1	
2	Altered resting state EEG microstate dynamics in acute-phase pediatric
3	mild traumatic brain injury
4	Sahar Sattari ¹ , Samir Damji ² , Julianne McLeod ³ , Maryam S. Mirian ^{4,5} , Lyndia C. Wu ^{1,6} , Naznin
5	Virji-Babul ^{5,7*}
6	¹ School of Biomedical Engineering, Faculty of Applied Science, University of British Columbia,
7	Vancouver, British Columbia, Canada
8	² Graduate Program in Neuroscience, Faculty of Medicine, University of British Columbia,
9	Vancouver, British Columbia, Canada
10	³ Department of Rehabilitation Sciences, Faculty of Medicine, University of British Columbia,
11	Vancouver, British Columbia, Canada
12	⁴ Pacific Parkinson's Research Centre, University of British Columbia, Vancouver, British
13	Columbia, Canada
14	⁵ Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver,
15	British Columbia, Canada
16	⁶ Department of Mechanical Engineering, Faculty of Medicine, University of British Columbia,
17	Vancouver, British Columbia, Canada
18	⁷ Department of Physical Therapy, Faculty of Medicine, University of British Columbia,
19	Vancouver, British Columbia, Canada
20	* Corresponding author
21	E-mail: nvb31@mail.ubc.ca (NVB)

22 Abstract

Objective: Sport-related concussion presents significant diagnostic and monitoring
challenges, especially in youth populations. This study investigates the potential of EEG
microstate analysis as a tool for assessing acute-phase brain activity changes in adolescent male
athletes following a concussion. We analyzed resting-state EEG data from 32 participants in a
between-subjects design, comparing participants with acute concussion (within two weeks of
injury) to an age- and sex- matched sample with no reported history of concussion.

Methodology: We applied a modified k-means clustering algorithm to group restingstate EEG topographical maps into seven clusters, with each cluster represented by one of the canonical microstate classes (A-G). Average duration, occurrence rate, and time coverage for each microstate were extracted.

Results: Statistically significant differences in mean duration, occurrence rate, and time coverage of microstates B and E were observed. Specifically, the mean duration, occurrence and time coverage of microstate E showed a significant decrease in the concussed cohort in comparision to the controls (p < 0.001). In addition, the mean duration, occurrence rate and time coverage was higher in the concussed cohort in comparision with the healthy cohort (p = 0.003). A significant negative linear relationship was found between microstate E and symptom severity (p = 0.006, F = 15.72).

Discussion: These results suggest that mild traumatic brain injury may disrupt the
 dynamic interaction of large-scale brain networks, hinting at potential biomarkers of injury. This
 study may help to inform future work on objective, brain-based tools for diagnosis and recovery
 assessment in concussed adolescents. Further research in larger, more diverse populations is
 necessary to validate these potential biomarkers.

45 Introduction

46 The neurophysiological and neurobiological changes due to mild traumatic brain injury (mTBI: hereafter used synonymously with concussion) are not well understood. Concussion 47 48 results from biomechanical forces exerted to the skull and the subsequent shearing and stretching 49 of brain tissue; although this injury is particularly prevalent, concussion diagnosis remains a 50 challenge due to its largely subjective nature and the absence of definitive clinical tests [1]. 51 There is a need for objective, brain-based diagnostic methods that can reliably indicate the 52 occurrence of a concussion [2,3]. The dynamic and complex nature of concussion injury is of 53 particular concern in adolescents due to the complexity and heterogeneity of neurodevelopment. 54 In Canada, concussion incidence is highest among adolescents aged 12 to 19 compared to 55 other age brackets across the lifespan; this result is corroborated by a similar epidemiological 56 investigation of the US population [4,5]. It should be noted that most concussion injuries in this age group are sustained during sport (hockey, rugby, and football in particular) and other 57 58 physical activities [4]. 59 Adolescents are particularly vulnerable to the neuropsychological consequences that stem 60 from sustaining a concussion. A growing base of structural and functional brain imaging studies 61 indicate that adolescence is an extremely dynamic period of brain development [6,7], and a 62 mTBI superimposed on a rapidly developing brain and body undergoing puberty reportedly leads 63 to more severe and persistent symptoms in comparison to younger children and adults [8,9]. 64 The current methods used in concussion detection, monitoring, and return-to-play clearance for adolescent athletes are suboptimal; these tools tend to lack either objectivity, utility, or 65

67 have begun to provide some clues regarding potential biomarkers of concussion. Resting-state

feasibility for this population [10,11]. Functional brain imaging studies on concussion to date

3

68 fMRI studies have examined the disruptions to brain functional connectivity caused by 69 concussion. This research has shown alterations in both dynamic (i.e., spending less time in a 70 frontotemporal default mode/limbic brain state) and static measures of functional connectivity 71 (i.e., altered interhemispheric connectivity, as well as hyperconnected frontal nodes and 72 hypoconnected posterior nodes in the salience and fronto-parietal networks) relative to healthy 73 control subjects [12,13].

74 While these findings are informative, the feasibility of performing multiple MRI tests on young athletes as an objective diagnosis/recovery assessment is not ideal and often not available 75 76 in remote regions. Portable and feasible neuroimaging techniques are required that are sensitive 77 to changes in the brain post impact. Electroencephalography (EEG) has the potential to serve as 78 an objective diagnostic tool and has been extensively studied in recent years for this purpose. 79 EEG microstate analysis describes the discrete functional cortex-wide states that occur in the resting-state brain. During the resting state, the brain does not exist in one such state, but shifts 80 81 dynamically between 4-7 different EEG "microstate topographies" that are usually stable for 82 approximately 30-120 milliseconds (ms) before shifting to another state [14–16]. These 83 microstates show high reliability [17]. This resting-state analysis method provides a lens for 84 understanding altered dynamics of large scale brain networks, rather than isolated features of the 85 EEG signal [18]. Importantly, EEG microstates have been shown to have strong associations 86 with large-scale fMRI resting state brain networks (RSNs) [19]. In addition, EEG microstates 87 capture subtle temporal dynamics about these functional brain areas and networks that cannot be captured with resting-state fMRI alone, given the limited temporal resolution of the BOLD 88 89 signal. To our knowledge, EEG microstate analysis has been applied by only one group to the 90 study of concussion [20]. However, this group investigated adults (mean age of 40 yrs) who had

sustained a concussion several years prior (0.3 – 16 yrs) and there was no healthy control
comparison group. In this chronic adult group with chronic neurophysiological impairement the
duration of the four canonical microstates (A-D) were negatively correlated with a
neuropsychological impairment index. Only four microstate classes were investigated and only
average duration of each microstate was investigated. To our knowledge, this is the first study
that has examined whether or not EEG microstate dynamics are altered in a adolecent sample in
acute phase of mTBI injury.

In the present study, we investigate acute-phase changes in EEG microstate features in a 98 99 cohort of adolescent males, with the primary aim to enhance our current understanding of neural 100 dynamic changes associated with concussion. We hypothesized that mTBI will be associated 101 with differences in microstate sequences as an indicator of acute changes in temporal dynamic 102 interplay of large scale brain networks in comparison to healthy controls, and that these changes 103 will be correlated with concussion symptoms. This pilot study is part of a larger project that aims 104 to identify potential microstate markers of concussion to enhance the objectivity and reliability 105 of concussion diagnoses and recovery assessments.

106 Materials and methods

107 Participants

Male athlete participants with normal or corrected to normal vision between the ages of 109 10 and 18 years were recruited for this study. Concussed participants received a diagnosis of 110 concussion from a physician or team doctor. Any participants exhibiting focal neurologic 111 deficits, pathology and/or those on prescription medications for neurological or psychiatric 112 conditions were excluded from the study. All participants provided written assent and their 113 parents gave written informed consent as per the guidelines of the Human Ethics Review Board

of the University of British Columbia. The study was approved by the University of British
Columbia Clinical Research Ethics Board (Approval number: H17-02973). The recruitment
period of the study started on March 15, 2019 and is ongoing. The current ethics protocol was
renewed and is valid until October 24, 2025. Eight of the concussed participants had completed
either the Sport Concussion Assessment Tool 3 (SCAT3) or the Child SCAT3 (for children aged
12 and under), a standardized tool widely used for the evaluation of athletes suspected of having
sustained a concussion [21].

121 *EEG data collection and preprocessing*

122 Five minutes of eyes closed, resting-state EEG data were collected from individuals using 123 a 64-channel HydroCel Geodesic Sensor Net (EGI, Eugene, OR). After obtaining assent and 124 informed consent, participants were seated in an experimental room with controlled lighting 125 levels and fitted with the EEG cap. They were instructed to minimize movement and remain 126 seated with their eyes closed. Before initiating the data collection, the electrode-scalp resistance 127 was ensured to be below 50 k Ω . The signals were referenced to the vertex (Cz) and recorded at a 128 sampling rate of 250 Hz. The collected EEG data were imported into Python for preprocessing. 129 The initial step involved re-referencing the data from the current reference (Cz) to the average of 130 all channels. This was followed by applying a 4th order Butterworth filter with zero phase shift 131 for band-pass filtering, and a frequency range of 1-50 Hz was kept. This frequency range was 132 chosen due to excessive noise at frequencies above 50 Hz, which interfered with subsequent 133 preprocessing steps. Independent component analysis (ICA) was performed using the *infomax* 134 algorithm and the MNE preprocessing package in Python [22]. To identify and remove noise-135 contaminated components, we employed the ICLabel algorithm, an automated labeling tool for 136 independent components [22]. We retained only those components that the algorithm labeled as

"Brain" (i.e. probability of the component presenting brain source was higher than artifactualsource).

139 *Microstate analysis*

140 For the microstate analysis, we utilized the *Pvcrostates* package in Python [23]. Global 141 Field Potential (GFP) peaks were extracted from individual data sets using the functions 142 available in the *Pycrostates* package. This process includes calculating the standard deviation of 143 all channel values at each time point, generating a time series of GFP values. Subsequently, 144 peaks in this time series were identified using the *Scipy* integrated function, with the minimum 145 distance between peaks set to the default value of 2 samples. We identified the minimum number 146 of GFP peaks across all subjects. We then resampled the GFP series for each subject to match 147 this minimum number. This step was performed to prevent the clustering results from being biased by variations in the number of peaks across individual data sets and to ensure uniformity. 148 149 The GFP peaks were clustered using a modified k-means algorithm, aligning with established 150 microstate literature.

151 The number of microstates used in previous studies typically ranges from 4 to 7 152 microstates; at present, there is no consensus on how to determine the optimal number of classes 153 to use for EEG microstate analysis [19]. Although 4 canonical microstate templates (A to D) 154 were initially prevalent in the EEG microstate literature, it has been argued that 7 distinct 155 microstate templates (A to G) best capture the scope of spontaneous electrophysiological activity 156 topographies that are observed in resting-state EEG studies [24]. We decided that 7 clusters 157 optimally fit our cohort's data, given that the 7-class solution maximized Global Explained 158 Variance (GEV), and the alignment of the 7 identified microstates' topographical shapes with 159 extant literature [25].

160 We aligned the 7 microstates with the established microstate labels (A to G) in the 161 literature, based on the topographical shapes of the states. We then assigned each time point of 162 the original dataset to one of the 7 microstates, using a "winner takes all" strategy, a smoothing 163 parameter of 6 samples (24 ms), and correlation between topographical maps as the similarity 164 measure. This involved correlating each time point with each microstate and assigning the time 165 point to the state with the highest correlation, taking smoothing parameter into account but not 166 taking polarity into account. The smoothing was applied to reduce the influence of rapid 167 fluctuations, potentially caused by artifacts, and to prevent false identification of microstate 168 changes.

169 Three measures were extracted from each individual's microstate sequence: average 170 duration, occurrence rate, and time coverage. First, the average duration of each microstate was 171 calculated by determining how long the microstate remained unchanged in each sequence. These 172 individual durations were averaged for each subject, and group-level means, and standard 173 deviations were extracted. The same method was applied to calculate the occurrence rate of each 174 microstate (i.e., the number of occurrences per second) and the time coverage (i.e., the 175 proportion of time each microstate occupied relative to the total recording time). Lastly, we also 176 extracted the average transition rate per second between microstates for each individual, as the 177 stability of staying in one state or frequently transitioning between states may be a characteristic 178 of mTBI. Fig. 1 presents the pipeline.

179

180 Fig 1. Study pipeline for microstate extraction from resting-state EEG to differentiate between

181 groups. The Global Field Power (GFP) time series is computed, and microstate maps are

182 identified by clustering maps from the local peaks of the GFP. Seven distinct microstates are

183 identified. The maps are labeled based on their shape and the labeling method introduced in [25].

The original EEG time series is backfitted to these microstates using a "winner-takes-all"
algorithm. Key metrics such as microstate duration, occurrence rate, and time coverage are then
extracted for group comparison.

187 Statistical analysis of microstate features

188 We used permutation statistics for all comparisons. For each feature, we conducted seven tests (one corresponding to each microstate). We applied a significance level of $\frac{0.05}{7}$ according to 189 190 the Bonferroni method to adjust for multiple comparisons. During each test, we randomized the 191 subjects from both groups and generated new groups of the same size as the original ones. We 192 then conducted a t-test on these newly formed random groups. If the t-value from the random 193 groups exceeded that of the original groups, we added the error rate by one. This procedure was 194 repeated 10,000 times. The total error rate was then calculated and divided by the number of 195 permutations. Features and microstates falling below the adjusted significance threshold are 196 reported.

We conducted a multiple regression analysis to understand the relationship between 197 198 symptom severity and quantity (as assessed by the SCAT) and microstate feature measures. 199 Specifically, we used the mean duration and occurrence rate of each microstate as independent 200 variables, while the reported quantity and severity of symptoms were used as the dependent 201 variables. To account for multiple comparisons, the significance threshold was adjusted to 0.007. 202 We did not include time coverage in this analysis, as it can be derived from the average duration 203 and occurrence rate. If these two features show a significant relationship with symptoms, time 204 coverage information is unnecessary.

205 Results

206 *Demographics*

207 34 male athletes were recruited: n = 14 with concussion and n = 20 with no history of

208 concucssion. Data from two acutely concussed males were removed due to excessive noise

remaining on EEG after preprocessing, leaving a final sample of 32 male adolescent athletes

between the ages of 13 and 18 (Table 1). The total number of symptoms and symptom severity

211 from the SCAT 3 were reported for 8 participants.

212

213 Table 1. Demographic and Clinical Characteristics of Study Participants.

214

	Mean age (min:max)	SCAT total symptoms (mean ± SD)	SCAT symptom severity (mean ± SD)
Healthy Controls (N = 20)	16.0 (14:18)	N/A	N/A
Acutely Concussed (N = 12)	15.1 (13:18)	$9.1 \pm 6.8 \ (n = 8)$	$21.0 \pm 20.4 \ (n = 8)$

215

216 *Microstate duration, occurrence rate and time coverage*

Seven microstates with topographies similar to those identified in the meta-analysis by Koenig et al. (2024) were extracted from each subject (Fig. 1). The variance explained by each microstate series was calculated for individual datasets following the backfitting process (Fig. 2B). The total global explained variance (GEV) was $58 \pm 12\%$. The analysis revealed several significant differences. First, microstate E was significant shorter in the concussed cohort compared to controls (p < 0.001) as shown by significantly lower duration, occurrence rate and

223	time coverage of this microstate in the concussed group (<i>Cohen's D</i> = 1.26 , 1.08 , 1.24
224	respectively). Additionally, microstate B exhibited a significantly higher mean duration,
225	occurrence rate, and time coverage (<i>p</i> = 0.003, 0.003, < 0.001; <i>Cohen's D</i> = 1.06, 1.07, 1.14
226	respectively) in the concussed group, i.e. there were fewer time points where the topographical
227	map of the concussed data resembled microstate E, and more time points where it resembled
228	microstate B (Fig.2B, C and D). We visualized these findings in two ways. In Fig. 2B, the
229	average duration of microstates B and E is shown for each participant as individual data points,
230	and by a bar plot comparing the group averages. In Fig. 3, we present the distribution of the
231	duration for microstates B (Fig. 3 left) and E (Fig. 3 right). Specifically, each time one of these
232	microstates occurred, its duration was recorded across data from all subjects. We then tested
233	whether the distributions of these duration values significantly differed between the two cohorts
234	using the Kolmogorov-Smirnov test and observed significant differences ($p < 0.001$).

235

236 Fig. 2. Microstate global variance, and comparative metrics in healthy vs. concussed participants: 237 (A) Total global explained variance (GEV) of each microstate sequence, with each dot representing 238 individual datasets. (B), (C) and (D) Bar plots representing the three key microstate measures: average 239 duration, average occurrence rate, and time coverage, respectively. Healthy (blue, n = 20) and concussed 240 (orange, n = 12) groups are compared, with dots representing individual data points. Significant 241 differences in mean duration, occurrence rate, and time coverage of microstates B and E are observed. 242 Overall, microstate E had a significantly lower duration, occurrence rate and time coverage in the 243 concussed group compared to controls (p < 0.001). Additionally, microstate B exhibited a significantly 244 higher mean duration, occurrence rate, and time coverage in the concussed group (p = 0.003). AC = Acute 245 concussed; HC = Healthy Controls. 246

247	Fig 3. Distribution of microstate B and E durations across all occurrences. The distributions for
248	microstate B and E durations are shown for both the HC (healthy control) and AC (acutely concussed)
249	groups. Significant differences were observed between the duration distributions, with $p < 0.001$ (D =
250	0.01 for microstate B, $D = 0.06$ for microstate E).
251 252	<i>Microstate transition rate</i> Transitions were recorded for each second of the data. The total number of transitions per
253	second was averaged over the entire data length for each individual. Interestingly, we observed
254	higher variability in the number of transitions among the control group, though their average and
255	median values were lower compared to the concussed group. However, this difference was not
256	statistically significant (Fig. 4).
257	
258	Fig 4. Rate of transitions in microstate sequences. (A) Shows a spike for each transition from one
259	microstate to another within a 4-second time window for example subgroups. (B) Shows a box plot
260	comparing the transition rates per second between the healthy control (HC) and acutely concussed (AC)
261	groups, with individual data points representing the average number of transitions per second for each
262	participant in the cohort. The difference between the two groups was not statistically significant.
263	
264 265	Concussion symptoms related to microstate features Among the eight concussed athletes who completed the SCAT, microstate E had a
266	significant negative linear relationship with the SCAT symptom severity measure ($p = 0.006$, F
267	= 15.72). This model indicates that increased symptom severity is associated with shorter
268	duration and lower occurrence rates of microstate E.
269	

Fig 5. Multiple regression model. The average duration and occurrence rate of microstate E are used as independent variables, and symptom severity is used as the dependent variable. Data from eight concussed participants were included in this model due to the absence of SCAT information from other concussed individuals. The model demonstrates a statistically significant linear relationship between the duration and occurrence rate of microstate E and the reported symptom severity (p = 0.006, F = 15.72).

275 Discussion

In the present study, we applied EEG microstate analysis to study the changes associatedwith acute sport-related concussion in a small cohort of adolescent male athletes.

278 As hypothesized, we observed disturbed functional dynamics in the resting-state brain microstate

279 of concussed adolescent participants relative to healthy control adolescent participants. Our

280 findings show specific changes in microstates E and B: there was a significant decrease in the

281 presence of microstate E and a significant increase in the presence of microstate B in our

concussed sample across three microstate measures examined. Specifically, microstate B occured
more frequently, lasted longer, and as a result covered a higher proportion of the total time; in

contrast, this pattern was the opposite for microstate E. Interestingly, we also observed that the
decreased occurrence rate and average duration of microstate E was associated with increased
symptom severity within the concussed group. We did not find any significant differences in
microstates transition rate between the groups.

EEG microstate analysis offers insights into temporal changes in the spatial organization of brain activity during rest. Specifically, this type of analysis relies on dynamic patterns of brain connectivity as opposed to a static measure of brain activity. Essentially, resting-state EEG scalp topographies tend to aggregate into a limited number of prototypical spatial distributions. These distributions are replicable both within and across individuals and have been extensively studied in the literature [17,25]. Typically, these topographical distributions are labeled A to G based on

their maps. While labeling systems may vary across studies, we compared our results with those
from studies that utilized similar topographical maps, though these studies report different labels.
Our labeling system is based on the unifying work of Tarailis et al. [26] which facilitates crossstudy comparisons.

298 Our findings indicate that the concussed group exhibits a reduced duration, occurrence rate 299 and time coverage of microstate E. Microstate E is characterized by a topography with a centro-300 parietal maximum [26] and has been associated with activity in the default mode network (DMN) 301 and the salience network (SN) [24,26]. Additionally, the scores of symptom severity showed a 302 significant negative correlation with the duration and occurrence of this microstate. Given the 303 critical role of the networks associated with microstate E in modulating both external and internal 304 attention, we also observed a similar relationship in the concussed group where symptom severity 305 which was linked to acute cognitive and somatic impairments were associated with characteristics 306 of microstate E. Previous research has shown that sleep deprivation [27] and acute drug 307 intoxication [28] are similarly associated with higher transition probability from microstate E to 308 other microstates and reduced duration and coverage of microstate E respectively. Self-reported 309 comfort at rest, on the other hand, has been associated with an increased duration of microstate E, 310 which may be due to increased self-referential processing associated with the DMN [29]. These 311 findings align well with what we observe in duration, occurrence and time coverage of microstate 312 E. Although not statistically significant, we also observed a decrease in the duration, occurrence 313 rate and time coverage of microstate D in the concussed group relative to the healthy control group. 314 Microstate D has been shown to primarily be associated with the frontoparietal executive control network (ECN) (e.g., activity in the right inferior parietal lobe and the right middle and 315

superior frontal gyri) [24,30]. Given that the salience network is thought to modulate the dynamic

interaction between the ECN and DMN [32], it is interesting that we observed a reduced presence of both microstates E and D in the concussed cohort, indicating potential impairment in key resting state networks related to attention regulation and cognitive control. Finally, a reduced duration of microstate E is consistent with the results of resting-state fMRI studies that have studied disruptions to dynamic functional connectivity in the brain as a result of concussion. Concussed pediatric participants have been observed to spend less time in a frontotemporal default mode/limbic brain state relative to their healthy control peers [13].

In contrast, microstate B characterized by a topography with a left frontal to right 324 posterior configuration has been shown to be associated with the visual processing regions of the 325 326 brain and particularly of activity in the left and right occipital cortices, including primary visual 327 cortex [24,26]. It is possible that an altered balance of sensory processing activity in the brain 328 (i.e., more activity in the visual imagery RSN), as a result of concussion, relates to acute 329 cognitive impairments associated with concussion. Future research should explore, how 330 concussion intervention alters the processing of various sensory inputs in the brain, and whether 331 or not an imbalance in brain activity related to the processing of different sensory inputs (e.g., 332 compensatory hyperactivity in one sensory modality) is a cause or effect of common concussion 333 symptoms such as mental fatigue, attentional deficits, and memory impairment. It is also worth 334 noting that in the aforementioned sleep deprivation study [27] and acute drug intoxication study 335 [28], not only was there a decreased presence of microstate E, but also a higher transition 336 probability from microstate E to microstate B. Perhaps concussion creates a state of impairment 337 that prompts a similar neurobiological compensatory response to sleep deprivation and alcohol 338 intoxication, where the brain is forced to prioritize bottom-up sensory processing (i.e., microstate

B) over internally-oriented networks or higher-order top-down cognitive networks (i.e.,

340 microstates E and D) [28].

341 We also analyzed the number of transitions between microstates for each participant. 342 This measure reflects the brain's ability to switch between different functional states, which is 343 crucial for maintaining efficient cognitive functioning. In a previous fMRI study, we observed 344 reduced transitions between functional states in individuals with a concussion, suggesting 345 cognitive rigidity [33]. However, we did not replicate this finding in the current study. A higher 346 transition rate could potentially indicate neural instability or increased metabolic cost, which 347 may also be linked to concussion pathology. Nevertheless, this observation did not reach 348 statistical significance. To draw more definitive conclusions, future studies should explore this 349 further with larger sample sizes and greater statistical power.

350 Limitations and methodological considerations

Our study results have limited generalizability as we were only able to include a small cohort of male adolescent participants. Future studies will build upon the results of the present study by performing microstate analysis on a larger and more diverse sample and in a longitudinal design to address normal within-subject variabilities. An additional limitation is that only 8 of the 12 concussed subjects had completed the SCAT3, reducing the power of the correlation analysis performed between SCAT symptom quantity/severity scores and microstate features.

358 Conclusion

In conclusion, using resting-state EEG in conjunction with microstate analysis, we observed significant disruptions in the dynamic interplay of large-scale brain networks in acutely concussed adolescent male athletes compared to their healthy control peers. EEG microstates

362	provide a readily accessible and replicable method to study the effects of concussion on whole-
363	brain network dynamics with high temporal resolution. This study aids in the effort to identify
364	an objective biomarker for concussion, which is direly needed for the most accurate diagnosis
365	and monitoring of this subtle and multifaceted injury.
366	
367	Acknowledgements
368	We would like to express our gratitude to the participants of this study, whose
369	cooperation and commitment made this research possible.
370	
371	
372	
373	
374	
375	
376	
377	
378	
379	
380	
381	
382	
383	
384	

c	ο	ᄃ
J	o	υ

386

387

388	References
389	1. Sussman ES, Pendharkar AV, Ho AL, Ghajar J. Mild traumatic brain injury and concussion:
390	terminology and classification. Handb Clin Neurol. 2018;158:21-4.
391	2. D'Lauro C, Jones E, Swope L, Anderson M, Broglio S, Schmidt J. Under-representation of
392	female athletes in research informing influential concussion consensus and position statements:
393	an evidence review and synthesis. British Journal of Sports Medicine. 2022;56:bjsports-2021.
394	3. Rubin TG, Lipton ML. Sex Differences in Animal Models of Traumatic Brain Injury. J Exp
395	Neurosci. 2019;13:1179069519844020.
396	4. Champagne AS, Yao X, McFaull SR, Saxena S, Gordon KR, Babul S, et al. Self-reported
397	concussions in Canada: A cross-sectional study. Health Rep. 2023;34:17-28.
398	5. Zhang AL, Sing DC, Rugg CM, Feeley BT, Senter C. The Rise of Concussions in the
399	Adolescent Population. Orthopaedic Journal of Sports Medicine. 2016;4:2325967116662458.
400	6. Ferschmann L, Bos MGN, Herting MM, Mills KL, Tamnes CK. Contextualizing adolescent
401	structural brain development: Environmental determinants and mental health outcomes. Current
402	Opinion in Psychology. 2022;44:170–6.
403	7. Smit DJA, Boersma M, Schnack HG, Micheloyannis S, Boomsma DI, Pol HEH, et al. The
404	Brain Matures with Stronger Functional Connectivity and Decreased Randomness of Its

405 Network. PLOS ONE. 2012;7:e36896.

- 406 8. Baillargeon A, Lassonde M, Leclerc S, Ellemberg D. Neuropsychological and
- 407 neurophysiological assessment of sport concussion in children, adolescents and adults. Brain Inj.
 408 2012;26:211–20.
- 409 9. Davis GA, Anderson V, Babl FE, Gioia GA, Giza CC, Meehan W, et al. What is the
- 410 difference in concussion management in children as compared with adults? A systematic review.
- 411 Br J Sports Med. 2017;51:949–57.
- 412 10. Daly E, Pearce AJ, Finnegan E, Cooney C, McDonagh M, Scully G, et al. An assessment of
- 413 current concussion identification and diagnosis methods in sports settings: a systematic review.
- 414 BMC Sports Science, Medicine and Rehabilitation. 2022;14:125.
- 415 11. Damji F, Babul S. Improving and standardizing concussion education and care: a Canadian
 416 experience. Concussion. 2018;3:CNC58.
- 417 12. Shafi R, Crawley AP, Tartaglia MC, Tator CH, Green RE, Mikulis DJ, et al. Sex-specific
- 418 differences in resting-state functional connectivity of large-scale networks in postconcussion
- 419 syndrome. Sci Rep. 2020;10:21982.
- 420 13. van der Horn HJ, Ling JM, Wick TV, Dodd AB, Robertson-Benta CR, McQuaid JR, et al.
- 421 Dynamic Functional Connectivity in Pediatric Mild Traumatic Brain Injury. Neuroimage.
- **422** 2024;285:120470.
- 423 14. Lehmann D. Brain Electric Microstates and Cognition: The Atoms of Thought. In: John ER,
- 424 Harmony T, Prichep LS, Valdés-Sosa M, Valdés-Sosa PA, editors. Machinery of the Mind: Data,
- 425 Theory, and Speculations About Higher Brain Function [Internet]. Boston, MA: Birkhäuser;

- 426 1990 [cited 2024 Apr 24]. p. 209–24. Available from: https://doi.org/10.1007/978-1-4757-1083427 0_10
- 428 15. Lehmann D, Pascual-Marqui RD, Michel C. EEG microstates. Scholarpedia. 2009;4:7632.
- 429 16. Katayama H, Gianotti LRR, Isotani T, Faber PL, Sasada K, Kinoshita T, et al. Classes of
- 430 Multichannel EEG Microstates in Light and Deep Hypnotic Conditions. Brain Topogr.
- 431 2007;20:7–14.
- 432 17. Khanna A, Pascual-Leone A, Michel CM, Farzan F. Microstates in resting-state EEG: current
- 433 status and future directions. Neurosci Biobehav Rev. 2015;49:105–13.
- 18. Michel CM, Koenig T. EEG microstates as a tool for studying the temporal dynamics of
- 435 whole-brain neuronal networks: A review. NeuroImage. 2018;180:577–93.
- 436 19. da Cruz JR, Favrod O, Roinishvili M, Chkonia E, Brand A, Mohr C, et al. EEG microstates
- 437 are a candidate endophenotype for schizophrenia. Nat Commun. 2020;11:3089.
- 438 20. Corradini PL, Persinger MA. Spectral power, source localization and microstates to quantify
- 439 chronic deficits from 'mild' closed head injury: Correlation with classic neuropsychological
- 440 tests. Brain Injury. 2014;28:1317–27.
- 441 21. McCrory P, Meeuwisse WH, Aubry M, Cantu B, Dvorák J, Echemendia RJ, et al. Consensus
- 442 statement on concussion in sport: the 4th International Conference on Concussion in Sport held
- 443 in Zurich, November 2012. Br J Sports Med. 2013;47:250–8.

- 444 22. Li A, Feitelberg J, Saini AP, Höchenberger R, Scheltienne M. MNE-ICALabel:
- 445 Automatically annotating ICA components with ICLabel in Python. Journal of Open Source
- 446 Software. 2022;7:4484.
- 447 23. Férat V, Scheltienne M, Brunet D, Ros T, Michel C. Pycrostates: a Python library to study
- 448 EEG microstates. Journal of Open Source Software. 2022;7:4564.
- 449 24. Custo A, Van De Ville D, Wells WM, Tomescu MI, Brunet D, Michel CM.
- 450 Electroencephalographic Resting-State Networks: Source Localization of Microstates. Brain
- 451 Connect. 2017;7:671–82.
- 452 25. Koenig T, Diezig S, Kalburgi SN, Antonova E, Artoni F, Brechet L, et al. EEG-Meta-
- 453 Microstates: Towards a More Objective Use of Resting-State EEG Microstate Findings Across
- 454 Studies. Brain Topogr. 2024;37:218–31.
- 455 26. Tarailis P, Koenig T, Michel CM, Griškova-Bulanova I. The Functional Aspects of Resting
- 456 EEG Microstates: A Systematic Review. Brain Topogr [Internet]. 2023 [cited 2024 Feb 14];
- 457 Available from: https://doi.org/10.1007/s10548-023-00958-9
- 458 27. Ke M, Li J, Wang L. Alteration in Resting-State EEG Microstates Following 24 Hours of
- 459 Total Sleep Deprivation in Healthy Young Male Subjects. Front Hum Neurosci [Internet]. 2021
- 460 [cited 2024 Aug 24];15. Available from: https://www.frontiersin.org/journals/human-
- 461 neuroscience/articles/10.3389/fnhum.2021.636252/full
- 462 28. Schiller B, Heinrichs M, Beste C, Stock A-K. Acute alcohol intoxication modulates the
- temporal dynamics of resting electroencephalography networks. Addict Biol. 2021;26:e13034.

- 464 29. Tarailis P, Šimkutė D, Koenig T, Griškova-Bulanova I. Relationship between Spatiotemporal
- 465 Dynamics of the Brain at Rest and Self-Reported Spontaneous Thoughts: An EEG Microstate
- 466 Approach. Journal of Personalized Medicine. 2021;11:1216.
- 467 30. Britz J, Van De Ville D, Michel CM. BOLD correlates of EEG topography reveal rapid
- 468 resting-state network dynamics. NeuroImage. 2010;52:1162–70.
- 469 31. Seitzman BA, Abell M, Bartley SC, Erickson MA, Bolbecker AR, Hetrick WP. Cognitive
- 470 manipulation of brain electric microstates. Neuroimage. 2017;146:533–43.
- 471 32. Menon V, Uddin LQ. Saliency, switching, attention and control: a network model of insula
- 472 function. Brain Struct Funct. 2010;214:655–67.
- 473 33. Muller AM, Virji-Babul N. Stuck in a State of Inattention? Functional Hyperconnectivity as
- 474 an Indicator of Disturbed Intrinsic Brain Dynamics in Adolescents With Concussion: A Pilot
- 475 Study. ASN Neuro. 2018;10:1759091417753802.

Group comparison Modified k-means algorithm Resting-state EEG Microstate classes מנות מיילאים היא ליייייירו ליייי היו ביואיי מייי הייי איין אייין אייין אייין אייין אייין אייין אייי and and the second and a second and the second and Duration 14/14/12/2012 Jan MAN Markey Markey Markey Markey 10/1 Mayorken he your mangar man Mangar Minory with Minor Man with the for the market and the state of the state of the section and the section of the s anone phan por states and an are anima phall particular and particular winning the physic way was an wing the deal deal and and the set of the se many martinely why have a some many more the 10114Marthallerty and mound Murde ABCDEFG many in all the way of a state and the work In March Harry and March March Occurrence MILINIA MARCHARMAN MANAMANIA CAM with a short we will want a will be with GFP Maps ABCDEFG Α в C E F D G

Figure 1



Figure 2



(A)

(B)



Figure 4

Microstate E vs Symptoms Severity



Figure 5