

Emergence of a Novel Reassortant Clade 2.3.2.1c Avian Influenza A/H5N1 Virus Associated with Human Cases in Cambodia

Jurre Y. Siegers^{1*}, Ruopeng Xie^{2,3*}, Alexander M.P. Byrne^{4*}, Kimberly M. Edwards^{2,3}, Shu Hu^{2,3}, Sokhoun Yann¹, Sarath Sin¹, Songha Tok¹, Kimlay Chea¹, Sreyviseth Horm¹, Chenthearath Rith¹, Seangmai Keo¹, Leakhena Pum¹, Veasna Duong¹, Heidi Auerswald¹, Yisuong Phou¹, Sonita Kol¹, Andre Spiegel⁴, Ruth Harvey⁵, Sothyra Tum⁶, San Sorn⁶, Bunnary Seng⁶, Yi Sengdoeur⁷, Chau Darapheak⁸, Chin Savuth⁸, Makara Hak⁹, Vanra Ieng¹⁰, Sarika Patel¹⁰, Han Di¹¹, Charles Todd Davis¹¹, Alyssa Finlay¹², Borann Sar¹², Peter Thielen¹³, Filip F. Claes¹⁴, Nicola S. Lewis⁵, Ly Sovann⁷, Vijaykrishna Dhanasekaran^{2,3#}, Erik A. Karlsson^{1#}

¹ Virology Unit, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

² School of Public Health, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China.

³ HKU-Pasteur Research Pole, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong SAR, China.

⁴ Direction, Institut Pasteur du Cambodge, Phnom Penh, Cambodia

⁵ Worldwide Influenza Centre, The Francis Crick Institute, London, UK.

⁶ National Animal Health and Production Research Institute, Phnom Penh, Cambodia.

⁷ Communicable Disease Control Department, Ministry of Health, Phnom Penh, Cambodia.

⁸ National Institute of Public Health, Ministry of Health, Phnom Penh, Cambodia

⁹ Food and Agriculture Organization of the United Nations (FAO) Country Office, Phnom Penh, Cambodia.

¹⁰ World Health Organization Country Office, Phnom Penh

¹¹ United States Centers for Disease Control and Prevention, Atlanta, GA, USA

¹² United States Centers for Disease Control and Prevention Country Office, Phnom Penh, Cambodia

¹² Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA

¹³ FAO Emergency Centre for Transboundary Animal Diseases, Regional Office for Asia and the Pacific, Bangkok, Thailand

*These authors contributed equally

#**Address for Correspondence:** For correspondence regarding avian influenza in Cambodia and Southeast Asia, please contact Dr. Erik Karlsson at ekarlsson@pasteur-kh.org. For inquiries related to influenza ecology and evolution, please contact Dr. Vijaykrishna Dhanasekaran at veej@hku.hk.

Keywords: avian influenza, H5N1, Cambodia, Southeast Asia, One Health, genomic surveillance

44 I. Abstract

45 After nearly a decade without reported human A/H5N1 infections, Cambodia faced a sudden
46 resurgence with 16 cases between February 2023 and August 2024, all caused by A/H5 clade
47 2.3.2.1c viruses. Fourteen cases involved a novel reassortant A/H5N1 virus with gene segments from
48 both clade 2.3.2.1c and clade 2.3.4.4b viruses. The emergence of this novel genotype underscores
49 the persistent and ongoing threat of avian influenza in Southeast Asia. This study details the timeline
50 and genomic epidemiology of these infections and related poultry outbreaks in Cambodia.

51

52 II. Introduction

53 Avian influenza viruses (AIVs), particularly highly pathogenic avian influenza (HPAI) A/H5N1 strains,
54 have caused significant economic losses in the poultry industry and pose a serious zoonotic threat
55 (1). Monitoring AIV circulation and evolution in high-risk regions, such as the Greater Mekong
56 Subregion (GMS), is crucial due to the region's heavy reliance on agriculture, particularly backyard
57 poultry farming, which often operates with minimal biosafety or biosecurity measures.

58

59 Cambodia, a lower middle-income country in the GMS has a large socio-economic dependence on
60 agriculture and faces ongoing circulation of AIVs in poultry. Longitudinal surveillance at key live bird
61 markets (LBMs) in Cambodia has shown year-round co-circulation of AIVs of agricultural and zoonotic
62 concern including HPAI A/H5N1, low pathogenic avian influenza (LPAI) A/H9N2, and to a lesser
63 extent, A/H7 viruses¹⁻⁴. In addition, there has been infrequent detection of wild-bird-associated LPAI
64 subtypes, including A/H1, A/H2, A/H3, A/H4, A/H6, A/H10, A/H11, and even A/H14, the latter marking
65 the first detection of this subtype in Asia⁵⁻⁷. Along with the high prevalence of AIV in poultry,
66 Cambodia has faced numerous instances of zoonotic spillover between 2005 and 2014, resulting in
67 21 human cases of A/H5N1, with 19 deaths (case-fatality rate [CFR]: 90.5%) by the end of 2012. In
68 January 2013, a clade 1.1.2 reassortant virus emerged in poultry in the GMS, containing the
69 hemagglutinin (HA) and neuraminidase (NA) genes from a clade 1.1.2 genotype Z virus and internal
70 genes from a clade 2.3.2.1a virus. Following the emergence of this novel reassortant, 26 new human
71 cases were reported in 2013 alone, with 15 deaths (CFR: 57.9%)^{5,8}. The outbreak continued into the
72 first quarter of 2014, resulting in 8 confirmed cases and 4 deaths (CFR: 50%)⁹. This clade 1.1.2

73 reassortant virus was replaced by clade 2.3.2.1c viruses in poultry by March 2014, with only one
74 clade 2.3.2.1c human infection occurring thereafter³. The reassortant 1.1.2 virus has not been
75 detected since. HPAI A/H5 clade 2.3.2.1c viruses have continued to circulate in the GMS since then¹⁰;
76 however, clade 2.3.4.4b viruses were first detected in Cambodian LBMs in 2021, co-circulating with
77 clade 2.3.2.1c viruses¹¹.

78

79 After almost a decade without a human case, a cluster of A/H5N1 cases involving two related
80 individuals was detected in February 2023 caused by the historically circulating clade 2.3.2.1c viruses
81 (1 death, CFR: 50%). Between October 2023 and end of August 2024, Cambodia reported 14 more
82 A/H5N1 cases (6 deaths, CFR: 42.8%). Critically, this recent surge in human A/H5N1 cases is linked
83 to the emergence of a novel reassortant genotype, which combines elements from multiple AIV
84 lineages. This study aims to elucidate the timeline, transmission dynamics, and evolutionary patterns
85 of these A/H5N1 viruses across human and avian populations in Cambodia between February 2023
86 and August 2024. Here, we seek to trace viral origins, identify transmission pathways, and assess
87 potential for increased risks to human health.

88

89 **III. Materials and Methods**

90 **Ethical approval**

91 All H5 sequences and information (e.g. collection date, location) included in this analysis were
92 obtained as part of human H5 case testing and One Health response as described below. All data
93 included in this study is available in the public record. All patient samples were de-identified and no
94 other individual-specific information was used in this study. Analyses in this study have been
95 approved by the Cambodian National Ethics Committee for Health Research (#365NECHR/2024).

96

97 **Sample collection**

98 ***Human case detection***

99 The Cambodian Influenza-like Illness (ILI) and Severe Acute Respiratory infections (SARI)
100 surveillance systems are part of the WHO Global Influenza Surveillance and Response System and
101 are a public health activity managed by the Ministry of Health (MoH) and Cambodia's Communicable

102 Disease Control (CCDC) Department. This surveillance system has existed since 2006, with
103 increased capacity during the COVID-19 pandemic. Cases were also detected through event-based
104 surveillance and active case finding.

105 ***Case investigation***

106 For each human case, a collaborative One Health investigation was conducted by MoH-CCDC, the
107 National Animal Health and Production Research Institute (NAHPRI), Ministry of Environment, and
108 provincial authorities, with assistance from the United States Centers for Disease Control and
109 Prevention (US CDC), the World Health Organization (WHO), and the Food and Agriculture
110 Organization of the United Nations (FAO).

111 ***Animal sampling***

112 Animal sampling was conducted by NAHPRI under the direction of the General Directorate for Animal
113 Health and Production (GDAHP), Cambodian Ministry of Agriculture, Forestry and Fisheries (MAFF)
114 along with provincial officials as part of One Health investigations surrounding each human case. In
115 addition, Institut Pasteur du Cambodge (IPC) and NAHPRI/GDAHP collaborate on active disease
116 surveillance activities focusing on LBMs¹¹, with assistance from the Food and Agriculture
117 Organization of the United Nations (FAO).

118 **RNA extraction and qRT-PCR**

119 Viral RNA was extracted from samples using the QIAamp Viral RNA Mini Kit (Qiagen, Maryland, USA)
120 according to the manufacturer's protocol. Quantitative real-time reverse transcription PCR (qRT-PCR)
121 was performed to screen for the matrix (MP) gene of influenza A virus, as described previously^{6,7}.
122 Samples with a cycle threshold (Ct) value <40 were deemed positive. Both original samples and
123 isolates were subtyped using influenza RT-PCR assays to test for H1, H3, H5, H7 and H9 HA genes
124 and N1, N2, N6, N8 and N9 NA genes^{6,7}.

125

126 **Virus isolation**

127 Virus isolation was attempted for all qRT-PCR positive human and poultry samples from each
128 outbreak cluster¹². Ten-day-old embryonated hens' eggs were inoculated via the allantoic route and
129 allantoic fluid was harvested three days post inoculation. For each sample, three blind passages were

130 performed. Additionally, virus isolation from human samples was performed using Madin-Darby
131 canine kidney (MDCK) cells stably transfected with human CMP-*N*-acetylneuraminase:β-galactoside
132 α-2,6-sialyltransferase (SIAT1), which leads to overexpression of the α-2,6-linked sialic acid (α2,6-SA)
133 receptor¹³. The presence of influenza virus in the allantoic fluid and cell culture supernatant was
134 tested using qRT-PCR as described above.

135

136 **Sequencing**

137 Whole viral genomes were amplified from patient nasal swabs, poultry samples, and/or isolates using
138 a modified multi-segment RT-PCR method that includes integrated molecular indices (iiMS-PCR)¹⁴.
139 The iiMS-PCR products were then pooled, prepared for sequencing with the ligation sequencing kit
140 SQK-LSK114 (R10 chemistry, Oxford Nanopore Technologies (ONT), Oxford, UK) and sequenced on
141 the GridION platform (ONT) using R10 flow cells as described previously¹⁵. Sequencing reads were
142 de-multiplexed, quality trimmed, and filtered using Porechop software
143 (<https://github.com/rrwick/Porechop>). Consensus sequences were generated using CDC's Iterative
144 Refinement Meta Assembler (IRMA) v1.1.4 using the default "IRMA FLU-minion" settings¹⁶.
145 Consensus sequences were manually inspected for errors such as insert-deletion mutations (INDELs)
146 and mixed bases and corrected if required. A minimum coverage depth of 10x was set for all genes.
147 Genomic sequencing was completed for twelve of the fifteen reported human cases, with one HA
148 sequence (A/Cambodia/RL240007/2024) being partial. This study generated a total of 83 gene
149 segments obtained from 12 human A/H5N1 and 329 segments from 51 avian A/H5N1 influenza A
150 viruses were deposited in GISAID. Accession numbers can be found in supplemental table 2.

151

152 **Phylogenetic analysis**

153 All available full-genome H5Nx influenza A virus sequences (N = 11,765), including the internal
154 protein coding genes were obtained from GISAID EpiFlu¹⁷ (26) databases (accessed on 2024-03-21).
155 The full genome sequences were concatenated using SeqKit¹⁸ and aligned prior to the inference of
156 preliminary maximum-likelihood (ML) phylogenetic trees with Augur¹⁹. This tree was then down-
157 sampled to maintain phylogenetic representatives using PARNAS²⁰. Sequences generated in this
158 study, along with publicly available sequences from Asia since 2020-01-01, were added to this

159 dataset to enhance regional representation and maximize phylogenetic diversity. Sequences
160 identified as duplicates (based on strain name), laboratory-derived, mixed subtypes, or with coverage
161 <90% of full length were excluded from further analysis. Each gene dataset was aligned with MAFFT
162 v.7.490²¹, and optimized manually. Large-scale ML phylogenetic trees were inferred with FastTree
163 v2²² using the GTR nucleotide substitution model. Focused ML trees were generated in IQ-TREE
164 v.2.2.0²³ using the best-fit nucleotide substitution model (HKY+Γ+G4) chosen according to
165 Bayesian Information Criterion. Trees were visualized and annotated using FigTree v.1.4.4
166 (<http://tree.bio.ed.ac.uk/software/figtree/>) and ggTree²⁴ in R (version 4.2.2).

167

168 **Reassortment analysis**

169 To investigate reassortment events among major clades (2.3.2.1c, 2.3.4.4b, 2.3.4.4x, 2.3.2.1a, 2.3.4,
170 1.1.2) and minor clades since 1996, we first constructed a ML tree for HA gene sequences using IQ-
171 TREE v.2.1.4. We subsampled 1,000 genomes, ensuring all Cambodian sequences were retained,
172 using the Phylogenetic Diversity Analyzer tool v.1.0.3 (<http://www.cibiv.at/software/pda>).
173 Subsequently, we constructed ML trees for the remaining seven segments, including polymerase
174 basic 2 (PB2), polymerase basic 1 (PB1), polymerase acidic (PA), neuraminidase (NA), nucleoprotein
175 (NP), matrix (MP), and non-structural (NS) gene segments of the 1,000 subsampled genomes. Baltic
176 v.0.1.5 (<https://github.com/evogytis/baltic>) was used to visualize the incongruence between the
177 phylogenetic trees across the eight genes. The HA tree was rooted using clade 0, while the other
178 trees were rooted at the midpoint.

179

180 **Molecular marker analysis**

181 Molecular markers were analyzed using a pipeline described previously²⁵ and Flutile
182 (<https://github.com/flu-crew/flutile>).

183

184

185

186

187 **IV. Results**

188 **Direct poultry-to-human transmission with no evidence of human-to-human spread**

189 Viruses sequenced from human cases in February 2023 in Prey Veng province were closely related
190 to A/H5N1 viruses from poultry in Cambodian LBMs between 2022 and 2023, particularly from Phnom
191 Penh and Kandal provinces (Figure 1a-c). Similarly, viruses from human cases in October 2023 (Svay
192 Rieng and Prey Veng province), November 2023 (Kampot province), January 2024 (Siem Reap
193 province), February 2024 (Kratie province), and early July 2024 (Svay Rieng province) were
194 genetically similar to poultry viruses collected at their respective outbreak sites. Although the January
195 2024 Prey Veng human case lacked direct poultry-associated sequences, it was related to viruses
196 detected from LBMs in Prey Veng and Kandal in January 2024. In February 2024, no viral sequences
197 could be obtained from neither the Kampot human case nor poultry associated cases. In July 2024,
198 no human viruses could be sequenced from the two Takeo province cases, although the two poultry-
199 associated viruses were closely related to human and poultry viruses from the October 2023 Svay
200 Rieng and one duck virus from the October 2023 Prey Veng outbreak. Investigations into the August
201 2024 cases Prey Vengcase showed that this virus was related to human and poultry viruses from
202 October 2023 (Prey Veng and Svay Rieng) and July 2024 (Kampot). These findings consistently
203 suggest zoonotic transmission from poultry to humans, with no evidence of human-to-human
204 transmission (Figure 1c).

205

206 **Genotype replacement by a novel reassortant clade 2.3.2.1c virus in late 2023**

207 Phylogenetic analysis of the HA gene segment reveals that all A/H5 viruses sequenced from human
208 cases belong to clade 2.3.2.1c (Figure 1c). The virus responsible for the initial cases in February 2023
209 was closely related to clade 2.3.2.1c A/H5N1 viruses circulating in Cambodian poultry and wild birds
210 since 2013¹. These early viruses retained complete gene cassettes within clade 2.3.2.1c viruses
211 (Figure 2). However, starting in late 2023 (October), HA sequences from both humans and poultry in
212 Cambodia displayed significant divergence from earlier Cambodia strains. These HA genes clustered
213 with two duck samples from Vietnam during July and August 2023, and were derived from a
214 sublineage of clade 2.3.2.1.c continuously detected in Laos during October 2020 and May 2023.


215 Together, this indicates an introduction of a sublineage of clade 2.3.2.1c viruses with a common
216 ancestor detected in poultry in Vietnam and Laos, but the exact introduction pathway cannot be
217 pinpointed due to gaps in available surveillance data..

218 Phylogenetic analyses of each of the gene segments revealed that A/H5N1 viruses in Cambodia,
219 detected from October 2023 onwards in both humans and poultry, represent a novel genotype
220 resulting from reassortment. This reassortment combines segments from clade 2.3.2.1c (HA, NP, and
221 NA) and clade 2.3.4.4b (PB2, PB1, PA, MP, and NS) viruses (Figure 2). In Cambodia, this novel
222 2.3.2.1c genotype has completely replaced the endemic 2.3.2.1c genotype that has dominated in
223 poultry for the last decade.

224 **Molecular markers of risk in novel reassortant genotype Cambodian A/H5N1 viruses have the** 225 **potential for increased mammalian adaptation**

226 Gene segments from the Cambodian human A/H5N1 viruses were analyzed for adaptive mutations
227 affecting receptor binding affinity, HA fusion and/or stability, polymerase activity, virus replication in
228 mammalian cells, and pathogenicity (Supplemental Table 1)²⁶. All human A/H5N1 viruses isolated
229 since 2023 are classified asHPAI due to the presence of a multi-basic cleavage site in the HA gene²⁷.
230 Three different cleavage sites were present: PQKERRKR↓GLF in human cases in February 2023,
231 PQRERRKR↓GLF in cases from October 2024 to January 2024 and from July 2024 onwards, and
232 KERRRKKR↓GLF, containing two additional basic amino acids from cases in February 2024.


233 Among the minimal set of five substitutions that confer airborne transmissibility in ferrets for the HPAI
234 A/H5N1 virus strain A/Indonesia/5/2005 (E627K in PB2; H99Y in PB1; H103Y, T156A, and either
235 Q222L or G224S in HA)^{28,29}, only HA 156A was consistently observed in all A/H5N1 viruses from
236 Cambodian human cases. Notably, a subset of the novel genotype forms a monophyletic lineage in
237 the HA tree (arrow in Fig 1c, bootstrap value 92), and contains PB2 E627K, a key molecular
238 determinant for host range, cross-species transmission and airborne transmission (PMID: 8445709
239 and 22723413)None of the human viruses from Cambodia contained previously defined HA
240 substitutions (Q222L and G224S (H5 HA numbering) that switch receptor specificity from avian α 2,3-
241 SA to human α 2,6-SA³⁰. The presence of PB2 E627K and HA T156A substitutions, along with the
242 potential for other functionally similar mutations, underscores the need for close monitoring of these
243 A/H5N1 strains for signs of potential increased mammalian adaptation³¹.



244

245 **Figure 1. Geographic location, timeline of events, and phylogenetic relationships of human A/H5N1 infections and**
 246 **poultry-associated clade 2.3.2.1c A/H5N1 viruses in Cambodia, February 2023 - July 2024. a)** Map of human A/H5N1
 247 cases. Circles indicate clinical outcome, with smaller circles for recovered individuals and larger circles for fatal cases, colored
 248 by outbreak cluster. **b)** Timeline of A/H5N1 human cases. Colors represent outbreak clusters; dotted circles indicate
 249 epidemiologically linked cases; mortality symbols indicate fatal cases; chicken and duck silhouettes denote associated poultry
 250 viruses; and major Cambodian festivals are labeled. **c)** Maximum likelihood phylogenetic tree of the A/H5 HA genes, with
 251 human and poultry-associated A/H5N1 viruses colored by outbreak cluster. The color-coded right y-axis represent the E/K
 252 amino acid in the PB2 gene at position 627 (blue/orange), country of isolation (shades of green) and host (shades of red),
 253 respectively. Bootstrap values >80% are indicated. Scale bar denotes nucleotide substitutions per site. Tree is rooted to the
 254 clade 2.3.2.1c vaccine strain, A/duck/Vietnam/NCVD-1584/2012. Names are provided as strain_name | collection_date.

255



256

257

Figure 2. Genotypic reassortment patterns of clade 2.3.2.1c A/H5N1 viruses causing human infections in Cambodia since February 2023. Tanglegram of ML phylogenetic trees for each genomic segment. For each virus, links have been displayed to connect phylogenetic positioning of individual viruses across all eight genomic segments. Phylogenetic links are color coded based on relevant A/H5 clade numbering: clade 2.3.2.1c in orange, 2.3.2.1a in dark blue, 2.3.4.4 in light blue, 2.3.4.4b in green, 2.3.4 in purple, 1.1.2 in brown, minor A/H5N1 clades in grey and recent Cambodian human A/H5N1 sequences in bright red.

263

264

265

266

267

268

269

270

271

272

273

274

275

276 V. Discussion

277 Reassortment events leading to the exchange of gene segments between viruses can significantly
278 alter viral properties and potentially increase the risk for spillover and human infections³²⁻³⁵. AIVs with
279 novel gene combinations have been associated with increases in zoonotic human cases, such as with
280 A/H5N1 (Cambodia clade 1.1) A/H7N9³⁴, A/H3N8³³, and even human pandemics³². Clade 2.3.4.4b
281 A/H5N1, which has spread globally since its emergence in 2020, has undergone further reassortment
282 during its dissemination.

283 The resurgence of zoonotic A/H5N1 cases in Cambodia underscores the complex dynamics of AIV
284 evolution and potential for zoonotic transmission. Phylogenetic analysis identified two distinct spillover
285 phases. The first phase, in February 2023, involved spillover of a clade 2.3.2.1c virus genotype, which
286 had been dominant in Cambodian poultry since 2014³. The second phase, starting in October 2023, is
287 marked by a novel reassortant virus combining genes from clades 2.3.2.1c and 2.3.4.4b. The exact
288 origins of this reassortment are unclear, but likely facilitated by high-density poultry farming, wild bird
289 migration, and cross-border poultry trade in the region, highlighting the ongoing risk of zoonotic AIV
290 transmission in Southeast Asia.

291 While the phenotypic contributions of newly introduced clade 2.3.4.4b internal gene segments have
292 yet to be elucidated, the presence of amino acid mutations in both human and poultry viruses such as
293 E627K in the 2.3.4.4b-origin PB2 gene segment suggests enhanced capacity for mammalian infection.
294 To better understand the zoonotic risk that these viruses pose, further risk assessment *in silico*, *ex*
295 *vivo*, *in vivo*, and *in vitro* is critical. In addition, the detection of the PB2 627K mutation in poultry is
296 also a concern, as it may become established in widespread circulation.

297 The success of the Cambodian One Health response to these recent reassortant A/H5N1 cases
298 highlights the importance of coordinated efforts between human and animal health sectors in
299 managing zoonotic threats. In addition, rapid genomic surveillance has been critical in understanding
300 the dynamics of virus spillover events and has provided clear evidence linking human cases to
301 infected poultry from the same household. This integration of genomic data with field investigations
302 allowed for timely identification of transmission pathways and underscores the importance of

303 ongoing surveillance in high-risk areas. These efforts exemplify the strength of the One Health
304 approach in Cambodia, where collaboration across disciplines and rapid data sharing have proven
305 essential for understanding, responding and controlling human AIV outbreaks in the country.

306 Given the success of Cambodian One Health response efforts being able to link cases to direct
307 contact with infected poultry, public health strategies should prioritize reducing human exposure,
308 particularly in high-risk rural areas. Reinforcing public education on the risks associated with direct
309 contact with infected birds has proven effective during other AIV outbreaks, but significant challenges
310 remain in achieving consistent behavioral changes. Most Cambodians in rural areas are still at high-
311 risk for potential exposure to AIV³⁶. In conjunction with human case investigations, community-based
312 education campaigns have been implemented in Cambodia, but their reach and effectiveness
313 vary^{37,38}. Only 32% of survey participants reported receiving information about AIV from healthcare
314 providers, 10.6% from village health support groups, and 2% from village animal health workers
315 (VAHW)³⁷. Furthermore, only 49% of participants reported poultry illness and deaths to local
316 authorities, and 23% of participants reported that they cook sick or dead poultry for consumption. This
317 is particularly crucial given that 68.3% of participants raised chickens in their backyards and 10.2%
318 raised ducks, generating a high level of potential exposure to infected birds³⁷. Further research is
319 necessary to identify, understand and overcome the barriers in risk perception that hinder
320 individuals from adopting safer behaviors, and to more effectively tailor behavioral messages to
321 specific target audiences³⁹. Educational interventions on the necessity of early access to healthcare,
322 recognizing AIV infection as a life-threatening disease, and awareness of the risks associated with
323 handling sick poultry have proven to be a powerful tool in increasing awareness and modifying health
324 behaviors^{40,41}.

325 Training and funding for VAHW and primary healthcare workers (PHW) are critical for early detection
326 and outbreak response⁴⁰⁻⁴⁴. Educational interventions can help VAHW and PHW identify poultry
327 deaths and exposure history, while also supporting event-based surveillance in both human and
328 animal sectors. Strengthening VAHW reporting and raising clinical suspicion among healthcare
329 providers will enhance early detection and case management. Equipping healthcare workers to

330 manage cases effectively and increasing awareness among high-risk individuals about early
331 healthcare access and poultry-related risks are essential to improving preparedness^{40,41}.

332 This study is limited by potential biases in human sample collection due to potential underreporting of
333 cases from rural areas with limited healthcare access. Additionally, sustainable funding for active AIV
334 surveillance is lacking across the GMS and broader Asia, resulting in inconsistent data collection and
335 limited genomic information available for in-depth analysis. Moreover, there is an urgent need to
336 develop a classification system that better reflects genotypes, as the conventional HA and NA
337 subtype nomenclature fails to capture the full complexity of reassortant AIV genomes and associated
338 evolutionary context.

339 In conclusion, these recurrent zoonotic infections caused by a novel reassortant A/H5N1 viruses in
340 Cambodia serve as a reminder of the ever-present threat of AIV to global health security. Despite the
341 recent focus on global dissemination and expanded host range of clade 2.3.4.4b⁴⁵⁻⁴⁷, clade 2.3.2.1c
342 viruses remain a significant concern, particularly in Asia, where the two clades co-circulate. A
343 coordinated, regional approach is essential for effectively monitoring the threat of AIV and ensuring
344 preparedness and response to emerging viral threats. Indeed, this study underscores the critical need
345 and effectiveness of a unified, One Health approach to combat the evolving landscape of AIV. The
346 urgency of this collaborative effort cannot be overstated as these viruses continue to adapt and
347 reassort, increasing the risk of a strain evolving the capacity for efficient human-to-human
348 transmission. To stay ahead of this threat, we must prioritize sustainable funding for long-term
349 surveillance, enhance laboratory capacity for rapid whole genome sequencing, and foster open, trust-
350 based information sharing across borders. Our collective preparedness today will determine our ability
351 to protect global health tomorrow.

352

353

354

355 VI. Acknowledgments

356 The investigators thank everyone involved in the critical discussions and review of this manuscript.
357 They also thank everyone involved in influenza surveillance and response in the Kingdom of
358 Cambodia including teams at the National Institute for Public Health, Cambodian Communicable
359 Disease Control Department, Ministry of Health, National Animal Health and Production Institute,
360 General Directorate of Animal Health and Production, Ministry of Agriculture, Forestry, and Fisheries,
361 World Health Organization, United States Centers for Disease Control and Prevention, provincial
362 health directors, province rapid response teams, and the Virology Unit at Institut Pasteur du
363 Cambodge who contributed to this study. We gratefully acknowledge the authors from the originating
364 laboratories responsible for obtaining the specimens and the submitting laboratories where genetic
365 sequence data were generated and shared via the GISAID Initiative, on which this research is based
366 (full list of GISAID acknowledgments are available in supplemental material). We also thank all
367 National Influenza Centers and laboratories that have supplied influenza viruses to the WHO
368 Collaborating Centers for Reference and Research on Influenza for analysis. The text as published
369 does not necessarily represent the official view of US-CDC, WHO, or FAO.

370

371 VII. Funding Statement

372 Avian influenza work in the Virology Unit at Institut Pasteur du Cambodge was funded, in part, by
373 the Food and Agriculture Organization of the United Nations' Global Health Security Program
374 (funded by USAID), the World Health Organization, and the Bill and Melinda Gates Foundation. H.A.
375 is supported, in part, by the German Centre for International Migration and Development. P.M.T was
376 supported by Johns Hopkins APL internal research and development. R.X., K.M.E., S.H. and V.D.
377 were funded in part by the National Institute of Allergy and Infectious Diseases, National Institutes of
378 Health, United States Department of Health and Human Services, under Contract No.
379 75N93021C00016. The funders had no role in study design, data collection and interpretation, or the
380 decision to submit the work for publication.

381

382 **VIII. Author Contributions**

383 Conceptualization: J.Y.S., R.X., A.M.P.B., K.M.E., N.S.L., V. Dhanasekaran, and E.A.K. Data
384 curation: J.Y.S., R.X., A.M.P.B., K.M.E., S.H., S.Y., S. Sin, S. Tok, K.C., S.V.H., C.R., S. Keo, L.P.,
385 H.A., Y.P., S. Kol, R.H., C.D., C.S., V.I., S.P., A.F., B. Sar, V. Dhanasekaran, and E.A.K. Formal
386 analysis: J.Y.S., R.X., A.M.P.B., K.M.E., S.H., S.Y., S. Sin, S. Tok, K.C., S.V.H., C.R., S. Keo, L.P.,
387 H.A., Y.P., S. Kol, R.H., N.S.L., V. Dhanasekaran, and E.A.K. Funding acquisition: V. Duong, A.S.,
388 V.I., S.P., A.F., B. Sar, F.F.C., N.S.L., V. Dhanasekaran, and E.A.K. Investigation: J.Y.S., R.X.,
389 A.M.P.B., K.M.E., S.H., S.Y., S. Sin, S. Tok, K.C., S.V.H., C.R., S. Keo, L.P., Y.P., S. Kol, R.H., S.
390 Tum, S. Sorn, B. Seng, Y.S., C.D., C.S., M.H., V.I., S.P., H.D., T.D., A.F., B. Sar, P.T., F.F.C., N.S.L.,
391 L.S., V. Dhanasekaran, and E.A.K. Methodology: J.Y.S., R.X., A.M.P.B., K.M.E., H.A., P.T., N.S.L., V.
392 Dhanasekaran, and E.A.K. Project administration: V. Duong, H.A., A.S., S. Tum, S. Sorn, B. Seng,
393 C.D., C.S., M.H., V.I., S.P., F.F.C., N.S.L., L.S., V. Dhanasekaran, and E.A.K. Resources: V. Duong,
394 H.A., S. Tum, S. Sorn, B. Seng, C.D., C.S., M.H., V.I., S.P., H.D., T.D., A.F., B. Sar, P.T., F.F.C.,
395 N.S.L., L.S., V. Dhanasekaran, and E.A.K. Software: P.T. and V. Dhanasekaran Supervision: V.
396 Duong, H.A., A.S., S. Tum, S. Sorn, B. Seng, Y.S., C.D., C.S., M.H., V.I., S.P., H.D., T.D., A.F., B.
397 Sar, F.F.C., N.S.L., L.S., V. Dhanasekaran, and E.A.K. Validation: J.Y.S., R.X., A.M.P.B., K.M.E.,
398 S.Y., S.V.H., V. Dhanasekaran, and E.A.K. Visualization: J.Y.S., R.X., A.M.P.B., K.M.E., V.
399 Dhanasekaran, and E.A.K. Writing - original draft: J.Y.S., R.X., A.M.P.B., K.M.E., P.T., F.F.C., N.S.L.,
400 V. Dhanasekaran, and E.A.K. Writing - review & editing: J.Y.S., R.X., A.M.P.B., K.M.E., S.H., S.Y., S.
401 Sin, S. Tok, K.C., S.V.H., C.R., S. Keo, L.P., V. Duong, H.A., Y.P., S. Kol, A.S., R.H., S. Tum, S. Sorn,
402 B. Seng, Y.S., C.D., C.S., M.H., V.I., S.P., H.D., T.D., A.F., B. Sar, P.T., F.F.C., N.S.L., L.S., V.
403 Dhanasekaran, and E.A.K.

404

405 **IX. Conflict Of Interest**

406 The authors declare no conflicts of interest.

407

408 X. References

- 409 1 Horwood, P. F. *et al.* Co-circulation of Influenza A H5, H7, and H9 Viruses and Co-infected
410 Poultry in Live Bird Markets, Cambodia. *Emerg Infect Dis* **24**, 352-355,
411 doi:10.3201/eid2402.171360 (2018).
- 412 2 Suttie, A. *et al.* The evolution and genetic diversity of avian influenza A(H9N2) viruses in
413 Cambodia, 2015 - 2016. *PLoS One* **14**, e0225428, doi:10.1371/journal.pone.0225428 (2019).
- 414 3 Suttie, A. *et al.* Diversity of A(H5N1) clade 2.3.2.1c avian influenza viruses with evidence of
415 reassortment in Cambodia, 2014-2016. *PLoS One* **14**, e0226108,
416 doi:10.1371/journal.pone.0226108 (2019).
- 417 4 Suttie, A. *et al.* Detection of Low Pathogenicity Influenza A(H7N3) Virus during Duck
418 Mortality Event, Cambodia, 2017. *Emerg Infect Dis* **24**, 1103-1107,
419 doi:10.3201/eid2406.172099 (2018).
- 420 5 Horm, S. V. *et al.* Intense circulation of A/H5N1 and other avian influenza viruses in
421 Cambodian live-bird markets with serological evidence of sub-clinical human infections.
422 *Emerg Microbes Infect* **5**, e70, doi:10.1038/emi.2016.69 (2016).
- 423 6 Siegers, J. Y. *et al.* Genetic and Antigenic Characterization of an Influenza A(H3N2) Outbreak
424 in Cambodia and the Greater Mekong Subregion during the COVID-19 Pandemic, 2020. *J*
425 *Virology* **95**, e0126721, doi:10.1128/JVI.01267-21 (2021).
- 426 7 Siegers, J. Y. *et al.* Detection and phylogenetic analysis of contemporary H14N2 Avian
427 influenza A virus in domestic ducks in Southeast Asia (Cambodia). *Emerg Microbes Infect* **13**,
428 2297552, doi:10.1080/22221751.2023.2297552 (2024).
- 429 8 Rith, S. *et al.* Identification of molecular markers associated with alteration of receptor-
430 binding specificity in a novel genotype of highly pathogenic avian influenza A(H5N1) viruses
431 detected in Cambodia in 2013. *J Virology* **88**, 13897-13909, doi:10.1128/jvi.01887-14 (2014).

- 432 9 Ly, S. *et al.* Seroprevalence and Transmission of Human Influenza A(H5N1) Virus before and
433 after Virus Reassortment, Cambodia, 2006-2014. *Emerg Infect Dis* **23**, 300-303,
434 doi:10.3201/eid2302.161232 (2017).
- 435 10 Suttie, A. *et al.* Avian influenza in the Greater Mekong Subregion, 2003-2018. *Infect Genet*
436 *Evol* **74**, 103920, doi:10.1016/j.meegid.2019.103920 (2019).
- 437 11 Edwards, K. M. *et al.* Detection of Clade 2.3.4.4b Avian Influenza A(H5N8) Virus in Cambodia,
438 2021. *Emerg Infect Dis* **29**, 170-174, doi:10.3201/eid2901.220934 (2023).
- 439 12 Brauer, R. & Chen, P. Influenza virus propagation in embryonated chicken eggs. *J Vis Exp*,
440 doi:10.3791/52421 (2015).
- 441 13 Matrosovich, M., Matrosovich, T., Carr, J., Roberts, N. A. & Klenk, H. D. Overexpression of
442 the alpha-2,6-sialyltransferase in MDCK cells increases influenza virus sensitivity to
443 neuraminidase inhibitors. *J Virol* **77**, 8418-8425, doi:10.1128/jvi.77.15.8418-8425.2003
444 (2003).
- 445 14 Thielen, P. *Influenza Whole Genome Sequencing with Integrated Indexing on Oxford*
446 *Nanopore Platforms*, <<https://dx.doi.org/10.17504/protocols.io.kxygxm7yzl8j/v1>> (2022).
- 447 15 Ratcliff, J. D. *et al.* Improved Resolution of Highly Pathogenic Avian Influenza Virus
448 Haemagglutinin Cleavage Site Using Oxford Nanopore R10 Sequencing Chemistry. *bioRxiv*,
449 2023.2009.2030.560331, doi:10.1101/2023.09.30.560331 (2023).
- 450 16 Shepard, S. S. *et al.* Erratum to: Viral deep sequencing needs an adaptive approach: IRMA,
451 the iterative refinement meta-assembler. *BMC Genomics* **17**, 801, doi:10.1186/s12864-016-
452 3138-8 (2016).
- 453 17 Khare, S. *et al.* GISAID's Role in Pandemic Response. *China CDC Wkly* **3**, 1049-1051,
454 doi:10.46234/ccdcw2021.255 (2021).
- 455 18 Shen, W., Le, S., Li, Y. & Hu, F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File
456 Manipulation. *PLOS ONE* **11**, e0163962, doi:10.1371/journal.pone.0163962 (2016).

- 457 19 Huddleston, J. *et al.* Augur: a bioinformatics toolkit for phylogenetic analyses of human
458 pathogens. *J Open Source Softw* **6**, doi:10.21105/joss.02906 (2021).
- 459 20 Markin, A. *et al.* PARNAS: Objectively Selecting the Most Representative Taxa on a
460 Phylogeny. *Syst Biol* **72**, 1052-1063, doi:10.1093/sysbio/syad028 (2023).
- 461 21 Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software Version 7:
462 Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**, 772-780,
463 doi:10.1093/molbev/mst010 (2013).
- 464 22 Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2 – Approximately Maximum-Likelihood
465 Trees for Large Alignments. *PLOS ONE* **5**, e9490, doi:10.1371/journal.pone.0009490 (2010).
- 466 23 Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective
467 Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology
468 and Evolution* **32**, 268-274, doi:10.1093/molbev/msu300 (2014).
- 469 24 Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T.-Y. ggtree: an r package for visualization and
470 annotation of phylogenetic trees with their covariates and other associated data. *Methods in
471 Ecology and Evolution* **8**, 28-36, doi:https://doi.org/10.1111/2041-210X.12628 (2017).
- 472 25 Byrne, A. M. P. *et al.* Investigating the Genetic Diversity of H5 Avian Influenza Viruses in the
473 United Kingdom from 2020-2022. *Microbiol Spectr* **11**, e0477622,
474 doi:10.1128/spectrum.04776-22 (2023).
- 475 26 Suttie, A. *et al.* Inventory of molecular markers affecting biological characteristics of avian
476 influenza A viruses. *Virus Genes* **55**, 739-768, doi:10.1007/s11262-019-01700-z (2019).
- 477 27 de Bruin, A. C. M. *et al.* Hemagglutinin Subtype Specificity and Mechanisms of Highly
478 Pathogenic Avian Influenza Virus Genesis. *Viruses* **14**, doi:10.3390/v14071566 (2022).
- 479 28 Herfst, S. *et al.* Airborne transmission of influenza A/H5N1 virus between ferrets. *Science*
480 **336**, 1534-1541, doi:10.1126/science.1213362 (2012).

- 481 29 Linster, M. *et al.* Identification, characterization, and natural selection of mutations driving
482 airborne transmission of A/H5N1 virus. *Cell* **157**, 329-339, doi:10.1016/j.cell.2014.02.040
483 (2014).
- 484 30 Matrosovich, M. N. *et al.* Avian Influenza A Viruses Differ from Human Viruses by
485 Recognition of Sialyloligosaccharides and Gangliosides and by a Higher Conservation of the
486 HA Receptor-Binding Site. *Virology* **233**, 224-234,
487 doi:<https://doi.org/10.1006/viro.1997.8580> (1997).
- 488 31 Russell, C. A. *et al.* The potential for respiratory droplet-transmissible A/H5N1 influenza virus
489 to evolve in a mammalian host. *Science* **336**, 1541-1547, doi:10.1126/science.1222526
490 (2012).
- 491 32 Shu, L. P. *et al.* Genetic reassortment in pandemic and interpandemic influenza viruses.
492 *European Journal of Epidemiology* **12**, 63-70, doi:10.1007/BF00144430 (1996).
- 493 33 Bao, P. *et al.* Human infection with a reassortment avian influenza A H3N8 virus: an
494 epidemiological investigation study. *Nature Communications* **13**, 6817, doi:10.1038/s41467-
495 022-34601-1 (2022).
- 496 34 Gao, R. *et al.* Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J*
497 *Med* **368**, 1888-1897, doi:10.1056/NEJMoa1304459 (2013).
- 498 35 Horwood, P. F. *et al.* Transmission experiments support clade-level differences in the
499 transmission and pathogenicity of Cambodian influenza A/H5N1 viruses. *Emerg Microbes*
500 *Infect* **9**, 1702-1711, doi:10.1080/22221751.2020.1792353 (2020).
- 501 36 Ly, S., Van Kerkhove, M. D., Holl, D., Froehlich, Y. & Vong, S. Interaction between humans
502 and poultry, rural Cambodia. *Emerg Infect Dis* **13**, 130-132, doi:10.3201/eid1301.061014
503 (2007).
- 504 37 Vang, D., Chau, D., Vutha, K. & Um, S. Knowledge, Attitudes, and Practices Related to Avian
505 Influenza (H5N1) After the Outbreak in Rural, Cambodia. *medRxiv*,
506 2023.2009.2025.23296059, doi:10.1101/2023.09.25.23296059 (2023).

- 507 38 Waisbord, S. R., Michaelides, T. & Rasmuson, M. Communication and social capital in the
508 control of avian influenza: lessons from behaviour change experiences in the Mekong
509 Region. *Glob Public Health* **3**, 197-213, doi:10.1080/17441690801887620 (2008).
- 510 39 World Health Organization. *Addressing the low risk perception of avian flu in Cambodia*,
511 <[https://www.who.int/westernpacific/news-room/feature-stories/item/addressing-the-low-](https://www.who.int/westernpacific/news-room/feature-stories/item/addressing-the-low-risk-perception-of-avian-flu-in-cambodia)
512 [risk-perception-of-avian-flu-in-cambodia](https://www.who.int/westernpacific/news-room/feature-stories/item/addressing-the-low-risk-perception-of-avian-flu-in-cambodia)> (2024).
- 513 40 Manabe, T. *et al.* Impact of Educational Intervention Concerning Awareness and Behaviors
514 Relating to Avian Influenza (H5N1) in a High-Risk Population in Vietnam. *PLOS ONE* **6**,
515 e23711, doi:10.1371/journal.pone.0023711 (2011).
- 516 41 Manabe, T. *et al.* Knowledge, attitudes, practices and emotional reactions among residents
517 of avian influenza (H5N1) hit communities in Vietnam. *PLoS One* **7**, e47560,
518 doi:10.1371/journal.pone.0047560 (2012).
- 519 42 Koh, G. C. *et al.* Avian influenza and South Jakarta primary healthcare workers: a controlled
520 mixed-method study. *Tropical medicine & international health : TM & IH* **14**, 817-829,
521 doi:10.1111/j.1365-3156.2009.02297.x (2009).
- 522 43 Tzeng, H.-M. & Yin, C.-Y. Nurses' Fears and Professional Obligations Concerning Possible
523 Human-to-Human Avian Flu. *Nursing Ethics* **13**, 455-470, doi:10.1191/0969733006nej893oa
524 (2006).
- 525 44 Kreslake, J. M. *et al.* The Intersection of Care Seeking and Clinical Capacity for Patients With
526 Highly Pathogenic Avian Influenza A (H5N1) Virus in Indonesia: Knowledge and Treatment
527 Practices of the Public and Physicians. *Disaster medicine and public health preparedness* **10**,
528 838-847, doi:10.1017/dmp.2016.81 (2016).
- 529 45 World Health Organization. *Genetic and antigenic characteristics of clade 2.3.4.4b A(H5N1)*
530 *viruses identified in dairy cattle in the United States of America*,
531 <[https://www.who.int/publications/m/item/genetic-and-antigenic-characteristics-of-clade-](https://www.who.int/publications/m/item/genetic-and-antigenic-characteristics-of-clade-2.3.4.4b-a(h5n1)-viruses-identified-in-dairy-cattle-in-the-united-states-of-america)
532 [2.3.4.4b-a\(h5n1\)-viruses-identified-in-dairy-cattle-in-the-united-states-of-america](https://www.who.int/publications/m/item/genetic-and-antigenic-characteristics-of-clade-2.3.4.4b-a(h5n1)-viruses-identified-in-dairy-cattle-in-the-united-states-of-america)> (2024).

533 46 Burrough, E. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Clade 2.3.4.4b Virus Infection
534 in Domestic Dairy Cattle and Cats, United States, 2024. *Emerging Infectious Disease journal*
535 **30**, 1335, doi:10.3201/eid3007.240508 (2024).

536 47 Uyeki, T. M. *et al.* Highly Pathogenic Avian Influenza A(H5N1) Virus Infection in a Dairy Farm
537 Worker. *New England Journal of Medicine* **390**, 2028-2029, doi:doi:10.1056/NEJMc2405371
538 (2024).

539

540 **IX. Supplemental Figures and Tables**

541

542 **Supplemental Table 1.** Molecular marker analysis and amino acid variation between human A/H5N1
543 virus in Cambodia, 2023-2024.

544 **Supplemental Table 2.** GISAID accession numbers.

545