

1 **Fascioliasis in north-central Vietnam: assessing community** 2 **knowledge, attitudes, and practices**

3 Vinh Hoang Quang^{1,2,3,4}, Bruno Levecke², Dung Do Trung¹, Binh Vu Thi Lam¹, Le Thuy Dung¹,
4 Nguyen Duc Thuy¹, Tran Thi Tuyen¹, Nguyen Thi Thu Hien¹, Nguyen Ngoc Ha¹, Brecht
5 Devleeschauwer^{2,5}, Kathy Goossens³, Theodorus de Jong³, Linda Paredis³, Katja Polman⁶, Steven
6 Callens⁴, Pierre Dorny^{2,3}, Veronique Dermauw^{3*}

7 ¹Department of Parasitology, National Institute of Malariology, Parasitology and Entomology,
8 Hanoi, Vietnam

9 ²Department of Translational Physiology, Infectiology and Public Health, Ghent University,
10 Merelbeke, Belgium

11 ³Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium

12 ⁴Department of Internal Medicine and Pediatrics, Faculty of Medicine and Health Sciences, Ghent
13 University, Ghent, Belgium

14 ⁵Department of Epidemiology and Public Health, Sciensano, Brussels, Belgium

15 ⁶Department of Public Health, Institute of Tropical Medicine, Antwerp, Belgium

16 *Corresponding author: vdermauw@itg.be

17 **Abstract**

18 **Background:** Fascioliasis, caused by *Fasciola hepatica* and *Fasciola gigantica*, is a zoonotic
19 disease that significantly impacts public health in agricultural communities, particularly in Vietnam.
20 This study aims to assess the knowledge, attitudes, and practices (KAP) regarding fascioliasis
21 among residents in a rural community in Vietnam.

22 **Methodology/Principal Findings:** A cross-sectional study was conducted in Dong Thanh
23 commune, north-central Vietnam. A random sample of 621 households was selected, and 1,398
24 individuals participated in this study. All participants were interviewed for individual KAP to assess
25 KAP regarding fascioliasis. In contrast, household heads were also interviewed about household
26 practices, including life cycle knowledge, health-seeking behavior, water and sanitation practices,
27 livestock and crop management, and dietary habits. Descriptive statistics were used to assess KAP,
28 and generalized linear models were applied to examine the associations between socio-demographic
29 variables and KAP. Awareness of fascioliasis was low, with 85% (1,193/1,398) of respondents
30 reporting no prior knowledge. Detailed understanding of transmission, symptoms, and prevention
31 was limited. Only 9% (124/1,398) of participants could accurately identify the symptoms, while
32 12% (168/1,398) were knowledgeable about preventive measures. A high percentage of households
33 treated drinking water (99%, 613/619), and consumption of raw vegetables was widespread, with
34 93% (1,083/1,168) of individuals and 95% of households reporting this practice. Males were less
35 likely to engage in non-risky practices than females (odds ratio: 0.696; 95% confidence interval:
36 0.591-0.819). Most households (85%, 522/617) sourced plants from their parcels, and 67%
37 (395/588) used animal manure as fertilizer.

38 **Conclusion/Significance:** The study reveals significant gaps in knowledge, attitudes, and practices
39 related to fascioliasis in Dong Thanh commune. There is a pressing need for targeted educational

40 programs to enhance community awareness and promote safer practices to mitigate the risk of
41 fascioliasis transmission. Future interventions should emphasize gender-specific education and
42 broader community involvement to address these gaps effectively.

43 **Author Summary**

44 Fascioliasis, caused by *Fasciola hepatica* and *Fasciola gigantica*, is a zoonotic foodborne disease
45 emerging in Vietnam. This study investigates the knowledge, attitudes, and practices (KAP) related
46 to fascioliasis among residents of a commune in north-central Vietnam, where the disease is highly
47 prevalent in livestock. A cross-sectional survey of 1,398 individuals and 621 households assessed
48 life cycle knowledge, health-seeking behaviors, dietary habits, water and sanitation practices, and
49 management of both livestock and crops. The results show that 85% of respondents were unaware
50 of fascioliasis, with inadequate knowledge about its transmission, symptoms, and prevention. Water
51 treatment is commonly practiced, but consumption of raw vegetables persists. Socio-demographic
52 factors, especially gender, males more likely to engage in risky practices. The findings highlight
53 critical gaps in KAP, emphasizing the need for targeted education to improve awareness and
54 promote safer practices, incorporating gender-specific strategies and community engagement.

55 **1. Introduction**

56 Fascioliasis is an ubiquitous parasitic disease caused by the liver flukes *Fasciola hepatica*
57 and *Fasciola gigantica* [1]. It primarily affects ruminants causing economic losses due to
58 significant reductions in milk and meat production [2-5]. Fascioliasis represents a significant
59 public health concern due to its substantial health risks to humans [6, 7]. The World Health
60 Organization (WHO) classifies fascioliasis as a neglected tropical disease, with approximately
61 50 million individuals infected globally and about 180 million at risk [8]. During the last
62 decades, fascioliasis has become an emerging public health problem in many areas of the world,
63 with an estimated annual loss of 35,000 disability adjusted life years [9]. The disease is acquired
64 through ingesting water or plants contaminated with metacercariae, the parasite's infective stage,
65 which thrives in environments where human and livestock activities intersect [10].

66 The northern and central regions of Vietnam are highly endemic for both animal and
67 human fascioliasis [11, 12]. Traditional practices, such as the consumption of raw aquatic plants,
68 facilitate the transmission of the disease to humans [11, 13]. In a comprehensive study by De et
69 al., which analyzed over 53,000 cases of fascioliasis in Vietnam from 1995 to 2019, a notable
70 upward trend in the incidence of the disease was observed. The study documented a rise in the
71 number of cases from a mere two cases reported in 1995 to a peak of 4,026 cases identified in
72 2019 [12].

73 In 2020, the General Statistics Office of Vietnam estimated the population of cattle and
74 buffalos at 6.23 and 2.33 million, respectively [14]. In Vietnam, cattle and buffalos are typically
75 managed by small-scale farmers, each maintaining herds of fewer than 30 animals [15],
76 allowing them to freely graze in fields [11]. This practice significantly elevates the risk of
77 infection among these animals, as they are exposed to contaminated environments for the
78 majority of the year [16]. The prevalence of fascioliasis in livestock is high, especially in the

79 Red River Delta area and in Central Vietnam, with infection rates in livestock reaching up to
80 72% [17].

81 Despite the clear public health significance of fascioliasis, there is limited information on
82 the community knowledge, attitudes, and practices (KAP) regarding the disease in Vietnam.
83 Understanding these factors is crucial for designing effective intervention strategies. Studies in
84 Iran have shown that awareness and knowledge of fascioliasis are often low in affected
85 communities, leading to behaviors that increase the risk of infection, such as the consumption of
86 raw or improperly treated vegetables and untreated water [18]. Moreover, the utilization of
87 livestock feces as fertilizer and the climatic conditions favoring the presence of suitable snail
88 hosts further perpetuate the persistence and transmission of the disease [1, 19]. Previous
89 research highlighted the critical role of health education and community engagement in
90 controlling zoonotic diseases [17, 20]. Given the rising number of documented fascioliasis cases
91 [12] and the existing gap in comprehensive studies on the disease's epidemiology and control,
92 this study aims to evaluate the KAP related to fascioliasis among residents of Dong Thanh, a
93 commune in Nghe An province (north-central Vietnam).


94 **2. Methods**

95 **2.1. Ethics statement**

96 This study was approved by the institutional review boards of both the National Institute
97 of Malariology, Parasitology, and Entomology (NIMPE) in Vietnam (Approval No. 02-
98 2022/HDDD) and the Ghent University Hospital/Faculty of Medicine and Health Sciences,
99 Ghent University in Belgium (Approval No. BC-08915). Prior to commencing the fieldwork, the
100 study's aims and procedures were communicated to provincial and district health officers, as
101 well as the Dong Thanh Commune People's Committee, which included medical doctors and
102 veterinarians. These local authorities subsequently granted their approval for the study. During
103 the initial visit, detailed information about the study's objectives and procedures was provided to
104 all participants. Signed informed consent was obtained from each participant aged 18 years and
105 older before they were enrolled in the study. For minors (i.e., participants between 5 and 1 years
106 old), verbal assent was solicited and written informed consent was secured from their parents or
107 guardians.

108 **2.2. Study site**

109 This study was conducted in Dong Thanh commune in Yen Thanh district, Nghe An
110 province in north-central Vietnam (**Figure 1**). As of March 2022, this commune has 8,490
111 registered habitants across 2,036 households. The main occupation of the residents is farming.
112 The selection of this study site was based on (i) historical data on the presence of human
113 fascioliasis, (ii) culinary behaviors that favor disease transmission (e.g. consumption of raw
114 vegetables), (iii) the presence of both cattle and buffalos that have access to crop fields, and (iv)
115 the accessibility of the study site.



116

117 **Fig 1. Map of Vietnam showing the locations of Nghe An province, Yen Thanh district, and Dong Thanh**
118 **commune.** The map was created using QGIS Desktop version 3.36.2 software

119

120 **2.3. Study design and data collection**

121 Between April and May 2022, a cross-sectional study was conducted to assess the current
122 KAP related to fascioliasis within the Dong Thanh commune. This study was part of a multi-
123 host study with the primary goal to estimate the prevalence of fascioliasis across all hosts in the
124 life cycle. From a list with the commune's households provided by the local authorities,
125 households with a minimum of three members were extracted from this list. Subsequently, a
126 random sampling method was applied to this subset to select 621 households. During household
127 visits, if more than three individuals were present, a maximum of three eligible individuals were
128 randomly chosen to participate. In cases when the household comprised three or fewer
129 individuals, all members were invited to participate. After the eligibility of the selected
130 individuals was confirmed, written informed consent was sought for the selected adults, while
131 for children aged 5 to 17, verbal assent combined with written informed consent from their
132 parents or guardians was sought. All selected individuals were invited to participate in the
133 individual questionnaire. The household questionnaire was specifically administered to the

134 household head or the member who possessed the most comprehensive knowledge of household
135 activities. Finally, for all participants, general demographic details were collected including
136 gender, age, occupation, education level, and marital status.

137 **2.4. Questionnaires**

138 The individual questionnaire consisted of three comprehensive sections, each designed to
139 capture specific aspects of participants' KAP regarding fascioliasis. The first section contained
140 questions on general demographic information, including age, gender, education level, and
141 distance from the nearest health clinic. The second section focused on knowledge and awareness
142 of fascioliasis, featuring questions on participants' familiarity with the disease, its symptoms,
143 modes of transmission, and prevention strategies. The third section explored participants'
144 attitudes and practices related to fascioliasis, including health-seeking behaviors, dietary habits,
145 and the frequency of consuming raw vegetables. This section also included questions on
146 healthcare access and reasons for not seeking formal medical treatment.

147 The household questionnaire was structured into four sections (**Info S1**). The first section
148 collected demographic data, including age, gender, occupation, educational attainment, marital
149 status, and participants' familiarity with fascioliasis. Additionally, the GPS coordinates of each
150 household were recorded in this section. The second section encompassed 11 questions related
151 to household water and sanitation practices. The third section, consisting of 15 questions,
152 focused on livestock and crop management practices. The final section included 8 questions
153 aimed at assessing culinary habits, specifically food preparation methods and the types of plants
154 consumed.

155 Prior to field implementation, the research team received training on the questionnaire
156 content, conducted pilot interviews among team members, and refined the questions to align
157 with the local language, optimize interview duration, and enhance the clarity and

158 comprehensibility of the phrasing for the local population. Both questionnaires were generated
159 in English and then translated into Vietnamese. The questions were completed by interviewers
160 using a printed version. After the field data collection, the data were entered manually into the
161 RedCap application [21, 22]. Templates of both questionnaires are provided in **Info S1**. All
162 original data are available in **Info S2**.

163 **2.5. Statistical data analysis**

164 The data were exported from the RedCap application to an Excel file and consequently
165 analysed by the statistical software R [23]. First, all responses were presented using descriptive
166 statistics. Proportions were shown for categorical variables, whereas mean, median, and range
167 were presented for continuous variables.

168 The study examined relationships between socio-demographic factors and responses from
169 individual and household questionnaires. For each question, a new binary variable was created:
170 responses indicating non-risky practices or correct knowledge scored 1, while risky practices or
171 incorrect knowledge scored 0 (details in **Info S1**). New variables were created to sum up the 1s
172 and 0s, resulting in a total score for individual practices, household practices, and individual
173 knowledge separately.

174 Associations between socio-demographic variables (gender, age, education and
175 employment) and individual practice, household practice, and individual knowledge scores were
176 analysed. For individual scores, a generalized linear mixed model with household as a random
177 factor was used. Household practice scores were analysed using a generalized linear model. The
178 full model (i.e., with all the socio-demographic variables included as fixed effects) was
179 simplified by iteratively removing the least significant variable, comparing Akaike's
180 information criterion values to find the most parsimonious model. The significance level was set
181 at 5%.

182 **3. Results**

183 **3.1. Community KAP of fascioliasis in north-central Vietnam**

184 *3.1.1. Socio-demographic characteristics of individual participants*

185 A total of 1,398 individuals were included in the study, with a slight majority of males
186 (55.22%, 770/1,398). The age of the participants ranged from 5 to 92 years, with a mean age of
187 38 and a median age of 42. Farming was the predominant occupation, accounting for
188 (828/1,397) of respondents. School children and students made up 26.77% (374/1,397) of the
189 sample, while smaller proportions were workers (5.87%, 82/1,397) and government employees
190 (3.44%, 47/1,397). In terms of education, most participants (53.86%, 753/1,398) had completed
191 secondary school. Primary school education was the highest level achieved by 17.38%
192 (243/1,398) of respondents, and 21.75% (304/1,398) had completed high school. A small
193 percentage (5.22%, 73/1,398) had achieved a university degree, and a minority (1.79%,
194 25/1,398) had not attended school.

195 *3.1.2. Knowledge and attitudes on fascioliasis among individual participants*

196 Knowledge and awareness of fascioliasis among the participants were limited. An important
197 majority (85.34%, 1,193/1,398) had never heard of fascioliasis, while only a small fraction
198 (13.02%, 182/1398) recognized it as a disease affecting humans and livestock. Additionally,
199 1.57% (22/1,398) of respondents were unsure. Only participants who had heard about the
200 disease or were unsure were asked further knowledge questions. Most of these participants were
201 aware that fascioliasis impacts the liver (77.07%, 158/205). They perceived fascioliasis as
202 somewhat serious (69.95%, 142/203), both at a personal and national level (71.57%, 146/204).
203 Commonly reported symptoms included abdominal pain (44.39%, 91/205) and fever (15.61%,
204 23/205).

205 A total of 61.46% (126/205) correctly identified fascioliasis as a parasitic infection, and
206 60.78% (124/204) understood that it is transmitted through contaminated plants or vegetables.
207 Preventive measures noted included not eating raw water plants (42.44%, 87/205) and cooking
208 vegetables (24.88%, 51/205). There was a strong belief (75.61%, 155/205) that anyone could be
209 infected, and 78.54% (161/205) were aware that specific treatment from health centers could
210 cure the disease. Information sources were primarily television (40.49%, 83/205) and health
211 workers (33.66%, 69/205), while experience with the disease amongst direct contacts, such as
212 neighbors (6.34%; 13/205) and were family members (1.95%; 4/205) were rare. Among 204
213 respondents, 58.33% (119/204) believed they could contract fascioliasis, 6.37% (13/204)
214 believed they could not, and 35.29% (72/204) were unsure. The majority of the 205 respondents
215 reported that they would primarily feel afraid if given a potential diagnosis (83.90%; 172/205),
216 followed by feeling normal (4.88%; 10/205), surprise (1.95%; 4/205), and sadness or
217 hopelessness (0.49%; 1/205). None of the respondents indicated shame, and 8.78% (18/205)
218 were uncertain about how they would react. More details on the study outcomes can be found in
219 **Info S3**. Because of the low number of participants having responded to more than one
220 knowledge questions (i.e. participants who had heard about fascioliasis), it was impossible to fit
221 a model investigating the association between the socio-demographic characteristics and the
222 individual knowledge scores.

223 *3.1.3. Practices on fascioliasis among individual participants*

224 **Table 1** summarizes the healthcare-seeking behavior among 1,398 respondents. A vast
225 majority (95.85%; 1,340/1390) expressed their intention to use health facilities for general
226 health problems, with a small fraction (3.29%; 46/1,398) preferring a pharmacy. A negligible
227 percentage (0.14%; 2/1,398) opted to rest at home, while (3.29%; 46/1,398) of the respondents
228 were uncertain about where they would seek treatment. The frequency of seeking health care
229 was also examined; among 1,340 respondents, (84.55%; 1,133) sought health care at least once a

230 year. Of the 1,392 respondents, (90.88%; 1,265/1392) consumed specific vegetables (shown in
 231 pictures), and of the 1,391 respondents, (83.47%; 1,161/1391) indicated they prepare these
 232 plants at home. Among 1,168 respondents, (89.81%; 1,049/1,168) consumed raw lettuce, and
 233 56.59% (661/1,168) consumed raw fish mint. Moreover, 60.96% (712/1,168) of these
 234 respondents indicated they consume these plants at least once a week. Regarding herbal drink
 235 consumption, out of 1,398 respondents, 45.70% (639/1,398) consumed green tea or herbal tea,
 236 while 53.29% (745/1,398) did not consume any herbal drinks. Additionally, only 3.22%
 237 (45/1,398) of respondents chewed leaves, grass, or other plants found outdoors. When
 238 investigating the association between the individual practice scores and the socio-demographic
 239 variables, the model containing only gender was best explained the data, with male participants
 240 having lower odds for a 1 score, indicating more risky practices, as compared to female
 241 participants (odds ratio (OR): 0.696; 95% confidence interval (95%CI): (0.591-0.819)).

242 **Table 1. Practices related to fascioliasis among the household participants in Dong Thanh commune.**

Question	n (multiple answers possible)	% n
Do you consume one of these plants? (The participants were shown in pictures in Info S4)	N = 1,392	
Yes	1,265	90.88
No	101	7.26
Don't know	26	1.87
Which of the following plants do you consume raw?	N = 1,168	
Water spinach, water morning glory	75	6.42
Water cress	18	1.54
Rice Paddy Herb	118	10.10
Lettuce	1,049	89.81
Sweet Cabbage	40	3.42
Lotus	5	0.43
Fish mint, lettuce mint	661	56.59
Water dropwort	16	1.37
Other non-water plant	88	7.53
Other	0	0.00
I don't consume raw plants/vegetables	77	6.59
Don't know	8	0.68
If you consume these raw, how often? (last year)	N = 1,168	
Daily	22	1.88
At least once a week	690	59.08
At least once a month	335	28.68
At least once a year	41	3.51
Don't know	32	2.74

If you prepare these at home (either raw or cooked), how often? (last year)	N = 1,168	
Daily	395	33.82
At least once a week	473	40.50
At least once a month	228	19.52
At least once a year	32	2.74
Don't know	38	3.25
Where do you mainly consume these plants?	N = 1,168	
At home	1,119	95.80
At home of other people in same village	5	0.43
At home of other people in different village	1	0.09
At the market	44	3.77
In a restaurant	6	0.51
Other	0	0.00
Don't know	12	1.03
Do you ever consume herbal drinks?	N = 1,398	
Yes, tea	636	45.49
Yes, others	3	0.21
No	745	53.29
Don't know	15	1.07

243

244 **3.2. Household practices relating to fasciolosis in north-central, Vietnam**

245 *3.2.1. Socio-demographic characteristics of household participants*

246 A total of 621 individuals were interviewed about their household practices. Of the 620
 247 respondents, 64.68% (401/620) were household heads, while 35.32% (219/620) were other
 248 household members. The gender distribution among the 621 participants was nearly balanced,
 249 with 50.89% (316/621) being male. The age of the participants ranged from 17 to 92 years with
 250 a mean of 50 and median age of 52 years. Farming was the primary occupation, representing
 251 84.68% (525/620) of the surveyed individuals. Other occupations included workers (6.94%;
 252 43/620), civil servants or government cadres (3.87%; 24/620), others (4.52%; 28/620), and
 253 miscellaneous professions (3.87%). Among the 621 individuals surveyed, education levels
 254 varied, ranging from no formal schooling (0.48%; 3/621) to university training (4.67%; 29/621).
 255 The largest group (65.70%; 408/621) had completed secondary school, followed by high school
 256 graduates (26.09%; 162/621), while 3.06% (19/621) had finished primary education.

257 3.2.2. *Water and sanitation practices in households*

258 Water and sanitation practices are described in **Table 2**. Households primarily relied on tube
 259 wells/boreholes (241/621 households, 38.81%) and rainwater collection (239/621 households,
 260 38.49%) for their drinking water, while tube wells/boreholes (339/621 households, 54.59%) and
 261 protected dug wells (234/621 households, 37.68%) were water sources used for other purposes
 262 in the households. Nearly all households (613/619 households, 98.71%) treated their drinking
 263 water, with the majority using filters or water machines (501/615 households, 81.46%). Most
 264 households (450/621 households, 72.46%) had access to flush/pour flush toilets, and the
 265 wastewater was predominantly managed through septic tanks (454/619 households, 73.34%).
 266 Additionally, a vast majority of households (609/617 households, 98.70%) had private toilet
 267 facilities.

268 **Table 2. Household water and sanitation practices in Dong Thanh commune.**

Question	n (multiple answers possible)	% n
What is the source of drinking water for members of your household?	N = 621	
Piped water into dwelling	1	0.16
Tube well/borehole	241	38.81
Protected dug well	171	27.54
Rainwater collection	239	38.49
Bottled water	0	0.00
Surface water	5	0.81
Other	0	0.00
		0.00
What is the source of water used by your household for other purposes, such as washing vegetables, cooking and hand washing?	N = 621	
Piped water into dwelling	0	0.00
Tube well/ borehole	339	54.59
Protected dug well	234	37.68
Rainwater collection	59	9.50
Bottled water	2	0.32
Surface water	5	0.81
Other	0	0.00
Don't know	0	0.00
Do you treat your water in any way to make it safer to drink?	N = 619	
Yes	613	98.71
No	6	0.97
Don't know	0	0.00
How do you treat the water to make it safer to drink?	N = 615	
Boil	115	18.70

Using a chemical	0	0.00
Use filter, water machine	501	81.46
Use ceramic filter	37	6.02
Let it stand and settle	1	0.16
Don't know	0	0.00
Other	0	0.00
What kind of toilet facility do members of your household use?	N = 621	
Flush / pour flush	450	72.46
Composting toilet (single tank)	19	3.06
Composting toilet (double tank)	134	21.58
Hanging toilet	0	0.00
Dug pit latrine	18	2.90
Bush or field	1	0.16
No toilet	1	0.16
Where does wastewater from your toilet go? (two missing)	N = 619	
Piped sewer system	29	4.68
Septic tank	454	73.34
Pond/lake/canal	7	1.13
Composting latrine	122	19.71
Other	2	0.32
Don't know	1	0.16
Is the toilet facility shared with other households?	N = 617	
Yes	7	1.13
No	609	98.70
Don't know	1	0.16

269

270 Related to livestock and crop management (**Table 3**), most households (588/621
 271 households, 94.69%) owned agricultural parcels, with a significant portion (411/585 households,
 272 70.26%) being actively engaged in farming activities. These parcels were predominantly used
 273 for crop cultivation, particularly rice, which was the primary crop for most households (565/587
 274 households, 96.25%). Livestock ownership was prevalent, with 567/584 households (96.40%)
 275 owning livestock. In terms of livestock interactions with water bodies used for vegetable
 276 cultivation, 40/567 respondents (7.05%) reported that their livestock frequently accessed these
 277 areas, while 107/567 (18.87%) indicated occasional access. Conversely, 281/567 respondents
 278 (49.56%) stated that their livestock never came into contact with such water bodies.
 279 Furthermore, 139/567 respondents (24.51%) were uncertain about their livestock's interaction
 280 with these water sources. Among these, 119/567 households (20.99%) owned cattle, and 78/567
 281 households (13.75%) owned buffalo. Manure was commonly utilized as fertilizer (395/588
 282 households, 67.18%), with composting being the preferred treatment method of animal feces

283 (388/395 households, 98.23% of those using manure). Pesticide use was widespread among
 284 households (469/587 households, 79.90%), indicating a reliance on chemical inputs for crop
 285 management.

286 **Table 3. Crop and livestock management practices of households in the Dong Thanh commune.**

Question	n (multiple answers possible)	% n
What are the feed sources for your livestock?	N = 573	
Cut and carry, grown in or near waterbodies	29	5.06
Cut and carry, grown elsewhere	93	16.23
Tethering	23	4.01
Factory product	93	16.23
Free roaming/grazing in or near waterbodies	3	0.52
Free roaming/ grazing elsewhere	17	2.97
Other processed feed source	330	57.59
Other	2	0.35
Don't know	6	1.05
Do your livestock come in or near waterbodies where vegetables for human consumption are being grown?	N = 567	
Often	40	7.05
Sometimes	107	18.87
Never	281	49.56
Don't know	139	24.51
Does your livestock come in or near water supply for crop water irrigation?	N = 567	
Often	12	2.12
Sometimes	184	32.45
Never	233	41.09
Don't know	138	24.34
What is the source of drinking-water for your livestock?	N = 573	
Piped water into dwelling	1	0.17
Tube well/ borehole	297	51.83
Protected dug well	201	35.08
Rainwater collection	47	8.20
Surface water	14	2.44
Other	0	0.00
Don't know	15	2.62
Do you use manure of cattle/buffalo/goat/horse/pig/human as fertilizer of your parcel?	N = 588	
Yes	395	67.18
No	191	32.48
Don't know	2	0.34
How do you treat manure before the use as fertiliser?	N = 395	
No treatment	7	1.77
Composting before use	388	98.23
Don't know	0	0.00
Others	0	0.00
Did you use pesticides on your parcel the last year?	N = 587	
Yes	469	79.90
No	56	9.54
Don't know	62	10.56

287

288 **Table 4** illustrates the culinary practices among households in the Dong Thanh commune.

289 Most households engaged in the consumption and preparation of various plants. Nearly all
 290 households (98.06%; 608/620) consumed these plants, and an equal percentage prepare them at
 291 home. Commonly consumed raw plants included lettuce (92.07%; 569/617), fish mint/lettuce
 292 mint (64.56%; 399/617), and rice paddy herb (13.43%; 83/617), while the most frequently
 293 cooked plants were water spinach/water morning glory (95.30%; 588/617) and sweet cabbage
 294 (85.58%; 528/617). Washing practices were prevalent, with 99.67% (616/617) of households
 295 washing these plants before use, primarily with water alone (59.48%; 367/617) or with water
 296 and salt (40.52%; 250/617). Only (1.3%; 8/617) washed the plants with vinegar and water. Most
 297 households (84.60%; 522/617) obtained these plants from their own parcels, while a smaller
 298 percentage (17.50%; 108/617) purchased them from local markets (more details can be found in
 299 the **Info S3**). None of the investigated socio-demographic variables was found to be
 300 significantly associated with the household practice scores.

301 **Table 4. Culinary practices among the households in the Dong Thanh commune**

Question	n (multiple answers possible)	% n
Does your household consume one of these plants? (pictures of vegetables are shown, see Info S4)	N = 620	
Yes	608	98.06
No	12	1.94
Don't know	0	0.00
Does your household prepare one of these plants at home? Is it either raw or cooked? (pictures of vegetables are shown in Info S4)	N = 618	
Yes	606	98.06
No	12	1.94
Don't know	0	0.00
Does your household wash these plants before use?	N = 610	
Yes	608	99.67
No	2	0.33
Don't know	0	0.00
How does your household wash these plants?	N = 617	
With water	367	59.48
With water, and vinegar	8	1.30
With water, and salt	250	40.52
Other	0	0.00

Don't know	1	0.16
Where does your households obtain these plants?	N = 617	
Local market	108	17.50
Cultivated from own parcel	522	84.60
Cultivated from another parcel	2	0.32
Supermarket	0	0.00
Restaurant	0	0.00
Other/Do not know	0	0.00

302

303

304 **4. Discussion**

305 Fascioliasis, caused by *F. hepatica* and *F. gigantica*, significantly impacts the health of
306 both humans and animals, mainly as the result of liver damage. This study provides crucial
307 insights into the KAP related to fascioliasis among residents of a rural community in the north-
308 central coast region of Vietnam.

309 **Lack of knowledge about fascioliasis**

310 A substantial knowledge deficit regarding fascioliasis was revealed in our study, with
311 about 85.34% of participants indicating no prior awareness of the disease. This awareness level
312 is considerably lower compared to findings from other regions. For example, a study in northern
313 Vietnam reported that only 11.50% of participants had never heard of liver fluke [13], consistent
314 with results from Central Vietnam [17, 24, 25]. Another study by Lugong et al., (2024) in the
315 Philippines reported a higher level of awareness, with 81.62% of participants indicating they had
316 heard of fascioliasis [26]. A detailed understanding of fascioliasis's transmission, symptoms, and
317 prevention among our respondents was notably deficient. This aligns with findings from regions
318 where fascioliasis was prevalent but underreported, indicating a global need for enhanced public
319 health education [1].

320 **Common practices of consuming raw vegetables and misconceptions about safe** 321 **handling practices**

322 Our study reveals that the consumption of raw vegetables was frequently practiced within
323 the community. Of the 1,168 respondents, 1,083 (92.72%) indicated consumption of at least one
324 type of raw vegetable. Additionally, 712 participants (61.82%) noted consuming raw vegetables
325 at a minimum frequency of once per week. These findings align with a study by Phi et al. in
326 northern Vietnam, where 91.2% reported consuming raw vegetables [13]. Conversely, Quy et

327 al.'s research found lower rates in central Vietnam, with consumption ranging from 28.20% to
328 33.80% [17].

329 In Vietnam, there is a common belief that washing raw vegetables with clean water or
330 soaking them in a saltwater or vinegar solution after washing, makes them safe to eat. Our
331 results showed that 99.84% of households wash vegetables, 59.48% using water, 40.52%
332 saltwater, and 1.30% vinegar water. However, Phuong's study demonstrated that washing
333 vegetables, whether by rinsing them under running water or scrubbing them by hand, was
334 largely ineffective at removing metacercariae, with less than 2% of the larvae being detached.
335 Additionally, exposure to a 5% acetic acid solution, comparable to commercial vinegar, resulted
336 in 93.60% to 96.00% of metacercariae surviving after less than 20 minutes of treatment.
337 Similarly, soaking vegetables in saltwater for less than 20 minutes failed to effectively kill or
338 detach the metacercariae [27]. These findings are further corroborated by Hassan et al. (2008),
339 who reported that acetic acid solutions below 5% were ineffective at killing *Fasciola*
340 metacercariae, even after 30 minutes of exposure [28].

341 **Modifying livestock and crop farming practices to reduce fascioliasis transmission**

342 During the KAP study, we concurrently investigated the prevalence of *Fasciola* infection
343 in hosts within the same area. The results revealed a notable discrepancy: animal (buffalo and
344 cattle) infection rates were 51.52% by copro-microscopy and 54.12% by antibody ELISA, while
345 human infection rates were 0% by copro-microscopy and only 0.07% by antibody ELISA [6]. At
346 the same time, this KAP study showed that conditions appear highly conducive to the potential
347 transmission of *Fasciola*. Indeed, this is due to the widespread practice of free-grazing buffalo
348 and cattle in the commune, and people frequently consume raw vegetables, including aquatic
349 plants.

350 This low infection rate in humans could be explained by several key observations. Firstly,
351 among the raw vegetables consumed, non-aquatic varieties such as lettuce (92.07%) and herbs
352 (64.56%) were the most preferred. Secondly, 84.60% of the vegetables consumed were sourced
353 from household gardens, which included some aquatic plants like fish mint, morning glory, and
354 rice paddy herb. Additionally, households primarily used agricultural land (including submerged
355 land) for rice cultivation (96.25%), with significant portions also allocated to corn (30.66%) and
356 potatoes/sweet potatoes (14.65%). Only a small fraction (3.24%) of the land was dedicated to
357 the cultivation of aquatic vegetables. Furthermore, the study found that out of 395 households
358 using livestock manure in agriculture, 388 households (98.23%) composted the manure before
359 use. In Dong Thanh commune, animal manure is typically collected and piled for 3 to 6 months
360 before being used in agriculture. Composting livestock manure before use helps reduce the risk
361 of *Fasciola* transmission. A study by Moazeni et al. (2010) demonstrated that composting at
362 high temperatures and for a sufficient duration effectively inactivates *Fasciola* eggs and reduces
363 the formation of miracidia [29].

364 The infection rate of *Fasciola* in the livestock population remains very high, exceeding
365 50%. This can be attributed to the practice of free grazing of livestock near agricultural fields,
366 which is a significant risk factor for *Fasciola* transmission. Free grazing allows cattle to
367 consume contaminated water and aquatic plants, simultaneously dispersing feces into the
368 agriculture fields. This aligns with previous research by Nguyen et al., which indicated a high
369 risk of infection in livestock due to extensive grazing practices [16].

370 In general, the practices observed in Dong Thanh commune appear to present significant
371 risks for fascioliasis transmission, particularly due to the high prevalence of infection in
372 livestock, widespread free-grazing, and the consumption of raw vegetables. Combined with the
373 low level of knowledge about the disease, this creates an apparently "dangerous" situation.
374 However, the accompanying prevalence study shows that human infection rates remain very low

375 [30], suggesting that transmission to humans is occurring at a minimal level. Upon closer
376 examination, certain practices, such as the composting of livestock manure, may play a role in
377 reducing the survival of *Fasciola* eggs and limiting transmission risks. Additionally, the practice
378 of growing vegetables in household gardens, with a preference for non-aquatic varieties, may
379 further mitigate potential exposure. Nonetheless, further research is required to gain deeper
380 insights into the specific factors that influence the transmission dynamics and to determine
381 which practices are most effective in reducing the risk of fascioliasis transmission to humans.

382 **Conclusion**

383 This study reveals significant gaps in KAP related to fascioliasis among residents of Dong
384 Thanh commune, highlighting a substantial lack of awareness. Risky behaviors, such as the
385 frequent consumption of raw vegetables in humans, and free grazing of animals in fields, are
386 prevalent. Although the current prevalence in humans in the area is reported to be low, it is key
387 to stay alert about the potential for transmission, suggesting that certain practices, such as
388 manure composting and the preference for non-aquatic vegetables, may help mitigate
389 transmission. To reduce the risk of fascioliasis transmission, targeted interventions are required.
390 These should include educational programs aimed at raising community awareness about the
391 disease and promoting safer practices in livestock management and vegetable consumption. The
392 role of household gardens in limiting exposure should also be emphasized, alongside improved
393 composting techniques to ensure the inactivation of *Fasciola* eggs. It is essential to incorporate
394 gender-specific strategies and community involvement to enhance the effectiveness of these
395 interventions. Further investigations should elucidate the factors hampering transmission to
396 humans.

397 **5. Acknowledgments**

398 This research was funded by the EmFaVie project, supported by the Flemish
399 Interuniversity Council - University Development Cooperation (VLIR-UOS, grant
400 VN2020SIN317A103 to BL and DDT), and the FasciCoM project, jointly financed by the
401 Research Foundation - Flanders (FWO, grant G0E2921N to BL) and the Vietnamese National
402 Foundation for Science and Technology Development (NAFOSTED, grant FWO.108.2020.01
403 to DDT). The funders had no role in the study's design, data collection, analysis, publication
404 decisions, or manuscript preparation. We extend our gratitude to the study participants, the
405 dedicated field team members Nguyen Huu Truong, and Dinh Cong Think, and the support from
406 the Nghe An Province Center for Disease Control, Yen Thanh Health District Center, and Dong
407 Thanh commune health stations.

408 **6. References**

- 409 1. Mas-Coma S, Bargues MD, Valero MA. Fascioliasis and other plant-borne trematode
410 zoonoses. *International Journal for Parasitology*. 2005;35(11):1255-78. doi:
411 <https://doi.org/10.1016/j.ijpara.2005.07.010>.
- 412 2. Charlier J, Duchateau L, Claerebout E, Williams D, Vercruysse J. Associations between
413 anti-*Fasciola hepatica* antibody levels in bulk-tank milk samples and production parameters
414 in dairy herds. *Preventive Veterinary Medicine*. 2007;78(1):57-66. doi:
415 <https://doi.org/10.1016/j.prevetmed.2006.09.010>.
- 416 3. Schweizer G, Braun U, Deplazes P, Torgerson PR. Estimating the financial losses due to
417 bovine fasciolosis in Switzerland. *Veterinary Record*. 2005;157(7):188-93.
- 418 4. Rioux MC, Carmona C, Acosta D, Ward B, Ndao M, Gibbs BF, et al. Discovery and
419 validation of serum biomarkers expressed over the first twelve weeks of *Fasciola hepatica*
420 infection in sheep. *International Journal for Parasitology*. 2008;38(1):123-36. doi:
421 <https://doi.org/10.1016/j.ijpara.2007.07.017>.
- 422 5. Copeman D, Copland R, Gray GD. Overcoming liver fluke as a constraint to ruminant
423 production in South-East Asia: Australian Centre for International Agricultural Research;
424 2008.
- 425 6. Mas-Coma S. Epidemiology of fascioliasis in human endemic areas. *Journal of*
426 *helminthology*. 2005;79(3):207-16.
- 427 7. Mas-Coma S. The importance of emerging and re-emerging zoonotic diseases: recognition,
428 monitoring and control. Edited by NE Odongo, M Garcia & GJ Viljoen. 2010:277.
- 429 8. Nyindo M, Lukambagire A-H. Fascioliasis: An Ongoing Zoonotic Trematode Infection.
430 *BioMed Research International*. 2015;2015:1-8. doi: 10.1155/2015/786195.

- 431 9. Fürst T, Keiser J, Utzinger J. Global burden of human food-borne trematodiasis: a
432 systematic review and meta-analysis. *The Lancet Infectious Diseases*. 2012;12(3):210-21.
433 doi: [10.1016/S1473-3099\(11\)70294-8](https://doi.org/10.1016/S1473-3099(11)70294-8).
- 434 10. Cwiklinski K, O'Neill S, Donnelly S, Dalton J. A prospective view of animal and human
435 Fasciolosis. *Parasite immunology*. 2016;38(9):558-68.
- 436 11. Bui TD, Doanh PN, Saegerman C, Losson B. Current status of fasciolosis in Vietnam: an
437 update and perspectives. *Journal of Helminthology*. 2016;90(5):511-22. Epub 2015/11/13.
438 doi: [10.1017/S0022149X15000929](https://doi.org/10.1017/S0022149X15000929).
- 439 12. De NV, Minh PN, Le TH, Dung DT, Duong TT, Tuan BV, et al. A multidisciplinary
440 analysis of over 53,000 fascioliasis patients along the 1995–2019 countrywide spread in
441 Vietnam defines a new epidemiological baseline for One Health approaches. *One Health*.
442 2024;19:100869. doi: <https://doi.org/10.1016/j.onehlt.2024.100869>.
- 443 13. Phi NTT, Nguyen TTB, Le TTH, Do DT, Lenaerts M, Losson B, et al. Foodborne zoonotic
444 trematode infections in Yen Bai, Vietnam: a situational analysis on knowledge, attitude, and
445 practice (KAP) and risk behaviors. *J Prev Med Hyg*. 2022;63(2):E310-e9. Epub 20220731.
446 doi: [10.15167/2421-4248/jpmh2022.63.2.2104](https://doi.org/10.15167/2421-4248/jpmh2022.63.2.2104). PubMed PMID: 35968061; PubMed
447 Central PMCID: PMCPMC9351405.
- 448 14. Office GS. Statistical summary book of Vietnam. Statistical Publishing House 2020. 313 p.
- 449 15. Geurden T, Somers R, Thanh NTG, Vien LV, Nga VT, Giang HH, et al. Parasitic infections
450 in dairy cattle around Hanoi, northern Vietnam. *Veterinary Parasitology*. 2008;153(3):384-
451 8. doi: <https://doi.org/10.1016/j.vetpar.2008.01.031>.
- 452 16. Lan Anh NT, Thanh DTH, Hoan DH, Thuy DT, Khong NV, Anderson N. The transmission
453 of *Fasciola* spp. to cattle and contamination of grazing areas with *Fasciola* eggs in the Red
454 River Delta region of Vietnam. *Tropical Animal Health and Production*. 2014;46(4):691-6.
455 doi: [10.1007/s11250-014-0542-4](https://doi.org/10.1007/s11250-014-0542-4).
- 456 17. Quy T, Yeatman H, Flood VM, Chuong N, Tuan B. Prevalence and risks of fascioliasis
457 among adult cohorts in Binh Dinh and Quang Ngai provinces-central Viet Nam. 2015.
- 458 18. Sazmand A, Alipoor G, Zafari S, Zolhavarieh SM, Alanazi AD, Sargison ND. Assessment
459 of knowledge, attitudes and practices relating to parasitic diseases and anthelmintic
460 resistance among livestock farmers in Hamedan, Iran. *Frontiers in veterinary science*.
461 2020;7:584323.
- 462 19. Nguyen TGT, Le TH, Dao THT, Tran TLH, Praet N, Speybroeck N, et al. Bovine
463 fasciolosis in the human fasciolosis hyperendemic Binh Dinh province in Central Vietnam.
464 *Acta Tropica*. 2011;117(1):19-22. doi: <https://doi.org/10.1016/j.actatropica.2010.09.003>.
- 465 20. Francisco I, Jiz M, Rosenbaum M, Baltazar P, Steele JA. Knowledge, attitudes and
466 practices related to schistosomiasis transmission and control in Leyte, Philippines. *PLOS*
467 *Neglected Tropical Diseases*. 2019;13(5):e0007358. doi: [10.1371/journal.pntd.0007358](https://doi.org/10.1371/journal.pntd.0007358).
- 468 21. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. A metadata-driven
469 methodology and workflow process for providing translational research informatics support.
470 *J Biomed Inform*. 2009;42(2):377-81.
- 471 22. Harris PA, Taylor R, Minor BL, Elliott V, Fernandez M, O'Neal L, et al. The REDCap
472 consortium: building an international community of software platform partners. *Journal of*
473 *biomedical informatics*. 2019;95:103208.
- 474 23. Chongsuvivatwong V. epiDisplay: epidemiological data display package. R package
475 version. 2018;3(0.1):2018.
- 476 24. Nguyen V, Nguyen V, Bui V, Tran M. Trial model of fascioliasis control in two communes
477 of Phu Cat district, Binh Dinh province. *Vietnam Journal of Practical Medicine*.
478 2011;796:152-6.

- 479 25. Nguyen TH. Study on some characteristics and treatment effectiveness of fascioliasis with
480 triclabendazole in Tinh Ky and Nghia Son communes, Quang Ngai province. National
481 Institute of Malariology □ Parasitology and Entomology: Ha Noi 2012:p. 163.
- 482 26. Lugong R, Nicolas K, Bermudez J. Knowledge, attitude and practices of small-hold cattle
483 and carabao owners towards fasciolosis in Northeastern Luzon, Philippines.
- 484 27. Nguyen Thi Xuan P. Foodborne zoonotic trematode infections and integrated control in
485 Vietnam: Aspects related to snails and vegetables 2022.
- 486 28. Hassan AAA, Shoukary NM, El-Motayam M, Morsy ATA. Efficacy of five chemicals on
487 *Fasciola gigantica* encysted metacercariae infectivity. J Egypt Soc Parasitol.
488 2008;38(3):919-28. PubMed PMID: 19209774.
- 489 29. Moazeni M, Ansari-Lari M, Masoodfar M, Hosseinzadeh S, Mootabi Alavi A. Lethal effect
490 of high temperatures on the eggs of *Fasciola hepatica*. Iranian Journal of Veterinary
491 Research. 2010;11(2):168-73.
- 492 30. Hoang VQ, Dermauw V, Do DT, Vu BTL, Nguyen HTT, Nguyen HN, et al. Prevalence and
493 risk factors related to *Fasciola* spp. transmission in Northern Vietnam: A cross-sectional
494 study in multiple hosts. medRxiv. 2024:2024.07. 10.24310192.

495

496 7. Supplementary Files

497 **Info S1** Individual and Household Questionnaires

498 **Info S2** Original data

499 **Info S3** Data analysis results

500 **Info S4** Vegetable pictures

