

Implementation of Point of Care HIV Viral load monitoring for people living with HIV in Low and Middle-income Countries: A systematic review on implementation research outcomes.

Perry Msoka^{1,2,3}, Iraseni Swai^{1,3,4}, Kennedy Ngowi¹, Ria Reis^{3,5}, Andreja Lekic², Blandina T. Mmbaga^{1,6,7}, Anita Hardon^{3,8}, Marion Sumari-de Boer^{1,3,6,8}

Affiliations:

1. Kilimanjaro Clinical Research Institute, Moshi, Tanzania
2. Amsterdam Institute for Social Science Research, Amsterdam, the Netherlands
3. Amsterdam Institute for Global Health and Development, Amsterdam, the Netherlands
4. UMC Amsterdam, location AMC, Amsterdam, the Netherlands
5. The Children's Institute, University of Cape Town, Cape Town, South Africa
6. Kilimanjaro Christian Medical University College, Moshi, Tanzania
7. Kilimanjaro Christian Medical Centre, Moshi, Tanzania
8. Knowledge, Technology and Innovation Chair group, Social Sciences Department, Wageningen University & Research, the Netherlands

Corresponding author: Perry Cyril Msoka. Kilimanjaro Clinical Research Institute
Moshi, Kilimanjaro, Tanzania
p.msoka@kcri.ac.tz

Abstract

Background

Viral load monitoring has rapidly increased among people living with HIV(PLHIV) in low- and middle-income countries (LMICs), resulting in an increased laboratory workload. The use of innovative Point of Care (PoC) or near Point of Care (n)PoC HIV Viral Load (HIV VL) monitoring has enabled improved patient care, a reduction in laboratory workload, improved clinic retention and reduced turnaround time of results. However, implementation bottlenecks of such methods are uncertain, especially when PoC or (n)PoC is implemented in remote areas in low-volume clinics. The main aim of this study was to review implementation research outcomes of (n)PoC HIV VL monitoring for PLHIV in LMICs.

Methodology

We qualitatively synthesised peer-reviewed papers to explore implementation research outcomes (IROs) of (n)PoC HIV VL monitoring. We identified studies published between January 2013 and June 2024. We used the IROs described by Proctor et al., which are acceptability, adoption, appropriateness, cost, feasibility, fidelity, penetration and sustainability. We searched using the following Mesh terms: Point of care testing, HIV, viral load, acceptability, patient acceptance of health care, adoption, facilities and services utilisation, appropriateness, cost, feasibility, fidelity, penetration, coverage, sustainability and continuity of patient care through PubMed, Cochrane and Scopus. The PRISMA diagram in Figure 1 presents the selection process of included papers.

Results

Twenty-five studies reported implementation outcomes of PoC or (n)PoC HIV VL monitoring. Near PoC HIV VL monitoring using GeneXpert is considered acceptable to patients and healthcare providers. Point of care HIV VL monitoring using mPIMA was feasible as patients received the results the same day. From a health service provider's perspective, PoC HIV VL monitoring was acceptable because it influenced patients to accept the illness and adhere to medication. Additionally, there was high testing coverage in routine PoC HIV VL monitoring centres. Fidelity was questionable in some settings due to (n)PoC HIV VL monitoring results not being delivered as intended. Additionally, we found in several studies that the (n)PoC costs are higher than standard of care test, USD 54.93 per test, at low testing volume clinics conducting 20VL tests per month compared to costs of USD 24.25 at high testing volume clinics conducting

100VL tests per month. However, costs are expected to be lower when (n)PoC HIV VL monitoring is scaled up and targeted for those at risk.

Conclusion

Implementation of PoC or (n)PoC testing for HIV viral load monitoring is acceptable and feasible and can reach a vast population. However, higher costs, limited fidelity, lower penetration and limited sustainability may hinder using (n)PoC testing in improving patient care and health outcomes. More knowledge and training should be implemented to overcome these challenges.

Keywords: Point of Care HIV Viral Load monitoring; People living with HIV; systematic review; Implementation research outcomes; Low and middle-income countries

Introduction

Globally, an estimated 39.9 million people were living with Human Immunodeficient Virus (HIV) in 2023, of whom 77% of adults and 57% of children accessed antiretroviral therapy (ART).(1) The Joint United Nations Programme on HIV and AIDS (UNAIDS) aims to reach 95-

95-95 targets by 2030 that 95% of people living with HIV (PLHIV) know their HIV status, 95% of diagnosed PLHIV receive antiretroviral therapy (ART), and 95% of PLHIV receiving ART attain viral suppression below 1,000 copies/ml.(2) To reach these goals, the World Health Organization (WHO) emphasizes that all PLHIV receive ART and routine HIV viral load (VL) testing to monitor treatment response, adherence and virologic suppression.(3)

However, monitoring the VL of PLHIV on ART in resource-limited settings is still challenging due to limited infrastructure, shortage of skilled staff, loss of samples, an inefficient system of providing results and patients being lost to follow-up.(4) To overcome those challenges, WHO has advocated for integrating Point of care HIV Viral load (PoC HIV VL) monitoring in HIV care services.(5,6)

Point of Care HIV VL monitoring entails testing at the point where healthcare is provided. The results can be given on the same day. Also, it can usually be performed by non-trained staff.(7) Near Point of Care HIV VL monitoring is testing in laboratories close to treatment facilities, while still improving the turnaround time from sampling to result.(8) The current PoC HIV VL assays that are commercially available or in late development include the PoC HIV assay of mPIMA by Abbott, and the (n)PoC HIV assays SAMBA by Diagnostics for the Real World and GeneXpert HIV 1 by Cepheid.(9)

Several studies have been conducted on the performance of PoC HIV VL monitoring. In Mozambique, mPIMA was found to have a high sensitivity of 95% and specificity of 96.5% to identify virological failure at the threshold of 1000 copies/ml when a standard conventional plasma test was used as the golden standard.(10) In addition, a study evaluating the performance of SAMBA in Malawi and Uganda found it to be accurate in differentiating patients who had a VL below or above 1000 copies/ml.(11)

Furthermore, a systematic review study from developed and developing countries found that PoC testing is preferable to centralized testing, with a sensitivity of 93.3%- 100% and specificity of 99.5%- 100% for Early Infant Diagnosis (EID), acute HIV infection diagnosis, or VL monitoring.(12)

Despite the good performance of PoC HIV VL monitoring in previous studies, several challenges exist. A systematic review of the performance and clinical utility of POC HIV viral load testing

in resource-limited settings observed that there is a need for constant electricity for the machine to function and a temperature control room for sample processing and storage.(12)

Other reported challenges of PoC HIV VL monitoring are device operation by untrained staff, routine device maintenance, and the high costs of a PoC test. All these may hamper the implementation of PoC HIV VL monitoring.

Given the above-reported challenges, future studies should explore solutions to overcome possible implementation challenges. Proctor and colleagues (2011) designed a framework that describes eight distinct implementation outcomes to guide the successful implementation of innovations: acceptability, adoption, appropriateness, feasibility, fidelity, implementation costs, penetration and sustainability. Acceptability is the perception among stakeholders that a given treatment, service, practice, or innovation is agreeable, palatable, or satisfactory. Secondly, adoption is the intention, initial decision, or action to try or employ an innovation or evidence-based practice. Thirdly, appropriateness is the perceived fit of the innovation to address a particular issue or problem. Fourthly, the cost is the financial effect of an implementation effort. It may include charges for treatment delivery, implementation strategy and use of the service setting. Fifthly, feasibility is how a new treatment or innovation can be successfully used or carried out within a given agency or setting. Sixthly, fidelity is defined as the extent to which an intervention was implemented as prescribed in the original protocol or as the program developers intended. Seventhly, penetration is integrating practice within a service setting and its subsystems. Lastly, sustainability is how a newly implemented intervention is maintained or institutionalised within a service setting's ongoing, stable operations. These implementation outcomes can give insights into challenges and assist in developing strategies to be investigated in future pragmatic trials.(13)

In this systematic review, we use the Proctor framework to explore implementation research outcomes of (n)PoC HIV VL monitoring for PLHIV in LMICs through a qualitative synthesis of peer-reviewed papers. The specific objectives were to review existing evidence on (1) the acceptability, (2) the adoption, (3) the appropriateness, (4) the cost, (5) the feasibility, (6) the fidelity, (7) the penetration and (8) the sustainability of (n)PoC HIV VL monitoring.

Methodology

Using the reporting items in systematic review guideline(14) we qualitatively synthesised scientific papers to explore the implementation research outcomes (IROs) of PoC HIV VL monitoring for PLHIV. We used the Proctor framework of IROs and pre-defined topics to define the search terms.(13)

Literature search

The initial broad relevant MESH terms were defined based on the terms ‘PoC testing’, ‘HIV’ and ‘Viral load’. We viewed the proper entry terminology for the IROs: Acceptability, Adoption, Appropriateness, Cost, Feasibility, Fidelity, Penetration and Sustainability. After defining these terms, we searched for the terms in June 2024 using a combination of the MESH and entry terms. A detailed search strategy is presented in the S1 Table.

Three databases were searched for publications published between January 2013 and June 2024: (1) PubMed, (2) Cochrane and (3) Scopus. After this, we scanned the reference list of each obtained publication found through the above search for additional publications that could fit our search criteria.

Study selection

Publications were included based on the following criteria:

- Quantitative, including trials or qualitative studies.
- Original papers.
- Focus on (n)PoC VL monitoring.
- A publication should at least describe one of the IROs as described by Proctor.

We used Covidence software <https://support.covidence.org/help/how-can-i-cite-covidence> to select and organise manuscripts and data extraction. The publications were identified and imported with a librarian (A) into the software for selecting eligible publications. After collecting publications from PubMed, Cochrane and Scopus, the titles and abstracts of identified studies were screened by two reviewers (PM, MS) for relevance. Any disagreements during the screening of titles and abstracts were resolved through discussion. In case of doubt that a publication met the criteria, it was included for full-text assessment to assess the eligibility. The full-text assessment of the quality and risk of bias of all included publications was performed by

two reviewers (PM, MS). We used the Newcastle Ottawa scale (NOS) to assess the risk of bias. (15) The tool assessed the bias in three categories which are: selection, comparability, and outcome or exposure. A star system was used: a score of seven to nine stars was considered a low risk of bias, four to six stars were considered an “unclear risk of bias”, and three or fewer stars were considered a “high risk of bias”. The study was considered of low quality if we found it to have a high risk of bias.

All manuscripts included in the full-text assessment were judged based on the quality of the paper using twenty-seven criteria selected from the PRISMA guidelines for reporting systematic reviews.(16) The PRISMA checklist can be found in the S2 Table. The twenty selected criteria covered the following items:

- The abstract was comprehensive.
- The objective did clearly state the specific aims of the study.
- The rationale was based on a well-described literature review.
- The methodology clearly described qualitative or quantitative methods, research setting and participants, methods and data collection procedures.
- Ethical issues were assessed by judging informed consent procedures, voluntary participation, no harm, confidentiality and anonymity.
- The analysis and results had to be realistic to achieve the required study objective.

Exclusion criteria were:

- Studies with outcomes that were not relevant to our review.
- Studies describing interventions outside of our area of interest.
- Studies and reviews of papers published before 2013 in which PoC VL testing was not widely available or implemented.
- Protocol studies.

Data extraction and management

One reviewer (PM) extracted the data. The following information was extracted from the included studies:

- Country where the study was performed

- Study design classified as clinical trials, cohort, qualitative or modelling studies
- Study funding sources
- Population description
- Study period, start and end date
- Inclusion and exclusion criteria
- Methods of recruitment of the participants
- Implementation of research outcomes (IROs) investigated

Ethical Approval and Informed Consent

The review was registered on the PROSPERO registry for systematic reviews. (PROSPERO 2023 CRD42023394668). The review did not involve collecting new data from human subjects. Therefore, ethical approval and informed consent were not required.

Results

Three hundred and thirteen studies were found through the initial search of PubMed, Scopus and Cochrane, and an additional five studies were found through scanning the reference lists. Forty-four duplicate studies were removed from the 318 studies. As a result, 274 studies were screened for eligibility based on title and abstract. From that, 240 irrelevant studies were excluded because the publications focused on something other than (n) PoC VL monitoring. Thirty-four studies were assessed on eligibility by reading the full text of the paper. We found that nine studies were not eligible: three had irrelevant outcomes for our review, three described interventions outside our area of interest, and three had wrong study designs, as detailed in the S3 Table. This led to 25 studies being included in the final review, as shown in Table 1. The PRISMA diagram showing detailed information on search results and the selection process is presented in S1 Figure.

Study characteristics

Twenty-five papers were included in this systematic review.(17–41)Among the studies, seventeen were quantitative studies,(18,19,21,22,24–28,30–32,34,36–38,41) six were qualitative studies,(20,29,33,35,39,40) and two were mixed methods studies.(17,23)

268

269

270 **Table 1. Characteristics of included studies.**

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Boyce et al. (2023) Uganda	Open-label, pilot study in lower-level health centre in rural Uganda	Quantitative	GeneXpert						(n)PoC test was more successful than centralized testing due to a reduction in time to results			
Reif et al. (2022) Haiti	Randomized control trials included HIV AIDS clinic	Quantitative method	GeneXpert		The majority of patients receive results the same day and immediate counselling							
Tembo et al. (2022) Lusaka, Zambia	Purposive sampling included six health facilities	Qualitative method		mPIMA	Patients were influenced to accept the illness and adhere to medication			Low costs of PoC test as the test was done on the spot				

271

272

273

274

275

276

277

278

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Boeke et al. (2021) Cameroon, Congo, Senegal, Kenya, Tanzania, Zimbabwe, Malawi	Retrospective Study included 57 Public- sector health facilities	Quantitative	GeneXpert	mPIMA			High perceived fit of PoC VL monitoring results available at the PoC or (n)PoC					PoC or (n)PoC test program to be maintained, there is a need for clear protocol diagnostic pathways
Bulterys et al (2021) Kenya	Observation study included five Ministry of Health Facilities	Mixed Methods	GeneXpert					High (n)PoC test costs USD 54.93 in low- volume clinics and a low-cost USD 24.25 in high- volume clinics	Low feasibility in some settings that shared the same instrument caused delays in VL results			

279

280

281

282

283

284

285

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Gueguen et al. (2021) Malawi and Uganda	Cohort study included four HC, one DH, one RH	Quantitative	SAMBA 1 VL			Turnaround time within a short period from blood draw to results was reduced			(n)PoC test was more successful than centralized testing		Manpower insufficient and high workload	(n)PoC test in clinics with challenges of staff availability and power
Sharma et al. (2021) South Africa	Randomized control trials included large public clinic	Quantitative method	GeneXpert		The majority of patients receive results the same day and immediate counselling							(n)PoC test increased in healthcare settings, providing adequate care according to patient needs
Wang et al. (2022) Malawi and Zimbabwe	Quantitative study in 18 health facilities	Clinic register and patient charts	GeneXpert						Multidisea se tool was highly feasible to scale up HIV testing	Lack of training to implement well (n)PoC test		

286

287

288

289

290

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Broucker et al. (2021) Southern Zambia	Cohort study included healthcare facilities in Southern Zambia	Quantitative	GeneXpert	mPIMA				Low costs in PoC or (n)PoC test including the test to confirm treatme nt failure and shifting to second- line ART regimen				
Ganesh et al. (2021) Malawi	Observation study included two large health facilities	Quantitative method	GeneXpert						(n)PoC test was more successful than centralized testing			

291

292

293

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Villa et al.(2020) Ghana	Prospective pilot study included large hospitals	Quantitative	GeneXpert		Same-day result reduces drug resistance							
Drain et al. (2020) South Africa	Randomized control trials included large urban clinic	Quantitative method	GeneXpert	mPIMA	PoC or (n)PoC test improves HIV care and increases ART adherence monitoring				PoC or (n)PoC test shows no delays in diagnosis and sufficient manpower			
Kufa et al. (2020) South Africa	Prospective Feasibility included four tertiary obstetric units	Qualitative method	GeneXpert	mPIMA					Multidisease tool was highly feasible to scale up HIV testing		Nurses are supported in managing stable patients through guidelines	
Msimango et al. (2020) South Africa	Randomized control trial included research clinic	Qualitative method	GeneXpert		(n)PoC test minimised the frequency of clinic visits for high-risk patients		Low appropriatenes of (n)PoC test in public health facilities with large numbers of people					

294

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Girdwood et al. (2020) Kenya	Quantitative study included all facilities	Geospatial cost outcome	GeneXpert	mPIMA				Low costs of PoC test to confirm treatment failure				Low sustainability of (n)PoC test in clinics with several challenges of staff availability and power supply
Vasconcellos et al. (2020) Brazil	Quantitative study included hospitals	Quantitative method		mPIMA					High feasibility of PoC due to collection of the low volume of plasma by fingerstick in sites without laboratory infrastructure			
Girdwood et al. (2019) Kenya	Quantitative study included remote healthcare facilities	Geospatial method	GeneXpert			(n)PoC test is adopted well since it is used for multi-disease testing	(n)PoC test produced results immediately					

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			nPoC	PoC								
Simeon et al. (2019) South Africa	Randomized control trial included research and urban government clinic	Mixed method	GeneXpert	mPIMA				Low costs of PoC test in high testing volume clinics				
Nicholas et al. (2019) Malawi	Retrospective Cohort included four ART clinics and a district hospital	Quantitative method	SAMBA 1 VL		(n)PoC test improved HIV care and reduced loss of follow-up							
De Necker et al. (2019) Kenya	Cohort modelling study	Quantitative method	GeneXpert					Low costs of (n)PoC test in monitori ng treatme nt failure				
Ndlovu et al. (2018) Zimbabwe	Prospective feasibility study in district hospital and rural primary healthcare facility	Quantitative method	GeneXpert			Multi- disease testing increases system efficiency	High appropriate for multi-disease testing	High costs of (n)PoC fail to manage samples in some clinics				

296

297

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriateness	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			(n)PoC	PoC								
Rasti et al. (2017) Western Uganda	Qualitative study included HC, regional and referral hospital	By Phone	GeneXpert		(n)PoC test improves the rapid treatment of patients after they receive results	Knowledge gaps in interpreting the results			(n)PoC test has been feasible as it strengthens the health system and improves healthcare	(n)PoC test not implemented well due to distrust in test accuracy/lack of training		Low sustainability in clinic with poor documentation of results
Engel et al. (2017) South Africa	Purposive sampling included home, community, clinic, peripheral lab, hospital, a diagnostic company	Qualitative method	GeneXpert		Non- laboratory healthcare workers conducted the (n)PoC test					(n)PoC test was observed that HCWs were not aware and familiar with the guideline		Low sustainability of (n)PoC test in clinics with several challenges of staff availability and power supply
Engel et al. (2015) South Africa	Purposive sampling in two urban and one rural setting	Qualitative method	GeneXpert		Patient adheres to medication. Multi-disease test evaluates multiple health conditions	Immediate provision of results and counselling to a patient		Low costs of (n)PoC test. The test was done on the same day	No delays in diagnosis and sufficient staffing to run the (n)PoC test		Nurses are supported in managing stable patients through guidelines. Manpower insufficient	

Author/Year/ Country	Study design and setting	Data Collection method	Type of POC test		Acceptability	Adoption	Appropriatenes s	Cost (High/ low)	Feasibility	Fidelity	Penetration	Sustainability
			(n)PoC	PoC								
Estill et al. (2013) South Africa	Mathematical Modelling included township ART programs	Quantitative	GeneXpert					Low costs in (n)PoC test included the test to confirm treatme nt failure and shifting to second- line ART regimen s				Sustainable when (n)PoC test costs are lowered

301 nPoC describes near point of care

302 PoC VL describes point of care viral load

303 HC describes health Centre

304

DH describes district hospital

RH describes referral hospital

ART describes Antiretroviral Therapy

Overall results

We assessed the risk of bias with the Newcastle Ottawa Scale.(15) Fourteen studies had a low risk of bias (scored 7-9).(17,18,21–25,28–32,35,38) while ten studies had an unclear risk of bias (scored 4-6).(19,20,26,27,33,34,36,37,39,40) Only one study had a high risk of bias and scored 3; this was the prospective feasibility study conducted in Zimbabwe that highlighted potential bottlenecks.(41).The Newcastle Ottawa Scale completed in detail is available in the S4 Table.

Each study described at least one of the eight IROs defined by Proctor et al.(13)

Acceptability

Ten studies described the acceptability of (n)PoC VL monitoring.(18–21,25,29,33,35,38,40) A study in Malawi, Uganda and South Africa (SA) on SAMBA I HIV, reported high satisfaction with (n)PoC VL monitoring, reduced turnaround times from blood draw to results and at the same time minimizing clinic visits for high-risk patients.(38,40) Further, in the STREAM study in SA, a retrospective cohort study on SAMBA 1 VL in Malawi, and a qualitative study in Southwestern Uganda, PoC or (n)PoC VL monitoring was highly acceptable to patients as it improves HIV care and ART adherence monitoring.(18–20)

In addition, studies conducted in Ghana and Haiti using GeneXpert showed that (n)PoC VL monitoring was highly acceptable to patients with the added benefit of results on the same day, reduced drug resistance and immediate counselling.(21,25) Likewise, a qualitative study in SA and Zambia showed that (n)PoC or PoC VL monitoring was highly acceptable by health providers as it helps patients to accept their illness and adhere to medication.(29,33) However, low acceptability of (n)PoC VL monitoring was observed in the SA studies as nonlaboratory healthcare workers conducted (n)PoC tests.(29,35)

Adoption

Four studies described the adoption of (n)PoC VL monitoring.(20,29,30,41) Medical professionals and patients observed a high uptake of (n)PoC VL monitoring in studies conducted in SA, Kenya and Zimbabwe for its immediate provision of results and counselling to patients, including multi-disease testing, which increases system efficiency.(29,30,41) However,

in a qualitative study conducted in Southwestern Uganda, health staff observed low adoption of (n)PoC VL monitoring, coupled with limited knowledge regarding interpreting the results.(20)

Appropriateness

Appropriateness was touched upon in four studies.(30,32,40,41)

A retrospective study in seven Sub-Saharan countries and the STREAM study in SA showed high appropriateness of (n)PoC VL monitoring as results were available at the near point of care.(32,40)

In addition, a quantitative study in Kenya and the prospective feasibility study conducted in Zimbabwe revealed that (n)PoC testing with GeneXpert was highly appropriate for multi-disease testing as it provides immediate results(30,41) On the contrary, the second qualitative study done in SA observed a low appropriateness of (n)PoC VL monitoring in public health facilities with large numbers of people.(40)

Costs

Costs were described in nine studies.(17,23,27–29,31,33,36,41)

An observational study of (n)PoC VL monitoring conducted in Kenya revealed high costs in low-testing volume clinics.(17) Researchers compared the (n)PoC costs with centralised testing, which were 54.93 USD per test and 25.65 USD per test respectively. The costs include personal time for a blood draw, clinical supplies, lab processing, training, lab equipment, electricity, rental space and cartridges. Furthermore, the feasibility study in Zimbabwe showed that (n)PoC VL monitoring had high costs due to failure to manage samples in some clinics.(41)

A cohort study in SA and Zambia and quantitative studies in Kenya observed low costs of PoC VL monitoring that included both the confirmation of treatment failure and the transition to second-line ART regimens.(27,28,31,36) Also, qualitative studies in SA and Zambia showed low costs of PoC tests, as it was performed on the same day and on the spot.(29,33) In addition, a mixed method study in SA showed low costs of (n)PoC VL monitoring in high testing volume clinics.(23)

Feasibility

Feasibility was described in ten studies.(17,18,20,24,26,29,34,37–39)

The SAMBA and feasibility studies in Malawi and Uganda using GeneXpert found that (n)PoC VL monitoring was more successful due to a reduction in time to results than centralised testing.(34,37,38)

Similarly, a prospective study on the implementation of PoC maternal VL and Early infant diagnosis conducted in SA and the feasibility study in Malawi and Zimbabwe showed that GeneXpert used as a multi-disease testing tool, was highly feasible to scale up HIV testing.(26,39) Furthermore, the STREAM and the qualitative studies in SA and Southwestern Uganda found high feasibility of (n)PoC VL monitoring, enabling rapid diagnosis, which, when coupled with sufficient staffing, contributes to the improvement of healthcare.(18,20,29)

Similarly, the quantitative study on developing and validating a simple and rapid way to generate a low volume of plasma used in PoC VL technology observed high feasibility in sites without laboratory Infrastructure.(24) However, low feasibility was observed in the Kenyan observational study in some settings that shared the same instrument, causing delays in VL results.(17)

Fidelity

Fidelity was described in three studies. (20,26,35)

The qualitative study in Southwestern Uganda and the feasibility study in Malawi and Zimbabwe revealed that (n)PoC VL monitoring was not implemented accurately. This was due to clients' distrust in the accuracy of the tests and insufficient training for HCWs.(20,26) Also, the qualitative study in SA showed a low fidelity of (n)PoC VL monitoring as HCWs were not aware or familiar with the guidelines.(35)

Penetration

Three studies reported penetration.(29,38,39)

The qualitative, prospective studies in SA reported high penetration of PoC or (n)PoC services when nurses are supported in managing stable patients through guidelines.(29,39) On the contrary, a qualitative study in SA and the SAMBA study in Malawi and Uganda observed low

penetration of the (n)PoC VL monitoring programme due to insufficient manpower and a high workload.(29,38)

Sustainability

Sustainability was described in six studies.(20,22,32,35,36,38)

A retrospective study in Sub-Saharan African countries, STREAM and cost-effective studies in SA observed high sustainability of PoC or (n)PoC VL monitoring, by having clear diagnostic pathways in healthcare settings, ensuring increased utilisation and adequate patient care, especially when lowering PoC test costs.(22,32,36)

On the other hand, the qualitative studies in Southwestern Uganda and SA and SAMBA studies in Malawi and Uganda observed low sustainability of (n)PoC VL monitoring in clinics, which is attributed to poor documentation of results, staff availability challenges and power supply.(20,35,38)

Discussion

This review has accumulated valuable insights into how PoC VL monitoring was implemented in various settings in LMICs and how specific implementation bottlenecks may limit proper implementation. Out of the twenty-five studies, fifteen studies specifically used GeneXpert for (n)PoC VL monitoring, six studies employed both GeneXpert and mPIMA, two studies used SAMBA 1 VL and only two studies focused on mPIMA PoC VL monitoring.

The implementation outcomes of (n)PoC VL monitoring have been widely positive in most of the studies, with reported high satisfaction among patients and HCWs. The benefits reported include reduced turnaround time, minimised clinic visits for high-risk patients, improved HIV care and increased adherence monitoring to antiretroviral therapy (ART), which significantly contribute to the acceptance and uptake of (n)PoC VL monitoring.

The short turnaround time of (n)PoC VL monitoring, providing same-day results was a key factor in its high acceptability. This led to early detection of treatment failure, minimised clinic visits, improved patient outcomes and minimised drug resistance. This was observed in a study conducted by Msimango et al., that (n)PoC VL monitoring was preferred and satisfactory over central laboratory testing as it reduced the frequency of clinic visits. (41)

This review shows that (n)PoC VL monitoring has improved the quality of HIV care from the patient's perspective. Additionally, a study by Nicholas et al., with SAMBA 1 VL test, showed that the same-day VL load test results improved HIV care and reduced follow-up loss.(19)

Moreover, PoC or (n)PoC VL monitoring has shown to lead to increased adherence monitoring to ART. This was evident from a study in South Africa by Drain et al., in which patients highlighted the advantages of MPIMA PoC or GeneXpert (n)PoC VL monitoring and that it enhances the monitoring of adherence to Antiretroviral Treatment (ART).(18)

However, the adoption of PoC VL monitoring in various studies has been hindered by several implementation bottlenecks. Challenges include issues related to non-laboratory HCWs not knowing how to interpret the results well, making it difficult to integrate (n)PoC VL monitoring into healthcare practices. This was also noted in a study by Wang et al., where HCWs lacked training to implement (n)PoC VL monitoring tests effectively.(26)

Challenges also arose in terms of appropriateness in public health facilities, with large patient volumes leading to delays in clinic flow. According to a study by Msimango et al., challenges arose for the client, as (n)PoC tests may cause delays in clinic flow due to a high number of people in the facility.(40)

Costs emerged as a barrier, particularly in low-volume clinics due to the high costs of (n)PoC VL monitoring. The study by Simeon et al., noted that (n)PoC VL monitoring reduced its costs in low-volume clinics when used for other disease testing as well. (23)

However, the complexity of multi-disease testing was identified as a challenge too in the review studies. This was also reported by Bulterys et al., indicating that clinics with multi-disease testing for (n)PoC VL monitoring using GeneXpert are accompanied by several challenges, including multiple patient queues causing delays in results and extended clinic visits.(17)

Implementation of (n)PoC VL monitoring challenges including test accuracy, were identified. Research by Rasti et al., found that patients were concerned about the diagnostic accuracy of (n)PoC VL monitoring.(20)

Insufficient manpower and high workloads were recognised as obstacles. Engel et al. reported that nurses perceived (n)PoC implementation as an extra workload, leading to patients' delays in waiting for the results.(29)

To overcome these implementation bottlenecks, clear diagnostic pathways and proper documentation of results were identified as essential measures to ensure the successful implementation of PoC or (n)PoC VL monitoring.

Also, studies showed that (n)PoC VL monitoring was acceptable and well adopted by patients and HCWs due to multi-disease testing and delivery of same-day results.(21,30) On the other hand, PoC VL monitoring was shown to be cost-effective, appropriate and feasible when conducted at the point of care.(24,32,33)

This review has several limitations. The author considered only peer-reviewed published articles, did not include unpublished reviews and the period for the included studies was restricted. Studies published between 2013-2024 and limited databases were covered. The lack of data from geographic regions outside of low- and middle-income countries raises a concern about bias and the generalizability of the findings.

The strength of this review is that we used the Proctor framework, which gives a comprehensive overview of implementation research outcomes. Gaps could suggest further research that can provide better insights into using PoC or (n)PoC VL monitoring in healthcare settings, human resources, continuous staff training and sustainable supply chain of PoC or (n)PoC facilities.

Recommendations

Point of Care VL monitoring provides rapid results followed by adherence counselling compared to nPoC VL monitoring. Rapid results provided by PoC VL monitoring improved its acceptability, leading to early detection of treatment failure, minimized clinic visits and improved patient outcomes. To address the implementation challenges of (n)PoC VL monitoring, prioritisation of clear diagnostic pathways and proper documentation are needed, as identified in this review. Further research should address gaps in healthcare settings, human resources, staff training and the sustainable supply chain for facilities that use (n)PoC tests.

Conclusion

Implementing PoC or (n)PoC testing for HIV VL monitoring was acceptable, feasible and can reach a broad population. However, high costs, limited fidelity, and lack of penetration and sustainability may hinder the use of the (n)PoC test in improving patient care and health

outcomes. Our findings indicate different implementation bottlenecks, such as non-laboratory HCWs not knowing how to interpret results and clinic delays due to high patient volumes. Additional challenges include increased costs due to low attendance, complex multi-disease testing and test accuracy concerns. Insufficient manpower and high workloads also affect implementation. Point of care knowledge and training are needed to address problems related to implementation outcomes. This enables researchers, public health communities and policymakers to collaborate effectively in developing and testing evidence-based interventions.

Acknowledgements

The authors thank the members of the EAPOC-VL study. The study is a collaboration between Uganda National Health Research Organisation/Uganda Virus Research Institute (UNHRO/UVRI, Uganda), Karolinska Institute (Sweden), The Good Samaritan Foundation Kilimanjaro Christian Medical Centre (GSF KCMC, Tanzania), Stichting Amsterdam Institute for Global Health and Development (AIGHD, the Netherlands), London School of Hygiene and Tropical Medicine (MRC/UVRI & LSHTM Uganda Research Unit, Uganda), National Institute for Medical Research (NIMR, Tanzania), Kenya Medical Research Institute (KEMRI), Center for Global Health Research (CGHR), and the University of Rwanda (UR, Rwanda).

514

515 References

- 516 1. Joint United Nations Programme on HIV/AIDS (UNAIDS) 2024. Fact sheet 2024 - Latest global and
517 regional HIV statistics on the status of the AIDS epidemic. 2024; Available from:
518 <https://www.unaids.org/en>
- 519 2. Lebelonyane R, Bachanas P, Block L, Ussery F, Alwano MG, Marukutira T, et al. To achieve 95-95-
520 95 targets we must reach men and youth: High level of knowledge of HIV status, ART coverage,
521 and viral suppression in the Botswana Combination Prevention Project through universal test and
522 treat approach. PLoS One. 2021 Aug 1;16(8 August).
- 523 3. Carmona S, Peter T, Berrie L. HIV viral load scale-up: Multiple interventions to meet the HIV
524 treatment cascade. Vol. 12, Current Opinion in HIV and AIDS. Lippincott Williams and Wilkins;
525 2017. p. 157–64.
- 526 4. Bulage L, Ssewanyana I, Nankabirwa V, Nsubuga F, Kihembo C, Pande G, et al. Factors Associated
527 with Virological Non-suppression among HIV-Positive Patients on Antiretroviral Therapy in
528 Uganda, August 2014-July 2015. BMC Infect Dis. 2017 May 3;17(1).
- 529 5. Considerations for adoption and use of multidisease testing devices in integrated laboratory
530 networks: information note. CCBY-NC-SA 3.0 IGO licence [Internet]. 2017;(June). Available from:
531 <https://iris.who.int/handle/10665/255693>
- 532 6. Meeting Report. MOLECULAR DIAGNOSTICS INTEGRATION GLOBAL MEETING REPORT. African
533 Soc Lab Med [Internet]. 2019;(10-12 JULY). Available from: <http://apps.who.int/iris>
- 534 7. Vojnov L, Taegtmeyer M, Boeke C, Markby J, Harris L, Doherty M, et al. Performance of non-
535 laboratory staff for diagnostic testing and specimen collection in HIV programs: A systematic
536 review and meta-analysis. Vol. 14, PLoS ONE. Public Library of Science; 2019.
- 537 8. Tun NN, Smithuis F, Tun NL, Min M, Hlaing MMM, van Olmen J, et al. Near point-of-care HIV viral
538 load testing: Cascade after high viral load in suburban Yangon, Myanmar. PLoS One. 2023 Apr
539 1;18(4 April).
- 540 9. Drain PK, Dorward J, Bender A, Lillis L, Marinucci F, Sacks J, et al. Point-of-Care HIV Viral Load
541 Testing: an Essential Tool for a Sustainable Global HIV/AIDS Response [Internet]. 2019. Available
542 from: <https://journals.asm.org/journal/cmr>
- 543 10. Meggi B, Bollinger T, Zitha A, Mudenyanga C, Vubil A, Mutsaka D, et al. Performance of a True
544 Point-of-Care Assay for HIV-1/2 Viral Load Measurement at Antenatal and Postpartum Services
545 [Internet]. IMPLEMENTATION SCIENCE. 2021. Available from: www.jaids.com
- 546 11. Ritchie A V., Ushiro-Lumb I, Edemaga D, Joshi HA, De Ruiter A, Szumilin E, et al. SAMBA HIV
547 semiquantitative test, a new point-of-care viral-load-monitoring assay for resource-limited
548 settings. J Clin Microbiol. 2014;52(9):3377–83.
- 549 12. Agutu CA, Ngetsa CJ, Price MA, Rinke de Wit TF, Omosa-Manyonyi G, Sanders EJ, et al. Systematic
550 review of the performance and clinical utility of point of care HIV-1 RNA testing for diagnosis and
551 care. PLoS One. 2018 Nov 1;14(6).
- 552 13. Proctor E, Silmere H, Raghavan R, Hovmand P, Aarons G, Bunger A, et al. Outcomes for

553 implementation research: Conceptual distinctions, measurement challenges, and research
554 agenda. *Adm Policy Ment Heal Ment Heal Serv Res*. 2011 Mar;38(2):65–76.

555 14. Shamseer L, Moher D, Clarke M, Ghera D, Liberati A, Petticrew M, et al. Preferred reporting
556 items for systematic review and meta-analysis protocols (prisma-p) 2015: Elaboration and
557 explanation. Vol. 349, *BMJ* (Online). BMJ Publishing Group; 2015.

558 15. Wells G, Shea B, Robertson J, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale
559 (NOS) for Assessing the Quality of Nonrandomized Studies in Meta-Analysis.

560 16. Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020
561 explanation and elaboration: Updated guidance and exemplars for reporting systematic reviews.
562 Vol. 372, *The BMJ*. BMJ Publishing Group; 2021.

563 17. Bulterys MA, Oyaro P, Brown E, Yongo N, Karauki E, Wagude J, et al. Costs of point-of-care viral
564 load testing for adults and children living with hiv in kenya. *Diagnostics*. 2021 Jan 1;11(1).

565 18. Drain PK, Dorward J, Violette LR, Quame-Amaglo J, Thomas KK, Samsunder N, et al. Point-of-care
566 HIV viral load testing combined with task shifting to improve treatment outcomes (STREAM):
567 findings from an open-label, non-inferiority, randomised controlled trial. *Lancet HIV*. 2020 Apr
568 1;7(4):e229–37.

569 19. Nicholas S, Poulet E, Wolters L, Wapling J, Rakesh A, Amoros I, et al. Point-of-care viral load
570 monitoring: outcomes from a decentralized HIV programme in Malawi. *J Int AIDS Soc*.
571 2019;22(8):1–9.

572 20. Rasti R, Nanjebe D, Karlström J, Muchunguzi C, Mwanga-Amumpaire J, Gantelius J, et al. Health
573 care workers' perceptions of point-of-care testing in a low-income country - A qualitative study in
574 Southwestern Uganda. *PLoS One*. 2017 Jul 1;12(7).

575 21. Reif LK, Belizaire ME, Rouzier V, Seo G, Severe P, Bajo Joseph JM, et al. Point-of-care viral load
576 testing among adolescents and young adults living with HIV in Haiti: a randomized control trial.
577 *AIDS Care - Psychol Socio-Medical Asp AIDS/HIV*. 2022;34(4):409–20.

578 22. Sharma M, Mudimu E, Simeon K, Bershteyn A, Dorward J, Violette LR, et al. Cost-effectiveness of
579 point-of-care testing with task-shifting for HIV care in South Africa: a modelling study. *Lancet HIV*.
580 2021 Apr 1;8(4):e216–24.

581 23. Simeon K, Sharma M, Dorward J, Naidoo J, Dlamini N, Moodley P, et al. Comparative cost analysis
582 of point-of-care versus laboratory-based testing to initiate and monitor HIV treatment in South
583 Africa. *PLoS One*. 2019 Oct 1;14(10).

584 24. Vasconcellos I, Mariani D, Azevedo MCVM d., Ferreira OC, Tanuri A. Development and validation
585 of a simple and rapid way to generate low volume of plasma to be used in point-of-care HIV virus
586 load technologies. *Brazilian J Infect Dis*. 2020 Jan 1;24(1):30–3.

587 25. Villa G, Abdullahi A, Owusu D, Smith C, Azumah M, Sayeed L, et al. Determining virological
588 suppression and resuppression by point-of-care viral load testing in a HIV care setting in sub-
589 Saharan Africa. *EClinicalMedicine*. 2020 Jan 1;18.

590 26. Wang M, Boeke CE, Rioja MR, Maparo T, Banda C, Chavula C, et al. Feasibility and impact of near-
591 point-of-care integrated tuberculosis/HIV testing in Malawi and Zimbabwe. *AIDS*. 2021 Dec
592 1;35(15):2531–7.

- 593 27. Broucker G De, Salvatore PP, Mutembo S, Moyo N, Mutanga JN, Thuma PE, et al. The cost-
594 effectiveness of scaling-up rapid point-of-care testing for early infant diagnosis of HIV in southern
595 Zambia. PLoS One. 2021 Mar 1;16(3 March).
- 596 28. de Necker M, de Beer JC, Stander MP, Connell CD, Mwai D. Economic and public health impact of
597 decentralized HIV viral load testing: A modelling study in Kenya. PLoS One. 2019 Feb 1;14(2).
- 598 29. Engel N, Davids M, Blankvoort N, Pai NP, Dheda K, Pai M. Compounding diagnostic delays: A
599 qualitative study of point-of-care testing in South Africa. Trop Med Int Heal. 2015 Apr
600 1;20(4):493–500.
- 601 30. Girdwood SJ, Nichols BE, Moyo C, Crompton T, Chimhamhiwa D, Rosen S. Optimizing viral load
602 testing access for the last mile: Geospatial cost model for point of care instrument placement.
603 PLoS One. 2019 Aug 1;14(8).
- 604 31. Girdwood SJ, Crompton T, Sharma M, Dorward J, Garrett N, Drain PK, et al. Cost-effectiveness of
605 adoption strategies for point of care HIV viral load monitoring in South Africa. EclinicalMedicine.
606 2020 Nov 1;28.
- 607 32. Boeke CE, Joseph J, Atem C, Banda C, Coulibaly KD, Doi N, et al. Evaluation of near point-of-care
608 viral load implementation in public health facilities across seven countries in sub-Saharan Africa.
609 2021; Available from: <http://onlinelibrary.wiley.com/doi/10.1002/jia2.25663/full>
- 610 33. Tembo T, Dale H, Muttu N, Itoh M, Williamson D, Mwamba C, et al. “Testing Can Be Done
611 Anywhere”: A Qualitative Assessment of Targeted Community-Based Point-of-Care Early Infant
612 Diagnosis of HIV in Lusaka, Zambia [Internet]. 2022. Available from: www.ghspjournal.org
- 613 34. Boyce RM, Ndizeye R, Ngelese H, Baguma E, Shem B, Rubinstein RJ, et al. It takes more than a
614 machine: A pilot feasibility study of point-of-care HIV-1 viral load testing at a lower-level health
615 center in rural western Uganda. PLoS Glob Public Heal. 2023 Mar 1;3(3).
- 616 35. Engel N, Davids M, Blankvoort N, Dheda K, Pant Pai N, Pai M. Making HIV testing work at the
617 point of care in South Africa: A qualitative study of diagnostic practices. BMC Health Serv Res.
618 2017;17(1).
- 619 36. Estill J, Egger M, Blaser N, Vizcaya LS, Garone D, Wood R, et al. Cost-effectiveness of point-of-care
620 viral load monitoring of antiretroviral therapy in resource-limited settings: Mathematical
621 modelling study. AIDS. 2013 Jun 1;27(9):1483–92.
- 622 37. Ganesh P, Heller T, Chione B, Gumulira J, Gugsu S, Khan S, et al. Near Point-of-Care HIV Viral
623 Load: Targeted Testing at Large Facilities [Internet]. 2020. Available from: www.jaids.com
- 624 38. Gueguen M, Nicholas S, Poulet E, Schramm B, Szumilin E, Wolters L, et al. Implementation and
625 operational feasibility of SAMBA I HIV-1 semi-quantitative viral load testing at the point-of-care in
626 rural settings in Malawi and Uganda. Trop Med Int Heal. 2021 Feb 1;26(2):184–94.
- 627 39. Kufa T, Mazanderani AH, Sherman GG, Elie Mukendi A, Murray T, Moyo F, et al. Point-of-care HIV
628 maternal viral load and early infant diagnosis testing around time of delivery at tertiary obstetric
629 units in South Africa: a prospective study of coverage, results return and turn-around times. J Int
630 AIDS Soc 2020, 23e25487 [Internet]. 2020; Available from:
631 <http://onlinelibrary.wiley.com/doi/10.1002/jia2.25487/full>
- 632 40. Msimango L, Gibbs A, Shoji H, Ngobese H, Humphries H, Drain PK, et al. Acceptability of point-of-

care viral load testing to facilitate differentiated care: a qualitative assessment of people living with HIV and nurses in South Africa. BMC Health Serv Res. 2020 Dec 1;20(1).

41. Ndlovu Z, Fajardo E, Mbofana E, Maparo T, Garone D, Metcalf C, et al. Multidisease testing for HIV and TB using the GeneXpert platform: A feasibility study in rural Zimbabwe. PLoS One. 2018 Mar 1;13(3).

Supporting information

S1 Figure PRISMA Diagram

S1 Table Search Strategy

S2 Table PRISMA checklists

S3 Table Reasons for exclusions

S4 Table Newcastle Ottawa Scale

Authors' contributions

PM performed the literature search, screening, data extraction and analysis and constructed the manuscript. MS performed the literature search, screening, data extraction and data analysis, review of articles, corrected the manuscript, IS reviewed the study results and final draft, KN reviewed the introduction and final draft. RR reviewed and corrected the manuscript. AL performed the literature search, screening, data extraction and data analysis. AH corrected the final manuscript. BM reviewed and corrected the final draft.

medRxiv preprint doi: <https://doi.org/10.1101/2024.11.04.24316630>; this version posted November 5, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).

medRxiv preprint doi: <https://doi.org/10.1101/2024.11.04.24316630>; this version posted November 5, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](#).




Figure 1: PRISMA diagram of studies screening and selection