## Determinants of antenatal care service utilization among adolescent mothers of (10-19) years of age at Kanyama level one hospital in Lusaka district.

Authors: Kenny Saka<sup>1</sup>Alice Ngoma Hazemba<sup>2</sup>, Phoebe Bwembya<sup>3</sup>

<sup>1</sup>Department of Community and Family Medicine, School of Public Health, University of Zambia

#### Abstract

**Introduction:** The provision of adequate antenatal care services is crucial for promoting maternal and child health. Therefore, it is necessary to understand the factors influencing the use of these services, especially among special groups such as adolescents. Adolescents, defined as individuals between the ages of 10 and 19 years, face specific challenges and vulnerabilities during pregnancy and childbirth. Hence, it is vital to develop targeted interventions to improve their reproductive health outcomes. This study analyzed the factors associated with antenatal care utilization among adolescent mothers in Zambia. Methodology: The study used a cross sectional study design. Data was collected using a structured questionnaire from adolescents who were 8 months pregnant or more and adolescents who were already mothers at Kanyama level one hospital in Lusaka. Descriptive statistics of individual characteristics were developed to describe the data. Associations between ANC utilization and independent variables was tested using the Chi-Square test and multivariate logistic regression. Results: The proportion of antenatal care utilization in this study was 39.37%. At multivariate analysis, factors that were significantly associated with antenatal care utilization were past pregnant experience (OR = 0.35; 95% CI: 0.14 to 0.85) and partners education (OR = 1.38; 95% CI: 1.02 to 1.87). However, Partner education did not have a statistical significance between levels of education and ANC utilization. This could have been due to a small sample size. Other studies however, reviewed a significant statistical relationship between levels of education and adolescent ANC utilization. Conclusion: The findings underscore a low proportion of antenatal care utilization among adolescent mothers. Particularly low proportion of ANC utilization was found in adolescents mothers who had never experienced pregnancy in the past.

Keywords: Utilization of Antenatal care services, adolescent mothers

#### Introduction:

Antenatal care (ANC) is a crucial component of maternal health services, designed to ensure the well-being of both mother and child during pregnancy. Among adolescent mothers aged 10-19 years, the utilization of these services is particularly critical due to the increased health risks associated with adolescent pregnancies<sup>1</sup>.

Adolescent pregnancy is a global phenomenon<sup>1</sup>. Every year in the low and middle income countries, an estimated 21 million girls aged 15–19 years become pregnant. A further 1.5 per 1000 girls aged 10-14 experience child birth globally. In Central and Southern Africa, adolescent birth rates are well above the global rate at 9 and 5 births per 1,000 adolescent girls aged 15-19 and 10-14<sup>18</sup>. In Zambia, 29% of adolescents aged 15 to 19 have begun childbearing, 24% have given birth while 5% are pregnant with their first child<sup>2</sup>.

Adolescent pregnancy have remained a major contributor to maternal and child mortality. Complications relating to pregnancy and childbirth are the leading cause of death for girls aged 10-19 globally<sup>2</sup>. Adolescents are also at significantly increased risk of delivering premature and low-birth-weight offspring. The risk of neonatal and perinatal mortality is even higher among adolescents less than 15 years. It is therefore, essential to protect the health of adolescent girls and their unborn children through ANC<sup>3</sup>.

However, the uptake of antenatal care visits among adolescent mothers in Zambia is low<sup>4</sup>. The 2021 Annual Health Statistical Report highlight that the national contribution of adolescents to the total antenatal attendances stood at 37.9%<sup>5</sup>. Hence, it is important to understand determinants of ANC to develop a targeted interventions to improve maternal and neonatal outcomes within this vulnerable population.

<sup>&</sup>lt;sup>1</sup> ibid

<sup>&</sup>lt;sup>2</sup> 1

#### **Methods and Materials**

#### Study design

This was a cross sectional study. According to Thompson, cross sectional studies are one-time survey or observation of one or more groups of subjects. This is similar to taking a snap shot of these subjects at a single point in time. They are ideal for calculating simple prevalence rates. Investigators can also use this design to attempt to examine the natural history of phenomenon by performing cross-sectional observations<sup>6</sup>. Therefore, this study design was ideal for this study.

#### **Study population**

The study population were adolescent girls who were 8 or more months pregnant and adolescent who were already mothers.

#### **Inclusion criteria**

The study included adolescent who were already mothers and adolescents who were 8 or more months pregnant. The adolescents considered were only those found within the health facility premises.

#### **Exclusion criteria**

Adolescents that were admitted at the health facility, referred to the general hospital and those that did not consent to participate in the study were excluded.

#### **Sampling Procedure**

A simple random sampling method was used to collect data from participants. The study sample size was 475.

#### **Ethical approval**

The study was approved by the University of Zambia Biomedical Research Ethics Committee (REF. No. 3624-2023). Permission for data collection was obtained from the Lusaka district medical office. Further, consent and assent forms were obtained from both the parents and the study participants.

#### **Data collection**

Data were collected using a structured questionnaire on one on one interviews. A criterion of data completeness, timeliness, precision, integrity and validity was used and maximized to ensure data quality.

#### **Data Analysis**

STATA version 14.2 was used for data analysis. Frequencies and percentages were generated and used to present categorical variables. To determine the association between ANC utilization and each independent variable, chi squared test and fishers exact test were used. This was dependent on the chi square assumptions being met. Logistic regression model was used to check for the relationship between the dependent and the independent variables. At multivariate regression analysis, an investigator led best fit model selection approach was used to select the best predictors of ANC in the multiple logistic regression model. Selection of the predictor variables was done by checking p-values. Variables found with larger p-value and not statistically significant were removed until statistically significant variables were left in the model.

## Results

# Individual, Structural and Health facility characteristics of adolescents at Kanyama level one hospital in Lusaka.

#### Table 1: Individual, Structural and Health facility characteristics

	FREQUENCY	PERCENTAGE (%)
Maternal Age		
10-14	10	2.11
15-19	465	97.89
Past pregnant experience		
Never experienced	451	94.95
Experienced	24	5.05
Marital Status		
Married	59	12.42
Not Married	416	87.58
Residing with Partner		
Yes	111	23.37
No	364	76.63
<b>Residing with Parents/Guadians</b>		
Yes	360	75.79
No	115	24.21
Employment status of Partner		
Yes	435	91.58
No	40	8.42
Employment status of Parents/Guardians		
Yes	469	98.74
No	6	1.26
Partners Education		
Primary	40	8.42
Secondary	228	48.00
Tertiary	207	43.58
<b>Parents/Guardians Education</b>		
Primary	62	13.05
Secondary	259	54.53
Tertiary	154	32.42
Adolescent friendly space		
Yes	47	9.89
No	428	90.11
Distance To Health Facility		
Less than 5km	233	49.05
More than 5km	242	50.95

Table 1 summarizes the results of the sample (n = 475). The sample comprised of adolescent mothers in the age range 10-19. The majority of the adolescent were in the age range 15-19 representing 97.89%. Less than a quarter (12.42%) reported they were married and 75.79% of the adolescents reported they were residing with their parents or guardians.

#### ANC utilization among adolescent mothers

	Pregnant Adolescents (n=475)	Percentage (%)
ANC Utilization Proportion		
Utilized	187	39.37
Not Utilized	288	60.63

Table 2 shows the proportion of adolescent mothers who had utilized and who had not utilized ANC services. Utilized in this study is defined as having 4 or more contact visits to ANC services and not utilized is having 3 or less visits to ANC services among adolescent mothers who had 8 or more months pregnant or were already mothers. Only 39.37% of the adolescent mothers had utilized the ANC services.

## Factors associated with ANC utilization

#### Table 3: Factors associated with ANC utilization

	Utilization of ANC services		P-value
	Utilized	Not utilized	
Maternal age			
10-14	1 (0.53)	9 (3.13)	0.097
15-19	189 (99.47)	279 (96.88)	
Past pregnant experience			
Never experienced	171 (91.44)	280 (97.22)	0.005
Experienced	16 (8.56)	8 (2.78)	
Marital Status			
Married	30 (16.04)	29 (10.07)	0.054
Not married	157 (83.96)	259 (89.93)	
Residing with Partner			
Yes	50 (26.74)	61 (21.18)	0.162
No	137 (73.26)	227 (78.82)	
<b>Residing with Parents/Guardians</b>			
Yes	136 (72.73)	224 (77.78)	0.209
No	51 (27.27)	64 (22.22)	
<b>Employment status of Partner</b>			
Yes	173 (92.51)	262 (90.97)	0.555
No	14 (7.49)	26 (9.03)	
<b>Employment status of Parents/Guardians</b>			
Yes	185 (98.93)	284 (98.61)	1.000
No	2 (1.07)	4 (1.39)	
Partners Education			
Primary level	13 (6.95)	27 (9.38)	0.008
Secondary level	79 (42.25)	149 (51.74)	
More than secondary (Tertiary)	95 (50.80)	112 (38.89)	
Parents/Guardians education			
Primary level	29 (15.51)	33 (11.46)	0.407
Secondary level	101 (54.01)	158 (54.86)	
More than secondary (Tertiary)	57 (30.48)	97 (33.68)	
Adolescent friendly space			
Yes	17 (9.09)	30 (10.42)	0.636
No	170 (90.91)	258 (89.58)	
Distance to Health Facility			
Less than 5KM	93 (49.73)	140 (48.61)	0.811
More than 5KM	94 (50.27)	148 (51.39)	

Table 3 shows the chi square test of association between individual, structural, health facility factors and antenatal care utilization. There was an association between Past pregnant experience and antenatal care utilization at 95% CI, (0.005). There was also an association between Partners education and antenatal care utilization at 95% CI, (0.008). The rest of the variables were not statistically significant.

## **Bivariate logistic regression analysis**

#### Table 4: Predictors of ANC utilization among adolescent mothers

Characteristics/factors	Study sample (%)	Proportional Odds Ratio (95%CI)	P-value
Maternal age			
10-14	2.11	1.0	
15-19	97.89	5.99 (0.75-47.75)	0.090
Past pregnancy experience			
Experienced	5.05	1.0	
Never experienced	94.95	0.31 (0.13-0.73)	0.008
Marital Status			
Married	12.42	1.0	
Not Married	87.58	0.59 (0.34-1.01)	0.056
<b>Residing With Partner</b>			
No	76.63	1.0	
Yes	23.37	1.36 (0.88-2.09)	0.163
<b>Residing with Parents/Guardians</b>			
No	24.21	1.0	
Yes	75.79	0.76 (0.49-1.17)	0.210
<b>Employment status of Partner</b>			
Not Employed	8.42	1.0	
Employed	91.58	1.23 (0.62-2.41)	0.555
<b>Employment status Parent/Guardians</b>			
Not Employed	1.26	1.0	
Employed	98.74	1.30 (0.24-7.18)	0.761
Partners education			
Primary	8.42	1.0	
Secondary	48.00	1.10 (0.54-2.25)	0.792
Tertiary	43.58	1.76 (0.86-3.60)	0.121
<b>Parents/Guardians education</b>			
Primary	13.05	1.0	
Secondary	54.53	0.73 (0.42-1.27)	0.264
Tertiary	32.42	0.67 (0.37-1.21)	0.186
Adolescent friendly space			
No	90.11	1.0	
Yes	9.89	0.86 (0.46-1.61)	0.637
Distance To Health Facility			
Less than 5KM	49.05	1.0	
More than 5KM	50.95	0.96 (0.66-1.38)	0.811

The findings in table 4 shows that adolescents' mothers who had never experienced past pregnant had lower odds of utilizing the ANC services. Their odds were 0.31 less likely to utilize ANC services compared to adolescents' mothers who had experienced past pregnant (OR = 0.31; 95% CI: 0.13 to 0.73).

#### Factors associated with ANC utilization

In this study, antenatal care utilization is associated with a number of individual, structural and health facility factors. However, in order to assess the contribution of all these factors to the overall variance, it was important at this stage to control for confounding by conducting a multiple logistic regression analysis

Characteristics/factors	Study	Proportional	P-value
	sample N	<b>Odds Ratio</b>	
	(%)	(POR) (95%CI)	
Past pregnancy experience			
Experienced	24 (5.05)	1.0	
Never experienced	451(94.95)	0.35 (0.14-0.85)	0.021
Partners education			
Primary	40(8.42)	1.0	
Secondary	228(48.00)	1.19 (0.58-2.45)	0.630
Tertiary	207(43.58)	1.74 (0.85-3.58)	1.33

Table 5: Multivariate analysis on factors associated with ANC utilization

After adjusting for the effect of other factors, adolescents who had never experienced past pregnancy were less likely to utilize ANC services than adolescents who had experienced pregnancy before. Their odds were 0.35 less likely to utilize ANC services (OR = 0.35; 95% CI: 0.14 to 0.85).

#### Discussion

Although antenatal care utilization is considered a success in Zambia, a notably low proportion of adolescent mothers sought antenatal care at Kanyama Level One Hospital in Lusaka. This figure reflects an ongoing challenge in achieving effective antenatal care utilization among adolescents. This finding aligns with the Annual Health Statistical Report (2022), which indicates that the contribution of adolescents to first antenatal attendances varies across provinces. Luapula had the highest attendance rate, while Lusaka had the lowest. Additionally, only four out of ten provinces, excluding six, recorded a significant contribution of adolescent mothers to first antenatal attendances. Based on these figures, the national picture suggests a concerning trend in antenatal care utilization among adolescent mothers across the provinces<sup>7</sup>.

There are variety of reasons that lead to adolescent mothers absconding ANC services. According to Kayemba (2023), attributed the absent from ANC services due to fear of pregnancy disclosure, long distance to health facility, lack of knowledge of pregnancy and ANC, travel costs to health facility and preference for traditional care to ANC<sup>8</sup>. Another study by Rahman et al, (2022), sites care seeking behavior during pregnancy. This is complex and is shaped by multiple factors of availability, accessibility, and acceptability of ANC services. Perceived usefulness and quality of ANC services significantly determine adolescent's determination to return for subsequent visits. The study further sites that ANC services is more strongly associated with country's GDP per capita than with coverage. Antenatal care (ANC) utilization varies significantly between low-income and upper-middle-income countries. Among adolescents, the primary reasons for lower ANC utilization include reduced bargaining power, lower expectations of care, and poorer health literacy<sup>9</sup>.

The current study found that adolescent mothers who had never experienced pregnancy before were less likely to utilize antenatal care services compared to those who had previous pregnancy experience. This is consistent with earlier research, which showed that adolescent mothers who had never been pregnant before had a lower prevalence of ANC utilization<sup>10</sup>. Another study by Skep found a significant association between past pregnancy experience and the use of a minimum of four ANC visits among adolescent mothers. It found that past pregnancy experience was positively associated with ANC use, with adolescent mothers who had experienced a previous birth being more likely to have at least four ANC visits than those whose most recent birth was their first<sup>11</sup>.

Chaibva reported that adolescent mothers who had never experienced pregnancy before had a lower prevalence of ANC utilization. Among adolescents that were interviewed, some feared disclosing their pregnancies to their parents or schools and preferred their pregnancies to be discovered when they were in labor. Some adolescents disclosed that they felt well, while others lacked money to pay ANC fees. Other reported reasons included limited knowledge about the benefits of ANC, reliance on traditional birth attendants, lack of required documents such as Zimbabwe national identity cards to register for ANC, and religious factors. This preference among pregnant adolescents could detrimentally influence their non-utilization of ANC services, with potentially hazardous effects for both mothers and babies<sup>3</sup>. Another study by Kayemba reported fear of pregnancy disclosure, long distances to health facilities, lack of knowledge about pregnancy and ANC, travel costs to health facilities, and a preference for traditional care over antenatal care services<sup>4</sup>.

<sup>&</sup>lt;sup>3</sup> 10

<sup>&</sup>lt;sup>4</sup> 8

.Despite current study finding a significant association between partner's education and antenatal care utilization, there was no significant association between levels of education and ANC utilization. This could have been attributed by a small sample size in this study. However, other studies such as that of Awingura found that adolescent mothers whose partners had secondary or more of education had a higher prevalence of ANC utilization compared to adolescent mothers whose partners had less than a secondary level of education. It further suggests that education positively impacts healthcare service utilization and increment of knowledge about specific issues hence recommends empowering adolescent mothers through education and decision-making to increase the utilization of maternal healthcare service<sup>12</sup>.

A study conducted by Christiana (2021) found that adolescent mothers with partners had some form of education were more likely to fully utilize ANC services compared to those whose partners had no education. This was attributed to the fact that education attainment is associated to higher awareness of the importance of ANC. No education attainment was seen to be associated with lower awareness of the importance of ANC<sup>13</sup>. Another study by Aduragbemi found that partner's education was significant predictor for ANC utilization among adolescent mothers in Kenya. In this study, it was entirely conceivable that education was a significant factor for ANC utilization. However, it only suggested that education of partner enhances both resources and agency for ANC utilization in the interest of adolescent mothers as the two primary components of empowerment<sup>14</sup>.

Anaba 2022 noticed that partner's education enabled mothers to have more access to health information, appreciate the causes of adverse pregnancy outcomes and the importance of ANC to the wellbeing of the mother and the unborn baby. In addition, partner's education enabled adolescent mother to have greater autonomy to make decisions and financial access to quality healthcare<sup>15</sup>.

#### Conclusion

The findings highlight a low rate of antenatal care (ANC) utilization among adolescent mothers, with especially low usage among those experiencing their first pregnancy. Various factors, such as lack of funds to cover ANC fees and limited knowledge about the benefits of ANC, may contribute to this low utilization. Therefore, it is crucial to implement strategies like making health facilities more accessible to adolescent mothers by addressing geographical and financial barriers and conducting community awareness campaigns to inform about the benefits of ANC and its importance for a healthy pregnancy.

#### Author's contribution

The author was responsible for the Conceptualization, Methodology, Analyzing the data, interpretation of the findings and drafting the manuscript.

#### Acknowledgement

The author wish to send sincere appreciation and acknowledge the invaluable contribution made by the lecturers at the University of Zambia, school of public health in particular DR Alice Ngoma Hazemba for supervising the work, Lusaka district medical office and Kanyama level one hospital for granting the permission to collect data and UNZABREC who granted ethical approval.

#### **Funding statement**

The study was funded by the authors

#### **Ethical approval**

The study was approved by the University of Zambia Biomedical Research Ethics Committee (REF. No. 3624-2023). Prior to data collection, written permission was obtained from the Lusaka

District Medical Office. Additionally, consent and assent forms were provided to parents or next of kin, as well as to all study participants, to obtain their consent.

### **Competing interest**

The authors have no competing interest

#### References

<sup>1</sup>Plan International (2022) Teenage Pregnancy. PI 2022

<sup>2</sup> Zambia Demographic Health statistics. Demographic and Health Survey. ZDHS 2018

<sup>3</sup> United Nations Children Emergency Fund. Antenatal Care. UNICEF 2022

<sup>4</sup> Ministry of Health. Reproductive, Maternal, Newborn, Child and Adolescent Health and Nutrition Communication and Advocacy Strategy 2018-2021. MOH 2021

<sup>5</sup> Gambirani. Exploring solutions to improve antenatal care in resource-limited settings: an expert consultation. BMC child birth and pregnancy. 2022

<sup>6</sup> Thompson. Research Study Designs: Non-experimental. Air Medical Journal 26:1. 2006

<sup>7</sup> Ministry of Health. Annual Health Statistical Report 2017-2021. MOH 2021

<sup>8</sup> Kayemba, Kozuki. A systematic review of community-to-facility neonatal referral completion rates in Africa and Asia. BMC Public Health 15:989 DOI 10.1186/s12889-015-2330-0. 2023

<sup>9</sup> Rahman. Equity in antenatal care visits among adolescent mothers: An analysis of 54 country levels trend and projection of coverage from 2000 to 2030. PMCID: PMC8932365 PMID: 35356654. 2022

<sup>10</sup> Chaibva CN, Kitange H, Teuscher T. Assessment of process quality in Tanzanian primary care, Health Policy1993;26:119-3

<sup>11</sup> Furuta M. Women's position within the household as determinant of maternal health care use in Nepal. Int Fam Plan Perspect 2020,32(1):17–27

<sup>12</sup> Awingura. Enhancing focused antenatal care in Ghana: An exploration into perceptions of practicing midwives. International Journal of Africa Nursing Sciences. 2022

<sup>13</sup> Christiana. Correlates of antenatal care usage among adolescent mothers in Nigeria: a pooled data analysis. VOL. 61, WOMEN & HEALTH 2021, NO. 1, 38–49

<sup>14</sup> Aduragbemi. Maternal Health Services Utilization: Analysis of the Demographic Health Survey 2014. Kenyan Adolescent Mothers. 2016

<sup>15</sup> Anaba. Utilization of antenatal care among adolescent and young mothers in Ghana; analysis of the 2017/2018 multiple indicator cluster survey. BMC Pregnancy and Childbirth. 2022