

1 **Title:** Role of the gut microbiome in frequent gut colonization with extended-spectrum β -
2 lactamase-producing Enterobacterales among Peruvian children

3 **Running title:** Pediatric gut microbiome and ESBL susceptibility

4 **Authors:** Maya L. Nadimpalli^{1#}, Neha Sehgal¹, Luismarcelo Rojas-Salvatierra,² Robert H.
5 Gilman^{2,3}, Ashlee M. Earl⁴, Colin J. Worby⁴, Madison Schwab,⁵ Amy J. Pickering^{6,7,8}, Monica J.
6 Pajuelo^{2,3#}

7 **Author Information:** □

8 ¹Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory
9 University, Atlanta, GA, USA □

10 ²Laboratorio de Microbiología Molecular, Laboratorios de Investigación y Desarrollo, Facultad
11 de Ciencias e Ingeniería, Universidad Peruana Cayetano Heredia, Lima, Peru □

12 ³Department of International Health, Johns Hopkins Bloomberg School of Public Health,
13 Baltimore, Maryland, USA □

14 ⁴Infectious Disease and Microbiome Program, The Broad Institute of MIT and Harvard,
15 Cambridge, MA, USA □

16 ⁵Program in Population Biology, Ecology, and Evolution, Emory University, Atlanta, GA, USA

17 ⁶Department of Civil and Environmental Engineering, University of California, Berkeley, CA,
18 USA □

19 ⁷Blum Center for Developing Economies, University of California, Berkeley, CA, USA

20 ⁸Chan Zuckerberg Biohub - San Francisco, CA, USA

21

22

23 **#Correspondence:** Maya L. Nadimpalli, Claudia Nance Rollins Room 6009, 1518 Clifton Rd
24 NE, Atlanta GA 30322, USA. Email: maya.l.nadimpalli@emory.edu

25 Monica J. Pajuelo, Av. Honorio Delgado 430, San Martín de Porres 15102, Lima Peru. Email:
26 monica.pajuelo.t@upch.pe

27

28 **Abstract (150/150)**

29 We investigated whether differences in gut microbiome development could underlie differential
30 susceptibility to gut colonization with extended-spectrum β -lactamase-producing
31 Enterobacterales (ESBL-E) over time among children in peri-urban Lima. Weekly stool and daily
32 surveys were collected from 345 children <3 years old during a 2016-19 study of enteric
33 infections. A subset of children (n=12) were rarely gut-colonized with ESBL-E from 1-16 months
34 of age. We performed short-read metagenomic sequencing of stool collected at 3, 6, 7, 9, 12,
35 and 16 months from these children and a random subset of 42 frequently colonized children,
36 and characterized differences in antibiotic use, sociodemographic characteristics, and the gut
37 microbiome over time. No gut microbiome differences were identified, but rarely colonized
38 children were more likely to live in households of lower socioeconomic status and consumed
39 fewer antibiotics. Differences in exposures, rather than gut microbiome development, might
40 underlie patterns of ESBL-E gut colonization among children in peri-urban Lima.

41

42 **Introduction (3497/3500 words)**

43 Antimicrobial-resistant (AMR) pathogens are a public health threat, particularly in low- and
44 middle-income (LMIC) countries like Peru where the infectious disease burden remains high.
45 Gut colonization with extended-spectrum β -lactamase-producing Enterobacterales (ESBL-E),
46 which are resistant to 3rd generation cephalosporins, is common among children living in
47 shantytowns of LMIC cities and is a risk factor for infections, particularly in healthcare
48 settings.(1–5)

49 Previous studies in low-resource settings have sought to identify risk factors for
50 children’s ESBL-E acquisition.(6–8) However, it remains unknown why some children are rarely
51 or never colonized with ESBL-E, while others raised in the same settings are persistently gut-
52 colonized. Intestinal “colonization resistance” refers to the ability of gut microbiota to inhibit
53 colonization and overgrowth of exogenous bacteria.(9) Identifying microbial taxa that inhibit
54 children’s gut colonization with ESBL-E over time, as well as early-life exposures that support
55 the assembly and maintenance of these protective taxa, could be helpful in designing future
56 interventions to protect children’s health.

57 Our team recently led a prospective birth cohort study of enteric infections among 345
58 children living in a shantytown of peri-urban Lima. We previously used culture-based methods to
59 characterize children’s gut colonization with ESBL-producing *Escherichia coli* and ESBL-
60 producing *Klebsiella* spp., *Enterobacter* spp., and *Citrobacter* spp. (KEC) at or near 1, 3, 4, 5, 6,
61 7, 9, 12, and 16 months of age for a subset of 112 children enrolled in the cohort study.(4) We
62 noted that while most children were frequently gut-colonized with ESBL-E over time, a fraction
63 (n=12) were rarely or never colonized. Here, we investigated whether differences in gut
64 microbiome development could underlie differing patterns of susceptibility to ESBL-E gut-
65 colonization among this population.

66

67 **Methods**

68 Stool collection

69 Stool samples were collected during a prospective cohort study of childhood enteric infections in
70 Villa El Salvador, Lima, Peru (NIH R01A1108695-01A1).(4) Pregnant women were identified
71 through an initial population screening in the community and through community health centers.
72 Full-term babies were enrolled approximately within one month after birth from February 2016-
73 May 2017, and daily surveillance data were collected through at least two years of age. Starting
74 at recruitment and weekly thereafter, caregivers also kept a freshly soiled diaper for field worker
75 collection. Upon arrival at a local lab facility, stool was aliquoted and stored at -20°C for up to
76 48h, then transported to UPCH for long-term storage at -80°C.

77

78 Sample selection

79 As previously described,(4) Enterobacterales were cultured from weekly stool samples of all
80 newborns enrolled after December 2016 (112/345). For each sample, Enterobacterales colonies
81 were pooled then archived. Colonies collected at approximately 1, 3, 4, 5, 6, 7, 9, 12, and
82 16□months of age were later screened for ESBL production. We observed two distinct patterns:
83 children who were never or rarely colonized with ESBL-E (n=12), hereafter “*rarely colonized*,”
84 and children who were often or always colonized (n=100), hereafter “*frequently colonized*”
85 (**Figure S1**). Rarely colonized children first tested positive for ESBL-E at approximately 6
86 months or later, and were colonized at fewer than half of tested timepoints. Frequently colonized
87 children were colonized at more than half of timepoints.

88

89 Exposure assessment

90 As previously described,(4,10) socioeconomic variables were collected at recruitment and then
91 once yearly. Additionally, field staff visited infants every day except Sunday to collect survey
92 data from the previous 24 hours. If field workers were unable to visit an infants’ home (e.g., day

93 was Sunday), field workers collected survey data for the missing day at the next visit. At each
94 daily visit, caretakers were asked about children's milk consumption, antibiotic use, and diarrhea
95 in the past 24 hours. Caretakers were asked to provide medication packaging so trained field
96 workers could confirm it was an antibiotic. Because survey data were collected daily, this
97 packaging was typically available. A new antibiotic course began when caretakers reported
98 antibiotic consumption following two days of no exposure; a course ended when the child did
99 not consume antibiotics for two consecutive days. Children were considered to have had
100 diarrhea if they experienced ≥ 3 liquid or semiliquid stools in the past 24 hours.

101

102 Metagenomic sequencing

103 We performed short-read metagenomic sequencing on available frozen stool specimens
104 collected at approximately 3, 6, 7, 9, 12, and 16 months of age from all rarely colonized
105 children (n=12 children, n=71 stool samples), and a randomly selected subset of 42 frequently
106 colonized children (n=238 stool samples). Total DNA was extracted from approximately 0.25 g
107 of frozen feces at UPCH using the QIAamp PowerFecal Pro DNA Kit (Qiagen) according to
108 manufacturer's instructions and quantified using a Qubit 4 fluorometer (Invitrogen). Extraction
109 blanks were included with each batch of extractions; DNA concentrations were below the Qubit
110 level of detection (<0.1 ng/ μ L) for all. DNA extracts were sent to the Broad Institute for short-
111 read, paired-end 150 bp sequencing on an Illumina Novaseq 6000 System using SP4 flow cells
112 to achieve 6 Gb per sample.

113 Processing metagenomic data and profiling taxonomy

114 Fastq sequences were trimmed and filtered to remove sequencing adaptors and low quality
115 reads using bbmap.(11) We tabulated the number of raw reads for each sample and excluded
116 from further analysis any sample with more or less reads than two standard deviations from the
117 mean (n=11). Human contaminant sequences were removed from each sample by discarding
118 reads that mapped to a non-redundant version of the Genome Reference Consortium Human

119 Build 38 (GRCh38; www.ncbi.nlm.nih.gov) using Bowtie2 v2.5.1. Human sequence-filtered raw
120 reads were deposited in the Sequence Read Archive (SRA; <https://www.ncbi.nlm.nih.gov/sra>)
121 under the project number PRJNA1138246 (**Table S1**).

122 Taxonomic assignment of short reads was performed using MetaPhlan3 (with db
123 v31).(12) Alpha diversity metrics were derived from the MetaPhlan3 relative abundance output
124 using the vegan package in R [<https://cran.r-project.org/web/packages/vegan/>].

125

126 Data analysis

127 We examined differences in sociodemographic characteristics, diarrhea rates, and antibiotic use
128 between rarely and frequently gut-colonized children using t-tests and chi square tests for all
129 112 children for whom we characterized ESBL-E gut colonization, and separately for the 54
130 children whose stool samples were sequenced.

131 For the subset of 54 children with fecal metagenomes, we examined whether any taxa
132 were differentially abundant between rarely and frequently gut-colonized children across time
133 using mixed effects linear regression models. An arcsine square root transformation was
134 applied to relative abundance values. We also used mixed effects linear regression models to
135 examine differences in log₁₀ transformed species richness values and Shannon diversity over
136 time between groups. Transformed relative abundance values and alpha diversity metrics were
137 modeled as a function of subject (random effects term), colonization pattern, i.e., rarely versus
138 frequently colonized (binary fixed effects term), child age (continuous fixed effects term), and
139 hours before diaper retrieval (continuous fixed effects term). We also ran additionally
140 parametrized models adjusted for factors known to affect the gut microbiome, including child's
141 delivery mode (binary fixed effects term) and number of diarrhea episodes and antibiotic
142 courses in the past 30 days (continuous fixed effects terms), as well as all socioeconomic
143 characteristics that differed groups with $p < 0.20$. We did not adjust for breastfeeding as there
144 was minimal variation in this exposure over time (**Table S2**). All taxa with more than 10% non-

145 zero values were fitted using the nlme package in R. Significance of associations was
146 determined using Wald's test, and p-values were adjusted for multiple hypothesis testing using
147 the Benjamini-Hochberg correction at each taxonomic level.

148 For the subset of metagenomes for which we had matched ESBL-E gut colonization
149 status, we examined whether any taxa were differentially abundant between timepoints where
150 children were ESBL-E positive versus negative using mixed effects logistic regression models.
151 For any given taxa detected in at least 10% of samples, we modeled ESBL-E positivity (yes/no)
152 as a function of subject, child age, hours before diaper retrieval, number of recent diarrhea
153 episodes and antibiotic courses, and the transformed relative abundance value of that taxa. We
154 used the lme4 package in R and adjusted p-values using the Benjamini-Hochberg correction at
155 each taxonomic level.

156 Finally, we used the StrainGE tool(13) to characterize *E. coli* strains in children's fecal
157 metagenomes. A reference database was built using complete *E. coli* genomes from NCBI's
158 RefSeq database (~3,100 genomes downloaded March 5, 2024), and each genome was
159 assigned a Clermont phylotype using EzClermont v.0.70.(14)

160

161 Results

162 Children who were rarely gut-colonized with ESBL-E belonged to households of lower 163 socioeconomic status

164 There were no significant differences in exposures that are well-known to impact the gut
165 microbiome, including rates of vaginal birth, antibiotic consumption rates, children's age at time
166 of first antibiotic consumption, or numbers of diarrhea episodes per child month between rarely
167 versus frequently colonized children (**Table 1**). Rarely colonized children did consume slightly
168 fewer antibiotic courses overall (3.4 vs 4.5 courses/child-year, $p=0.2$), and fewer azithromycin
169 courses in particular (0.1 vs 0.5 courses/child-year, $p=0.14$), but differences were not

170 statistically significant. Only 2/112 children did not consume any antibiotics, yet both were
171 frequently colonized with ESBL-E.

172 Children who were rarely gut-colonized with ESBL-E did belong to households of lower
173 socioeconomic status (SES). Specifically, they were somewhat more likely to live in households
174 that relied on a public tap for drinking water ($p=0.06$) versus having an indoor connection, and
175 more likely to rely on pit latrines versus indoor flush toilets ($p=0.05$). Most households that relied
176 on water taps also used pit latrines, and vice versa. Rarely colonized children were also
177 somewhat less likely to be born to a mother who had completed high school ($p=0.11$).
178 Children's antibiotic use (total courses/child-year) was weakly correlated with household
179 reliance on public water taps ($r_{pb}=0.24$, $p<0.05$ by point-biserial correlation) and use of a pit
180 latrine ($r_{pb}=0.25$, $p<0.05$), but was not correlated with maternal education attainment ($r_{pb}=0.04$,
181 $p=0.67$).

182 Breastfeeding was exceptionally common. On average, children received breastmilk
183 nearly every day in the 30 days prior to a stool sample through 16 months of age (**Table S2**).
184 Notably, nearly 1/3 children who were frequently colonized with ESBL-E regularly consumed
185 formula early in life, compared to just 1 of 12 rarely colonized children (**Table 1**). However, this
186 difference was not significant ($p=0.29$).

187 The distributions of these exposures were similar among the subset of 54/112 children
188 whose fecal metagenomes were sequenced for this study (**Table S3**).

189
190 Peruvian children's gut microbiomes are dominated by *Bifidobacterium* spp. regardless of
191 ESBL-E colonization status

192 Samples included in our final analysis included 66 fecal metagenomes from 12 rarely colonized
193 children and 232 fecal metagenomes from 42 frequently colonized children ($n=298$ total, 4-6
194 samples per child).

195 As expected given the high proportion of breastfeeding in this setting, children's gut
196 microbiomes were largely dominated by *Bifidobacterium* spp., comprising 58% of bacterial
197 reads in average at 3 months, and 34% by 16 months (**Figure 1**). *B. longum* and *B. breve*
198 predominated earlier in life, while *B. bifidum* consistently comprised between 10-15% of
199 bacterial reads (**Figure 2**). The number of unique *Bifidobacterium* spp. in children's guts ranged
200 from 0-9 (median=4) and increased as children aged ($\beta=0.14$, $SE=0.019$, $p\text{-value}<0.0001$).
201 Pooling all timepoints together, there was no difference in the relative abundance of any
202 *Bifidobacterium* spp. between children who were rarely versus frequently gut-colonized with
203 ESBL-E. Likewise, the number of unique *Bifidobacterium* spp. did not differ between groups.

204
205 No differences in diversity or relative abundance of gut bacteria between rarely and frequently
206 colonized children

207 Besides *Bifidobacterium*, other dominant bacterial genera included *Blautia* (average across all
208 samples 3.1%), *Bacteroides* (2.1%), *Escherichia* (1.9%), *Lactobacillus* (1.1%), *Faecalibacterium*
209 (0.9%), and *Klebsiella* (0.8%) (**Figure 1**). Potential enteropathogens like *Klebsiella pneumoniae*
210 and *Enterococcus faecium* were detected among nearly all children (**Table S4**). After
211 adjustment for multiple comparisons, no taxa were differentially abundant between children who
212 were rarely versus frequently gut-colonized with ESBL-E over time in fully parameterized
213 models (corrected p-values all > 0.89). Results were similar for simpler models that only
214 adjusted for children's colonization pattern, child age, and hours before diaper retrieval (fdr-
215 corrected p-values all >0.66).

216 Like others,(15) we found the number of unique bacterial species in children's fecal
217 metagenomes increased as they aged, from 31 to 75, on average. The diversity of children's
218 fecal metagenomes also increased (0.99 Shannon diversity index on average at 3 months, to
219 2.31 at 16 months) (**Figure S4**). Pooling all timepoints together, we observed no differences in
220 diversity or richness between groups ($p\text{-values}> 0.62$). There were also no differences in

221 median richness or Shannon diversity at any specific sampling point ($p>0.52$ for all pairwise
222 differences by Wilcoxon rank sum tests).

223

224 No differences in the number and diversity of *E. coli* strains between rarely vs frequently
225 colonized children

226 Rarely colonized children appeared to harbor slightly fewer *E. coli* strains on average at each
227 timepoint (**Figure 3A**), but these differences were not statistically significant ($p>0.15$ for all by
228 Welch two-sample t-test). In general, children's stool samples were largely dominated by *E. coli*
229 phylotypes A and B1 (**Figure 3B**). There was no significant difference in the proportion of
230 strains belonging to any given *E. coli* phylotype between rarely and frequently colonized children
231 at any time point ($p>0.26$ for all pairwise comparisons by Fisher test).

232 We also examined differences in *E. coli* strains between sampling points where children
233 were gut-colonized with ESBL-E versus not (ESBL-E gut colonization status available for
234 263/298 fecal metagenomes). Overall, children harbored more unique *E. coli* strains at sampling
235 points where they were gut-colonized with ESBL-E (mean=2.3 across all age groups) versus
236 sampling points where they were not (mean=2.1, $p=0.05$ by Welch two-sample t-test), though
237 there were no differences for specific sampling points ($p>0.15$ for all by Welch two-sample t-
238 test). Overall, there was no difference in *E. coli* phylotype distribution at sampling points where
239 children were gut-colonized with ESBL-E versus not ($p=0.75$ by Fisher test).

240

241 Some differences in relative abundance of specific taxa at ESBL-E positive versus negative
242 timepoints

243 Regardless of whether children were rarely or frequently gut-colonized with ESBL-E over time,
244 we did note that the relative abundances of the phyla *Actinobacteria* (average across all
245 samples 45.9%; $\text{fdr-corrected } p\text{-value}=0.20$) and the genus *Slackia* (0.006%, $\text{fdr-corrected } p\text{-}$
246 $\text{val}=0.03$) in children's stool were higher at timepoints where children were not gut-colonized

247 with ESBL-E by culture, when controlling for child age, hours before diaper retrieval, number of
248 recent antibiotic courses and diarrhea episodes, and repeated measurements (**Figure S3**).
249 Relative abundances of the phyla *Proteobacteria* (3.4%, fdr-corrected p-value=0.20), the phyla
250 *Bacteroidetes* (2.9%, fdr-corrected p-value=0.20), and the class *Erysipelotrichia* (0.47%; fdr-
251 corrected p-val=0.15) were higher at timepoints where children colonized with ESBL-E by
252 culture.

253

254 Discussion

255 This is the first longitudinal study of the gut microbiome among South American children. As
256 previously reported, gut colonization with ESBL-E was prevalent among children in this peri-
257 urban setting and steadily increased as children aged.(4) Similar to recent studies among
258 healthy adults,(16–18) our findings suggest that differences in exposures, rather than
259 differences in gut microbiome development, might underlie patterns of ESBL gut colonization
260 over time among children in peri-urban Lima. To our knowledge, this is the first study to
261 examine this research question among young children who are living in a low-resource setting
262 and whose gut microbiomes are rapidly developing.

263 Three recent Western studies and two LMIC studies have investigated associations
264 between gut microbial taxa and ESBL-E gut colonization at single time points.(16–20) Two
265 case-control studies among Dutch(17) and French adults(18) found no differences in alpha
266 diversity or taxa abundance between healthy adults who were gut-colonized with ESBL-E
267 versus not, after controlling for age, sex, ethnicity, and recent travel. Another study among US
268 travelers(16) found no difference in the pre-travel gut microbiomes of individuals who acquired
269 ESBL-E while abroad versus those who did not, suggesting that differences in exposures while
270 traveling, rather than inherent differences in intestinal colonization resistance, are more
271 important determinants of ESBL-E acquisition. One study of healthy Thai adults(20) reported
272 differences in gut taxa between ESBL-E carriers and non-carriers, but because potential

273 confounders were not controlled for, it is challenging to attribute differences to ESBL-E
274 colonization status. A small study among indigenous persons in French Guyana identified some
275 differences in gut microbial taxa between 8 cases and 24 controls, but whether these
276 differences preceded or were a result of ESBL-E gut colonization could not be determined.(19)
277 To our knowledge, no previous study has examined whether differences in the gut microbiome
278 over time, rather than just at a single timepoint, may underlie ESBL-E gut colonization patterns.
279 Our study adds to the existing literature by indicating that patterns of rare versus frequent
280 ESBL-E gut colonization among children who are frequently exposed to enteropathogens and
281 antibiotics may not primarily be a function of intestinal colonization resistance.

282 Taxa that previous studies have identified to be associated with ESBL carriage or non-
283 carriage differed from what we observed. *Actinobacteria*, which typically dominate the guts of
284 breastfeeding children,(21) were somewhat more abundant at ESBL-negative timepoints (fdr-
285 corrected p-val=0.2), while *Proteobacteria*, *Bacteroidetes*, and *Erysipelotrichia*, all of which are
286 commonly isolated in children's stool,(21) were somewhat more abundant at ESBL-positive
287 timepoints (p-val between 0.15 and 0.2 for all). *Slackia*, which is not well-described among
288 children but has previously been isolated from both gut and oral infections in adults,(22) was
289 more abundant at ESBL-negative timepoints though overall abundance was low (0.006%).
290 Differences between our study and previous work(19,20) could be a result of our focus on the
291 pediatric microbiome, differences in our sequencing approach, differences in environmental
292 exposures, or other factors. We note that whether these differentially abundant taxa preceded or
293 followed ESBL-E gut colonization cannot be distinguished through our analytical approach.

294 Overall, the fact that we did not identify gut taxa that were differentially abundant
295 between rarely and frequently colonized children could be because such differences do not
296 exist; for example, perhaps the differential exposures we noted only affected the gut
297 metabolome. There are other gut microbiome differences that we were unable to explore, e.g.,
298 higher bacteriocin production among commensal strains harbored by rarely colonized children,

299 which could favor the exclusion of ESBL-E.(23) Or, we may have been underpowered to detect
300 differences. Given the high and increasing rates of ESBL-E gut colonization in community
301 settings in LMICs, we note that enrolling sufficient children to adequately investigate this
302 research question may be challenging in future studies.

303 While we did not identify differentially abundant gut taxa or differences in *E. coli* strains
304 between rarely and frequently colonized children, we surprisingly found that rarely colonized
305 children were more likely to belong to households of lower SES. Although these differences
306 were not statistically significant at the $p < 0.05$ level, the households of rarely colonized children
307 were somewhat more likely to lack access to indoor piped water and flush toilets, and their
308 mothers were less likely to have completed high school. A previous study of drug-resistant
309 urinary tract infections in women in Hyderabad, India, found that poorer women were most likely
310 to have an ESBL-producing infection, despite consuming fewer antibiotics than wealthier
311 women, suggesting environmental exposures unique to low SES populations might be
312 underlying elevated risks of exposure to AMR bacteria in this setting.(24) We find the opposite
313 trend here, with lower SES children having the lowest risk of being persistently colonized with
314 ESBL-E early in life. The role of the environment as a source of exposure to ESBL-E likely
315 differs among and perhaps within LMICs. We note that while multiple global analyses indicate
316 that antimicrobial resistance is more common in LMICs compared to high-income
317 settings,(25,26) many LMICs have high rates of income inequality and within-country
318 heterogeneity in acquisition risks have rarely been examined.(27)

319 This study had strengths and limitations. Our study represents the first longitudinal
320 analysis of the gut microbiome of South American children, whose fecal metagenomes are
321 under-represented in global databases.(28) Because daily surveys were collected, we were able
322 to control for time-varying exposures that are known to affect the gut microbiome, including
323 antibiotic use and recent diarrhea, with a low risk of recall bias. However, our study was
324 relatively small, and we may have been underpowered to detect differences in gut microbial

325 taxa between groups. Further, as previously noted,(4) we used culture-based methods with a
326 relatively high lower limit of detection (~2 log₁₀ CFU/g wet feces) to identify ESBL-E gut
327 colonization for this study. Rarely colonized children may have been gut-colonized with ESBL-E
328 at timepoints where they tested negative, just at concentrations below the limit of detection.
329 However, since higher concentrations of multidrug-resistant Enterobacterales in the gut are
330 associated with elevated risks of infection,(29,30) identifying whether specific gut taxa may
331 suppress the growth of ESBL-E remains a clinically relevant question.

332

333 Conclusion

334 This study examined whether differences in Peruvian children's gut microbiome development
335 could underpin differences in early-life susceptibility to ESBL-E gut colonization. We did not
336 identify any significant differences in species richness, Shannon diversity, *E. coli* strain diversity,
337 or the relative abundance of specific taxa between rarely versus frequently colonized children.
338 Nevertheless, rarely colonized children who were somewhat more likely to live in households of
339 lower SES. Regardless of gut colonization patterns over time, children harbored higher relative
340 abundances of *Proteobacteria*, *Bacteroidetes*, and *Erysipelotrichia* at timepoints where they
341 were ESBL-E positive and higher abundances of *Actinobacteria* and *Slackia* at timepoints where
342 they were negative. Our findings are in line with recent studies among healthy adults(16–18)
343 suggesting that differences in exposures, rather than differences in the gut microbiome, might
344 underlie differing patterns of susceptibility to ESBL-E gut-colonization. Our findings suggest that
345 preventing ESBL-E colonization among children in LMICs may require reducing their exposures
346 to environmental reservoirs of ESBL-E.

347

348 Funding

349 This work was supported by NIH R01AI108695-01A, NIH award KL2TR002545, and NIH award
350 U19AI110818. MLN was supported by Emory University and the MP3 Initiative. NS was

351 supported by NIH award 5T32ES012870. The content is solely the responsibility of the authors
352 and does not necessarily represent the official views of Emory University or the MP3 Initiative.

353

354 Acknowledgements

355 We thank all field workers and the community of Villa El Salvador for their collaboration in the
356 parent study.

357

358 Conflicts of interest

359 None to declare.

360

361 Data Availability

362 Children's fecal metagenomes are available in NCBI's Sequence Read Archive

363 (<https://www.ncbi.nlm.nih.gov/sra>) under BioProject number PRJNA1138246.

364

365 Ethics approval and consent to participate

366 The parent cohort study was approved by the Institutional Review Boards of Asociación

367 Benefica PRISMA, Universidad Peruana Cayetano Heredia, and Johns Hopkins University.

368 Written informed consent was obtained from caretakers of the infants for both study participation

369 and the use of collected specimens for subsequent research. The analysis of stool specimens

370 for this substudy was approved by the Institutional Review Boards of Asociación Benefica

371 PRISMA and Universidad Peruana Cayetano Heredia (no. 201592). Metagenomic sequencing

372 of stool specimens was determined to be exempt by the Tufts University Health Sciences IRB

373 (STUDY00001328).

374

375

376 References

- 377 1. Cheikh A, Belefquih B, Chajai Y, Cheikhaoui Y, El Hassani A, Benouda A.
378 Enterobacteriaceae producing extended-spectrum β -lactamases (ESBLs) colonization as a
379 risk factor for developing ESBL infections in pediatric cardiac surgery patients:
380 “retrospective cohort study.” *BMC Infect Dis.* 2017 29;17(1):237.
- 381 2. Ben-Ami R, Schwaber MJ, Navon-Venezia S, Schwartz D, Giladi M, Chmelnitsky I, et al.
382 Influx of extended-spectrum beta-lactamase-producing enterobacteriaceae into the hospital.
383 *Clin Infect Dis.* 2006 Apr 1;42(7):925–34.
- 384 3. Denis B, Lafaurie M, Donay JL, Fontaine JP, Oksenhendler E, Raffoux E, et al. Prevalence,
385 risk factors, and impact on clinical outcome of extended-spectrum beta-lactamase-
386 producing *Escherichia coli* bacteraemia: a five-year study. *Int J Infect Dis.* 2015 Oct;39:1–6.
- 387 4. Nadimpalli ML, Rojas Salvatierra L, Chakraborty S, Swarouth JM, Cabrera LZ, Pickering AJ,
388 et al. Effects of breastfeeding on children’s gut colonization with multidrug-resistant
389 Enterobacteriales in peri-urban Lima, Peru. *Gut Microbes.*
- 390 5. Nadimpalli ML, Marks SJ, Montealegre MC, Gilman RH, Pajuelo MJ, Saito M, et al. Urban
391 informal settlements as hotspots of antimicrobial resistance and the need to curb
392 environmental transmission. *Nature Microbiology.* 2020 Jun 1;5(6):787–95.
- 393 6. Kothari C, Gaiind R, Singh LC, Sinha A, Kumari V, Arya S, et al. Community acquisition of β -
394 lactamase producing Enterobacteriaceae in neonatal gut. *BMC Microbiol.* 2013 Jun
395 17;13:136.
- 396 7. Islam MA, Amin MB, Roy S, Asaduzzaman M, Islam MR, Navab-Daneshmand T, et al. Fecal
397 colonization with multidrug-resistant *E. coli* among healthy infants in rural Bangladesh.
398 *Frontiers in Microbiology.* 2019;10:640.
- 399 8. Tellevik MG, Blomberg B, Kommedal Ø, Maselle SY, Langeland N, Moyo SJ. High
400 Prevalence of Faecal Carriage of ESBL-Producing Enterobacteriaceae among Children in
401 Dar es Salaam, Tanzania. *PLOS ONE.* 2016 Dec;11(12):e0168024.
- 402 9. Lawley TD, Walker AW. Intestinal colonization resistance. *Immunology.* 2013 Jan;138(1):1–
403 11.
- 404 10. Pajuelo MJ, Noazin S, Cabrera L, Toledo A, Velagic M, Arias L, et al. Epidemiology of
405 enterotoxigenic *Escherichia coli* and impact on the growth of children in the first two years of
406 life in Lima, Peru. *Front Public Health.* 2024;12:1332319.
- 407 11. Brian Bushnell. BBMap [Internet]. Available from: <https://sourceforge.net/projects/bbmap/>
- 408 12. Truong DT, Tett A, Pasolli E, Huttenhower C, Segata N. Microbial strain-level population
409 structure and genetic diversity from metagenomes. *Genome Res.* 2017;27(4):626–38.
- 410 13. Van Dijk LR, Walker BJ, Straub TJ, Worby CJ, Grote A, Schreiber HL, et al. StrainGE: a
411 toolkit to track and characterize low-abundance strains in complex microbial communities.
412 *Genome Biol.* 2022 Dec;23(1):74.

- 413 14. Waters NR, Abram F, Brennan F, Holmes A, Pritchard L. Easy phylotyping of *Escherichia*
414 *coli* via the EzClermont web app and command-line tool. *Access Microbiol.*
415 2020;2(9):acmi000143.
- 416 15. Stewart CJ, Ajami NJ, O'Brien JL, Hutchinson DS, Smith DP, Wong MC, et al. Temporal
417 development of the gut microbiome in early childhood from the TEDDY study. *Nature.*
418 2018;562(7728):583–8.
- 419 16. Worby CJ, Sridhar S, Turbett SE, Becker MV, Kogut L, Sanchez V, et al. Gut microbiome
420 perturbation, antibiotic resistance, and *Escherichia coli* strain dynamics associated with
421 international travel: a metagenomic analysis. *The Lancet Microbe* [Internet]. 2023 Sep 13
422 [cited 2023 Sep 29];0(0). Available from:
423 [https://www.thelancet.com/journals/lanmic/article/PIIS2666-5247\(23\)00147-7/fulltext](https://www.thelancet.com/journals/lanmic/article/PIIS2666-5247(23)00147-7/fulltext)
- 424 17. Ducarmon QR, Zwitter RD, Willems RPJ, Verhoeven A, Nooij S, van der Klis FRM, et al.
425 Gut colonisation by extended-spectrum β -lactamase-producing *Escherichia coli* and its
426 association with the gut microbiome and metabolome in Dutch adults: a matched case-
427 control study. *Lancet Microbe.* 2022 Jun;3(6):e443–51.
- 428 18. Boyd A, El Dani M, Ajrouche R, Demontant V, Cheval J, Lacombe K, et al. Gut microbiome
429 diversity and composition in individuals with and without extended-spectrum β -lactamase-
430 producing *Enterobacteriales* carriage: a matched case–control study in infectious diseases
431 department. *Clinical Microbiology and Infection* [Internet]. 2024 Mar 23 [cited 2024 Jul 17];
432 Available from: <https://www.sciencedirect.com/science/article/pii/S1198743X24001460>
- 433 19. Gosalbes MJ, Vázquez-Castellanos JF, Angebault C, Woerther PL, Ruppé E, Ferrús ML, et
434 al. Carriage of *Enterobacteria* Producing Extended-Spectrum β -Lactamases and
435 Composition of the Gut Microbiota in an Amerindian Community. *Antimicrob Agents*
436 *Chemother.* 2016 Jan;60(1):507–14.
- 437 20. Piewngam P, Quiñones M, Thirakittiwatthana W, Yungyuen T, Otto M, Kiratisin P.
438 Composition of the intestinal microbiota in extended-spectrum β -lactamase-producing
439 *Enterobacteriaceae* carriers and non-carriers in Thailand. *Int J Antimicrob Agents.* 2019
440 Apr;53(4):435–41.
- 441 21. Milani C, Duranti S, Bottacini F, Casey E, Turrone F, Mahony J, et al. The First Microbial
442 Colonizers of the Human Gut: Composition, Activities, and Health Implications of the Infant
443 Gut Microbiota. *Microbiology and Molecular Biology Reviews.* 2017 Nov
444 8;81(4):10.1128/mnbr.00036-17.
- 445 22. Kim KS, Rowlinson MC, Bennion R, Liu C, Talan D, Summanen P, et al. Characterization of
446 *Slackia exigua* isolated from human wound infections, including abscesses of intestinal
447 origin. *J Clin Microbiol.* 2010 Apr;48(4):1070–5.
- 448 23. Majeed H, Gillor O, Kerr B, Riley MA. Competitive interactions in *Escherichia coli*
449 populations: the role of bacteriocins. *ISME J.* 2011 Jan;5(1):71–81.
- 450 24. Alsan M, Kammili N, Lakshmi J, Xing A, Khan A, Rani M, et al. Poverty and Community-
451 Acquired Antimicrobial Resistance with Extended-Spectrum β -Lactamase-Producing
452 Organisms, Hyderabad, India. *Emerging Infect Dis.* 2018;24(8):1490–6.

- 453 25. Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance
454 in 2019: a systematic analysis. *Lancet*. 2022 Feb 12;399(10325):629–55.
- 455 26. Hendriksen RS, Munk P, Njage P, van Bunnik B, McNally L, Lukjancenko O, et al. Global
456 monitoring of antimicrobial resistance based on metagenomics analyses of urban sewage.
457 *Nature Communications*. 2019 Mar 8;10(1):1124.
- 458 27. Blackmon S, Avendano E, Nirmala N, Chan CW, Morin RA, Balaji S, et al. Socioeconomic
459 status and the risk for colonization or infection with priority bacterial pathogens: a global
460 evidence map [Internet]. 2024 [cited 2024 Jun 5]. Available from:
461 <http://medrxiv.org/lookup/doi/10.1101/2024.04.24.24306293>
- 462 28. Fuhrmeister ER, Harvey AP, Nadimpalli ML, Gallandat K, Ambelu A, Arnold BF, et al.
463 Evaluating the relationship between community water and sanitation access and the global
464 burden of antibiotic resistance: an ecological study. *The Lancet Microbe*. 2023 Aug
465 1;4(8):e591–600.
- 466 29. Ruppé E, Lixandru B, Cojocaru R, Büke C, Paramythiotou E, Angebault C, et al. Relative
467 fecal abundance of extended-spectrum- β -lactamase-producing *Escherichia coli* strains and
468 their occurrence in urinary tract infections in women. *Antimicrob Agents Chemother*. 2013
469 Sep;57(9):4512–7.
- 470 30. Shimasaki T, Seekatz A, Bassis C, Rhee Y, Yelin RD, Fogg L, et al. Increased Relative
471 Abundance of *Klebsiella pneumoniae* Carbapenemase-producing *Klebsiella pneumoniae*
472 Within the Gut Microbiota Is Associated With Risk of Bloodstream Infection in Long-term
473 Acute Care Hospital Patients. *Clin Infect Dis*. 2019 May 30;68(12):2053–9.
- 474

475 Tables

476 **Table 1.** Sociodemographic characteristics and early-life exposures among 112 Peruvian
477 children who were rarely or frequently colonized with ESBL-E during the first 16 months of life.

	Frequently colonized (n=100)	Rarely colonized (n=12)	p-val
Male (n, %)	48 (48.0)	7 (58.3)	0.711
Birth mode (n, %)			0.234
Vaginal	68 (68.0)	10 (83.3)	
Cesarean section	31 (31.0)	1 (8.3)	
Missing	1 (1.0)	1 (8.3)	
Mother completed high school (n, %)	69 (69.0)	5 (41.7)	0.117
Water Source (n,%)			0.063
Other	5 (5.0)	0 (0.0)	
Public Tap	15 (15.0)	5 (41.7)	
Indoor water connection (24 hours)	80 (80.0)	7 (58.3)	
Toilet (n,%)			0.051
Other	2 (2.0)	0 (0.0)	
Pit latrine	14 (14.0)	5 (41.7)	
Indoor flush toilet	84 (84.0)	7 (58.3)	
Home has improved flooring ^a	97 (97.0)	11 (91.7)	0.906
Lives in household with more than six household members (n,%)	40 (40.0)	2 (16.7)	0.207
Received breastmilk 90% of days in first 3 months (n,%)	63 (63.0)	8 (66.7)	1
Received formula 50% of days in first 3 months (n,%)	27 (27.0)	1 (8.3)	0.29
Household owned chickens (n,%)	17 (17.0)	1 (8.3)	0.721
Total number of antibiotic courses per child-year (mean (SD))	4.45 (2.86)	3.36 (2.29)	0.207
Number of azithromycin courses per child-year (mean (SD))	0.47 (0.95)	0.06 (0.22)	0.144
Number of sulfa-trimethoprim courses per child-year (mean (SD))	1.03 (1.15)	0.98 (1.09)	0.888
Number of amoxicillin courses per child-year (mean (SD))	1.59 (1.41)	1.93 (2.36)	0.471
Number of cefalexin courses per child-year (mean (SD))	0.53 (0.90)	0.32 (0.61)	0.425
Age in months at time of first antibiotic use (mean (SD))	5.67 (4.49)	7.23 (3.75)	0.251
Number of diarrhea episodes/100-child-month (mean (SD))	6.88 (5.09)	5.28 (2.59)	0.288

478 ^aImproved flooring included cement, wood, tiles, and “falso piso.” Unimproved flooring included
479 dirt and sand.

480

481 Figures

482

483 **Figure 1.** Proportion of bacterial genera with at least 1% average relative abundance in stool

484 samples collected from Peruvian children aged 3-16 months.

485

486 **Figure 2.** Relative abundance of dominant *Bifidobacterium spp.* in stool samples collected from

487 Peruvian children aged 3-16 months.

488 Note: Fifteen *Bifidobacterium spp.* were detected among all children; species with at least 1%

489 average relative abundance are depicted here.

490

491 **Figure 3.** Distribution of *E. coli* strains in stool samples from Peruvian children aged 3-16


492 months who were frequently vs rarely gut-colonized with ESBL-E, including A) mean number of

493 unique *E. coli* strains, and B) Clermont phylotype.


494

495


496



Proportional Abundance (%)



A)



B)

