

1 **Full Title:** “Around the clock”. Exploring health care professionals’ experience of discharge of
2 older people during out of hours from the emergency department: a qualitative study.

3 **Short Title:** Older people and out of hours emergency department discharge.

4

5 **Mary Dunnion^{1*}¶, Debbie Goode²¶, Assumpta Ryan¹¶, Sonja McIlfatrick²¶,**

6 ¹School of Nursing and Paramedic Science, Faculty of Life and Health Sciences, Ulster
7 University, Northland Road, Derry~Londonderry, United Kingdom.

8 ²School of Nursing and Paramedic Sciences, Faculty of Life and Health Science, Ulster
9 University, 2-24 York Street, Belfast, United Kingdom.

10 ***Corresponding Author:** Mary Dunnion. Email: Dunnion-m1@ulster.ac.uk

11 **ORCID:** Mary Dunnion <https://orcid.org/0000-0003-2648-5125>

12

13 ¶¶These authors contributed equally to this work.

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29 **Abstract**

30 **Background**

31 Older adults constitute a considerable number of attendances at emergency departments
32 (EDs). Whilst many require hospital admission, a greater focus now is on admission avoidance
33 with older adults being subsequently discharged from EDs. Little is known, however, about
34 the experiences out of hours (OOH) when specialist older person support services are
35 unavailable.

36 **Aim**

37 To explore senior health care professionals' experiences of discharging older people during
38 out of hours from EDs.

39 **Method**

40 A qualitative study involving individual semi-structured interviews was used to explore the
41 experiences of healthcare professionals. Fourteen participants in total responded from a wide
42 geographical spread. Data analysis was undertaken using Braun and Clarke's (2022) six-step
43 framework for Thematic Analysis.

44 **Results**

45 One overarching theme was identified focusing on risks and safety for the older person whilst
46 being discharged OOH from the ED. Three sub-themes included "Should they stay, or should
47 they go?", "Bright lights and noise", and "New ways of working", which included risks in
48 discharging an older person OOH from the ED, risks in delaying discharge, and
49 recommendations for future practice. Significant differences were reported between office
50 hours and OOH discharge of older adults from the ED. Diversity in practice assessments, and
51 resources available was also evident across EDs. Significant adverse consequences were
52 identified for older adults having to stay for prolonged periods in the ED.

53 **Conclusions**

54 There have been many welcome developments in healthcare services for older people who
55 attend and are subsequently discharged from the ED. However, further innovative practice
56 and collaboration with senior decision makers is needed to meet the healthcare needs of a
57 rapidly ageing population. Safe, equitable and effective discharge practice 24/7 should be a
58 norm for all older people in every ED regardless of location. Gaps in care identified must be
59 addressed taking cognisance of the relevant recommendations for future practice.

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75 Introduction

76 Countries worldwide are witnessing an unprecedented population increase in both the number
77 and the proportion of older people (1). The United Nations General Assembly (2) projected
78 that between 2019 and 2030, the number of persons aged 60 years or over will grow by 38
79 per cent. According to Eliopoulos (3), older adults are generally defined as 65 years of age
80 and older; however, several categories are also recognised, including, young old (65 to 74
81 years); old (75 to 84 years) and oldest-old (85 years and over) taking cognisance of the
82 diversity between these age groups. Alongside the demographic growth, a current focus on
83 healthy ageing is advocated (4). However, by nature many older adults become unwell which
84 often necessitates a visit to an emergency department (ED). This can often be a difficult
85 experience for some older people who attend the ED with an atypical presentation and in an
86 acutely unwell state. According to Cassarino et al (5) and Corcoran et al (6) attendance at the
87 ED can result in high levels of adverse outcomes for older people. Issues with communication,
88 privacy, and personal care for older people in the ED have been highlighted (7), whilst the ED
89 has been reported as a complex, stressful and unpredictable area (8; 9). Although some older
90 adults may require subsequent admission to hospital from the ED (10), many are discharged
91 home from the department. Whilst the latter option may be the most preferable for older
92 people, Cetin-Sahin (11) and Gettel et al (12) reported on several transitional care needs for
93 patients following discharge.

94 It is imperative that the discharge process from the ED is safe, comprehensive, effective, and
95 timely. However, outside of office hours (after 5pm until 8am and at all weekends and public
96 holidays) many older person specialist teams are inaccessible in the ED or in the community
97 which poses the question as to the experiences and risks for this cohort being discharged late
98 at night or at weekends (13). Whilst some attention has been given to the discharge of older
99 people from hospital at night (14), a review of the literature has identified a dearth of research
100 regarding out of hours (OOH) discharge of older people from the ED. The aim of this paper

101 was to explore the experiences of Health Care Professionals (HCP) regarding the discharge
102 of older people OOH versus during office hours from the ED.

103

104 **Methodology**

105 **Aim and Objectives**

106 This paper was part of an overall study exploring the discharge of older adults out of hours
107 from the ED. This paper focused on the experiences of senior Health Care Professionals'
108 when discharging older people during OOH versus office hours from the ED.

109 Objectives:

- 110 • To determine the experiences of HCPs when discharging older people from the ED
- 111 • To examine the differing practices, if any, between OOH and office hours discharge
112 of older adults from the ED
- 113 • To identify recommendations for future practice when discharging older people OOH
114 from the ED

115 **Study design**

116 In this study a qualitative methodology was used consisting of semi-structured interviews. The
117 questions for the interviews were constructed by the researchers following a review of relevant
118 literature, taking cognisance of the Integrated Care Theoretical Framework ([15](#)), the
119 biopsychosocial aspects of health, and the Integrated Care Programmes for Older People
120 (ICPOP). ICPOP was introduced in Ireland with an aim to improve quality of life for older
121 people and to shift from acute hospital, episodic care to community based longitudinal,
122 coordinated, and integrated care ([16](#)). This links in with the aim, objectives and theoretical
123 framework of this study. The interview schedule (see Appendix 1) was developed by the

124 authors and reviewed by an ED expert group. A pilot study with one participant was also
125 undertaken to ensure further rigour of the semi-structured questions.

126 **Population and Recruitment**

127 This study was undertaken in Ireland. Inclusion criteria for participants stipulated that they
128 were required to be senior healthcare professionals working in an ED. Fourteen participants
129 in total responded which included clinical nurse managers, consultants in emergency
130 medicine, advanced nurse practitioners, clinical nurse specialists in frailty, clinical education
131 facilitators and physiotherapists. Participants included a broad range of healthcare
132 professionals and originated from a diversity of urban and rural geographical areas and
133 models of hospitals which added to the rigour of the study.

134 **Ethical approval**

135 Ethical approval was obtained from Ulster University Research Ethics Filter committee
136 (Project Number: FCNUR-22-017) as well as from the individual hospital Research Ethics
137 Committees (RECs). No national ethics committee currently exists in Ireland; therefore,
138 approval had to be sought from individual hospital RECs. In many of the sites, the Consultant
139 in Emergency Medicine acted as gatekeeper for the research participants as stipulated by the
140 hospital REC. Following confirmation of ethical approval, assistance to identify and invite
141 participants to take part in the study was provided by members of The National Clinical
142 Programme for Emergency Medicine and The Irish Association for Emergency Medicine. All
143 participants were provided with a participation information sheet providing the purpose, risks,
144 benefits of the study, the right to withdraw, and the assurance of anonymity. This was
145 explained by one of the research team to each of the respondents and a consent form was
146 subsequently signed by participants prior to commencement. Consent was also obtained for
147 the possible dissemination of the study findings. Due to the small number of urban and rural
148 EDs in Ireland, it was vital not to breach General Data Protection Regulations (GDPR) by

149 identifying any of the hospitals and/or the participants who contributed to the study. Therefore,
150 careful selection and reporting of the narrative exemplars was undertaken to take cognisance
151 of and avoid the risk of this possibility.

152 **Data collection**

153 The recruitment period for this study was between February 17th and November 24th, 2023.
154 One-to-one interviews were conducted. Written informed consent was obtained by the
155 researcher and consent was reconfirmed at commencement of the interviews with all the
156 participants. All interviews took place via video conferencing and at a time to suit the
157 participants following individual consent to use this option. Due to the busyness of the ED
158 environment, time constraints for participants and the distance between each site for the
159 researcher, this method was deemed to be the most appropriate. Interviews lasted 50 to 60
160 minutes in duration and were recorded using a digital recorder and via the computer to reduce
161 the risk of loss of the interview data. Data saturation was reached following the final interview
162 as no new data was noted and recruitment then ceased. The recordings were transcribed by
163 one member of the research team and checked for accuracy prior to analysis (MD).

164 **Data analysis**

165 Data analysis was undertaken using Braun and Clarke's (17) six-step framework for Thematic
166 Analysis (TA) which included familiarisation with the data set; coding; generating initial
167 themes; developing and reviewing themes; refining, defining, and naming themes; and writing
168 up. TA is a highly effective method to use when looking to analyse and understand a collection
169 of experiences, thoughts, or behaviours (18), therefore taking cognisance of the aim of this
170 study, this framework was selected as appropriate to use in this data analysis. These six steps
171 involved a collaborative approach with the four members of the research team (MD, AR, DG,
172 SMcl). Several meetings took place to interpret the data, identify categories, codes, provisional
173 themes, edit and refine themes to ensure rigour, consensus, and a richer interpretation of the

174 data. These collaborative meetings with the experienced research team at all stages of the
175 study are advocated to ensure reflexive practices in qualitative research (19; 20). The
176 Standards for Reporting Qualitative Research (SRQR) (21) checklist was also completed to
177 ensure rigour and a transparent reporting of the research process (see Appendix 2).

178

179 Findings

180 Following data analysis, several themes and sub-themes were identified. One recurrent theme
181 highlighted the issues of risks and safety for older people being discharged from the ED OOH.
182 In addition to this over-arching theme, several sub-themes were also identified (see Table 1).
183 Significant differences were reported by the respondents between office hours and OOH
184 discharge of older adults from the ED.

185

186 Table 1: Risks and Safety Theme and Sub-themes

Theme	Sub-themes
Risks and safety during older person	Should they stay or should they go?
Emergency Department discharge out of hours	Bright lights and noise
	New ways of working

187

188

189 Should they stay or should they go?

190 Participants reported on the risks in OOH ED discharge for the older person. The safety
191 decision to discharge an older adult OOH from the ED or to detain them until reviewed by

192 specialist older people services during office hours was influenced by several factors, including
193 the time of discharge and distance to home. However, if there was any concern that the older
194 person required more support following discharge OOH, it was reported that s/he would be
195 detained or “boarded for review” in ED until 8am when older person specialist staff are
196 available. This was despite staff being aware of the pressures of not having people lying on
197 trolleys.

198 *“It’s two or four in the morning, but they are a very vulnerable high-risk patient and they’re*
199 *going to go home to an empty house where there is no one. There’s not a light on... we*
200 *would keep them until the morning.” (Clinical Nurse Manager No. 2)*

201 Participants reported being conscious of the need to ensure safe discharge and refer older
202 people to the specialist supports in the ED and the community OOH. They also reported the
203 benefits of having these specialist older person teams as a great form of support. However,
204 participants also identified difficulties with the limited availability of these specialists OOH
205 whilst trying to run a 24-hour service in the department and the need to make a quick decision.
206 During OOH, it was identified that admission avoidance became a difficulty due to a lack of
207 resources for the older person both within the ED and in the community as these specialist
208 staff are not on duty. OOH discharge was not deemed to be as safe as office hours. Often
209 older adults would end up having to stay in ED or be admitted due to an identified risky
210 discharge OOH, especially at weekends.

211 *“There is a huge differential in you coming and being seen by a Frailty Intervention Team*
212 *during office hours on Monday to Friday and you coming to the emergency department at*
213 *any other time.” (Consultant in Emergency Medicine No.1)*

214 Respondents also reported a level of risk based on the variances of when assessments were
215 undertaken, and their suitability for assessing older people attending the ED. This included
216 triage and frailty assessments following ED presentation. Variances existed in relation to if,
217 and when, these assessments took place, and the scales used. Some participants reported

218 that frailty assessments were not undertaken at the initial triage assessment and not at all
219 during OOH due to demands on and availability of specialist older person team members, and
220 the numbers of attendees to the ED. Others reported it was the first important step and always
221 tried to get older people assessed as quickly as possible. Again, variances occurred between
222 office hours and OOH practice.

223 *“We do the clinical frailty scale.... The triage nurses are overwhelmed.... They can’t keep up*
224 *with the tsunami of patients coming through, so that would be done post triage.” (Consultant*
225 *in Emergency Medicine No. 1)*

226 Several respondents reported strategies to reduce risk and ensure safety for older people
227 when being discharged home OOH from the ED. These included arranging transport,
228 providing a staff escort for support and to ensure safety, and using a discharge lounge which
229 was only available during office hours. Practices varied between hospitals. Many hospitals
230 witnessed changing variances in staff profiles and skill mix with staff turnover being significant
231 in some areas. This change in personnel following the employment of many international
232 nurses to Ireland was reported to be a difficulty for the older adult in the ED, with cultural and
233 language differences evident. This was identified as a specific issue OOH when no specialist
234 frailty team support was available to support these new international staff members who may
235 be unfamiliar with the specific needs of older people on discharge from the ED. Thus, adding
236 to a potential risk on OOH discharge. Other challenges identified included the increasing
237 numbers of international older adults with growing health care needs living in inappropriate
238 places such as hotels, and associated difficulties in accessing support on discharge from the
239 ED.

240 *“There are issues with new staff from outside of Ireland, and the biggest problem is*
241 *communication particularly for the elderly patient. Language is a big thing, same with the*
242 *doctor. They just have a different culture. They just presume that there is going to be*
243 *somebody at home to look after that patient, because that's what they would do.” (Clinical*
244 *Education Facilitator No. 4)*

245 **Bright lights and noise**

246 Participants identified several risks for the older adults who were delayed or detained in the
247 ED OOH whilst awaiting discharge, often because of an overstimulated environment resulting
248 in adverse consequences. These included significant biological and psychological effects such
249 as sleep deprivation, anxiety, delirium, and medication and mobility issues. Older people also
250 faced challenges within the ED environment, including being exposed to frightening incidents,
251 and the risks of acquiring infections and pressure ulcers. A drop in the level of supervision
252 and care from HCPs was also a concern once an older person was deemed fit for discharge
253 but awaiting review by the specialist team.

254 *“Older people staying in an ED can.... get sicker, pressure sores and anxiety..... you have to*
255 *weigh up the pros and cons and what is safest in the long run and having to keep them in to*
256 *be seen by the team the next morning.” (Clinical Nurse Specialist in Frailty, No. 2)*

257 Keeping an older person in ED OOH for follow-up by the older person services although
258 medically discharged was reported to result in an increase in patient experience times and a
259 reduction in patient flow within the ED. This was identified as a cause of stress for staff and
260 an increase in workload, due to overcrowding within the ED but also due to a reflection of poor
261 ED performance because of prolonged patient experience times for those over 75 years of
262 age.

263 *“Out of hours, where our colleagues notice an issue with a patient that would benefit from*
264 *the multi-disciplinary team input, is when we would see that knock-on effect on time.”*
265 *(Consultant in Emergency Medicine No.2)*

266 It was reported that older people also faced challenges within the ED environment OOH, as
267 often no specific older person area was available within the ED due to the volume of
268 attendances, overcrowding and a lack of space. Respondents reported that this could lead to
269 increased adverse outcomes for the older person because of difficulties in meeting

270 fundamental needs, from having to stay overnight on a chair or trolley and often on a busy
271 corridor.

272 *“.... for a patient who has poly morbidities, their risks of having a negative outcome are much
273 higher. There is also the risk of them not getting appropriate medications, and meals as well
274 as not getting the toilet” (Consultant in Emergency Medicine No.2)*


275

276 **New ways of working**

277 Participants interviewed recognised the difficulties in providing a 24-hour service in the ED,
278 when their counterparts in other areas of the hospital and community services were not
279 available. They were very forthcoming in providing recommendations for future practice to
280 improve services for older people whilst being discharged from the ED OOH (see Figure 1).
281 This included various aspects of discharge and care within the department and extending
282 services to have specialist older person HCPs working 24/7 to ensure safe older person
283 discharge from the ED OOH.

284

285



286

287

Fig 1 New ways of working- recommendations for practice

288

289 Many respondents identified the need to improve experiences OOH. This included ED staffing
 290 OOH, and the need for more older person specialist health care professionals 24/7 in the ED
 291 including physicians in older person emergency medicine and employment of a nurse to
 292 ensure any required follow-up is organised for older people following ED
 293 discharge. Respondents also highlighted the need for upskilling of all ED staff. With the
 294 increasing older person demographics, it was identified that a need existed to ensure that ED
 295 staff had the knowledge and skills to be able to provide a 24/7 service safely and effectively
 296 for this specialist cohort. This could be in the form of a national education programme.

297 *“..... that the ED staff themselves need a greater skills set which could be provided by a 2-*
 298 *day course with online learning we wouldn't contemplate having staff in the emergency*
 299 *department if they didn't have ACLS or ATLS training and yet, there isn't much of our work*
 300 *which is now cardiac arrest. There's far more of our work that is older adult now.”*

301

(Consultant in Emergency Medicine No.1)

302 Participants in this study recommended a greater and earlier focus on frailty on a 24/7 basis,
303 not just at triage in the ED, but also pre-attendance by ambulance personnel and at primary
304 care centres. This was identified as being very important considering the growth in older
305 people demographics internationally and the associated risk of an increased incidence of
306 frailty.

307 *“Ideally frailty should be scored in triage from the minute the person presents to the door. I*
308 *think everybody should be thinking frailty..... to start scoring them coming through the door*
309 *from triage. And even before that, the ambulance staff, the GP surgery.” (Clinical Nurse*
310 *Specialist in Frailty No. 1)*

311 Recommendations from participants also included a need to focus on specific documentation
312 which could identify and highlight the risk of frailty for the older person presenting to the ED.
313 It was suggested that a national discharge checklist could help to ensure that there is
314 uniformity in discharging the older person from the ED. A focus on written instructions and
315 patient education for older people in the ED was also suggested in order to have a record of
316 treatment provided and further care required. Improvements to the ED environment were also
317 suggested. These included specific frailty units and separate treatment areas for older people
318 as seen internationally. It was suggested that these could help reduce the risk of adverse
319 reactions for older adults presenting to the ED by the avoidance of frightening experiences in
320 a busy, highly stimulated ED.

321 *“What I would say needs to happen is to have an older person's unit, like a frailty unit part of*
322 *the ED. In America some of the hospitals are just older people EDs because that's the*
323 *population they're getting. Their whole focus in the 24 hours is older people. And that's the*
324 *way it's going to be. It's not going to go away.” (Clinical Nurse Specialist in Frailty, No. 1)*

325

326 **Discussion**

327 This paper identified the safety and risk aspects of HCP's experiences of discharging older
328 people during OOH versus office hours from the Emergency Department. Patient safety is
329 defined as "the reduction of risk of unnecessary harm associated with healthcare to an
330 acceptable minimum" (22). However, it is estimated that around one in ten patients are harmed
331 during health care, resulting in substantial consequences for patients (23). Discharge
332 processes are an important aspect of a patient's health care journey. Many older adults who
333 attend the ED for assessment are subsequently discharged. Participants in this study
334 highlighted risks for both discharging older people home OOH without organised supports and
335 in keeping older patients in the ED until services are available. The results identified important
336 but diverse practices regarding older people ED discharge occurring in different geographical
337 areas and between OOH and office hours. Despite its unique and challenging clinical
338 environment, the principles of patient-centred care and shared decision making are evident
339 within the ED (24) with a holistic problem-solving, person-centred approach to discharge
340 needed (14). Participants in this study were cognisant of the wishes of the older person;
341 however, this was alongside their concern over patient safety if an adverse event were to
342 occur post discharge or indeed in the ED OOH whilst waiting until further assessment could
343 take place during office hours.

344 Whilst some developments in ED care are not new, such as an admission avoidance (AA)
345 team (25), in more recent years many welcome developments have been realised in older
346 person ED care and follow up in the community. These aim to expand community healthcare
347 services to be closer to where older people reside and to reduce the demand on hospital
348 services by admission avoidance and transition from hospital to home (26), recognising the
349 risks of hospitalisation for this cohort. A study exploring the impact of Home FIRsT (Frailty
350 Intervention & Response Team) interventions on older patients' experience times in the ED
351 (27), identified a need for further research in the area. The authors recognised that
352 assessments by the Home FIRsT can be lengthy and in EDs where there are waiting criteria
353 (e.g. 4-hour National Health Service target), it would be particularly challenging to run such a

354 service. Although the study excluded patients attending out of hours, the authors did suggest
355 a future potential to deliver continuous Home FIRsT-like services 24 hours a day, seven days
356 a week, both in the ED and in community services. Interestingly, ED PLUS, an initiative
357 introduced to link the transition of care between the ED visit and the community, was shown
358 to result in a lower incidence of functional decline and improvement in quality of life for the
359 older adult post ED (28). Whilst these developments are very welcome, the findings in this
360 study showed that many older person specialist teams are inaccessible in the ED or in the
361 community when discharge occurs OOH which impacted on the decision to discharge.

362 Point-of-care identification is needed for older adults presenting to the ED who may be at risk
363 of adverse outcomes (29). Assessments including triage, comprehensive geriatric
364 assessment, frailty and delirium assessments are regularly undertaken with the older person
365 attending the ED. Zachariasse et al (30) and Brouns et al (31) reported that the validity of the
366 Manchester Triage System in emergency care was moderate to good; however, lowest
367 performance was in older patients over 65 years of age. Questions over the use of the
368 Manchester Triage System with older adults in the ED also emerged in this study, with some
369 participants expressing concerns over its suitability. Potential risks for older adults with frailty
370 in the ED have been recognised. Frailty is often assessed using the Clinical Frailty Scale on
371 a scale from 1-9, assessing domains such as comorbidity, function, and cognition (32).
372 Significant findings from a cross-sectional study in 14 European countries, identified that 40%
373 of older people attending the ED had a score of 5 or more when assessed using Rockwood
374 et al's (33) Version 1.2 Clinical Frailty Scale (34). The use of a clinical frailty scale in ED may
375 assist with prompt decision making at triage (35), timely access to appropriate care for older
376 patients attending (36; 37), and in obtaining a more holistic clinical assessment (38).
377 Additionally, early assessment and intervention in ED can increase patient satisfaction levels
378 (5). Although not OOH focused, O'Shaughnessy et al (39), also identified that comprehensive
379 geriatric assessment may help improve patient outcomes by achieving admission avoidance
380 and thus avoiding any possible adverse consequences. However, the ED environment is not

381 appropriate for performing such in-depth assessments, with a need for quick and user-friendly
382 tools (40) as frailty screening should be undertaken within four hours of attending the ED and
383 take less than five minutes to complete (41). Daily encounters with older people with frailty will
384 be a norm for almost all health and social care professionals (42). Therefore, staff must be
385 cognisant of increased frailty especially during long department waits in overcrowded and
386 sensory heightened areas present in many EDs. Despite the reported benefits of using a
387 Clinical Frailty Scale at triage, this study identified that frailty assessment is not performed at
388 triage in some hospitals and often never undertaken OOH, resulting in delayed or missed
389 identification of some older adults with frailty.

390 Whilst early ED frailty screening at triage can transform care through fast tracking to required
391 care pathways, education for all professionals is essential to ensure engagement in and
392 recognition of frailty (43). Despite the prevalence of frailty, there is a lack of training in medical
393 and nursing curricula and a need exists for education and skills development for HCPs (44).
394 Emergency services must be equipped to provide care for older patients who present with
395 complex health and care needs (34). Assessments can also be compounded by language
396 barriers in nurse/patient communication which Ali and Watson (45) reported can have adverse
397 effects on the provision of appropriate and effective care. Indeed, the World Health
398 Organisation (46) highlighted the increase in the international migration and mobility of health
399 workers. In this present study, participants identified communication issues between staff and
400 older patients, and cultural and decision-making concerns during older person ED discharge
401 OOH. Health education is an important aspect of ensuring safe discharge re treatment and
402 follow-up. However, participants reported it as a difficulty for many patients being discharged
403 from the ED. A need exists to ensure a greater adherence to discharge instructions and to
404 improve the transition from ED to home (47), possibly using a multimodal approach in
405 delivering discharge instructions (48). Participants in this study highlighted issues surrounding
406 ED discharge information for older adults and recommended the need for specific older patient
407 discharge information leaflets.

408 ED discharge is associated with increased risk with up to six months decline in functional
409 status for the older adult living in the community (49). Concerningly almost 80% of older people
410 are discharged from the ED with a minimum of one unaddressed health concern (Memedovich
411 et al (50). Condon et al (51) explored the transition of older adults to the community from the
412 ED. Whilst the qualitative review was not OOH focused, findings identified fragmented care,
413 communication difficulties, reports of unplanned discharges with transport not being organised
414 for older adults. Participants in this study identified that they had to consider many factors to
415 ensure a safe older person ED discharge including time, distance, and transport issues.
416 Discharging an older adult living alone in a rural area may not be a satisfactory start to the
417 discharge process. In Ireland, around 5% of the population live 50km or more from an adult
418 emergency department while 0.8% of the population have a travel distance of 75km or more
419 (52). Time of discharge can be challenging for older people, especially at night when support
420 networks and services in the community are less available (14).

421 This study also highlighted the risks in being detained in ED whilst awaiting further
422 assessments prior to discharge. Older person ED discharge needs to be safe and effective on
423 a 24/7 basis. However, although not specific to OOH, some of the challenges caring for older
424 people in the ED, included the unsuitable physical environment and equipment,
425 communication issues, a limited time factor, and a lack of geriatric-specific knowledge (53).
426 Attendance at an ED can have adverse consequences for the older patient with dementia (54;
427 8), which can be very distressing but avoidable for patients and their families. Healthcare
428 systems benefit from providers delivering a positive patient experience (55). However, long
429 waits, difficult waiting environments, lack of communication, privacy, and personal care have
430 been identified previously as areas of concern for older people in the ED (7). Additionally,
431 many healthcare needs of older adults are not being met either in the ED or by interventions
432 initiated by the ED (50). Similarly, in this study, participants recognised the potentially serious
433 risks in keeping an older adult for extended periods of time in the ED, which included risk of
434 infection, delirium, anxiety, and possible neglect. Often this delay OOH was due to the

435 perceived risks for the older person, and the unavailability of specialist staff who only work
436 during office hours. This is a major area of concern as these serious risks are preventable if
437 adequate resources were in place.

438 ED overcrowding results in decreased access to an appropriate treatment space, and a
439 decrease in adherence to the National Emergency Access Target (NEAT) (56). An increase
440 in older people attendances with complex and chronic conditions are causes of overcrowding
441 and it is recognised that there is need for efficient ED patient flow, using initiatives to meet
442 timed patient disposition targets, as well as expanded primary care (57). Internationally, older
443 adults have advocated for older adult services such as a geriatric ED to meet specific health
444 needs for their cohort which could help to reduce overcrowding and wait times (58; 7. Similar
445 difficulties were also highlighted in this study including increased wait times and older people
446 having to frequently stay overnight on a chair in an over stimulated ED corridor, due to
447 overcrowding, with a recommendation for specific older person areas.

448 Lowthian et al (59) reported a need for further research around safe older person ED discharge
449 as a lack of consensus existed as to what forms evidence-based best practice for patients
450 discharged from the ED. Enhancement of integrated care is required especially with regards
451 to improved communications, sharing of responsibility between care providers, additional
452 resources and further research evaluating care practices (60). Recognising the need to
453 improve care for older patients in the ED, Lucke et al (61) developed expert clinical
454 recommendations for Geriatric Emergency Medicine (GEM). Mooijaart et al (62) also
455 recognised the rapid development of GEM in response to ageing demographics, proposing
456 that further work, research and innovation is needed to provide holistic care for the older adult,
457 including organisation of care, identification of patient preferences and needs, and improving
458 HCP's knowledge and expertise. Respondents in this study also identified a need for improved
459 documentation, staff training, and national discharge guidelines to ensure uniformity of
460 discharge practices for older people from all EDs on a 24/7 basis.

461

462 **Strengths and weaknesses**

463 This study had several strengths and weaknesses. Firstly, whilst considerable research has
464 been undertaken on ED matters in recent years, a dearth still exists in relation to out of hours
465 ED discharge experiences. This study is the first of its kind to explore this important topic and
466 has addressed the gap in research which existed. It is likely that the practices, experiences,
467 and concerns regarding out of hours discharge seen in this study also pertain to EDs globally
468 although there is a need for further international research. Secondly, participants included
469 HCPs from a variety of backgrounds and from a wide geographical spread which enhanced
470 the rigour of the study. This study identified diverse practices in healthcare, taking cognisance
471 of the availability of resources and follow-up care, the geographical landscape, demographics,
472 and cultural aspects in caring for older family members. However, additional participants such
473 as occupational therapists and social workers may have generated additional insightful
474 experiences.

475

476 **Conclusions**

477 Whilst the older adult cohort are one of the main attenders at EDs, many are subsequently
478 discharged home. However, this study highlighted that if deemed not for admission but unfit
479 for discharge home, older adults may experience a delayed discharge OOH. This principally
480 occurs if a perceived biological/psychological/social risk is identified and/or if further
481 assessment is required by members of the multi-disciplinary team, yet the service is
482 unavailable OOH. Due to the multiple contesting factors within a dynamic emergency
483 department, attendance and long waits therein can lead to a risk of serious adverse effects for
484 the older adult including discomfort from inappropriate waiting areas, a lack of sleep, anxiety,
485 and delirium. Patient experience times and ED flow can also be impacted. To ensure a safe

486 and effective discharge home 24/7 in every ED regardless of location, there is a need to
487 address gaps in care identified. With the significant increase in older person demographics,
488 there is an urgent need for healthcare managers and relevant policy makers to develop best
489 practice guidelines, ensure greater availability of older people services and teams both in the
490 ED and in the community, increase explicit older person ED treatment areas, and develop
491 specific education courses and upskilling for ED staff. These recommendations can assist in
492 achieving a safe and appropriate older person ED discharge around the clock.

493

494 **Supporting information:**

495 S1 Appendix 1: Interview schedule (DOC)

496 S2 Appendix 2 Standards for reporting qualitative research checklist (SRQR) (DOC)

497

498 **Acknowledgements**

499 The authors acknowledge the participating health care professionals for their interest, time,
500 and honesty in this study, along with support and guidance received from members of the
501 National Clinical Programme for Emergency Medicine and the Irish Association for Emergency
502 Medicine.

503

504 **Conflict of interest**

505 The authors declare no conflict of interest.

506

507 **Author Contributions**

508 **Conceptualization:** Mary Dunnion, Assumpta Ryan, Debbie Goode, Sonja McIlpatrick.
509 **Formal analysis:** Mary Dunnion, Assumpta Ryan, Debbie Goode, Sonja McIlpatrick.
510 **Investigation:** Mary Dunnion.
511 **Methodology:** Mary Dunnion, Assumpta Ryan, Debbie Goode, Sonja McIlpatrick.
512 **Project administration:** Mary Dunnion.
513 **Supervision:** Assumpta Ryan, Debbie Goode, Sonja McIlpatrick.
514 **Writing– original draft:** Mary Dunnion
515 **Writing– review & editing:** Mary Dunnion, Assumpta Ryan, Debbie Goode, Sonja McIlpatrick.

516

517 **References**

- 518 1. World Health Organisation. Ageing and health. 2022. Available:
519 [https://www.who.int/news-room/fact-sheets/detail/ageing-and](https://www.who.int/news-room/fact-sheets/detail/ageing-and-health#:~:text=Every%20country%20in%20the%20world%20is%20experiencing%20growth,from%201%20billion%20in%202020%20to%201.4%20billion.)
520 [health#:~:text=Every%20country%20in%20the%20world%20is%20experiencing%20](https://www.who.int/news-room/fact-sheets/detail/ageing-and-health#:~:text=Every%20country%20in%20the%20world%20is%20experiencing%20growth,from%201%20billion%20in%202020%20to%201.4%20billion.)
521 [growth,from%201%20billion%20in%202020%20to%201.4%20billion.](https://www.who.int/news-room/fact-sheets/detail/ageing-and-health#:~:text=Every%20country%20in%20the%20world%20is%20experiencing%20growth,from%201%20billion%20in%202020%20to%201.4%20billion.) Accessed:
522 07/06/2024
- 523 2. United Nations General Assembly. United Nations Decade of Healthy Ageing (2021-
524 2030): resolution/adopted by the General Assembly. 2020. United Nations, New York.
525 Available: [https://policycommons.net/artifacts/8967746/united-nations-decade-of-](https://policycommons.net/artifacts/8967746/united-nations-decade-of-healthy-ageing-2021-2030/9835673/)
526 [healthy-ageing-2021-2030/9835673/](https://policycommons.net/artifacts/8967746/united-nations-decade-of-healthy-ageing-2021-2030/9835673/) Accessed: 07/06/2024.
- 527 3. Eliopoulos C. Gerontological Nursing (Tenth edition). China, Wolters Kluwer; 2021.
- 528 4. World Health Organization. Decade of Healthy Ageing Baseline Report. World Health
529 Organization, Geneva; 2020.
- 530 5. Cassarino M, Robinson K, Trépel D, O’Shaughnessy Í, Smalle E, White S, Devlin C,
531 Quinn R, Boland F, Ward ME. and McNamara R. Impact of assessment and

- 532 intervention by a health and social care professional team in the emergency
533 department on the quality, safety, and clinical effectiveness of care for older adults: A
534 randomised controlled trial. *PLoS Medicine*, 2021; 18(7), p.e1003711.
- 535 6. Corcoran G, Bernard P, Kenna L, Malone A, Horgan F, O'Brien C, Ward P, Howard
536 W, Hogan L, Mooney, R., Masterson, S. "Older People Want to Be in Their Own
537 Homes": A Service Evaluation of Patient and Carer Feedback after Pathfinder
538 Responded to Their Emergency Calls. *Prehospital Emergency Care*. 2023 ;27(7):866-
539 874.
- 540 7. Mwakilasa, M. T., Foley, C., O'Carroll, T., Flynn, R., & Rohde, D. Care experiences of
541 older people in the emergency department: a concurrent mixed-methods study.
542 *Journal of Patient Experience*, 2021; 8, doi:10.1177/23743735211065267.
- 543 8. Goode, D., Ryan, A., Melby, V., & Slater, P. Care experiences of older people with
544 mental health needs and their families in emergency medical services settings.
545 *International Journal of Older People Nursing*, 2023; 18, e12500.
546 <https://doi.org/10.1111/opn.12500>
- 547 9. Rowe, A., & Knox, M. The impact of the healthcare environment on patient experience
548 in the Emergency Department: a systematic review to understand the implications for
549 patient-centered design. *HERD: Health Environments Research & Design Journal*,
550 2023; 16(2), 310-329.
- 551 10. Greenwald, P.W., Estevez, R.M., Clark, S., Stern, M.E., Rosen, T., Flomenbaum, N.
552 The ED as the primary source of hospital admission for older (but not younger) adults.
553 *The American Journal of Emergency Medicine*, 2016, 34, 6, pgs. 943-947. ISSN 0735-
554 6757. <https://doi.org/10.1016/j.ajem.2015.05.041>
- 555 11. Cetin-Sahin D, Ducharme F, McCusker J, Veillette, N., Cossette, S., Minh Vu, T.,
556 Vadeboncoeur, A., Lachance, P., Mah, R., Berthelot, S. Experiences of an Emergency
557 Department Visit Among Older Adults and Their Families: Qualitative Findings from a
558 Mixed-Methods Study. *Journal of Patient Experience*, 2020; 7(3):346-356.
559 doi:10.1177/2374373519837238

- 560 12. Gettel, C.J., Serina, P.T., Uzamere, I., Hernandez-Bigos, K., Venkatesh, A.K., Rising,
561 K.L., Goldberg, E.M., Feder, S.L. Cohen, A.B., Hwang, U. Emergency department-to-
562 community care transition barriers: A qualitative study of older adults. *J Am Geriatr*
563 *Soc.*, 2022; 70(11): 3152-3162.
- 564 13. Dunnion, M., Ryan, A., Goode, D., McIlfratrick, S. Supporting older people following out
565 of hours discharge from the Emergency Department: An integrative review of the
566 literature. *International Journal of Older People Nursing*, 2023; 18(3): e12529.
- 567 14. Hyslop, B. 'Not safe for discharge'? Words, values, and person-centred care. *Age and*
568 *Ageing*, 2020; 49(3), 334-336.
- 569 15. World Health Organisation. World report on ageing and health. Geneva: World Health
570 Organization.2015.Available:[https://iris.who.int/bitstream/handle/10665/326843/WHO](https://iris.who.int/bitstream/handle/10665/326843/WHO-FWC-ALC-19.1-eng.pdf?sequence=17)
571 [-FWC-ALC-19.1-eng.pdf?sequence=17](https://iris.who.int/bitstream/handle/10665/326843/WHO-FWC-ALC-19.1-eng.pdf?sequence=17) Accessed: 19/08/2024.
- 572 16. Integrated Care Programme for Older People Steering Group. Making a start in
573 integrated care for older persons a practical guide to the local implementation of
574 integrated care programmes for older persons. 2017. Available:
575 [https://www.hse.ie/eng/services/publications/clinical-strategy-and-programmes/a-](https://www.hse.ie/eng/services/publications/clinical-strategy-and-programmes/a-practical-guide-to-the-local-implementation-of-integrated-care-programmes-for-older-persons.pdf)
576 [practical-guide-to-the-local-implementation-of-integrated-care-programmes-for-older-](https://www.hse.ie/eng/services/publications/clinical-strategy-and-programmes/a-practical-guide-to-the-local-implementation-of-integrated-care-programmes-for-older-persons.pdf)
577 [persons.pdf](https://www.hse.ie/eng/services/publications/clinical-strategy-and-programmes/a-practical-guide-to-the-local-implementation-of-integrated-care-programmes-for-older-persons.pdf) Accessed 22/7/2024.
- 578 17. Braun, V. & Clarke, V. Thematic Analysis. A Practical Guide. Sage Publications; 2022.
- 579 18. Braun, V. & Clarke, V. One size fits all? What counts as quality practice in (reflexive)
580 thematic analysis? *Qualitative Research in Psychology*, 2021; 18:3, 328-352.
- 581 19. Jamieson, M. K., Govaart, G. H., & Pownall, M. Reflexivity in quantitative research: A
582 rationale and beginner's guide. *Social and Personality Psychology Compass*, 2023;
583 17(4), e12735.
- 584 20. Olmos-Vega, F. M., Stalmeijer, R. E., Varpio, L., & Kahlke, R. A practical guide to
585 reflexivity in qualitative research: AMEE Guide No. 149. *Medical teacher*, 2023; 45(3),
586 241-251.

- 587 21. O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for reporting
588 qualitative research: a synthesis of recommendations. *Academic Medicine*, 2014; 89
589 (9):1245-1251. DOI: 10.1097/ACM.0000000000000388
- 590 22. World Health Organisation Patient Safety and World Health Organization. Conceptual
591 framework for the international classification for patient safety version 1.1: final
592 technical report January 2009 (No. WHO/IER/PSP/2010.2). Report No.: 606940937X.
593 Geneva, World Health Organization, 2010.
- 594 23. Slawomirski, L., Aaraaen, A., Klazinga, N.S. "The economics of patient
595 safety: Strengthening a value-based approach to reducing patient harm at national
596 level", *OECD Health Working Papers*, 2017, No. 96, OECD Publishing,
597 Paris. <https://doi.org/10.1787/5a9858cd-en>.
- 598 24. Hess, E. P., Grudzen, C. R., Thomson, R., Raja, A. S., & Carpenter, C. R. Shared
599 decision-making in the emergency department: respecting patient autonomy when
600 seconds count. *Academic Emergency Medicine*, 2015; 22(7), 856-864.
- 601 25. Hardy C, Whitwell D, Sarsfield B, Maimaris, C. Admission avoidance and early
602 discharge of acute hospital admissions: an accident and emergency-based scheme.
603 *Emergency Medicine Journal*, 2001; 18:435-440.
- 604 26. Health Service Executive. Enhanced Community Care. 2024. Available:
605 [https://www.hse.ie/eng/services/list/2/primarycare/enhanced-community-](https://www.hse.ie/eng/services/list/2/primarycare/enhanced-community-care/#:~:text=The%20ECC%20programme%20has%20delivered:%201.%2096%20o)
606 [care/#:~:text=The%20ECC%20programme%20has%20delivered:%201.%2096%20o](https://www.hse.ie/eng/services/list/2/primarycare/enhanced-community-care/#:~:text=The%20ECC%20programme%20has%20delivered:%201.%2096%20o)
607 [f](https://www.hse.ie/eng/services/list/2/primarycare/enhanced-community-care/#:~:text=The%20ECC%20programme%20has%20delivered:%201.%2096%20o) Accessed 17/07/2024.
- 608 27. O'Shaughnessy, Í., Romero-Ortuno, R., Edge, L., Dillon, A., Flynn, S., Briggs, R.,
609 Shields, D., McMahon, G., Hennessy, A., Kennedy, U. and Staunton, P. Home FIRsT:
610 interdisciplinary geriatric assessment and disposition outcomes in the emergency
611 department. *European journal of internal medicine*, 2021; 85, pp.50-55.
- 612 28. Conneely, M., Leahy, S., O'Connor, M., Corey, G., Gabr, A., Saleh, A., Okpaje, B.,
613 O'Shaughnessy, Í., Synnott, A., McCarthy, A. and Holmes, A. A physiotherapy-led
614 transition to home intervention for older adults following emergency department

- 615 discharge: a pilot feasibility randomised controlled trial (ED PLUS). *Clinical*
616 *Interventions in Aging*, 2023; pp.1769-1788.
- 617 29. Carpenter, C. R., Shelton, E., Fowler, S., Suffoletto, B., Platts-Mills, T. F., Rothman,
618 R. E., & Hogan, T. M. Risk factors and screening instruments to predict adverse
619 outcomes for undifferentiated older emergency department patients: a systematic
620 review and meta-analysis. *Academic Emergency Medicine*, 2015; 22(1), 1-21.
- 621 30. Zachariasse, J.M., Seiger, N., Rood, P.P., Alves, C.F., Freitas, P., Smit, F.J.,
622 Roukema, G.R. and Moll, H.A. Validity of the Manchester Triage System in emergency
623 care: A prospective observational study. *PloS ONE*, 2017; 12(2), p.e0170811.
- 624 31. Brouns, S. H., Mignot-Evers, L., Derkx, F., Lambooj, S. L., Dieleman, J. P., & Haak,
625 H. R. Performance of the Manchester triage system in older emergency department
626 patients: a retrospective cohort study. *BMC Emergency Medicine*, 2019; 19, 1-11.
- 627 32. Church, S., Rogers, E., Rockwood, K., & Theou, O. A scoping review of the Clinical
628 Frailty Scale. *BMC geriatrics*, 2020; 20, 1-18.
- 629 33. Rockwood, K., Song, X., MacKnight, C., Bergman, H., Hogan, D.B., McDowell, I.,
630 Mitnitski, A. A global clinical measure of fitness and frailty in elderly people. *Canadian*
631 *Medical Association Journal*, 2005; 173(5):489-495.
- 632 34. European Taskforce on Geriatric Emergency Medicine (ETGEM) Collaborators.
633 Prevalence of Frailty in European Emergency Departments (FEED): an international
634 flash mob study. *European Geriatric Medicine*, 2024; 15(2):463-470.
- 635 35. Elliott, A., Taub, N., Banerjee, J., Aijaz, F., Jones, W., Teece, L., van Oppen, J. and
636 Conroy, S. Does the clinical frailty scale at triage predict outcomes from emergency
637 care for older people? *Annals of emergency medicine*, 2021; 77(6), pp.620-627.
- 638 36. Blomaard, L.C., Speksnijder, C., Lucke, J.A., de Gelder, J., Anten, S., Schuit, S.C.,
639 Steyerberg, E.W., Gussekloo, J., de Groot, B. and Mooijaart, S.P. Geriatric screening,
640 triage urgency, and 30-day mortality in older emergency department patients. *Journal*
641 *of the American Geriatrics Society*, 2020; 68(8), pp.1755-1762.

- 642 37. Ng, C.J., Chien, L.T., Huang, C.H., Chaou, C.H., Gao, S.Y., Sherry, Y.H.C., Hsu, K.H.
643 & Chien, C.Y. Integrating the clinical frailty scale with emergency department triage
644 systems for elder patients: A prospective study. *The American Journal of Emergency*
645 *Medicine*, 2023; 66, pp.16-21.
- 646 38. Elliott, A., Phelps, K., Regen, E., & Conroy, S. P. Identifying frailty in the Emergency
647 Department—feasibility study. *Age and ageing*, 2017; 46(5), 840-845.
- 648 39. O’Shaughnessy, Í., Robinson, K., Whiston, A., Barry, L., Corey, G., Devlin, C.,
649 Hartigan, D., Synnott, A., McCarthy, A., Moriarty, E. and Jones, B. Comprehensive
650 geriatric assessment in the emergency department: a prospective cohort study of
651 process, clinical, and patient-reported outcomes. *Clinical Interventions in Aging*, 2024;
652 pp.189-201.
- 653 40. van Dam, C.S., Hoogendijk, E.O., Mooijaart, S.P., Smulders, Y.M., de Vet, R.C.,
654 Lucke, J.A., Blomaard, L.C., Otten, R.H., Muller, M., Nanayakkara, P.W. and
655 Trappenburg, M.C. A narrative review of frailty assessment in older patients at the
656 emergency department. *European Journal of Emergency Medicine*, 2021; 28(4),
657 pp.266-276.
- 658 41. Moloney, E., O’Donovan, M.R., Carpenter, C.R., Salvi, F., Dent, E., Mooijaart, S.,
659 Hoogendijk, E.O., Woo, J., Morley, J., Hubbard, R.E. and Cesari, M. Core
660 requirements of frailty screening in the emergency department: an international Delphi
661 consensus study. *Age and ageing*, 2024; 53(2), p. afae013.
- 662 42. Conroy, S., Carpenter, C., Banerjee, J. Silver Book II Quality Care for Older People
663 with Urgent Care Needs [Online]. 2021. Available: [Silver Book II | British Geriatrics](https://www.bgs.org.uk/silver-book-ii)
664 [Society \(bgs.org.uk\)](https://www.bgs.org.uk) Accessed: 07/06/2024.
- 665 43. van Oppen, J. D., & Suzanne, M. Frailty screening in the Emergency Department: why
666 does it matter? *Age and Ageing*, 2024; 53(4), afae056.
- 667 44. Hammel, I. S., & Chapman, E. N. Frailty training among healthcare professionals. *The*
668 *Journal of nutrition, health and aging*, 2024; 28 (6), 100258.

- 669 45. Ali, P. A., & Watson, R. Language barriers and their impact on provision of care to
670 patients with limited English proficiency: Nurses' perspectives. *Journal of Clinical*
671 *Nursing*, 2018; 27(5-6), e1152-e1160.
- 672 46. World Health Organisation. Addressing the international migration of health workers.
673 2024. Available: [Addressing the international migration of health workers \(who.int\)](#)
674 Accessed: 13/8/2024.
- 675 47. Benjenk, I., DuGoff, E.H., Jacobsohn, G.C., Cayenne, N., Jones, C.M., Caprio, T.V.,
676 Cushman, J.T., Green, R.K., Kind, A.J., Lohmeier, M. and Mi, R. Predictors of older
677 adult adherence with emergency department discharge instructions. *Academic*
678 *Emergency Medicine*, 2021; 28(2), pp.215-225.
- 679 48. Hoek, A. E., Anker, S. C., van Beeck, E. F., Burdorf, A., Rood, P. P., & Haagsma, J.
680 A. Patient discharge instructions in the emergency department and their effects on
681 comprehension and recall of discharge instructions: a systematic review and meta-
682 analysis. *Annals of Emergency Medicine*, 2020; 75(3), 435-444.
- 683 49. Nagurney, J. M., Fleischman, W., Han, L., Leo-Summers, L., Allore, H. G., & Gill, T.
684 M. Emergency department visits without hospitalization are associated with functional
685 decline in older persons. *Annals of emergency medicine*, 2017; 69(4), 426-433.
- 686 50. Memedovich, A., Asante, B., Khan, M., Eze, N., Holroyd, B. R., Lang, E., Kashuba, S.,
687 Clement, F. Strategies for improving ED-related outcomes of older adults who seek
688 care in emergency departments: a systematic review. *International Journal of*
689 *Emergency Medicine*, 2024; 17, 16. <https://doi.org/10.1186/s12245-024-00584-7>.
- 690 51. Condon, B., Griffin, A., Fitzgerald, C., Shanahan, E., Glynn, L., O'Connor, M., Hayes,
691 C., Manning, M., Galvin, R., Leahy, A. and Robinson, K. Older adults experience of
692 transition to the community from the emergency department: a qualitative evidence
693 synthesis. *BMC geriatrics*, 2024; 24(1), p.233.
- 694 52. Central Statistics Office. Measuring Distance to Everyday Services in Ireland [Health](#)
695 [Services Measuring Distance to Everyday Services in Ireland - Central Statistics](#)

- 696 [Office. 2024. Available: Introduction Measuring Distance to Everyday Services in](#)
697 [Ireland - Central Statistics Office](#) Accessed 19/07/202
- 698 53. Lennox, A., Braaf, S., Smit, D. V., Cameron, P., & Lowthian, J. A. (2019) Caring for
699 older patients in the emergency department: health professionals' perspectives from
700 Australia—the safe elderly emergency discharge project. *Emergency Medicine*
701 *Australasia*, 2019; 31(1), 83-89.
- 702 54. Watkins, S., Murphy, F., Kennedy, C., Graham, M., & Dewar, B. Caring for older people
703 with dementia in the emergency department. *British Journal of Nursing*, 2020; 29(12),
704 692-699.
- 705 55. Aleksandrovskiy, I., Ganti, L., Simmons, S. The Emergency Department Patient
706 Experience: In Their Own Words. *Journal of Patient Experience*, 2022; 20;9,1-4.
- 707 56. Smith, B., Bouchoucha, S., & Watt, E. 'Care in a chair'—The impact of an overcrowded
708 Emergency Department on the time to treatment and length of stay of self-presenting
709 patients with abdominal pain. *International Emergency Nursing*, 2016; 29, 9-14.
- 710 57. Morley, C., Unwin, M., Peterson, G. M., Stankovich, J., & Kinsman, L. Emergency
711 department crowding: a systematic review of causes, consequences, and solutions.
712 *PLoS ONE*, 2018; 13(8), e0203316.
- 713 58. American College of Emergency Physicians. Policy Statement. Safe Discharge from
714 the Emergency Department. Dallas, American College of Emergency
715 Physicians. 2019.
- 716 59. Lowthian, J. A., McGinnes, R. A., Brand, C. A., Barker, A. L., & Cameron, P. A.
717 Discharging older patients from the emergency department effectively: a systematic
718 review and meta-analysis. *Age and ageing*, 2015; 44(5), 761-770.
- 719 60. Dunlea, S., McCombe, G., Broughan, J., Carroll, Á., Fawsitt, R., Gallagher, J., Melin,
720 K. and Cullen, W. Priorities in integrating primary and secondary care: a multimethod
721 study of GPs. *Journal of Integrated Care*, 2022; 31(5), pp.1-14.
- 722 61. Lucke JA, Mooijaart SP, Heeren P, Singler K, McNamara R, Gilbert T, Nickel CH,
723 Castejon S, Mitchell A, Mezera V, Van der Linden L, Lim SE, Thaur A, Karamercan

724 MA, Blomaard LC, Dundar ZD, Chueng KY, Islam F, de Groot B, Conroy S. Providing
725 care for older adults in the Emergency Department: expert clinical recommendations
726 from the European Task Force on Geriatric Emergency Medicine. *European Geriatric*
727 *Medicine*, 2022; 13(2):309-317. doi: 10.1007/s41999-021-00578-1. Epub 2021 Nov 5.
728 PMID: 34738224; PMCID: PMC8568564.

729 62. Mooijaart, S. P., Carpenter, C. R., & Conroy, S. P. Geriatric emergency medicine—a
730 model for frailty friendly healthcare. *Age and Ageing*, 2022; 51(3), afab280.