perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

# Transdiagnostic alterations in white matter microstructure associated with suicidal thoughts and behaviours in the ENIGMA Suicidal Thoughts and Behaviours consortium

5 Running title: White matter microstructure and suicidality

# 7 Authors:

4

6

Laura S. van Velzen<sup>1,2</sup>, Lejla Colic<sup>3,4,5</sup>, Zuriel Ceja<sup>6,7</sup>, Maria R. Dauvermann<sup>8,9</sup>, Luca M. Villa<sup>3</sup>, 8 Hannah S. Savage<sup>2,10</sup>, Yara J. Toenders<sup>1,2,11</sup>, Niousha Dehestani<sup>10,12</sup>, Alyssa H. Zhu<sup>13</sup>, Adrian I. 9 Campos<sup>14</sup>, Lauren E. Salminen<sup>13</sup>, Ingrid Agartz<sup>15,16,17</sup>, Nina Alexander<sup>18,50</sup>, Rosa Ayesa-Arriola<sup>19</sup>, 10 Elizabeth D. Ballard<sup>20</sup>, Nerisa Banaj<sup>21</sup>, Carlotta Barkhau<sup>22</sup>, Zeynep Başgöze<sup>24</sup>, Jochen Bauer<sup>23,</sup> 11 Francesco Benedetti<sup>25</sup>, Klaus Berger<sup>26</sup>, Bianca Besteher<sup>4,5</sup>, Katharina Brosch<sup>18,50,86</sup>, Manuel Canal-Rivero<sup>27,28</sup>, Simon Cervenka<sup>17,29</sup>, Romain Colle<sup>30</sup>, Colm G. Connolly<sup>31</sup>, Emmanuelle Corruble<sup>30</sup>, Philippe Courtet<sup>32,33</sup>, Baptiste Couvy-Duchesne<sup>34,35</sup>, Benedicto Crespo-Facorro<sup>27,28,36</sup>, Kathryn R Cullen<sup>24</sup>, Udo Dannlowski<sup>22</sup>, Jeremy Deverdun<sup>37,38</sup>, Ana M. Diaz-Zuluaga<sup>87,88</sup>, Lorielle M.F. Dietze<sup>39</sup>, 12 13 14 15 Jennifer W Evans<sup>20</sup>, Negar Fani<sup>40</sup>, Kira Flinkenflügel<sup>22</sup>, Naomi P. Friedman<sup>41</sup>, Ian H. Gotlib<sup>42</sup>, Nynke A. Groenewold<sup>43</sup>, Dominik Grotegerd<sup>22</sup>, Tomas Hajek<sup>39</sup>, Alexander S. Hatoum<sup>44</sup>, Marco Hermesdorf<sup>26</sup>, Ian B. Hickie<sup>89</sup>, Yoshiyuki Hirano<sup>90,91</sup>, Tiffany C. Ho<sup>92,93</sup>, Yuki Ikemizu<sup>90,94</sup>, Frank Iorfino<sup>89</sup>, Jonathan C. Ipser<sup>43</sup>, Yuko Isobe<sup>90,91</sup>, Andrea P. Jackowski<sup>45,46</sup>, Fabrice Jollant<sup>47,48,49,30</sup>, Tilo 16 17 18 19 Kircher<sup>18,50</sup>, Melissa Klug<sup>22</sup>, Sheri-Michelle Koopowitz<sup>43</sup>, Anna Kraus<sup>22</sup>, Axel Krug<sup>18,51</sup>, Emmanuelle Le Bars<sup>37,38</sup>, Elisabeth J. Leehr<sup>22</sup>, Meng Li<sup>4,5</sup>, Elizabeth T.C. Lippard<sup>52,53</sup>, Carlos Lopez-Jaramillo<sup>88</sup>, Ivan I. Maximov<sup>15,54</sup>, Andrew M. McIntosh<sup>55</sup>, Katie A. McLaughlin<sup>56,57</sup>, Sean R. McWhinney<sup>39</sup>, 20 21 22 Susanne Meinert<sup>22,58</sup>, Elisa Melloni<sup>25</sup>, Philip B. Mitchell<sup>59</sup>, Benson Mwangi<sup>60,95</sup>, Igor Nenadić<sup>18,50</sup>, Stener Nerland<sup>15,16</sup>, Emilie Olie<sup>32,33</sup>, Victor Ortiz-García de la Foz<sup>19</sup>, Pedro M. Pan<sup>46</sup>, Fabricio Pereira<sup>61,62</sup>, Fabrizio Piras<sup>21</sup>, Federica Piras<sup>21</sup>, Sara Poletti<sup>25</sup>, Andrew E. Reineberg<sup>63</sup>, Gloria Roberts<sup>59</sup>, Rafael Romero-García<sup>9,27</sup>, Matthew D. Sacchet<sup>64</sup>, Giovanni A. Salum<sup>65,66</sup>, Anca-Larisa 23 24 25 26 Sandu<sup>67</sup>, Carl M. Sellgren<sup>17,68</sup>, Eiji Shimizu<sup>90,91,94</sup>, Harry R. Smolker<sup>69</sup>, Jair C. Soares<sup>60,95</sup>, 27 Gianfranco Spalletta<sup>21,70</sup>, J. Douglas Steele<sup>71</sup>, Frederike Stein<sup>18,50</sup>, Dan J. Stein<sup>72</sup>, Benjamin Straube<sup>18,50</sup>, Lea Teutenberg<sup>18,50</sup>, Florian Thomas-Odenthal<sup>18,50</sup>, Paula Usemann<sup>18,50</sup>, Romain Valabregue<sup>73,74</sup>, Johanna Valencia-Echeverry<sup>88</sup>, Gerd Wagner<sup>4,75</sup>, Gordon Waiter<sup>67</sup>, Martin Walter<sup>4,5,76,77</sup>, Heather C. Whalley<sup>55,78</sup>, Mon-Ju Wu<sup>60,95</sup>, Tony T. Yang<sup>79</sup>, Carlos A. Zarate<sup>20</sup>, Andre Zugman<sup>80</sup>, Giovana B. Zunta-Soares<sup>60,95</sup>, Kees van Heeringen<sup>96</sup>, Sanne J.H. van Rooij<sup>40</sup>, Nic van der Wee<sup>81,82</sup>, Steven van der Werff<sup>81,82</sup>, Paul M. Thompson<sup>13</sup>, Hilary P. Blumberg<sup>3,83,84</sup>, Anne-Laura van Harmelen<sup>9,85</sup>, Miguel E. Rentería<sup>6,7</sup>, Neda Jahanshad<sup>13</sup>, ENIGMA Suicidal Thoughts and Pabaviaura Consertium<sup>\*</sup> and Lippae Schmaol<sup>1,2</sup> 28 29 30 31 32 33 34 35 Behaviours Consortium\* and Lianne Schmaal<sup>1,2</sup>

# 36 Affiliations:

- 37 <sup>1</sup> Orygen, Parkville, VIC, Australia
- 38 <sup>2</sup> Centre for Youth Mental Health, University of Melbourne, Melbourne, VIC, Australia
- 39 <sup>3</sup> Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA
- 40 <sup>4</sup> Department of Psychiatry and Psychotherapy, Jena University Hospital, Jena, Germany
- 41 <sup>5</sup> DZPG (German Center for Mental Health), partner site Halle/Jena/Magdeburg
- 42 <sup>6</sup> QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia
- 43 <sup>7</sup> School of Biomedical Sciences, Faculty of Medicine, The University of Queensland, Brisbane, QLD,
- 44 Australia
- 45 <sup>8</sup> Institute for Mental Health, School of Psychology, University of Birmingham, Birmingham, UK
- 46 <sup>9</sup> Department of Psychiatry, University of Cambridge, Cambridge, UK
- <sup>10</sup> Melbourne Neuropsychiatry Centre, Department of Psychiatry, The University of Melbourne, Victoria,
   Australia
- 49 <sup>11</sup> Department of Psychology, Education and Child Studies, Erasmus University Rotterdam, the Netherlands
- 50 <sup>12</sup> School of Psychology, Deakin University, Victoria, Australia
- <sup>13</sup> Imaging Genetics Center, Mark and Mary Stevens Neuroimaging and Informatics Institute, Keck School of
   Medicine, University of Southern California, Marina del Rey, CA, USA
- 53 14 Rage and should not be used to guide clinical practice.
- <sup>15</sup> Norwegian Centre for Mental Disorders Research (NORMENT), Institute of Clinical Medicine, University of Oslo, Oslo, Norway

- <sup>16</sup> Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, Norway
- <sup>17</sup> Centre for Psychiatry Research, Department of Clinical Neuroscience, Karolinska Institutet, & Stockholm
   Health Care Services, Region Stockholm, Stockholm, Sweden
- <sup>18</sup> Department of Psychiatry and Psychotherapy, University of Marburg, Marburg, Germany
- 60 <sup>19</sup> Department of Psychiatry. Marqués de Valdecilla University Hospital, IDIVAL, School of Medicine,
- 61 University of Cantabria, Santander, Spain
- Experimental Therapeutics and Pathophysiology Branch, National Institute of Mental Health, National
   Institutes of Health, Bethesda MD, USA
- <sup>21</sup> Laboratory of Neuropsychiatry, Clinical Neuroscience and Neurorehabilitation Department, Santa Lucia
   Foundation IRCCS, Rome, Italy
- 66 <sup>22</sup> Institute for Translational Psychiatry, University of Münster, Münster, Germany
- 67 <sup>23</sup> Clinic for Radiology, University Hospital Münster, Münster, Germany
- <sup>24</sup> Department of Psychiatry and Behavioral Sciences, University of Minnesota Medical School, Minneapolis,
   MN, USA
- <sup>25</sup> Division of Neuroscience, Psychiatry and Psychobiology Unit, IRCCS San Raffaele Scientific Institute,
   Milan, Italy
- 72 <sup>26</sup> Institute of Epidemiology and Social Medicine, University of Münster, Münster, Germany
- 73 <sup>27</sup> Instituto de Biomedicina de Sevilla (IBiS)/HUVR/CSIC/Universidad de Sevilla. CIBERSAM (ISCIII)
- 74 <sup>28</sup> Mental Health Service, Hospital Universitario Virgen del Rocío, Sevilla, Spain
- <sup>29</sup> Department of Medical Sciences, Psychiatry, Uppsala University, Uppsala, Sweden
- 76 <sup>30</sup> MOODS Team, INSERM 1018, CESP, Université Paris-Saclay, Faculté de Médecine Paris-Saclay,
- 77 Service Hospitalo-Universitaire de Psychiatrie de Bicêtre, Assistance Publique-Hôpitaux de Paris, Hôpitaux
- 78 Universitaires Paris-Saclay, Hôpital de Bicêtre, Le Kremlin Bicêtre, F-94275, France
- 79 <sup>31</sup> Department of Biomedical Sciences, Florida State University, Tallahassee, FL
- <sup>32</sup> Department of Emergency Psychiatry and Acute Care, Lapeyronie Hospital, CHU Montpellier, Montpellier,
   France
- 82 <sup>33</sup> IGF, Univ. Montpellier, CNRS, INSERM, Montpellier, France
- 83 <sup>34</sup> Institute for Molecular Bioscience, the University of Queensland, St Lucia, QLD, Australia
- <sup>35</sup> Sorbonne University, Paris Brain Institute ICM, CNRS, Inria, Inserm, AP-HP, Hôpital de la Pitié
- 85 Salpêtrière, F-75013, Paris, France
- <sup>36</sup> Centro de Investigación Biomédica en Red de Salud Mental, Instituto de Salud Carlos III (CIBERSAM),
   Madrid, Spain
- <sup>37</sup> Institut d'Imagerie Fonctionnelle Humaine, I2FH, Department of Neuroradiology, Gui de Chauliac Hospital
   and University of Montpellier, Montpellier, France
- 90 <sup>38</sup> Department of Neuroradiology, Gui de Chauliac Hospital and University of Montpellier, Montpellier, France
- 91 <sup>39</sup> Department of Psychiatry, Dalhousie University, Halifax, NS, Canada
- 92 <sup>40</sup> Department of Psychiatry and Behavioral Sciences, Emory University, Atlanta, Georgia, USA
- <sup>41</sup> Institute for Behavioral Genetics and Department of Psychology and Neuroscience, University of Colorado
   Boulder, Boulder, CO, USA
- 95 <sup>42</sup> Department of Psychology, Stanford University, Stanford, CA 94305 USA
- 96 <sup>43</sup> Department of Psychiatry & Neuroscience Institute, University of Cape Town, Cape Town, South Africa
- 97 <sup>44</sup> Department of Psychological & Brain Sciences, Washington University in St. Louis
- 98 <sup>45</sup> Østfold University College Department of Education, ICT and Learning, Halden, Norway
- 99 <sup>46</sup> Universidade Federal de São Paulo, Brazil
- <sup>47</sup> Faculty of medicine, University Paris-Saclay & Bicetre hospital, APHP, Le Kremlin-Bicetre, France
- 101 <sup>48</sup> Department of psychiatry, CHU Nîmes, Nîmes, France
- <sup>49</sup> Department of psychiatry, McGill University, Montreal, Canada
- <sup>50</sup> Center for Mind, Brain, and Behavior (CMBB), University of Marburg, Marburg, Germany
- 104 <sup>51</sup> Department of Psychiatry, University Hospital Bonn, Bonn, Germany
- <sup>52</sup> Department of Psychiatry and Behavioral Sciences, Dell Medical School, University of Texas at Austin
- 106 <sup>53</sup> University of Texas at Austin
- 107 <sup>54</sup> Department of Health and Functioning, Western Norway University of Applied Sciences, Bergen, Norway
- <sup>55</sup> Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, United
- 109 Kingdom
- 110 <sup>56</sup> Ballmer Institute for Children's Behavioral Health, University of Oregon
- 111 <sup>57</sup> Department of Psychology, Harvard University
- <sup>58</sup> Institute for Translational Neuroscience, University of Münster, Münster, Germany
- <sup>59</sup> Discipline of Psychiatry and Mental Health, School of Clinical Medicine, University of New South Wales,

114 Sydney, Australia

- <sup>60</sup> Louis A. Faillace, MD, Department of Psychiatry and Behavioral Sciences, McGovern Medical School, The
   University of Texas Health Science Center at Houston
- <sup>61</sup> MIPA, Université de Nîmes, Nimes, France
- <sup>62</sup> Division for clinical research and innovation, University Hospital Center of Nimes, Nimes, France

- 119 <sup>63</sup> Institute for Behavioral Genetics, University of Colorado Boulder, Boulder, CO, USA
- 120 <sup>64</sup> Meditation Research Program, Department of Psychiatry, Massachusetts General Hospital, Harvard
- 121 Medical School, Boston, MA, USA
- 122 <sup>65</sup> Child Mind Institute, New York
- 123 <sup>66</sup> Universidade Federal do Rio Grande do Sul, Hospital de Clinicas de Porto Alegre - Porto Alegre, Brazil
- 124 <sup>67</sup> Aberdeen Biomedical Imaging Centre, University of Aberdeen, Aberdeen, UK
- 125 <sup>68</sup> Department of Physiology and Pharmacology, Karolinska Institutet, Stockholm, Sweden
- 126 <sup>69</sup> Institute for Cognitive Science, University of Colorado Boulder, Boulder, CO, USA
- 127 <sup>70</sup> Menninger Department of Psychiatry and Behavioral Sciences, Baylor College of Medicine, Houston, TX, 128 USA
- 129 <sup>71</sup> Division of Imaging Science and Technology, Medical School, University of Dundee, Dundee UK
- 130 <sup>72</sup> SAMRC Unit on Risk & Resilience in Mental Disorders, Department of Psychiatry & Neuroscience Institute, 131 University of Cape Town
- 132 <sup>73</sup> Institut du Cerveau et de la Moelle épinière, ICM, Inserm U 1127, CNRS UMR 7225, Sorbonne Université, 133 Paris, France
- 134 <sup>74</sup> Centre de Neuro-Imagerie de Recherche, CENIR, ICM, Paris, France
- 135 <sup>75</sup> Center for Intervention and Research on adaptive and maladaptive brain Circuits underlying mental health 136 (C-I-R-C), Jena-Magdeburg-Halle, Jena, Germany.
- 137 <sup>76</sup> Department of Psychiatry and Psychotherapy, University Tübingen, Tübingen, Germany
- 138 <sup>77</sup> Leibniz Institute for Neurobiology, Magdeburg, Germany
- 139 <sup>78</sup> Generation Scotland, Centre for Genomic and Experimental Medicine, Institute of Genetics and Molecular 140 Medicine, University of Edinburgh, Edinburgh, United Kingdom
- 141 <sup>79</sup> Department of Psychiatry and Behavioral Sciences, Division of Child and Adolescent Psychiatry, Weill
- 142 Institute for Neurosciences, University of California San Francisco, San Francisco, California, USA
- 143 <sup>80</sup> National Institute of Mental Health, National Institutes of Health, Bethesda, Maryland
- 144 <sup>81</sup> Department of Psychiatry, Leiden University Medical Center, Leiden, The Netherlands
- 145 <sup>82</sup> Leiden Institute for Brain and Cognition, Leiden University Medical Center, Leiden, The Netherlands
- 146 <sup>83</sup> Department of Radiology and Biomedical Imaging, Yale University School of Medicine, New Haven, CT, 147 USA
- 148 <sup>84</sup> Child Study Center, Yale School of Medicine, New Haven, CT, USA
- 149 <sup>85</sup> Education and Child Studies, Leiden University, Leiden, the Netherlands
- 150 <sup>86</sup> Institute of Behavioral Sciences, Feinstein Institutes for Medical Research, Manhasset, NY, USA
- 151 <sup>87</sup> Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Behavior David Geffen School 152 of Medicine, University of California Los Angeles
- 153 <sup>88</sup> Research Group in Psychiatry GIPSI, Department of Psychiatry, Faculty of Medicine, Universidad de
- 154 Antioquia, Medellin, Columbia.
- 155 <sup>89</sup> The University of Sydney, Sydney, Australia
- 156 <sup>90</sup> Research Center for Child Mental Development, Chiba University
- 157 <sup>91</sup> United Graduate School of Child Development, Osaka University, Kanazawa University, Hamamatsu 158 University, Chiba University and University of Fukui
- 159 <sup>92</sup> Department of Psychology, University of California, Los Angeles, Los Angeles, CA, USA
- 160 <sup>93</sup> Brain Research Institute, University of California, Los Angeles, Los Angeles, CA, USA
- 161 <sup>94</sup> Department of Cognitive Behavioral Physiology, Graduate School of Medicine, Chiba University
- 162 <sup>95</sup> UTHealth Houston School of Behavioral Health Sciences.
- 163 <sup>96</sup> Ghent University, Ghent, Belgium

### 164 165

#### 166 Corresponding author:

- 167 Laura S. van Velzen Name:
- 168 Affiliation: Orygen
- 169 Centre for Youth Mental Health, University of Melbourne
- 170 Address: 35 Poplar Road, Parkville, VIC 3052, Australia
- 171 Telephone: +31 6 205 28 057
- 172 Email: laurasvanvelzen@gmail.com
- 173
- 174
- 175
- 176
- 177
- Abstract 178

179

180 Previous studies have suggested that alterations in white matter (WM) microstructure are implicated 181 in suicidal thoughts and behaviours (STBs). However, findings of diffusion tensor imaging (DTI) 182 studies have been inconsistent. In this large-scale mega-analysis conducted by the ENIGMA 183 Suicidal Thoughts and Behaviours (ENIGMA-STB) consortium, we examined WM alterations 184 associated with STBs.

185 Data processing was standardised across sites, and resulting WM microstructure measures 186 (fractional anisotropy, axial diffusivity, mean diffusivity and radial diffusivity) for 25 WM tracts were pooled across 40 cohorts. We compared these measures among individuals with a psychiatric 187 188 diagnosis and lifetime history of suicide attempt (n=652; mean age=35.4±14.7; female=71.8%). 189 individuals with a psychiatric diagnosis but no STB (i.e., clinical controls; n=1871; mean 190 age=34±14.8; female=59.8%), and individuals with no mental disorder diagnosis and no STB (i.e., 191 healthy controls; n=642; mean age=29.6±13.1; female=62.9%). We also compared these measures 192 among individuals with recent suicidal ideation (n=714; mean age=36.3±15.3; female=66.1%), 193 clinical controls (n=1184; mean age=36.8±15.6; female=63.1%), and healthy controls (n=1240; 194 mean age= 31.6±15.5; female=61.0%).

195 We found subtle but statistically significant effects, such as lower fractional anisotropy associated 196 with a history of suicide attempt, over and above the effect of psychiatric diagnoses. These effects 197 were strongest in the corona radiata, thalamic radiation, fornix/stria terminalis, corpus callosum and 198 superior longitudinal fasciculus. Effect sizes were small (Cohen's d < 0.25). Recent suicidal ideation 199 was not associated with alterations in WM microstructure.

200 This large-scale coordinated mega-analysis revealed subtle regional and global alterations in WM 201 microstructure in individuals with a history of suicide attempt. Longitudinal studies are needed to 202 confirm whether these alterations are a risk factor for suicidal behaviour.

#### Introduction 203

204 Suicide is a worldwide public health concern, with more than 700,000 deaths by suicide occurring 205 worldwide annually [1]. For each death by suicide, it is estimated that there are more than 20 206 attempts, which has a cascading effect that negatively impacts families, friends, and communities 207 [1]. Despite widespread international efforts to reduce deaths by suicide, the number of deaths by 208 suicide continues to increase in several regions of the world [2]. Risk and protective factors for 209 suicidal thoughts and behaviours (STBs) have been identified [3]. Our current understanding of the 210 underlying pathophysiological mechanisms is limited. Therefore, it is crucial that we increase our 211 knowledge and understanding of these neurobiological mechanisms so that we can better target 212 prevention and intervention efforts for individuals with high suicide risk. 213

214 Many prior magnetic resonance imaging (MRI) studies have sought to identify neural correlates of 215 suicidal ideation and suicide attempt [for a review, please see 4, 5]. However, identifying robust and 216 reliable patterns of neural alterations associated with STBs has been hampered by methodological 217 heterogeneity across studies. Furthermore, due to the high clinical heterogeneity, associations with 218 brain alterations are likely subtle; hence, large samples are needed to increase statistical power and 219 identify neural correlates of STBs. The ENIGMA Suicidal Thoughts and Behaviours (ENIGMA-STB) 220 consortium was established to address these issues by pooling data across international research 221 groups to identify neural correlates of STBs. In our first study of subcortical and cortical grey matter 222 morphology and STBs in young people between 8 and 25 years of age, we identified a subtle association between lifetime history of suicide attempt and the surface area of the frontal pole, a 223 224 region in the prefrontal cortex [6], in a well-phenotyped subsample that was enriched for STBs and 225 more severe symptoms of major depressive disorder (MDD) or bipolar disorder (BD). In addition to 226 grey matter morphology, alterations in the microstructure of white matter (WM) tracts connecting 227 brain regions may also contribute to risk for STBs. Identifying these WM alterations could reveal new 228 treatment targets and increase the accuracy of monitoring or treatment response prediction.

229

230 Previous neuroimaging studies have used diffusion tensor imaging (DTI) to examine microstructural 231 alterations related to STBs [for a review, please see 7]. These studies have focused primarily on

232 fractional anisotropy (FA), which measures the coordinated directionality of water diffusion in WM 233 fibre tracts and may reflect the coherence and myelination of neuronal fibres [8, 9]. A lifetime history 234 of suicide attempt has been associated with lower FA values in various regions and WM tracts. 235 including in the prefrontal cortex (PFC) [10–13], corpus callosum [14–16], cingulum [11], internal 236 capsule [11, 17], uncinate fasciculus [16, 18], and inferior fronto-occipital fasciculus [16]. However, 237 other studies found higher FA in individuals with a history of suicide attempt [19, 20]. Fewer studies 238 have examined associations between FA measures and suicidal ideation, but findings from these 239 studies suggest that suicidal ideation is associated with lower FA in the corpus callosum [21, 22], 240 uncinate fasciculus [23], and corona radiata [21].

241

242 Thus, few studies have examined WM microstructural alterations related to suicidal ideation, and 243 findings on suicide attempt are inconsistent across studies. Prior studies are hampered by small 244 sample sizes [4], which decreases the likelihood of identifying true effects, increases the probability 245 of false-negative findings, and may cause inflation of effect sizes [24]. In addition, large samples may 246 be needed to identify subtle associations between STBs and WM microstructure. Finally, most prior 247 studies focused on FA and did not examine other WM diffusivity measures, including axial diffusivity 248 (AD), which is associated with axonal number and organisation; mean diffusivity (MD), which may 249 be an estimate of membrane density; and radial diffusivity (RD), which can provide insights into 250 myelination [25, 26]. Previous studies have also focused only on the presence or absence of suicidal 251 ideation or suicide attempt, and have not examined specific aspects of suicide-related thoughts, such 252 as the severity of suicidal ideation.

253

To address these limitations, we pooled data from 40 cohorts from the ENIGMA-STB consortium to examine associations between measures of WM microstructure (FA, AD, MD and RD) for 25 WM tracts and STBs in a large transdiagnostic sample. We examined WM microstructure alterations in people with a psychiatric diagnosis and lifetime history of suicide attempt compared to individuals with a diagnosis but no history of suicide attempt (i.e., clinical controls; CLC) and individuals with no disorder and no history of suicide attempt (healthy controls). In addition, we examined WM alterations

in people with recent suicidal ideation (within the last six months) but no history of suicide attempt,compared to CLC and HC.

262

Based on previous findings concerning suicidal behaviour, we expected that a lifetime history of suicide attempt would be associated with lower FA in WM tracts that connect the frontal lobe with limbic regions or connect different limbic regions, such as the cingulum, corpus callosum, internal capsule, and uncinate fasciculus [11, 14, 17, 18, 23].

267

In a subsample of participants for whom more in-depth assessment of STBs from the Columbia Suicide Severity Rating Scale (CSSRS) was available, we investigated associations between WM microstructure and the severity of suicidal ideation. Finally, in this sample, we were able to examine associations between WM microstructure and suicide attempt, and also distinguish among interrupted, aborted, and actual suicide attempt.

273

# 274 Patients and methods

275 <u>Cohorts</u>

276 We pooled data from 40 international cohorts from 15 countries (see Supplementary Figure 1) to 277 investigate the association between STBs and WM microstructure in a transdiagnostic sample, 278 including individuals diagnosed with major depressive disorder, obsessive-compulsive disorder, 279 bipolar disorder, post-traumatic stress disorder, psychotic disorders, generalised anxiety disorder, 280 panic disorder, or social anxiety disorder. Demographic characteristics of the different samples are 281 presented in **Table 1** and **Table 2**. The inclusion and exclusion criteria for the different studies are 282 shown in **Supplemental Table S1**. All cohorts obtained ethics approval from their local institutional 283 review boards and ethics committees. Participants who were 18 years old and over provided written 284 informed consent; those under the age of 18 years provided written informed assent in addition to 285 written informed consent from a parent/guardian at the local institution.

286

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

# 288 **Table 1. Descriptive statistics for sites included in the lifetime suicide attempt analyses**

Presented here are age (mean and standard deviation) and sex for the three groups (HC: healthy controls, CLC: clinical controls, group with lifetime suicide attempt) for the different sites included in the analysis on lifetime history of suicide attempt.

Site	Age HC	Age CLC	Age suicide attempt group	% female HC	% female CLC	% female suicide attempt group	Total N HC	Total N CLC	Total N suicide attempt group
BiDirect	-	48.6 (7.3)	49.5 (7.3)	-	56.3	66.9	0	316	124
CHU Montpellier BICS	42.2 (11.7)	45.3 (14.4)	44.2 (13.8)	71.4	86.4	90.0	21	22	20
CHU Montpellier IMPACT	-	38.3 (12.9)	38.6 (9.8)	-	94.4	85.7	0	18	7
CHU Montpellier UF8555	37.3 (7.9)	35.4 (8.4)	37.9 (9.5)	100.0	100.0	100.0	28	47	42
DCHS	30.9 (5.6)	29.7 (6.4)	32.3 (6.0)	100.0	100.0	100.0	22	20	9
MOODS team DEP-ARREST CLIN	35.5 (13.0)	33.2 (10.2)	34.8 (14.3)	67.7	70.8	63.0	31	24	27
ETPB-STB	33.9 (11.3)	34.3 (7.9)	34.8 (11.9)	65.0	50.0	63.6	20	12	11
Fondazione Santa Lucia _ Neuropsychiatr y Lab / (FSL_NPL)- OCD sample	39.3 (10.5)	39.9 (11.1)	38.1 (7.6)	35.4	34.9	36.4	82	43	11
FSL_NPL - Schizophrenia sample	-	40.7 (11.7)	43.9 (9.9)	-	32.7	44.4	0	49	18
Ghent	33.6 (13.4)	38.2 (11.7)	34.8 (11.9)	94.1	53.3	69.2	17	15	13
GIPSI BD	-	31.5 (10.3)	34.4 (11.3)	-	25.0	25.0	0	24	8
GIPSI SZ	-	39.7 (11.9)	40.0 (11.4)	-	62.2	68.0	0	45	25
Grady Trauma Project Emory University	39.4 (13.0)	39.3 (12.1)	36.1 (10.9)	97.8	100.0	100.0	46	51	14
Halifax	40.4 (13.7)	42.8 (17.8)	51.0 (16.5)	64.3	41.2	57.1	42	17	7

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Houston BD	24.1 (14.7)	29.1 (16.4)	30.2 (11.2)	43.1	48.4	64.3	51	64	14
LTS Colorado site 1	28.7 (1.1)	28.6 (0.9)	28.5 (0.5)	63.6	37.0	76.9	55	92	13
LTS Colorado site 2	28.6 (0.6)	28.7 (0.8)	28.6 (0.5)	73.3	62.5	66.7	30	72	9
McGill University	35.8 (8.0)	44.5 (11.3)	39.4 (10.6)	53.3	50.0	77.8	15	14	9
MR-IMPACT	-	15.0 (1.4)	15.2 (1.2)	-	67.1	91.4	0	70	35
PAFIP	-	29.8 (8.5)	27.3 (7.8)	-	36.4	26.7	0	77	15
San Raffaele Hospital	-	47.7 (10.9)	46.7 (10.2)	-	63.5	74.7	0	288	79
Sydney Bipolar Kids and Siblings	20.9 (3.6)	23.0 (4.8)	24.1 (4.0)	52.2	65.5	75.0	67	84	24
Sydney Brain and Mind Centre	-	19.5 (3.5)	18.1 (2.6)	-	65.1	68.8	0	169	32
UCSF Adolescent MDD	15.4 (1.4)	15.7 (1.3)	16.0 (1.1)	60.0	55.8	71.4	40	43	14
University of Minnesota Adolescent MDD	16.5 (2.2)	15.5 (1.8)	16.9 (2.1)	72.7	78.9	71.4	11	38	7
University of Texas- Austin - Bipolar Seed Program	20.5 (1.7)	21.0 (2.3)	22.4 (2.0)	68.4	73.3	57.1	19	15	7
University of Washington/ Harvard	12.0 (1.8)	12.4 (2.6)	14.4 (1.8)	46.7	35.5	60.0	45	31	5
Yale School of Medicine	-	19.3 (3.4)	19.0 (3.2)	-	60.4	73.6	0	111	53
Total	29.6 (13.1)	34.0 (14.8)	35.4 (14.7)	62.9	59.8	71.8	642	1871	652

<sup>293</sup> 294

295 296 297

Table 2. Descriptive statistics for studies included in the recent suicidal ideation analysis
 Presented here are age (mean and standard deviation) and sex for the three groups (HC: healthy

controls, CLC: clinical controls, group with recent suicidal ideation) for the different sites included in
 the analysis on suicidal ideation.

Site Age HC Age Age % % % Total Tota CLC suicidal female female female N HC N CL	Age HC Ag	e Age C suicidal	% female	% female	% female	N HC	N CLC	suicidal
---	-----------	---------------------	-------------	-------------	-------------	------	-------	----------

			ideation group	НС	CLC	suicidal ideation group			ideation group
BiDirect	-	48.7 (7.3)	48.3 (7.0)	-	57.0	65.2	0	405	46
Brazilian High Risk Cohort Porto Alegre	11.0 (2.0)	11.2 (1.9)	11.2 (1.9)	48.1	61.9	80.0	181	21	5
Chiba University	31.2 (8.4)	34.1 (6.3)	32.2 (7.8)	20.0	65.0	61.5	5	20	26
CHU Montpellier BICS	42.2 (11.7)	54.5 (10.7)	41.9 (14.3)	71.4	83.3	87.5	21	6	16
CHU Montpellier IMPACT	-	35.3 (12.7)	43.0 (12.6)	-	90.9	100.0	0	11	7
CHU Montpellier UF8555	38.6 (8.2)	36.3 (7.7)	33.9 (9.9)	100.0	100.0	100.0	10	32	14
MOODS team DEP- ARREST- CLIN	35.5 (13.0)	-	31.9 (10.4)	67.7	-	66.7	31	0	18
EPISCA (Leiden adolescents)	14.6 (1.6)	15.9 (2.2)	15.8 (1.7)	85.0	80.0	86.4	20	10	22
ETPB-STB	34.6 (11.8)	-	34.1 (9.0)	72.2	-	62.5	18	0	8
FSL_NPL- OCD sample	-	35.1 (12.3)	38.7 (7.9)	-	39.6	0.0	0	53	6
FOR2107 Marburg	38.1 (13.4)	35.9 (12.8)	35.0 (12.8)	65.4	81.5	57.9	243	27	38
FOR2107 Muenster	29.5 (11.2)	27.6 (9.8)	36.6 (12.7)	64.9	71.0	63.6	154	31	44
Grady Trauma Project Emory University	38.4 (12.4)	41.3 (11.4)	34.7 (12.7)	97.7	97.5	100.0	44	40	11
Jena-SB	-	39.8 (11.1)	37.6 (13.1)	-	53.5	81.8	0	15	22
KASP	-	26.3 (6.1)	26.1 (7.0)	-	52.2	12.5	0	23	8
McGill University	35.8 (7.9)	47.7 (9.6)	42.1 (12.6)	53.3	50.0	50.0	15	6	8
MR-IMPACT	-	15.1 (1.4)	14.8 (1.4)	-	72.5	62.1	0	40	29
Muenster	42.1	38.1	36.1	54.2	56.8	58.2	260	37	98

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

Neuroimaging Cohort	(10.9)	(12.0)	(12.4)						
San Raffaele Hospital	-	48.0 (10.0)	48.3 (10.7)	-	63.5	62.9	0	74	151
Stanford University FAA	30.6 (10.2)	35.7 (6.7)	35.5 (9.9)	100.0	100.0	100.0	18	6	8
STRADL site	-	52.5 (11.7)	54.4 (13.8)	-	65.0	71.4	0	20	14
STRADL site 2	-	56.3 (10.2)	54.9 (6.4)	-	85.0	75.0	0	20	12
Sydney Brain and Mind Centre	24.0 (4.3)	20.0 (4.3)	19.7 (4.9)	56.0	59.8	69.8	75	112	43
TIPS-Jena	47.4 (16.1)	46.8 (12.3)	42.7 (13.0)	51.5	56.0	50.0	66	25	16
UCSF Adolescent MDD	15.4 (1.4)	15.7 (1.3)	15.7 (1.4)	60.0	61.1	28.6	40	36	7
University of Minnesota Adolescent MDD	16.1 (1.9)	15.9 (1.9)	15.6 (2.0)	55.0	75.9	83.3	20	29	24
University of Texas- Austin -Bipolar Seed Program	20.5 (1.7)	21.1 (2.2)	20.8 (2.7)	68.4	70.0	80.0	19	10	5
Yale School of Medicine	-	19.2 (3.5)	18.1 (2.5)	-	57.3	75.0	0	75	8
Total	31.6 (15.5)	36.8 (15.6)	36.3 (15.3)	61.0	63.1	66.1	1240	1184	714

<sup>300</sup> 

302

# 303 Image processing

304 Scanner characteristics and acquisition parameters for all cohorts are provided in Supplemental 305 Table S2. Each site performed local preprocessing of diffusion-weighted images, including diffusion 306 tensor fitting. The pre-processed images were then processed using the ENIGMA-DTI protocol, 307 including quality control procedures. This protocol is freely available on the ENIGMA-GitHub 308 webpage (https://github.com/ENIGMA-git#enigma-dti-imaging) NITRC and 309 (http://www.nitrc.org/projects/enigma dti). For 24 tracts of interest, FA, AD, MD and RD measures 310 were extracted (please see Supplemental Table S3). We combined regions of interest (ROIs) across

<sup>301</sup> 

hemispheres by calculating the mean of the left and right hemispheres, weighted by the number of
voxels, in order to reduce the number of statistical tests. In addition, a global anisotropy or diffusivity
measure was created, leading to a total of 25 measures for FA, AD, MD, and RD.

314

The WM microstructure measures were harmonised across sites using the ComBat algorithm in R [27, 28]. An empirical Bayes approach was used to adjust for variability between scanners while preserving biological variability related to age, sex, and diagnosis. In line with our previous study [6], all DTI measures included in the analyses were ComBat-corrected. Finally, within-site outliers (measures greater than three standard deviations away from the mean of that region) were visually inspected and, if necessary, excluded from the analysis.

321

## 322 Statistical analysis

323 The main aim of this study was to examine associations among WM microstructure. lifetime history 324 of suicide attempt, and recent (in the last six months) suicidal ideation in a large transdiagnostic 325 sample. The cohorts included in this multi-study analysis administered different instruments to 326 assess recent suicidal ideation and lifetime history of attempt. We used a similar approach to our 327 previous study [6] to harmonise these measures across studies (see Supplemental Table S4). 328 Lifetime history of suicide attempt (coded yes/no) was determined using clinical interviews (e.g., the 329 Kiddie Schedule for Affective Disorders and Schizophrenia (K-SADS; [29]) or Structured Clinical 330 Interview for DSM-5 (SCID; [30])) or detailed clinical scales on STBs (e.g., the CSSRS [31] or Self-331 Injurious Thoughts and Behaviours Interview (SITBI; [32])). Recent suicidal ideation (yes/no in the 332 past six months or more recent) was assessed using items from depression severity rating scales 333 (e.g. the Hamilton Depression Rating Scale (HDRS; [33]), Beck Depression Inventory (BDI; [34, 35]), 334 detailed clinical scales on STBs (e.g., CSSRS or Beck Scale for Suicidal Ideation (SSI; [36, 37]) and 335 clinical interviews (e.g. SCID). Similar to our previous study [6], we conducted separate analyses for 336 suicidal ideation and suicide attempt to optimise the sample size for each analysis, as only 16 out of 337 the 40 cohorts had information on both suicidal ideation and suicide attempt. In total, we included 338 data from 28 cohorts in the analysis on lifetime suicide attempt and from 28 cohorts in the analysis 339 on recent suicidal ideation.

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

340

341 We assessed differences among groups in global and regional anisotropy or diffusivity measures using multiple linear regression models in R [38]. We included a group variable to compare 342 343 individuals with a lifetime history of suicide attempt (N=652) to either CLC (N=1871) or HC (N=642) 344 (two group comparisons). A group variable was also included to compare individuals with recent 345 suicidal ideation (N=714) to CLC (N=1184) and HC (N=1240) in separate analyses. In supplementary 346 analyses, we also compared the HC and CLC groups (please see Supplemental note 1 for a 347 description of the methods and results). Consistent with previous ENIGMA-DTI analyses [39], the 348 analyses were corrected for age, sex, and their linear and non-linear interactions (age-by-sex interaction, age<sup>2</sup> and age<sup>2</sup>-by-sex interaction). Effect sizes were assessed using the Cohen's d349 metric. All p-values were corrected for multiple testing (for the 25 tracts per anisotropic/diffusivity 350 351 measure) using the Benjamini-Hochberg correction in R, resulting in FDR<.05.

352

Finally, in supplementary analyses, we examined interactions between group and type of lifetime psychiatric diagnosis in a subsample of participants, as data on lifetime psychiatric diagnosis were not available for all participants (please see Supplemental **note 2** for a description of the methods and results).

357

# 358 Analysis in the CSSRS sample

We also examined associations between WM microstructure and more detailed phenotypes of suicide attempt and suicidal ideation in a subsample of 7 cohorts that used the CSSRS to assess STBs. The CSSRS was specifically developed to assess the intensity and severity of suicidal thoughts and suicidal behaviour [31]. C-SSRS has good validity, high sensitivity and specificity for suicide attempts [40].

364

For analyses of suicidal ideation, we examined how the nominal measures of recent or lifetime severity of suicidal ideation (coded 0-5; 0: no ideation; 1: passive ideation; 2: non-specific active ideation; 3: active ideation with a method, but no plan or intent; 4: active ideation with intent, but no

plan; 5: active ideation with a plan and intent) were associated with WM microstructure. For theseanalyses, the standardised beta was calculated as an effect size.

370

371 In line with our previous study, for analyses of suicide attempt, we compared anisotropy/diffusivity 372 measures between individuals with a lifetime history of any attempt (actual, aborted, or interrupted 373 attempt) and individuals with no lifetime history of any attempt [6]. In addition, we compared these 374 measures between individuals with a lifetime history of an actual (non-interrupted and non-aborted) 375 attempt and those without any lifetime suicide attempt. Finally, these measures were compared 376 between individuals with a history of suicidal ideation (but no history of an actual suicide attempt), 377 and those with a lifetime history of an actual attempt. Cohen's d metric was calculated as an effect 378 size for these analyses.

379

Similar to the main analyses, all analyses in the CSSRS sample included age, sex, age-by-sex, age<sup>2</sup>, and age<sup>2</sup>-by-sex as covariates, and all resulting p-values were corrected for multiple testing (for the testing (for the 25 tracts per anisotropic/diffusivity measure) using the Benjamini-Hochberg correction in R to lead to FDR<.05.

384

# 385 **Results**

## 386 Lifetime suicide attempt

## 387 Individuals with a lifetime history of suicide attempt versus HC

388 Global FA and regional FA were lower in the attempt group (N=652) compared to HC (N=642) in the 389 following tracts: anterior limb of the internal capsule (ALIC), body of the corpus callosum (BCC), 390 corpus callosum (CC), cingulate gyrus of the cingulum bundle (CGC), corona radiata (CR), external 391 capsule (EC), fornix/stria terminalis (FXST), genu of the corpus callosum (GCC), inferior fronto-392 occipital fasciculus (IFO), posterior corona radiata (PCR), posterior thalamic radiation (PTR), 393 retrolenticular part of the internal capsule (RLIC), splenium of the corpus callosum (SCC), superior 394 corona radiata (SCR), superior longitudinal fasciculus (SLF), sagittal stratum (SS), and uncinate 395 fasciculus (UNC) (Cohen's d range: 0.125-0.226; Figure 1.A and Supplemental table S6). In 396 addition, MD was higher in the UNC in the attempt group compared to HC (Cohen's d=0.220;

Supplemental table S7). Finally, RD was higher in the BCC, CC, CR, FXST, PCR, SCC, SLF, and
UNC in the attempt group compared to HC (Cohen's *d* range: 0.148-0.170; Figure 1.A and
Supplemental table S8). There were no significant differences in AD between the attempt group and
HC (Supplemental table S5).

401



402	
-----	--

403	Figure 1.
404	Cohen's <i>d</i> map represents significant ROIs without the global covariate and threshold of pFDR < 0.05. From left to right,
405	middle slices of sagittal, axial, and coronal views were used, except for C, where only the sagittal view was used.
406	The figure includes three sections: A) FA and RD for individuals with a lifetime history of suicide attempt versus HC, B)
407	FA for individuals with a lifetime history of suicide attempt versus CLC and C) FA and RD for individuals with recent
408	suicidal ideation versus HC.
409	Abbreviations: CLC= clinical control group ; FA= fractional anisotropy; HC= healthy control group; pFDRR= p value
410	corrected using Benjamini Hochberg correction; RD= radial diffusivity; ROI= region of interest; SA= individuals with a
411	lifetime history of suicide attempt; SI= individuals with recent suicidal ideation.
412	
413	Individuals with a lifetime history of suicide attempt versus CLC

414 Global and regional AD, MD, and RD did not differ between individuals with a lifetime history of

415 suicide attempt (*N*=652) and CLC (*N*=1871) (**Figure 1.B**, Supplemental **tables S9**, **S11**, and **S12**).

- 416 Regional FA was lower in the CR, FXST, PCR, PTR, SCC and SLF in the suicide attempt group
- 417 compared to CLC (Cohen's *d* range: 0.123-0.138; Supplemental **table S10**).
- 418

# 419 Recent suicidal ideation

- 420 Individuals with recent suicidal ideation versus HC
- 421 Global and regional AD and MD did not differ between the suicidal ideation group (*N*=714) and HC
- 422 (*N*=1240) (Supplemental **tables S13 and S15**). Global FA and regional FA in the CR, GCC and
- 423 superior fronto-occipital fasciculus (SFO) were lower in the suicidal ideation group compared to HC
- 424 (Cohen's *d* range: 0.133-0.158; Figure 1.C, Supplemental table S14). Global RD and regional RD
- 425 in the SCR were higher in the suicidal ideation group compared to HC (Cohen's *d*: 0.142-0.150;
- 426 Supplemental **table S16**).
- 427

# 428 Individuals with recent suicidal ideation versus CLC

There were no significant differences between the suicidal ideation group (*N*=714) and CLC
(*N*=1184) (Supplemental **tables S17-20**).

431

# 432 Deeper phenotyping in the C-SSRS sample

433 There were no significant associations of regional FA, AD, MD, RD, with severity of lifetime suicidal 434 ideation (N=299; Supplemental tables S21-24) or severity of recent suicidal ideation (N=338; 435 Supplemental tables S25-28). In addition, there were no significant differences in WM microstructure 436 between people with a lifetime history of an actual suicide attempt (N=134) and those without a 437 history of any suicide attempt (no interrupted, aborted, or actual suicide attempt; N=244) (Supplemental tables 29-32). Moreover, there were no significant differences between people with 438 439 an actual suicide attempt (N=134) and those with lifetime suicidal ideation, but without a history of 440 an actual suicide attempt (N=122; Supplemental tables 33-36). Finally, there were no differences in FA or diffusivity measures between people with a lifetime history of an interrupted, aborted, or actual 441 442 suicide attempt (N=177) compared to individuals without a lifetime history of any subtype of suicide attempt (N=244; Supplemental tables S37-40). 443

# 445 **Discussion**

In this large-scale multi-cohort transdiagnostic analysis, we examined associations between global and regional measures of WM microstructure and STBs in a pooled sample from 40 international cohorts. A lifetime history of suicide attempt was associated with subtle differences in global and regional FA and diffusivity (higher regional MD and RD), and recent suicidal ideation was associated with lower global and regional FA and higher global and regional RD.

451

452 Individuals with a lifetime history of suicide attempts had lower regional FA than did clinical controls. This analysis was undertaken to ascertain that the observed differences are not merely attributable 453 454 to microstructural variances associated with psychiatric disorders, but rather, are significantly 455 influenced by lifetime suicide attempt. The WM tracts that showed this difference between individuals 456 with a lifetime history of suicide attempt and CLC included the corona radiata (including the posterior 457 corona radiata), part of the fornix (fornix/stria terminalis), thalamic radiation (specifically the posterior 458 thalamic radiation), splenium of the corpus callosum, and the superior longitudinal fasciculus. The 459 corona radiata and the thalamic radiation connect the cortex with subcortical brain structures, 460 including the brainstem and thalamus. The corona radiata is part of the thalamic-cortical circuitry and 461 has been associated with perceptual, motor, emotional and cognitive function, including behavioural 462 regulation, and lower FA in the corona radiata has been associated with poorer executive functioning 463 [41–44]. In addition, previous single-cohort studies have reported that lower FA in the corona radiata 464 is associated with suicidal behaviour [16, 45], albeit inconsistently [20]. Moreover, lower FA in the 465 corona radiata predicted suicidal behaviour on average two years later [46]. The (posterior) thalamic 466 radiations also connect the thalamus with cortical regions, including the parietal and occipital lobes. These are important for cortical arousal and consciousness but may also be associated with 467 468 cognitive control [47]. While lower FA in this tract has not been associated with suicidal behaviour 469 before, FA in this tract was lower in individuals with BD [48], who are at higher risk of suicidal 470 behaviour [49]. The splenium of the corpus callosum connects the occipital-parietal and temporal 471 cortex from both hemispheres and has been associated with visuospatial functioning, reading, 472 language processing, and consciousness [50]. Lower FA in the splenium of the corpus callosum has 473 been associated with a higher number of suicide attempts in BD and MDD [14]. The fornix (including

474 the fornix/stria terminalis) connects the amygdala to the hypothalamus and plays an important role 475 in threat monitoring, regulation of the hypothalamic-pituitary-adrenal (HPA) axis and behavioural 476 inhibition [51]. In addition, previous investigators have speculated that impaired structural 477 connectivity in this tract may contribute to feelings of anhedonia [52], which may drive suicidality [53]. 478 Finally, the superior longitudinal fasciculus is an association tract that connects the parietal and 479 temporal cortex with the frontal cortex and may play a role in speech, spatial awareness, processing 480 speed, and attention [54]. FA in this region has inconsistently been associated with suicidality (see 481 [20, 55] and has been consistently linked to brooding and rumination [56], which are risk factors for 482 suicidal behaviour [57].

483

In this large sample, we were unable to replicate previous findings of lower FA in other WM tracts, including the cingulum [11], internal capsule [11, 17], uncinate fasciculus [16, 18], and inferior frontooccipital fasciculus [16]. This lack of replication may be related to the small sample size of these previous studies, related to sample characteristics, or be specific to certain psychiatric disorders.

488

489 We also found differences in global and regional RD between individuals with a history of suicide 490 attempt and HC but not CLC, suggesting that these findings were driven by psychiatric illness and 491 not by suicidal behaviour per se. Similarly, differences in FA and RD were identified between 492 individuals with recent suicidal ideation and HC but not CLC, which may also be related to mental 493 illness in general and not to suicidal ideation specifically. Consistent with this finding, similar effects 494 for FA were observed comparing CLC to HC; however, these findings were weaker and most did not 495 survive correction for multiple testing (see Supplemental **note 1**). We hypothesise that mental illness 496 may be more severe in the suicidal ideation group than in the CLC group, which may explain why 497 differences in FA were more pronounced.

498

Our findings provide evidence consistent with previous ENIGMA reports about differences in WM microstructure in major depressive disorder [39], bipolar disorder [58] and schizophrenia [59] with DTI data, where associations with global and regional FA were reported. The MDD study included 1305 individuals diagnosed with MDD and 1602 HC; the results showed lower global FA and regional

503 FA in 16 out of 25 WM tracts and higher RD in adult individuals with recurrent MDD; the largest FA 504 differences were observed in the corpus callosum and *corona radiata* [39]. On the other hand, the 505 schizophrenia study included 1963 individuals diagnosed with schizophrenia and 2359 HC, and the 506 results showed lower global FA and regional FA in 20 out of 25 ROIs; the larger differences were 507 reported in anterior *corona radiata* and corpus callosum [59]. Finally, the study on bipolar disorder 508 included 1482 individuals diagnosed with BD and 1551 HC and showed lower global FA and regional 509 FA in 29 out of 43 tracts in patients, most prominently in the corpus callosum and cingulum [58].

510

511 In a subsample of 7 cohorts, which had more deeply phenotyped suicidal thoughts and behaviours 512 using the CSSRS, we were able to examine the associations with more detailed phenotypes of 513 suicide attempt and suicidal ideation. In our previous study [6] on grey matter morphology and STB 514 in young people, lower surface area of the frontal pole was associated with a history of actual (non-515 interrupted or non-aborted) suicide attempt in a subsample of studies that had used the CSSRS to 516 assess suicidal behaviour (4 out of 7 cohort overlap, 53% total sample overlap with the CSSRS 517 sample in the current study). In this study, we did not find any associations between WM anisotropy or diffusivity measures and severity of suicidal ideation, nor did we find differences in these measures 518 519 related to suicide attempt in general (actual, interrupted or aborted suicide attempt), or actual suicide 520 attempt specifically (non-interrupted or non-aborted suicide attempts). It is possible that this sample 521 was not large enough to identify subtle associations between suicide attempt and WM 522 microstructure.

523

524 This study increases our understanding of the mechanisms underlying suicidal behaviour by 525 confirming findings from previous smaller studies that lower FA is implicated in suicidal behaviour 526 and showing which tracts are involved. However, the effect sizes for significant group differences 527 observed in this study were small (all d<0.25), especially for differences between the suicidal 528 ideation/suicide attempt groups and the CLC. High clinical heterogeneity may mask larger effects in 529 individual persons, which we were not able to detect with the group average approach used in the 530 current study. Nonetheless, the currently observed differences are too subtle to be clinically useful 531 in terms of prediction of risk for suicidal behaviour at the individual level. In addition, it is unclear

whether alterations in WM microstructure represent a risk factor for suicidal behaviour or are a consequence of a previous suicide attempt. Previous longitudinal studies have shown that lower regional FA at baseline predicts suicidal behaviour at follow-up in individuals with mood disorders [11, 46]. Further, FA has been found to be highly heritable [60], suggesting that alterations in WM microstructure may indeed represent a risk factor for suicidal behaviour. However, cellular brain damage following hypoxia/anoxia during a suicide attempt may also cause WM damage and affect WM microstructure [61].

539

540 A strength of this study is the large sample size, which allowed the examination of more detailed 541 phenotypes in subsamples. A second strength is the use of harmonised protocols for diffusion image 542 processing and quality control. However, we should also acknowledge several limitations of this 543 study, including heterogeneity across samples in how STBs were assessed. We have tried to 544 minimise this effect by using a detailed process to harmonise these measures across studies (in line 545 with the approach used in our previous study). Finally, we could not control for severity of psychiatric 546 symptoms, as this was a transdiagnostic analysis and the different cohorts had used different severity rating scales; therefore, we cannot rule out the possibility that higher symptom severity in 547 548 the suicide attempt group affected our findings. The scope of this study was limited to cross-sectional 549 findings. However, this presents an opportunity for future research to delve into longitudinal 550 analyses, shedding light on the evolution of the differences reported.

551

To conclude, in this large-scale multi-cohort study we found an association between a history of suicide attempt and regional FA, above and beyond the effect of psychiatric diagnosis. FA in several tracts that have been associated with (among others) behavioural regulation/inhibition, executive functioning, and brooding/rumination show a subtle association with suicidal attempt. Future longitudinal studies are needed to examine if altered FA may represent a risk factor for suicidal behaviour, for instance by contributing to lower resilience to severely stressful life events.

- 558
- 559
- 560

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

## 561 Acknowledgments

562 This work was supported by the MQ Brighter Futures Award MQBFC/2 (LS, LC, MD, LvV, ALvH, HB) and the 563 National Institute of Mental Health of the National Institutes of Health under Award Number R01MH117601 564 (LS, LvV, NJ). ALvH was funded by a MQ Brighter Futures Award MQBFC/2, a Royal Society Dorothy 565 Hodgkin Fellowship (DH15017), an MRC MRF emerging leaders award, the Leiden University Social 566 Resilience and Security fund, and an NWO VIDI award. HPB was supported by the MQ Brighter Futures 567 Award MQBFC/2, the National Center for Advancing Translational Science (Grant Number: UL1TR000142), 568 the National Institute of Mental Health (Grant Numbers: RC1MH088366, R01MH69747), Brain and Behavior 569 Research Foundation, International Bipolar Disorders Foundation, American Foundation for Suicide 570 Prevention SRG-1-10-119, and the John and Hope Furth Endowment. EDB is supported by the National 571 Institute of Mental Health Intramural Research Program (ZIA MH002857). NB is supported by the Italian 572 Ministry of Health, grant Ricerca Corrente RC 23. The University of Minnesota adolescent MDD study is 573 supported by the National Institute of Mental Health (K23MH090421), the National Alliance for Research on 574 Schizophrenia and Depression, the University of Minnesota Graduate School, the Minnesota Medical 575 Foundation, and the Biotechnology Research Center (P41 RR008079 to the Center for Magnetic Resonance 576 Research), University of Minnesota, and the Deborah E. Powell Center for Women's Health Seed Grant, 577 University of Minnesota. FB is supported by the Italian Ministry of Health, grant PNRR-MAD-2022-12375859. 578 BiDirect was funded by grants from the German Federal Ministry of Education and Research (BMBF; Grants 579 FKZ-01ER0816 and FKZ-01ER1506) to KB. BB and LC are supported by the Interdisciplinary Center of 580 Clinical Research of the Medical Faculty Jena. M.C.-R. acknowledges funding support from the Consejería 581 de Salud y Familias (Junta de Andalucía) 2020 grant, which covers his salary (RH-0081-2020). Funding for 582 the DEPARRESTCLIN cohort was provided by a national grant (ANR SAMENTA 2012) of the Agence 583 Nationale de la Recherche (ANR). BCD is funded by a NHMRC CJ Martin fellowship (1161356). BCF 584 acknowledges this work was supported by the Instituto de Salud Carlos III (PI14/00639 and PI14/00918) and 585 Fundación Instituto de Investigación Margués de Valdecilla (NCT0235832 and NCT02534363). UD 586 acknowledges this work was funded by the German Research Foundation (DFG), Udo Dannlowski (co-587 speaker FOR2107, DA 1151/5-1, DA 1151/5-2, grant DA1151/9-1, DA1151/10-1 and DA1151/11-1) and the 588 Interdisciplinary Center for Clinical Research (IZKF) of the medical faculty of Münster (grant Dan3/022/22 to 589 UD). MuensterNeuroimagingCohort: This work was funded by the German Research Foundation (SFB-590 TRR58, Project C09 to UD). JWE is supported by the National Institute of Mental Health Intramural Research 591 Program (ZIA MH002857). NPF acknowledges the LTS Colorado data acquisition was supported by grants 592 from the National Institutes of Health (NIH) in the United States (AG046938 and MH063207). NPF was 593 supported by NIH grants AG046938, , MH117131, MH124846, DA042742, DA046413, DA046064, and 594 DA051018. IHG is supported by the National Institute of Mental Health Grant R37MH101495. TH 595 acknowledges this study was supported by funding from the Canadian Institutes of Health Research (103703 596 , 106469 and 142255, 180449, 186254), Nova Scotia Health Research Foundation, Dalhousie Clinical 597 Research Scholarship to T. Hajek, Brain & Behavior Research Foundation (formerly NARSAD); 2007 Young 598 Investigator and 2015 Independent Investigator Awards to T. Hajek. This work was also supported by the 599 AMED Brain/MINDS Beyond Program (JP18dm0307002) and JSPS KAKENHI (JP19K03309 and 600 JP22H01090). TCH is supported in part by the National Institutes of Health (K01MH117442, R21MH130817, 601 R01MH127176). YI was supported by JSPS KAKENHI (JP23K07004). FI was supported by an NHMRC EL1 602 Investigator Grant (GNT2018157) and the Bill and Patricia Richie Foundation Fellowship. YI was supported 603 by JSPS KAKENHI (JP23K02956). FJ acknowledges this study was supported by an operating grant from 604 the Canadian Institutes for Health Research (CIHR #119288). TK acknowledges this work was funded by the 605 German Research Foundation (DFG grants FOR2107 KI588/14-1, and KI588/14-2, and KI588/20-1, 606 KI588/22-1 to Tilo Kircher, Marburg, Germany). Biosamples and corresponding data were sampled, 607 processed and stored in the Marburg Biobank CBBMR. This work was funded by the German Research 608 Foundation (DFG grants KR 3822/5-1, KR 3822/7-2 to Axel Krug). ETCL s funded in part by the American 609 Foundation for Suicide Prevention SRG-0-112-20. CLJ is supported by PRISMA U.T. AMM is supported by 610 the Wellcome Trust Grants (220857/Z/20/Z, 226770/Z/22/Z, 104036/Z/14/Z, 216767/Z/19/Z), and the 611 European Union Horizon 2020 Grant (Grant Agreement 847776). KAM acknowledges this study was funded 612 by the National Institute of Mental Health (R01-MH103291). PBM acknowledges this study was funded by 613 the Australian National Medical and Health Research Council (Program Grant 1037196 (PBM and MB); 614 Investigator Grant 1177991 (PBM)), the Lansdowne Foundation, Good Talk, and the Keith Pettigrew Family

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

615 Bequest (PBM). IN work is funded by the Deutsche Forschungsgemeinschaft (DFG) grants NE2254/1-2, 616 NE2254/2-1, NE2254/3-1, NE2254/4-1. PMP acknowledges BHRCS was supported with grants from the 617 National Institute of Development Psychiatric for Children and Adolescent (INPD) and the Grant Fapesp 2014/50917-0 - 2021/05332-8 CNPq 465550/2014-2. FP acknowledges support for the team in Nimes, 618 619 CINES grants access to HPC facilities (A0100311413). FaP is supported by the Italian Ministry of Health, 620 grant RF-2019-12370182. FeP is supported by the Italian Ministry of Health, grant Ricerca Corrente RC 23. 621 SP is supported by the Italian Ministry of Health, grant PNRR-MAD-2022-12375716, RF-2018-12367249. GR 622 acknowledges this study was funded by the Australian National Medical and Health Research Council 623 (Program Grant 1037196 (PBM and MB); Investigator Grant 1177991 (PBM)), the Lansdowne Foundation, 624 Good Talk, and the Keith Pettigrew Family Beguest (PBM). RRG was supported by EMERGIA Junta de 625 Andalucía program (EMERGIA20 00139), the VII Plan Propio of the University of Seville, the Plan de 626 Consolidación (CNS2023-143647) and the Proyectos de Generación de Conocimiento (PID2021-122853OA-627 100) from the Spanish Ministry of Science and Innovation. MDS is supported by the National Institute of 628 Mental Health (Project Number R01MH125850); Brain and Behavior Research Foundation (Grant Number 629 28972). HRS was supported by a grant from the National Institutes of Health in the United States 630 (DA051018). JCS acknowledges this work was partially supported by NIMH (1R01MH085667-01A1), John S. 631 Dunn Foundation (Houston, Texas), and Pat Rutherford Chair in Psychiatry (UTHealth Houston). GS is 632 supported by the Italian Ministry of Health, grant Ricerca Corrente RC 23. JDS is supported by the grants 633 MRC (MR/S010351/1, MR/W002388/1, MR/W002566/1) and EPSRC (EP/Y017544/1). The DCHS cohort is 634 funded by the Bill & Melinda Gates Foundation [INV-006732]. DJS is funded by the South African Medical 635 Research Council. BS acknowledges this work was funded by the German Research Foundation (DFG grant 636 as part of FOR2107: STR1146/18-1 to Benjamin Straube, Marburg, Germany). MW is supported by German 637 Center for Mental Health (DZPG); FKZ: 01EE2103; and NeuroMarKet: Neuroimaging and Blood Markers as 638 Indicators of Ketamine Efficacy in Treatment Resistant Depression; BMBF-EU-EraNet-Neuron, FKZ: 639 01EW2010A. HCW acknowledges this work is supported by the Chief Scientist Office of the Scottish 640 Government Health Directorates [CZD/16/6], Scottish Funding Council [HR03006], Wellcome Trust 641 [216767/Z/19/Z] & Wellcome Trust (Wellcome Trust Strategic Award "STratifying Resilience and Depression 642 Longitudinally" (STRADL) Reference 104036/Z/14/Z). CAZ is supported by the National Institute of Mental 643 Health Intramural Research Program (ZIA MH002857). TTY is supported by the National Center for 644 Complementary and Integrative Health (NCCIH) R21AT009173, R61AT009864, and R33AT009864 to TTY; 645 by the National Center for Advancing Translational Sciences (CTSI), National Institutes of Health, through 646 UCSF-CTSI UL1TR001872 to TTY; by the American Foundation for Suicide Prevention (AFSP) SRG-1-141-647 18 to TTY; by UCSF Weill Institute for Neurosciences to TTY; by UCSF Research Evaluation and Allocation 648 Committee (REAC) and J. Jacobson Fund to TTY; by the National Institute of Mental Health (NIMH) 649 R01MH085734 and the Brain and Behavior Research Foundation (formerly NARSAD) to TTY. PMT is 650 supported by NIH grants R01AG058854, R01MH116147 and R01MH129742. NJ is supported by 651 R01MH134004. MER receives Fellowship support from the Rebecca L Cooper Medical Research 652 Foundation (F20231230).

#### 654 **Conflicts of interest**

653

655 ALvH receives consultancy fees from the Augeo foundation. CAZ is a full-time US government employee. 656 He is listed as a co-inventor on a patent for the use of ketamine and its metabolites in major depression and 657 suicidal ideation. Dr. Zarate has assigned his patent rights to the U.S. government but will share a 658 percentage of any royalties that may be received by the government. HPB has consulted to the Milken 659 Institute. MW is a member of the following advisory boards and gave presentations to the following 660 companies: Bayer AG, Germany; Boehringer Ingelheim, Germany; and Biologische Heilmittel Heel GmbH, 661 Germany. MW has further conducted studies with institutional research support from HEEL and Janssen 662 Pharmaceutical Research for a clinical trial (IIT) on ketamine in patients with MDD, unrelated to this 663 investigation. MW did not receive any financial compensation from the companies mentioned above. IA 664 received speakers honorarium from Lundbeck. IBH is the Co-Director, Health and Policy at the Brain and 665 Mind Centre (BMC) University of Sydney. The BMC operates an early-intervention youth services at 666 Camperdown under contract to headspace. He is the Chief Scientific Advisor to, and a 3.2% equity 667 shareholder in, InnoWell Pty Ltd which aims to transform mental health services through the use of 668 innovative technologies. AC reports being currently an employee of the Regeneron Genetics Center, and 669 may own Regeneron stock or stock options. JCS acknowledge to be related with the next companies

#### perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

- 670 ALKERMES (Advisory Board), BOEHRINGER Ingelheim (Consultant), COMPASS Pathways (Research
- 671 Grant), JOHNSON & JOHNSON (Consultant), LIVANOVA (Consultant), RELMADA (Research Grant),
- 672 SUNOVION (Research Grant), Mind Med (Research Grant). PMP received payment or honoraria for lectures
- and presentations in educational events for Sandoz, Daiichi Sankyo, Eurofarma, Abbot, Libbs, Instituto
- 674 Israelita de Pesquisa e Ensino Albert Einstein, Instituto D'Or de Pesquisa e Ensino. All other authors report
- no biomedical financial interests, disclosures or potential conflicts of interest.

- ~~~

perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license .

#### References 702

- 703 WHO. Suicide worldwide in 2019: global health estimates. 2021. 2021. 1.
- 704 2. Martínez-Alés G, Jiang T, Keyes KM, Gradus JL. The Recent Rise of Suicide Mortality in the 705 United States. Annu Rev Public Health. 2022;43:99–116.
- 706 3. Wasserman D, Carli V, Iosue M, Javed A, Herrman H. Suicide prevention in childhood and 707 adolescence: a narrative review of current knowledge on risk and protective factors and 708 effectiveness of interventions. Asia Pac Psychiatry. 2021;13:e12452.
- 709 4. Schmaal L, van Harmelen A-L, Chatzi V, Lippard ETC, Toenders YJ, Averill LA, et al. Imaging suicidal thoughts and behaviors: a comprehensive review of 2 decades of neuroimaging 710 711 studies. Mol Psychiatry. 2020;25:408-427.
- 712 Auerbach RP, Pagliaccio D, Allison GO, Algueza KL, Alonso MF. Neural Correlates 5. 713 Associated With Suicide and Nonsuicidal Self-injury in Youth. Biol Psychiatry. 2021;89:119-714 133.
- 715 6. van Velzen LS, Dauvermann MR, Colic L, Villa LM, Savage HS, Toenders YJ, et al. Structural 716 brain alterations associated with suicidal thoughts and behaviors in young people: results from 717 21 international studies from the ENIGMA Suicidal Thoughts and Behaviours consortium. Mol 718 Psychiatry. 2022:1–11.
- Zanghì E, Corallo F, Lo Buono V. Diffusion tensor imaging studies on subjects with suicidal 719 7. 720 thoughts and behaviors: A descriptive literature review. Brain Behav. 2022;12:e2711.
- Podwalski P, Szczygieł K, Tyburski E, Sagan L, Misiak B, Samochowiec J. Magnetic 721 8. 722 resonance diffusion tensor imaging in psychiatry: a narrative review of its potential role in 723 diagnosis. Pharmacol Rep. 2021;73:43-56.
- 724 Jones DK, Knösche TR, Turner R. White matter integrity, fiber count, and other fallacies: the 9. 725 do's and don'ts of diffusion MRI. Neuroimage. 2013;73:239-254.
- 726 10. Olvet DM, Peruzzo D, Thapa-Chhetry B, Sublette ME, Sullivan GM, Oquendo MA, et al. A 727 diffusion tensor imaging study of suicide attempters. J Psychiatr Res. 2014;51:60-67.
- 728 11. Lippard ETC, Cox Lippard ET, Johnston JAY, Spencer L, Quatrano S, Fan S, et al. 729 Preliminary examination of gray and white matter structure and longitudinal structural changes 730 in frontal systems associated with future suicide attempts in adolescents and young adults 731 with mood disorders. Journal of Affective Disorders. 2019;245:1139–1148.
- 732 12. Mahon K, Burdick KE, Wu J, Ardekani BA, Szeszko PR. Relationship between suicidality and 733 impulsivity in bipolar I disorder: a diffusion tensor imaging study. Bipolar Disord. 2012;14:80-734 89.
- 735 13. Gosnell SN, Molfese DL, Salas R. Brain Morphometry: Suicide. In: Spalletta G, Piras F, Gili T, 736 editors. Brain Morphometry, New York, NY: Springer New York; 2018. p. 403-427.
- 737 14. Cyprien F, de Champfleur NM, Deverdun J, Olié E, Le Bars E, Bonafé A, et al. Corpus callosum integrity is affected by mood disorders and also by the suicide attempt history: A 738 739 diffusion tensor imaging study. J Affect Disord. 2016;206:115–124.
- 740 15. Lischke A, Domin M, Freyberger HJ, Grabe HJ, Mentel R, Bernheim D, et al. Structural 741 Alterations in the Corpus Callosum Are Associated with Suicidal Behavior in Women with 742 Borderline Personality Disorder. Front Hum Neurosci. 2017;11:196.
- 743 16. Wei S, Womer FY, Edmiston EK, Zhang R, Jiang X, Wu F, et al. Structural alterations 744 associated with suicide attempts in major depressive disorder and bipolar disorder: A diffusion 745 tensor imaging study. Prog Neuropsychopharmacol Biol Psychiatry. 2020;98:109827.
- 746 17. Jia Z, Huang X, Wu Q, Zhang T, Lui S, Zhang J, et al. High-field magnetic resonance imaging 747 of suicidality in patients with major depressive disorder. Am J Psychiatry. 2010;167:1381-748 1390.
- 18. Johnston JAY, Wang F, Liu J, Blond BN, Wallace A, Liu J, et al. Multimodal Neuroimaging of 749 750 Frontolimbic Structure and Function Associated With Suicide Attempts in Adolescents and 751 Young Adults With Bipolar Disorder. Am J Psychiatry. 2017;174:667–675.
- 752 19. Kim B, Oh J, Kim M-K, Lee S, Tae WS, Kim CM, et al. White matter alterations are associated 753 with suicide attempt in patients with panic disorder. J Affect Disord. 2015;175:139–146.
- 754 20. Lee S-J, Kim B, Oh D, Kim M-K, Kim K-H, Bang SY, et al. White matter alterations associated 755 with suicide in patients with schizophrenia or schizophreniform disorder. Psychiatry Res 756 Neuroimaging. 2016;248:23-29.
- 757 21. Reis JV, Vieira R, Portugal-Nunes C, Coelho A, Magalhães R, Moreira P, et al. Suicidal

- 758Ideation Is Associated With Reduced Functional Connectivity and White Matter Integrity in759Drug-Naïve Patients With Major Depression. Front Psychiatry. 2022;13:838111.
- Zhang R, Jiang X, Chang M, Wei S, Tang Y, Wang F. White matter abnormalities of corpus
  callosum in patients with bipolar disorder and suicidal ideation. Ann Gen Psychiatry.
  2019;18:20.
- Fan S, Lippard ETC, Sankar A, Wallace A, Johnston JAY, Wang F, et al. Gray and white
  matter differences in adolescents and young adults with prior suicide attempts across bipolar
  and major depressive disorders. J Affect Disord. 2019;245:1089–1097.
- Provide the second stress of the second st
- 25. Winklewski PJ, Sabisz A, Naumczyk P, Jodzio K, Szurowska E, Szarmach A. Understanding
  the Physiopathology Behind Axial and Radial Diffusivity Changes—What Do We Know? Front
  Neurol. 2018;9:92.
- Tae WS, Ham BJ, Pyun SB, Kang SH, Kim BJ. Current Clinical Applications of Diffusion Tensor Imaging in Neurological Disorders. J Clin Neurol. 2018;14:129–140.
- Fortin J-P, Cullen N, Sheline YI, Taylor WD, Aselcioglu I, Cook PA, et al. Harmonization of
   cortical thickness measurements across scanners and sites. Neuroimage. 2018;167:104–120.
- Radua J, Vieta E, Shinohara R, Kochunov P, Quidé Y, Green MJ, et al. Increased power by
  harmonizing structural MRI site differences with the ComBat batch adjustment method in
  ENIGMA. Neuroimage. 2020;218:116956.
- Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, et al. Schedule for Affective
  Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADSPL): Initial Reliability and Validity Data. Journal of the American Academy of Child &
  Adolescent Psychiatry. 1997;36:980–988.
- First MB. Structured clinical interview for DSM-IV Axis I disorders SCID-I: Clinician version,
   scoresheet. American Psychiatric Press; 1997.
- 785 31. Posner K, Brent D, Lucas C, Gould M, Stanley B, Brown G, et al. Columbia-suicide severity
   786 rating scale (C-SSRS). New York, NY: Columbia University Medical Center. 2008. 2008.
- Nock MK, Holmberg EB, Photos VI, Michel BD. Self-Injurious Thoughts and Behaviors
  Interview: development, reliability, and validity in an adolescent sample. Psychol Assess.
  2007;19:309–317.
- 33. Hamilton M. A rating scale for depression. J Neurol Neurosurg Psychiatry. 1960;23:56–62.
- 34. Beck AT, Ward C, Mendelson M, Mock J, Erbaugh J. Beck depression inventory (BDI). Arch
  Gen Psychiatry. 1961;4:561–571.
- 35. Beck AT, Steer RA, Brown GK, Others. Beck depression inventory-II. San Antonio.
  1996;78:490–498.
- 36. Beck AT, Kovacs M, Weissman A. Assessment of suicidal intention: the Scale for Suicide
   Ideation. J Consult Clin Psychol. 1979;47:343–352.
- 797 37. Beck AT, Steer RA, Ranieri WF. Scale for suicide ideation: Psychometric properties of a self 798 report version. J Clin Psychol. 1988;44:499–505.
- R core team. R: A language and environment for statistical computing. R Foundation for
   Statistical Computing, Vienna, Austria. Http://www R-Project Org/. 2013. 2013.
- 39. van Velzen LS, Kelly S, Isaev D, Aleman A, Aftanas LI, Bauer J, et al. White matter
  disturbances in major depressive disorder: a coordinated analysis across 20 international
  cohorts in the ENIGMA MDD working group. Mol Psychiatry. 2020;25:1511–1525.
- 40. Posner K, Brown GK, Stanley B, Brent DA, Yershova KV, Oquendo MA, et al. The ColumbiaSuicide Severity Rating Scale: initial validity and internal consistency findings from three
  multisite studies with adolescents and adults. Am J Psychiatry. 2011;168:1266–1277.
- 41. Jenkins LM, Barba A, Campbell M, Lamar M, Shankman SA, Leow AD, et al. Shared white
   matter alterations across emotional disorders: A voxel-based meta-analysis of fractional
   anisotropy. Neuroimage Clin. 2016;12:1022–1034.
- 42. Chen Z, Cui L, Li M, Jiang L, Deng W, Ma X, et al. Voxel based morphometric and diffusion
  tensor imaging analysis in male bipolar patients with first-episode mania. Prog
  Neuropsychopharmacol Biol Psychiatry. 2012;36:231–238.
- 43. Leunissen I, Coxon JP, Caeyenberghs K, Michiels K, Sunaert S, Swinnen SP. Task switching
   in traumatic brain injury relates to cortico-subcortical integrity. Hum Brain Mapp.

815		2014;35:2459–2469.
816	44.	Wallace EJ, Mathias JL, Ward L. The relationship between diffusion tensor imaging findings
817		and cognitive outcomes following adult traumatic brain injury: A meta-analysis. Neurosci
818		Biobehav Rev. 2018;92:93–103.
819	45.	Jiang X, Guo Y, Jia L, Zhu Y, Sun Q, Kong L, et al. Altered Levels of Plasma Inflammatory

45. Jang X, Guo Y, Ja L, Zhu Y, Sun Q, Kong L, et al. Altered Levels of Plasma inhammatory
 Cytokines and White Matter Integrity in Bipolar Disorder Patients With Suicide Attempts. Front
 Psychiatry. 2022;13:861881.

- 46. Colic L, Villa LM, Dauvermann MR, van Velzen LS, Sankar A, Goldman DA, et al. Brain grey
  and white matter structural associations with future suicidal ideation and behaviors in
  adolescent and young adult females with mood disorders. JCPP Adv. 2022;2.
- 47. Chaddock-Heyman L, Erickson KI, Voss MW, Powers JP, Knecht AM, Pontifex MB, et al.
  White matter microstructure is associated with cognitive control in children. Biol Psychol.
  2013;94:109–115.
- 48. Hu R, Stavish C, Leibenluft E, Linke JO. White Matter Microstructure in Individuals With and At
  Risk for Bipolar Disorder: Evidence for an Endophenotype From a Voxel-Based Meta-analysis.
  Biol Psychiatry Cogn Neurosci Neuroimaging. 2020;5:1104–1113.
- 49. Dome P, Rihmer Z, Gonda X. Suicide Risk in Bipolar Disorder: A Brief Review. Medicina .
  2019;55.
- Blaauw J, Meiners LC. The splenium of the corpus callosum: embryology, anatomy, function
   and imaging with pathophysiological hypothesis. Neuroradiology. 2020;62:563–585.
- 51. Clauss J. Extending the neurocircuitry of behavioural inhibition: a role for the bed nucleus of
   the stria terminalis in risk for anxiety disorders. Gen Psychiatr. 2019;32:e100137.
- 52. Harnett NG, Stevens JS, van Rooij SJH, Ely TD, Michopoulos V, Hudak L, et al. Multimodal
  structural neuroimaging markers of risk and recovery from posttrauma anhedonia: A
  prospective investigation. Depress Anxiety. 2021;38:79–88.
- 840 53. Bonanni L, Gualtieri F, Lester D, Falcone G, Nardella A, Fiorillo A, et al. Can Anhedonia Be
  841 Considered a Suicide Risk Factor? A Review of the Literature. Medicina. 2019;55:458.
- 54. Janelle F, Iorio-Morin C, D'amour S, Fortin D. Superior Longitudinal Fasciculus: A Review of
   the Anatomical Descriptions With Functional Correlates. Front Neurol. 2022;13:794618.
- 55. Davey DK, Jurick SM, Crocker LD, Hoffman SN, Sanderson-Cimino M, Tate DF, et al. White
  matter integrity, suicidal ideation, and cognitive dysfunction in combat-exposed Iraq and
  Afghanistan Veterans. Psychiatry Res Neuroimaging. 2021;317:111389.
- 56. Pisner DA, Shumake J, Beevers CG, Schnyer DM. The superior longitudinal fasciculus and its
   functional triple-network mechanisms in brooding. Neuroimage Clin. 2019;24:101935.
- 57. Horwitz AG, Czyz EK, Berona J, King CA. Rumination, Brooding, and Reflection: Prospective
  Associations with Suicide Ideation and Suicide Attempts. Suicide Life Threat Behav.
  2019;49:1085–1093.
- 58. Favre P, Pauling M, Stout J, Hozer F, Sarrazin S, Abé C, et al. Widespread white matter
  microstructural abnormalities in bipolar disorder: evidence from mega- and meta-analyses
  across 3033 individuals. Neuropsychopharmacology. 2019;44:2285–2293.
- 59. Kelly S, Jahanshad N, Zalesky A, Kochunov P, Agartz I, Alloza C, et al. Widespread white
  matter microstructural differences in schizophrenia across 4322 individuals: results from the
  ENIGMA Schizophrenia DTI Working Group. Mol Psychiatry. 2018;23:1261–1269.
- Kochunov P, Jahanshad N, Marcus D, Winkler A, Sprooten E, Nichols TE, et al. Heritability of
  fractional anisotropy in human white matter: a comparison of Human Connectome Project and
  ENIGMA-DTI data. Neuroimage. 2015;111:300–311.
- 61. DiPoce J, Guelfguat M, DiPoce J. Radiologic findings in cases of attempted suicide and other self-injurious behavior. Radiographics. 2012;32:2005–2024.
- 863 62. Correia MM, Carpenter TA, Williams GB. Looking for the optimal DTI acquisition scheme given
  864 a maximum scan time: are more b-values a waste of time? Magn Reson Imaging.
  865 2009;27:163–175.
- 866
- 867
- 868
- 869
- 870

#### 871 **ENIGMA Suicidal Thoughts and Behaviours Consortium**

872

Laura S. van Velzen<sup>1,2</sup>, Lejla Colic<sup>3,4,5</sup>, Zuriel Ceja<sup>6,7</sup>, Maria R. Dauvermann<sup>8,9</sup>, Luca M. Villa<sup>3</sup>, 873 Hannah S. Savage<sup>2,10</sup>, Yara J. Toenders<sup>1,2,11</sup>, Niousha Dehestani<sup>10,12</sup>, Alyssa H. Zhu<sup>13</sup>, Adrian I. 874 Campos<sup>14</sup>, Lauren E. Salminen<sup>13</sup>, Ingrid Agartz<sup>15,16,17</sup>, Nina Alexander<sup>18,50</sup>, Rosa Ayesa-Arriola<sup>19</sup>, 875 Elizabeth D. Ballard<sup>20</sup>, Nerisa Banaj<sup>21</sup>, Carlotta Barkhau<sup>22</sup>, Zeynep Başgöze<sup>24</sup>, Jochen Bauer<sup>23</sup>, 876 Francesco Benedetti<sup>25</sup>, Klaus Berger<sup>26</sup>, Bianca Besteher<sup>4,5</sup>, Katharina Brosch<sup>18,50,86</sup>, Manuel Canal-877 Rivero<sup>27,28</sup>, Simon Cervenka<sup>17,29</sup>, Romain Colle<sup>30</sup>, Colm G. Connolly<sup>31</sup>, Emmanuelle Corruble<sup>30</sup>, 878 Philippe Courtet<sup>32,33</sup>, Baptiste Couvy-Duchesne<sup>34,35</sup>, Benedicto Crespo-Facorro<sup>27,28,36</sup>, Kathryn R 879 Cullen<sup>24</sup>, Udo Dannlowski<sup>22</sup>, Jeremy Deverdun<sup>37,38</sup>, Ana M. Diaz-Zuluaga<sup>87,88</sup>, Lorielle M.F. Dietze<sup>39</sup>, 880 Jennifer W Evans<sup>20</sup>, Negar Fani<sup>40</sup>, Kira Flinkenflügel<sup>22</sup>, Naomi P. Friedman<sup>41</sup>, Ian H. Gotlib<sup>42</sup>, Nynke 881 A. Groenewold<sup>43</sup>, Dominik Grotegerd<sup>22</sup>, Tomas Hajek<sup>39</sup>, Alexander S. Hatoum<sup>44</sup>, Marco 882 Hermesdorf<sup>26</sup>, Ian B. Hickie<sup>89</sup>, Yoshiyuki Hirano<sup>90,91</sup>, Tiffany C. Ho<sup>92,93</sup>, Yuki Ikemizu<sup>90,94</sup>, Frank 883 lorfino<sup>89</sup>, Jonathan C. Ipser<sup>43</sup>, Yuko Isobe<sup>90,91</sup>, Andrea P. Jackowski<sup>45,46</sup>, Fabrice Jollant<sup>47,48,49,30</sup>, Tilo 884 Kircher<sup>18,50</sup>, Melissa Klug<sup>22</sup>, Sheri-Michelle Koopowitz<sup>43</sup>, Anna Kraus<sup>22</sup>, Axel Krug<sup>18,51</sup>, Emmanuelle 885 Le Bars<sup>37,38</sup>, Elisabeth J. Leehr<sup>22</sup>, Meng Li<sup>4,5</sup>, Elizabeth T.C. Lippard<sup>52,53</sup>, Carlos Lopez-Jaramillo<sup>88</sup>, 886 Ivan I. Maximov<sup>15,54</sup>, Andrew M. McIntosh<sup>55</sup>, Katie A. McLaughlin<sup>56,57</sup>, Sean R. McWhinney<sup>39</sup>, 887 Susanne Meinert<sup>22,58</sup>, Elisa Melloni<sup>25</sup>, Philip B. Mitchell<sup>59</sup>, Benson Mwangi<sup>60,95</sup>, Igor Nenadić<sup>18,50</sup>. 888 Stener Nerland<sup>15,16</sup>, Emilie Olie<sup>32,33</sup>, Victor Ortiz-García de la Foz<sup>19</sup>, Pedro M. Pan<sup>46</sup>, Fabricio 889 Pereira<sup>61,62</sup>, Fabrizio Piras<sup>21</sup>, Federica Piras<sup>21</sup>, Sara Poletti<sup>25</sup>, Andrew E. Reineberg<sup>63</sup>, Gloria 890 Roberts<sup>59</sup>, Rafael Romero-García<sup>9,27</sup>, Matthew D. Sacchet<sup>64</sup>, Giovanni A. Salum<sup>65,66</sup>, Anca-Larisa 891 Sandu<sup>67</sup>, Carl M. Sellgren<sup>17,68</sup>, Eiji Shimizu<sup>90,91,94</sup>, Harry R. Smolker<sup>69</sup>, Jair C. Soares<sup>60,95</sup>, 892 Gianfranco Spalletta<sup>21,70</sup>, J. Douglas Steele<sup>71</sup>, Frederike Stein<sup>18,50</sup>, Dan J. Stein<sup>72</sup>, Benjamin 893 Straube<sup>18,50</sup>, Lea Teutenberg<sup>18,50</sup>, Florian Thomas-Odenthal<sup>18,50</sup>, Paula Usemann<sup>18,50</sup>, Romain 894 Valabregue<sup>73,74</sup>, Johanna Valencia-Echeverry<sup>88</sup>, Gerd Wagner<sup>4,75</sup>, Gordon Waiter<sup>67</sup>, Martin 895 Walter<sup>4,5,76,77</sup>, Heather C. Whalley<sup>55,78</sup>, Mon-Ju Wu<sup>60,95</sup>, Tony T. Yang<sup>79</sup>, Carlos A. Zarate<sup>20</sup>, Andre 896 Zugman<sup>80</sup>, Giovana B. Zunta-Soares<sup>60,95</sup>, Kees van Heeringen<sup>96</sup>, Sanne J.H. van Rooij<sup>40</sup>, Nic van 897 der Wee<sup>81,82</sup>, Steven van der Werff<sup>81,82</sup>, Paul M. Thompson<sup>13</sup>, Hilary P. Blumberg<sup>3,83,84</sup>, Anne-Laura 898 van Harmelen<sup>9,85</sup>, Miguel E. Rentería<sup>6,7</sup>, Neda Jahanshad<sup>13</sup>, Lianne Schmaal<sup>1,2</sup> 899