medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

perpetuity. It is made available under a CC-BY-ND 4.0 International license .

#### What interventions are effective and cost-effective for supporting the health and well-being of people with obesity on healthcare waiting lists? A Rapid Review

**Authors:** Toby Ayres<sup>1</sup>, Jordan Everitt<sup>1</sup>, Alesha Wale<sup>1</sup>, Chukwudi Okolie<sup>1</sup>, Amy Fox-McNally<sup>1</sup>, Helen Morgan<sup>1</sup>, Hannah Shaw<sup>1</sup>, Jacob Davies<sup>2</sup>, Rhiannon Tudor Edwards<sup>2</sup>, Adrian Edwards<sup>3</sup>, Alison Copper<sup>3</sup>, Ruth Lewis<sup>4,5</sup>

1 Public Health Wales Evidence Service, Cardiff, United Kingdom

2 Centre for Health Economics and Medicines Evaluation, Bangor University, United Kingdom

3 Health and Care Research Wales Evidence Centre, Cardiff University, United Kingdom

4 Health and Care Research Wales Evidence Centre, Bangor University, United Kingdom

5 Bangor Institute for Medical and Health Research, Bangor University, United Kingdom

#### Abstract:

Currently, there is a significant demand for tier 3 weight management services, with individuals waiting between three and five years to access these services in parts of Wales. This rapid review aimed to identify and synthesise the evidence for the effectiveness of strategies for supporting the health and well-being of individuals with obesity on such waiting lists, with a focus on practical and resource-efficient interventions that can be implemented within current healthcare constraints.

Seven studies were included, and these were published between 2017 and 2024. Studies were conducted in a range of countries, and no relevant UK based study was identified. Studies investigated exercise, physical activity counselling, education and text message-based prehabilitation interventions in people awaiting surgery.

This review did not identify any studies assessing the effectiveness of interventions that could be feasibly implemented or scaled up within the resource limitations of typical tier 3 weight management services in Wales. Most studies required significant resource and input from healthcare professionals, and were delivered in-person at healthcare settings or remotely via teleconferencing. All studies assessed patients with obesity on a waiting list for surgery, but none included a patient population that matched those on waiting lists for tier 3 weight management services in Wales. None of the studies evaluated the cost-effectiveness of interventions.

Overall, we are not confident in the evidence. Most studies were of low quality, with significant methodological and reporting limitations affecting the reliability of their findings. Although we have little confidence in the evidence, there is some evidence from four studies, that suggest exercise interventions may support the quality of life and anthropometric measures of people with obesity waiting for surgery. This evidence could be cautiously considered to inform interventions in practice, but those designing interventions, including text message-based prehabilitation interventions, preoperative educational interventions and physical activity counselling interventions were reliant on findings from single low-quality studies. Some of these interventions showed improvements for participant's quality of life, mental well-being and anthropometric measures.

In relation to obesity weight management services, allocation of resources should allow for conducting and evaluating robust studies and economic evaluations investigating interventions for those awaiting obesity weight management services. Given the current healthcare resource constraints, it may be beneficial to consider the feasibility and scalability of interventions during their design.

**Funding statement:** The authors and their Institutions were funded for this work by the Health and Care Research Wales Evidence Centre, itself funded by Health and Care Research Wales on behalf of Welsh Government.

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.



What interventions are effective and costeffective for supporting the health and well-being of people with obesity on healthcare waiting lists? A Rapid Review

November 2024









### **Report Contributors**

*Review Team* Toby Ayres, Jordan Everitt, Alesha Wale, Chukwudi Okolie, Amy Fox-McNally, Helen Morgan, Hannah Shaw (Public Health Wales Evidence Service)

*Economic Considerations* Rhiannon Tudor Edwards and Jacob Davies (Centre for Health Economics & Medicines Evaluation, Bangor University)

*Methodological Advice* Ruth Lewis (Health and Care Research Wales Evidence Centre, Bangor University)

*Evidence Centre Team* Adrian Edwards, Ruth Lewis, Alison Copper, Micaela Gal and Elizabeth Doe involved in Stakeholder engagement, review of report and editing

*Public Partners* Praveena Pemmasani and Libby Humphris

*Stakeholders* Enzo Battista-Dowds, Maria Cole and Megan Elliot

Evidence need submitted to the Evidence Centre: December 2023

Initial Stakeholder Consultation Meeting: 26 June 2024

Final Report issued: November 2024

**The review should be cited as:** Health and Care Research Wales Evidence Centre. What interventions are effective and cost-effective for supporting the health and well-being of people with obesity on healthcare waiting lists? A Rapid Review (RR0033). November 2024

**Disclaimer:** The views expressed in this publication are those of the authors, not necessarily Health and Care Research Wales. The Health and Care Research Wales Evidence Centre and authors of this work declare that they have no conflict of interest

medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

perpetuity. It is made available under a CC-BY-ND 4.0 International license .

# What interventions are effective and cost-effective for supporting the health and well-being of people with obesity on healthcare waiting lists? A Rapid Review

Report Number RR0033 (November 2024)

# EXECUTIVE SUMMARY

#### What is a Rapid Review?

Our rapid reviews (RR) use a variation of the systematic review approach, abbreviating or omitting some components to generate the evidence to inform stakeholders promptly whilst maintaining attention to bias.

### Who is this Rapid Review for?

The question was proposed by the Cwm Taf Morgannwg University Health Board Weight Management Service.

#### **Background / Aim of Rapid Review**

Currently, there is a significant demand for tier 3 weight management services, with individuals waiting between three and five years to access these services in parts of Wales. This rapid review aimed to identify and synthesise the evidence for the effectiveness of scalable strategies for supporting the health and well-being of individuals on waiting lists, with a focus on practical and resource-efficient interventions that can be implemented within current healthcare constraints.

### **Results of the Rapid Review**

Recency of the evidence base

 The literature searches were conducted in July 2024, and the included studies were published between 2017 and 2024.

#### Extent of the evidence base

- Seven studies were included in the rapid review: two randomised controlled trials (RCTs), a
  non-randomised controlled trial, a before and after study (that also used the usual care
  comparator group from a previous study), and three uncontrolled before and after studies.
- Studies were conducted in a range of countries, including Canada (n=2), Turkey (n=2), Australia (n=1), Germany (n=1), and Spain (n=1).
- Studies investigated exercise, physical activity counselling, education and text message-based prehabilitation interventions in people awaiting surgery.

#### Key findings and certainty of the evidence

- This rapid review did not identify any studies assessing the effectiveness of interventions that could be feasibly implemented or scaled up within the resource limitations of typical tier 3 weight management services in Wales. Most studies required significant resource and input from healthcare professionals, and were delivered in-person at healthcare settings or remotely via teleconferencing.
- All studies assessed patients with obesity on a waiting list for surgery, none included a patient
  population with obesity that matched those on waiting lists for tier 3 weight management
  services.
- Overall, we are not confident in the evidence as most studies were of low methodological quality, most intervention types were only assessed by a single study, most studies included small numbers of participants and three interventions assessed the feasibility of implementing an intervention rather than its effectiveness.

medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

#### perpetuity. It is made available under a CC-BY-ND 4.0 International license .

- Exercise interventions: Findings from four studies (two RCTs, one non-randomised controlled trial and one before and after study (with comparator groups from a previous study)) showed some evidence for improving quality of life and anthropometric measures, and very little evidence supporting improvements in functional outcomes (i.e. fatigue and walking capacity). Study designs, interventions and outcome measurement tools differed greatly, which limits our overall confidence in the findings.
- Text message-based prehabilitation interventions, preoperative educational interventions and physical activity counselling interventions were reliant on findings from single low-quality studies. Some of these interventions showed improvements for participant's quality of life, mental well-being and anthropometric measures.
- We found no economic evaluations to help us establish the cost-effectiveness of interventions.

#### **Policy and Practice Implications**

- Although we have little confidence in the evidence, there is more evidence of better quality, including two RCTs, that suggest exercise interventions may support the quality of life of people with obesity waiting for surgery. This evidence could be cautiously considered to inform interventions in practice but those designing interventions should be mindful of the population and setting in which they are applied.
- In relation to obesity weight management services, allocation of resources should allow for conducting and evaluating robust studies and economic evaluations investigating interventions for those awaiting obesity weight management services. Given the current healthcare resource constraints, it may be beneficial to consider the feasibility and scalability of interventions during their design.

#### **Research Implications and Evidence Gaps**

- No UK-based studies were identified, making it unclear whether the findings are applicable to UK healthcare settings.
- No studies specifically focused on brief, low-resource interventions suitable for scalable implementation within current weight management service constraints.
- Most studies assessed patients awaiting bariatric surgery, not those on waiting lists for tier 3 weight management services.
- The evidence base for interventions effects on mental well-being is very limited
- None of the included studies reported participant engagement outcomes suitable for inclusion in this review.
- None of the included studies evaluated the cost-effectiveness of interventions.
- Most studies were of low quality, with significant methodological and reporting limitations affecting the reliability of their findings.

#### **Economic considerations**

- There is a lack economic evidence on the impact of implementing interventions for supporting the health and well-being of people with obesity on healthcare waiting lists.
- Obesity is a considerable and prevalent public health issue in Wales that incurs significant economic cost. Obesity costs NHS Wales £73 million per annum. If rates of overweight and obesity continue to rise in line with recent trends, by 2050, they will cost NHS Wales £465 million per year, with a cost to the wider society and economy of £2.4 billion.

The certainty of the evidence has been assessed using the GRADE (Grading of Recommendation, Assessment, Development and Evaluation) approach (<u>https://www.gradeworkinggroup.org/</u>)

# **TABLE OF CONTENTS**

| TABLE (  | OF CONTENTS  | 6  |
|----------|--|----|
| Abbrevia | tions  | 8  |
| 1. BAG   | CKGROUND   | 9  |
| 1.1      | Who is this review for?  | 9  |
| 1.2      | Background and purpose of this review                                    | 9  |
| 2. RES   | SULTS  | 10 |
| 2.1      | Overview of the Evidence Base  | 10 |
| 2.2      | Effectiveness of exercise interventions                                  | 13 |
| Qua      | ality of life outcomes   | 14 |
| Ant      | hropometric outcomes   | 14 |
| Fun      | ctional outcomes   | 15 |
| Bot      | tom line results for exercise interventions                              | 15 |
| 2.3      | Effectiveness of text message-based prehabilitation interventions        | 15 |
| Qua      | ality of life outcomes   | 16 |
| Ant      | hropometric outcomes   | 16 |
| Bot      | tom line results for text message-based prehabilitation interventions    | 16 |
| 2.4      | Effectiveness of group-based preoperative educational interventions      | 16 |
| Mer      | ntal well-being outcomes   | 16 |
| Qua      | ality of life outcomes   | 16 |
| Ant      | hropometric outcomes   | 17 |
| Bot      | tom line results for group-based preoperative educational interventions. | 17 |
| 2.5      | Effectiveness of physical activity counselling interventions             | 17 |
| Mer      | ntal well-being and quality of life outcomes                             | 17 |
| Bot      | tom line results for physical activity counselling interventions         | 17 |
| 3. DIS   | CUSSION  | 20 |
| 3.1      | Summary of the findings  | 20 |
| 3.2      | Strengths and limitations of the available evidence                      | 20 |
| 3.3      | Strengths and limitations of this Rapid Review                           | 21 |
| 3.4      | Implications for policy and practice                                     | 21 |
| 3.5      | Implications for future research   | 22 |
| 3.6      | Economic considerations*   | 23 |
| 4. RE    | FERENCES   | 24 |
| 5. RA    | PID REVIEW METHODS   | 26 |
| 5.1      | Eligibility criteria   | 26 |
| 5.2      | Literature search  | 26 |
| 5.3      | Reference management   | 27 |
| 5.4      | Study selection process  | 27 |
| 5.5      | Data extraction  | 27 |

| 5.6    | Study design classification        | 27 |
|--------|------------------------------------|----|
| 5.7    | Quality appraisal                  | 27 |
| 5.8    | Synthesis                          | 27 |
| 5.9    | Assessment of body of evidence     |    |
| 6. EVI | DENCE                              | 28 |
| 6.1    | Search results and study selection | 28 |
| 6.2    | Data extraction                    |    |
| 6.3    | Quality appraisal                  |    |
| 6.4    | Information available on request   | 41 |
| 7. ADI | DITIONAL INFORMATION               | 42 |
| 7.1    | Conflicts of interest              | 42 |
| 7.2    | Acknowledgements                   | 42 |
| 8. APF | PENDIX                             | 43 |
|        |                                    |    |

### **LIST OF FIGURES**

| Figure 1 | . PRSIMA flow diagram |  | 9 |
|----------|-----------------------|--|---|
|----------|-----------------------|--|---|

# **LIST OF TABLES**

| Table 1: Summary of interventions                                   | 12 |
|---|----|
| Table 2: Summary of the evidence                                    | 18 |
| Table 3: Eligibility Criteria                                       | 26 |
| Table 4: Summary of included studies                                | 30 |
| Table 5: Quality appraisal results for quasi-experimental studies   | 39 |
| Table 6: Quality appraisal results for randomised controlled trials | 40 |

### **Abbreviations**

| Acronym        | Full Description   |
|----------------|--|
| BMI            | Body Mass Index  |
| CI             | Confidence interval  |
| COVID-19       | Coronavirus disease  |
| CSEP           | core stabilisation exercise program                                    |
| EG             | Exercise group   |
| ES             | Effect size  |
| EQ-5D-3L       | EuroQol-5 dimension 3-level questionnaire                              |
| EWL            | Excess weight loss   |
| FACIT-F        | Functional Assessment of Chronic Illness Therapy – Fatigue             |
| FSS            | Fatigue Severity Scale   |
| GAD-7          | Generalized Anxiety Disorder Questionnaire - 7                         |
| GRADE          | Grading of Recommendations, Assessment, Development, and Evaluations   |
| HTE            | Health technology evaluation   |
| ICTRP          | International Clinical Trials Registry Platform                        |
| IWQOL-Lite     | Impact of Weight on Quality of Life-Lite                               |
| JBI            | Johanna Briggs Institute   |
| Kg             | Kilograms  |
| MD             | Mean difference  |
| MWT            | Minute walk test   |
| MVPA           | moderate-to-vigorous physical activity                                 |
| NICE           | National Institute for Health and Care Excellence                      |
| OECD           | Organisation for Economic Co-operation and Development                 |
| OSQOL          | Obesity Specific Quality of Life                                       |
| PAC            | physical activity counselling  |
| PHQ-9          | Patient Health Questionnaire - 9                                       |
| PICO           | Population, intervention, comparator, outcome                          |
| PMOABS         | people with morbid obesity awaiting bariatric surgery                  |
| PRISMA         | Preferred Reporting Items for Systematic reviews and Meta-<br>Analyses |
| PSADBE         | Pre-surgical aerobic dance-based exercise                              |
| PSQ-20         | Perceived Stress Questionnaire - 20                                    |
| PreSET         | Pre-Surgical Exercise Training intervention                            |
| QoL            | Quality of Life  |
| RCT            | Randomised controlled trial  |
| SD             | Standard deviation   |
| SF-12          | 12-Item Short Form Survey  |
| SF-36          | 36-Item Short Form Survey  |
| TelePreSET     | Telehealth Pre-Surgical Exercise Training intervention                 |
| TELE-BariACTIV | TELEhealth BARIatric behavioural intervention                          |
| UK             | United Kingdom   |
| WRQOL          | weight-related quality of life   |

# 1. BACKGROUND

#### 1.1 Who is this review for?

This rapid review was conducted as part of the Health and Care Research Wales Evidence Centre Work Programme. The question was proposed by Cwm Taf Morgannwg University Health Board Weight Management Service to address the need for feasible, low-resource interventions that could support individuals with obesity on waiting lists for tier 3 services. helping to prevent deterioration in health and well-being during prolonged waiting periods.

#### 1.2 Background and purpose of this review

Prevalence of obesity (body mass index [BMI] of ≥30kg/m<sup>2</sup>) has risen rapidly in recent decades and is considered one of the most significant public health challenges worldwide (Watkins, 2024). In Wales, latest data estimates that 25% of adults aged 16 and over are living with obesity (Watkins, 2024). By 2035, it is projected that 11% of the Welsh population will have severe obesity (BMI ≥40kg/m<sup>2</sup>), compared to 8% in England, and 5% in Scotland (Keaver et al, 2020). Currently the UK utilises a four-tier pathway for the management of services to support adults with overweight (BMI 25 - 29.9kg/m<sup>2</sup>) and obesity to lose weight. In Wales specifically, the All-Wales Weight Management Pathway sets out the key elements and principles underpinning the delivery of weight management services. Tier 1 includes brief advice and self-directed materials for overweight adults, Tier 2 includes multicomponent interventions for adults with obesity, Tier 3 includes specialist multi-disciplinary services for adults with severe obesity, and Tier 4 involve specialist surgical services (Welsh Government, 2021).

Obesity prevalence within the Cwm Taf Morgannwg University Health Board area (29% of adults) is higher than the national average (StatsWales, 2023) however, a weight management service operating at tiers 1 to 4 has only been available in the area since February 2023 (Life Sciences Hwb, 2023). Currently, there is a significant demand for tier 3 services, with individuals waiting between three and five years to access these services across parts of Wales (Wales NHS Confederation, 2024). During these prolonged waits, individuals with severe obesity face considerable risk of health deterioration. Due to the high demand on the service leading to extended wait times, and the lack of resources to increase capacity, there is an urgent need for effective interim support strategies for people with severe obesity on waiting lists for tier 3 weight management services in order to prevent their health and well-being deteriorating and promote 'waiting well'.

Stakeholders highlighted the need for scalable strategies that fit within current resource constraints to support people awaiting tier 3 services to 'wait well' and prevent deterioration in health and well-being during this period. Preliminary scoping work conducted for this rapid review identified little secondary or primary research specifically focused on evaluating interventions to support people on waiting lists to attend tier 3 multi-disciplinary services. Consequently, the focus of this rapid review has been broadened to identify primary research on interventions that could potentially support adults with obesity on any healthcare waiting list.

# 2. RESULTS

#### 2.1 Overview of the Evidence Base

The methods and eligibility criteria for the review are presented in Section 5. Eligibility was restricted to comparative primary studies and economic evaluations assessing the effectiveness or cost-effectiveness of interventions aimed at supporting adults with obesity on any waiting list (see section 5.1 for full eligibility criteria). Seven studies were eligible for inclusion. All studies investigated interventions designed to support patients with obesity on surgical waiting lists. None of the studies addressed the stakeholders initial research question which was to support people waiting for tier 3 obesity weight management services.

Two studies were randomised controlled trials (RCTs) while the remaining five studies included two uncontrolled before and after studies, one pilot uncontrolled before and after study, one before and after study that also utilised the comparator usual care group from a previous study, and one non-randomised controlled trial. No economic evaluations were identified. Studies were conducted in a range of countries, including Canada (n=2), Turkey (n=2), Australia (n=1), Germany (n=1), Spain (n=1), and were published between 2017 and 2024. Sample sizes across the studies ranged from six to 305, with six of the seven studies reporting a sample size less than 35. Intervention duration ranged from six-weeks to sixmonths. Six studies included people with severe obesity (BMI  $\ge$  40 kg/m<sup>2</sup> or  $\ge$  35 kg/m<sup>2</sup> with comorbidities) who were on a waiting list for bariatric surgery (n=5) or either bariatric or metabolic surgery (n=1). The final study included people with obesity (BMI  $\ge$  30 kg/m<sup>2</sup>) on waiting lists for any elective surgery.

Four studies evaluated the effectiveness of interventions, while three focused on assessing their feasibility of implementation. All participants were recruited from surgical waiting lists, either through referral by surgical teams (n=3) or direct contact from researchers (n=1), and three studies did not specify their recruitment methods. The interventions aimed to prepare participants for surgery by targeting improvements in quality of life, physiological, anthropometric, behavioural, functional, or mental health outcomes. In four studies, participants in the intervention group received an intervention in combination with usual care, and outcomes were compared to a control group who also received the usual care component. In these three studies, the usual care component was physical activity counselling (n=2) or nutritional and psychological counselling (n=1). One study compared the effectiveness of a group-based preoperative educational intervention delivered via teleconferencing to an in-person intervention. This study was conducted during the COVID-19 pandemic, when teleconferencing became an alternative to in-person meetings. In the remaining three studies, participants received an intervention, no usual care component was described and outcomes were compared before and after the intervention.

Interventions were based on exercise (n=4), physical activity counselling (n=1), group-based preoperative educational (n=1) and text message-based prehabilitation (n=1). Most interventions were delivered face-to-face or with significant direct healthcare professional input, suggesting a potential mismatch with the need for scalable, low-resource solutions. Notably, six interventions required direct input from healthcare professionals, with delivery methods including remote teleconferencing (n=2), face-to-face sessions (n=3), and one study that compared both approaches, often conducted in settings like hospitals and university physiotherapy departments. One intervention was delivered through text messaging, where patients received four-weekly text message prompts encouraging healthy

behaviours based on Australian physical activity and nutrition guidelines. Exercise interventions frequently required specialist equipment, which, in the case of home-based interventions, was provided to participants along with internet connectivity installation or upgrades where appropriate for use during the study period, adding to the resource demands. Remote interventions conducted via videoconferencing or telehealth required participants to have access to digital devices, internet connectivity, and the necessary skills to use these technologies. Greater detail of the specific interventions being investigated is shown in Table 1.

Included studies reported a variety of outcomes that were related to aspects of health and well-being. Commonly reported outcomes included quality of life (n=6) and anthropometric measures such as weight, BMI, fat mass, and fat-free mass (n=5). Mental health measures, including anxiety and depression, were reported in two studies, while fatigue (n=2), stress (n=1), and functional capacity (n=1) were less frequently assessed.

The methodological quality of included studies was assessed in a structured way using established and validated critical appraisal tools for RCTs (Barker et al, 2023) or quasiexperimental studies (Barker et al, 2024). The approach used is reported in more detail in Section 5. All studies had some risk of bias, with quasi-experimental studies rated as low quality (n=5) and RCTs as moderate (n=1) and low (n=1) quality. The moderate quality RCT assessed a Core Stabilisation Exercise Program plus physical activity counselling (usual care), compared with physical activity counselling alone (Arman et al, 2021). A common issue in six studies was the lack of appropriate statistical analysis, including the absence of power calculations and ineffective statistical methods for assessing intervention effectiveness. Three before and after studies did not include control groups and participant outcomes were assessed and comparisons were made before and after the intervention. One RCT lacked blinding, and the other raised concerns about allocation concealment, group similarity at baseline, measurement reliability, and follow-up reporting. Further details of the quality of individual studies are provided in Section 6.3.

A summary of the findings of the effectiveness of each intervention type is reported in the next section. Our certainty (or confidence) in the overall body of evidence on which the findings were based for each outcome measure was assessed according to: the relevance of the available evidence in addressing the review question (directness), the amount and quality of the evidence, the magnitude and direction of effects, similarity of the studies and consistency in the findings (in whether they favour the intervention or control). This information was used to classify the body of evidence for each outcome into one of four levels of certainty: high, moderate, low, and very low. These ratings indicate the degree of confidence we have in the findings, with a high rating indicating, that having assessed the potential problems with the available evidence we are very confident that the summary findings represent the true value of the intervention effect, whilst a very low rating indicates that we have very little confidence that our summary findings represents the true underlying intervention effect. The results of this assessment for the main outcomes, quality of life and mental well-being (anxiety or depression) are provided in Table 2.

#### Table 1: Summary of interventions

| Citation<br>(Country) and<br>Study design  | Intervention details   | Participants and setting  |
|--|--|---|
| etuaj aceign   | Exercise Interventions   |   |
| Arman et al.<br>2021   | Intervention: Core Stabilisation Exercise Program.<br>Consisting of a combination of strengthening,<br>endurance and balance exercises accompanied by<br>breathing   | <b>Participants:</b> Patients on a waiting list for bariatric surgery, referred by their surgical team. |
| RCT  | <b>Comparator group/ usual care intervention:</b> Both groups received physical activity counselling.  | <b>Setting:</b> University physiotherapy and rehabilitation centre.                                     |
|  | <b>Duration:</b> Twice weekly, for eight weeks.<br><b>Mode of delivery:</b> Supervised, in-person.   |   |
|  |  |   |
| Baillot et al.<br>2017   | <b>Intervention:</b> Endurance and strength training supervised via telehealth.  | <b>Participants:</b> Patients on a waiting list for bariatric surgery. Referral method not stated.      |
| (Canada)<br>Before and<br>after study with<br>comparator<br>group from a<br>previous study | Comparator group/usual care intervention:<br>Participants received individual lifestyle counselling<br>sessions every six to eight weeks. Historical<br>comparator groups included an in-person physical<br>activity programme and a comparator group that<br>received individual lifestyle counselling only.<br>Duration: Two, weekly 80-minute sessions. | Setting: Participants homes   |
|  | <b>Mode of delivery:</b> Supervised, via videoconferencing.  |   |
| Marc-<br>Hernández et<br>al. 2019  | <b>Intervention:</b> An exercise programme combining endurance and resistance training.  | <b>Participants:</b> Patients awaiting bariatric surgery, referred by surgical teams.                   |
| (Spain)  | <b>Comparator group/usual care intervention:</b> All participants followed usual presurgical care including nutritional and psychological counselling.   | Setting: Not stated.  |
| Non-<br>randomised<br>controlled trial   | <b>Duration:</b> 12-week programme, participants trained up to four times per week as the intervention progressed.   |   |
|  | Mode of delivery: Supervised, in-person.   |   |
| Tokgoz et al.<br>2022  | Intervention: Pre-surgical aerobic dance-based exercise  | <b>Participants:</b> Patients awaiting bariatric surgery. Referral method not stated                    |
| (Turkey)   | <b>Comparator group/usual care intervention:</b> All participants including the comparator group received  | Setting: Not stated.  |
| RCT  | Duration and frequency: 60-minutes twice-weekly for 8-weeks.   |   |
|  | Mode of delivery: Supervised, in-person group sessions   |   |
|  | Physical activity counselling interver   | itions  |
| Baillot et al.   | Intervention: Physical activity counselling sessions   | Participants: Patients on a waiting   |
| 2024   | via videoconferencing designed to increase   | list for metabolic or bariatric surgery.  |
|  | moderate-to-vigorous intensity physical activity.  | referred by a clinician.  |

medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

perpetuity. It is made available under a CC-BY-ND 4.0 International license .

| (Canada)                   |   |  |
|----------------------------|---|--|
| Uncontrolled<br>before and | <b>Comparator group/usual care intervention:</b> No usual care intervention was stated. | Setting: Participants homes.             |
| after study                | <b>Duration:</b> Six, weekly session lasting 45 minutes per session.                    |  |
|                            | <b>Mode of delivery:</b> Remote, videoconferencing delivered by a trained professional. |  |
|                            | Text message-based prehabilitation inter  | ventions                                 |
| Kulinski &                 | Intervention: Mobile phone-based automated text   | Participants: Participants on a          |
| Smith 2020                 | message prehabilitation programme including a bank                                      | waiting list for elective surgery were   |
| 011111 2020                | of 96 messages from the fields of nutrition exercise                                    | contacted by researchers asking to       |
| (Australia)                | psychology and medical management based on  | participate in the study. Specific       |
| (, luci alla)              | Australian quidelines.  | surgery types were not stated.           |
|                            |   |  |
| Pilot                      | Comparator group/usual care intervention: Not   | Setting: Text messages delivered         |
| uncontrolled               | stated.   | to participants mobile device.           |
| before and                 |   |  |
| after study                | <b>Duration:</b> Messages were sent four times per week                                 |  |
| ,                          | for six months. Links to local health resources were                                    |  |
|                            | also provided   |  |
|                            |   |  |
|                            | Mode of delivery: Text messaging  |  |
|                            |   |  |
|                            | Group-based preoperative educational inte   | erventions                               |
| Lau et al. 2024            | Intervention: Group-based preoperative Educational                                      | Participants: Patients on a waiting      |
|                            | Programme delivered in-person or online via   | list for metabolic or bariatric surgery. |
| (Germany)                  | videoconferencing. All sessions were group-based  | Referral method not stated.              |
|                            | and aimed to bridge the gap in preoperative   |  |
| Uncontrolled               | preparation.  | Setting: Hospital (in-person group)      |
| before and                 |   | or in participants homes (online         |
| after study                |   | group).                                  |
|                            | Comparator group/usual care intervention:   |  |
|                            | Intervention modes were compared (online vs in-   |  |
|                            | person). Usual care was not stated.   |  |
|                            |   |  |
|                            | <b>Duration:</b> The programme lasted three months and                                  |  |
|                            | consists of four theoretical and three exercise   |  |
|                            | sessions.   |  |
|                            |   |  |
|                            | Mada of delivery in person or constants   |  |
|                            | widessenferencing. Originally delivered in person                                       |  |
|                            | videoconierencing. Originally delivered in-person,                                      |  |
|                            | but conducted online via video conferencing due to                                      |  |
|                            |   |  |
|                            |   |  |
| 1                          |   | 1  |

### 2.2 Effectiveness of exercise interventions

Four studies (Arman et al, 2021; Baillot et al, 2017; Marc-Hernández et al, 2019; Tokgoz et al, 2022) investigated the effects or feasibility of exercise interventions on the health and well-being of patients on waiting lists for bariatric surgery. The types of exercise interventions varied, including combined resistance and endurance programmes (Baillot et al, 2017; Marc-Hernández et al, 2019), a core stability exercise programme (Arman et al, 2021), and aerobic dance-based exercise (Tokgoz et al, 2022). Participants received the exercise intervention in addition to usual care in all four studies. The studies included two RCTs (Arman et al, 2021; Tokgoz et al, 2022), one non-randomised controlled trial (Marc-Hernández et al, 2019) and one before and after study with comparator group from a

previous study (Baillot et al, 2017;). Overall, the evidence provided by these studies was of very low certainty.

#### Quality of life outcomes

- Arman et al (2021) assessed the impact of a supervised core stability exercise programme in people with obesity awaiting bariatric surgery. Both the intervention group and control group received physical activity counselling as part of usual care. After the eight-week intervention, a statistically significant difference in Obesity Specific Quality of Life Questionnaire (OSQOL) scores were observed in both groups and between-groups (p=0.001). The intervention significantly improved their OSQOL scores (ES= 0.41, p=0.005), whereas the control group significantly worsened (ES= -0.48, p=0.011).
- Two studies explored the effectiveness of supervised resistance and endurance exercise interventions in people with obesity awaiting bariatric surgery (Baillot et al, 2017; Marc-Hernández et al, 2019). Baillot et al (2017) assessed the feasibility of implementing a home-based videoconferencing intervention and found no significant difference in weight-related quality of life (WRQOL) scores compared to usual care (p≥0.4) or an in-person intervention (p=0.9) group from a previous study (descriptive statistics to indicate the direction of effect were not reported). Marc-Hernández et al (2019) used the 36-Item Short Form Survey (SF-36) questionnaire to assess the impact of a supervised in-person intervention. While the intervention group increased mean scores in all domains, significant between-group differences in favour of the intervention group were reported for only two of eight scales which included: physical functioning (p=0.026), and general health perception (p=0.005), a significant difference was apparent for the physical component summary (p=0.046) as well.
- A group-based aerobic dance intervention (Tokgoz et al, 2022) significantly improved Impact of Weight on Quality of Life-Lite (IWQOL-Lite) scores in people with morbid obesity awaiting bariatric surgery. The intervention group improved their scores immediately post-intervention (p>0.005), whereas control group scores significantly worsened following eight-weeks of physical activity counselling. The improvement in scores shown in the intervention group were greater than that of the control group post-intervention (mean difference: -14.46 ± 22.08 vs 1.64 ± 22.08, p=0.007).

#### Anthropometric outcomes

- Arman et al (2021) measured the effect of the intervention on a range of body composition measures including body weight, BMI, fat mass and fat free mass. No significant differences were reported for either group following the programme (p > 0.05).
- Baillot et al (2017) found no significant differences in BMI or fat mass in participants undergoing the videoconferencing intervention compared to the usual care or inperson programme (p>0.2).
- Tokgoz et al (2022) reported no significant between-group differences for body weight, BMI or fat mass post-intervention (p>0.05).

medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

- perpetuity. It is made available under a CC-BY-ND 4.0 International license .
- Results reported by Marc-Hernández et al (2019) showed the intervention group • obtained significant reductions in all anthropometric and body composition variables evaluated total weight (-7.3 ± 4.1 kg, p<0.01), BMI (- 2.6 ± 1.5, p=0.001), fat mass  $(-7.1 \pm 4.7 \text{ kg}, \text{ p} < 0.01)$ , and waist circumference  $(-5.3 \pm 2.1 \text{ cm}, \text{ p} < 0.01)$  except in the fat-free mass (kg), which remained constant. In addition, all variables excluding fat-free mass (kg), showed significant reductions that were superior to the control group.

#### Functional outcomes

- Arman et al (2021) found that the intervention group (IG) significantly improved their 6-Min Walk Test (MWT)-distance values after eight weeks (MD= 146 ± 42.33 [95% CI: 115.71 to 176.28]). The control group 6-MWT-distance values significantly worsened after eight-weeks (MD= -36.63 ± 50.91 [95% CI:-70.83 to -2.43]). There was a statistically significant difference in 6 MWT-distance scores between the groups after the programme in favour of the exercise group (p<0.05). Cohen's d indicated on all scores of 6 MWT-distance ranged from 0.03 to - 0.78 for the exercise group and - 0.05 to 0.26 for the control group.
- Tokgoz et al (2022) measured fatigue using the Fatigue Severity Scale (FSS), and • reported significant improvements from baseline to post intervention. These improvements were significantly greater than the control group (MD=  $-2.89 \pm -1.13$ vs -0.49 ± 1.26, p=0.001).

#### Bottom line results for exercise interventions

Overall, our certainty in the findings that exercise interventions are effective in improving the quality of life of people with obesity awaiting surgery is very low (Table 2). The evidence base was limited, with only four relevant studies identified three of which were rated as low methodological quality and one study assessed the feasibility of implementing their intervention. The one moderate quality RCT reported statistical improvements in quality of life following exercise interventions. However, the lack of blinding of participants, outcome assessors and those delivering the treatment limited the strength of these findings. The differences in study design, intervention characteristics such as exercise type, and mode of delivery among these studies limit our confidence in the findings. Studies also used a variety of measurement tools to assess quality of life outcomes, which complicates comparisons and further limits certainty. Evidence for the impact of exercise on anthropometric measures, such as body weight and BMI varied, with a majority of studies showing no significant effect of the intervention. There is also very little evidence supporting improvements in functional outcomes.

#### 2.3 Effectiveness of text message-based prehabilitation interventions

One study (Kulinski and Smith 2020) examined the feasibility of implementing of a text message-based prehabilitation intervention on patients with obesity awaiting elective surgery. This study was a before and after pilot study conducted in Australia. This study was judged to be of low quality and only provided descriptive analysis of outcomes relevant to this rapid review.

#### Quality of life outcomes

 Kulinski and Smith (2020) found that 61% (n=11) of participants reported that their overall health score, as indicated by the quality of life EQ-5D-3L visual analogue scale improved after completing the programme. Twenty-seven percent (n=5) of participants reported improved general health-related quality of life.

#### Anthropometric outcomes

Kulinski and Smith (2020) reported that 40% (n=8) of participants lost at least 2 kg over the course of the intervention. Those who did lose weight lost a mean of 6 kg over six months (range 2– 12 kg), with a mean weight loss of 1.7 kg for the entire cohort.

#### Bottom line results for text message-based prehabilitation interventions

The evidence for the effectiveness of text message-based prehabilitation interventions for improving quality of life, weight is of very low certainty, is reliant on a single low-quality study that lacked appropriate statistical analysis and reporting. Due to the significant methodological limitations and reliance on a single feasibility study, the effectiveness of this intervention remains unclear.

### 2.4 Effectiveness of group-based preoperative educational interventions

One study (Lau et al., 2024) examined the impact of a group-based educational intervention on mental health outcomes, comparing videoconferencing and face-to-face delivery modes of delivery. This low quality, before and after study conducted in Germany provided very low certainty evidence for the effectiveness of group-based preoperative educational interventions.

#### Mental well-being outcomes

Anxiety and depression scores, measured using the GAD-7 and PHQ-9 questionnaires respectively, were significantly improved following completion of the study (anxiety: -1.1 units ± 4.6, z = -3.914, p<0.001; depression: -0.9 units ± 4.6, z = -3.771, p<0.001). Participants also significantly reduced perceived stress scores (PSQ-20), overall stress reduced between by 4.6 ± 15.6 points (z = -4.976, p<0.001). In particular, the subscale "worries" showed an average improvement of -7.9 ± 22.7 points (z = -5.761, p<0.001). A mean reduction of five points was achieved for the subscale "tension" (z = -4.237, p<0.001) and for the subscale "demands" (z = -4.944, p<0.001). There were no significant changes on the "joy" subscale. There were no interaction effects between the two subgroups indicating that the mode of delivery (face-to-face vs. videoconference) did not significantly influence outcomes.</li>

#### Quality of life outcomes

• There was no significant change in the physical component of the SF-12 however, there was a trend toward improvement in the mental component of quality of life, with a mean increase of 2.3 points ± 10.8 (z = 3.022, p=0.003).

#### Anthropometric outcomes

No significant differences in weight or BMI were reported following completion of the intervention (weight: -0.3 kg ± 8.7, z = -0.561, p=0.575; BMI: -0.1 units, ± 3.1, z= -0.277, p=0.782).

#### Bottom line results for group-based preoperative educational interventions

The evidence for group-based educational interventions, comparing videoconferencing and face-to-face delivery modes, is of very low certainty, with findings from a single low-quality study. Results for both interventions indicate improvements in mental well-being outcomes, including anxiety, depression, and stress reduction. However, there were no significant changes in quality of life, with only a trend towards improvement in the mental component, and no significant changes in the physical component. The interventions showed no effect on anthropometric outcomes, such as weight or BMI. The mode of delivery (face-to-face vs. videoconference) did not significantly influence the results.

#### 2.5 Effectiveness of physical activity counselling interventions

One study (Baillot et al., 2024) evaluated the feasibility, acceptability, and effectiveness of a telehealth-based physical activity counselling intervention designed to increase moderate-to-vigorous physical activity in adults awaiting metabolic or bariatric surgery. This low quality, before and after study was conducted in Canada. The study's limited statistical analysis undermines the precision and clarity of the findings, making the effectiveness of the intervention unclear.

#### Mental well-being and quality of life outcomes

Baillot et al (2024) found at least one minimally important difference (improvement in RAND-36 score > 25% post-intervention) for psychosocial outcomes (anxiety and depressive symptoms, quality of life) was reached by 90.9% (n=10) of participants during and after the intervention. This low-quality study does not report inferential statistics for these outcomes; therefore, the statistical significance is unclear.

#### Bottom line results for physical activity counselling interventions

The evidence for the effectiveness of physical activity counselling interventions is of very low certainty and is limited to a single low quality feasibility study. While a high proportion of participants showed improvements in some mental well-being and quality of life domains, the poor reporting restricts the ability to gauge the true effectiveness of the intervention. The results suggest potential benefits in improving psychosocial outcomes, but the overall evidence is very low certainty and inconclusive.

### Table 2: Summary of the evidence

| Intervention<br><i>Outcome</i><br>[No. of studies]                                  | Findings   | Limitations   | Number of<br>participants                | Indirectness  | Inconsistency   | Certainty   |
|---|--|---|--|---|---|---|
|   | Intervention effect<br>based on<br>statistical<br>significance                                   | Study design and quality<br>from critical appraisal<br>checklist  | Participant<br>numbers across<br>studies | Does the evidence cover similar<br>people/ intervention/<br>comparators we were interested<br>in?   | Variation in findings across studies (eg CIs don't<br>overlap   | $\begin{array}{c} \oplus \ominus \ominus \ominus \\ \text{very low} \\ \oplus \oplus \ominus \ominus \\ \text{low} \\ \oplus \oplus \oplus \oplus \ominus \\ \text{moderate} \\ \oplus \oplus \oplus \oplus \\ \text{high} \end{array}$ |
|   |  |   | Q  | uality of life  |   |   |
| Exercise-based<br>interventions<br><i>Quality of life</i><br>N=4                    | Positive <sup>Arman 2021,</sup><br>Tokgoz 2022*<br>Neutral Baillot 2017†,<br>Marc-Hernandez 2019 | One moderate quality<br>RCT Arman 2021<br>One low quality RCT Tokgoz<br>2022<br>One low quality non-RCT <sup>,</sup><br>Marc-Hernandez 2019<br>One low quality before<br>and after study with<br>comparator group from a<br>previous study Baillot 2017 | Range: 6-34                              | Yes<br>Participants were on waiting lists<br>for metabolic or bariatric surgery.<br>Interventions primarily aimed to<br>prepare them for surgery. | Results favoured the intervention but was not<br>always statistically significant. Interventions<br>differed in terms of content, setting and mode of<br>delivery. Tools used to measure outcomes<br>varied across studies. | ⊕⊝⊝<br>very low   |
| Education<br>interventions<br><i>Quality of life</i><br>N=1                         | Neutral <sup>Lau 2024**</sup>  | One low quality<br>uncontrolled study <sup>Lau 2024</sup>   | 305                                      | Yes<br>Patients with obesity on a<br>waiting list for bariatric surgery.<br>The intervention primarily aimed<br>to prepare them for surgery.      | Results from one study. Findings favoured the intervention but no significant improvement was identified.   | ⊕⊝⊝<br>very low   |
| Text-messaging<br>prehabilitation<br>interventions<br><i>Quality of life</i><br>N=1 | Positive <sup>Kulinski</sup><br>2020***  | One low quality<br>uncontrolled study <sup>Kulinski</sup><br><sup>2020</sup>  | 22                                       | Yes<br>Patients with obesity on waiting<br>lists for elective surgery. The<br>intervention primarily aimed to<br>prepare them for surgery.        | Results from one study. Findings favoured the intervention but no significance values reported  | ⊕⊝⊝⊝<br>very low  |
| Physical activity<br>counselling<br>interventions                                   | Unclear Baillot 2024†  | One low quality<br>uncontrolled study <sup>Baillot 2024</sup>   | 12                                       | Yes   | Results from one study. which did not report<br>inferential statistics  | ⊕⊝⊝⊝<br>very low  |

| i                 | 1                     |  |     |                                     |  |                               |
|-------------------|-----------------------|--|-----|-------------------------------------|--|-------------------------------|
| Quality of life   |                       |  |     | Patients with obesity on a          |  |                               |
| N=1               |                       |  |     | waiting list for bariatric surgery. |  |                               |
|                   |                       |  |     | The intervention aimed to           |  |                               |
|                   |                       |  |     | prepare them for surgery.           |  |                               |
|                   |                       |  | Mei | ntal well-being                     |  |                               |
| Education         | Positive Lau 2024     | One low quality                        | 305 | Yes                                 | Results from one study. Findings favoured the    | $\Theta \Theta \Theta \Theta$ |
| interventions     |                       | uncontrolled study <sup>Lau 2024</sup> |     |                                     | intervention and were statistically significant  | very low                      |
| Anxiety           |                       |  |     | Patients with obesity on a          |  |                               |
| N=1               |                       |  |     | waiting list for bariatric surgery. |  |                               |
|                   |                       |  |     | The intervention primarily aimed    |  |                               |
|                   |                       |  |     | to prepare them for surgery.        |  |                               |
| Education         | Positive Lau 2024     | One low quality                        | 305 | Yes                                 | Results from one study. Findings favoured the    | $\Theta \Theta \Theta \Theta$ |
| interventions     |                       | uncontrolled study <sup>Lau 2024</sup> |     |                                     | intervention and were statistically significant  | very low                      |
| Depression        |                       |  |     | Patients with obesity on a          |  | -                             |
| N=1               |                       |  |     | waiting list for bariatric surgery. |  |                               |
|                   |                       |  |     | The intervention primarily aimed    |  |                               |
|                   |                       |  |     | to prepare them for surgery.        |  |                               |
| Education         | Positive Lau 2024     | One low quality                        | 305 | Yes                                 | Results from one study. Findings favoured the    | $\Theta \Theta \Theta \Theta$ |
| interventions     |                       | uncontrolled study <sup>Lau 2024</sup> |     | Patients with obesity on a          | intervention and were statistically significant  | very low                      |
| Stress            |                       |  |     | waiting list for bariatric surgery. |  | -                             |
| N=1               |                       |  |     | The intervention primarily aimed    |  |                               |
|                   |                       |  |     | to prepare them for surgery.        |  |                               |
| Physical activity | Unclear Baillot 2024† | One low quality                        | 12  | Yes                                 | Results from one study. Findings favoured the    | $\Theta \Theta \Theta \Theta$ |
| counselling       |                       | uncontrolled study Baillot             |     |                                     | intervention but no significance values reported | very low                      |
| interventions     |                       | 2024                                   |     | Patients with obesity on a          |  | -                             |
| Anxiety           |                       |  |     | waiting list for bariatric surgery. |  |                               |
| N=1               |                       |  |     | The intervention primarily aimed    |  |                               |
|                   |                       |  |     | to prepare them for surgery.        |  |                               |
| Physical activity | Unclear Baillot 2024† | One low quality                        | 12  | Yes                                 | Results from one study. Findings favoured the    | $\Theta \Theta \Theta \Theta$ |
| counselling       |                       | uncontrolled study Baillot             |     |                                     | intervention but no significance values reported | very low                      |
| interventions     |                       | 2024                                   |     | Patients with obesity on a          |  |                               |
| Depression        |                       |  |     | waiting list for bariatric surgery. |  |                               |
| N=1               |                       |  |     | The intervention primarily aimed    |  |                               |
|                   |                       |  |     | to prepare them for surgery.        |  |                               |

Footnotes:

\*Positive effect post-intervention not sustained five months post-operatively \*\*Despite no overall QoL improvement, there was a significant trend toward improvement in the mental component \*\*\*No significance values reported <sup>†</sup>Results cannot be sufficiently interpreted due to poor reporting

# 3. DISCUSSION

### 3.1 Summary of the findings

This rapid review aimed to identify and synthesise the evidence for the effectiveness of interventions to support the health and well-being of people with obesity on waiting lists, with a focus on identifying practical and resource-efficient interventions that can be implemented within the current healthcare constraints of tier 3 weight management services in Wales.

Seven primary studies were identified. These assessed the effectiveness or feasibility of implementing exercise, physical activity counselling, education or text messaging-based interventions. The findings of these studies indicated that these interventions can improve quality of life, anxiety, stress and depressive symptoms, as well as anthropometric measures (body weight, BMI, fat mass, fat-free mass) of patients awaiting surgery. However, the evidence supporting these findings was deemed to be of very low certainty.

Four studies investigating exercise interventions were identified, while the remaining intervention types were reliant on findings from a single study. Although exercise interventions demonstrated positive results for improving quality of life, interventions varied greatly in terms of 1) exercise type (e.g, a Core Stability Exercise Programme and an aerobic based dance intervention), 2) mode of delivery (e.g, supervised in-person or remote delivery via teleconferencing), 3) study design and aim (e.g, RCTs, non-RCTs and uncontrolled studies, and one study assessed the feasibility of implementing an intervention), and 4) quality of life outcomes and tools used to assess them. This, as well as the studies being predominantly of low methodological quality, means that we are not confident in the evidence.

The stakeholders of this rapid review aimed to use findings to inform support strategies for people with obesity on waiting lists for tier 3 weight management services. However, it is likely that the type of interventions assessed in the included studies would not be feasible for implementation within this context. The interventions were resource-intensive and most were delivered in-person, which contrasts with our stakeholder need for brief and low-resource solutions that are scalable to a large number of patients waiting for treatment. One study (Kulinski and Smith 2020) assessed an automated text message-based prehabilitation intervention, which may be better suited to weight management service waiting list constraints; however, this study assessed feasibility of implementation, not effectiveness.

### 3.2 Strengths and limitations of the available evidence

The evidence included in this rapid review was impacted by significant methodological limitations. Most studies were rated as low quality due to not including a control group, absence of blinding in RCTs, and inadequate statistical analysis to support the validity of the findings. These issues contributed to a very low certainty of evidence, which makes it challenging to draw robust conclusions about the effectiveness of the interventions evaluated. Furthermore, three studies assessed the feasibility of an intervention, so effectiveness could not be determined.

Several key evidence gaps further limit the applicability of the findings to the stakeholder context. Firstly, there were no UK-based studies included in the review, and the geographic

scope of the studies may affect the generalisability of the results, particularly given differences in healthcare systems, patient demographics, and service availability.

Additionally, most of the included studies focused on individuals on waiting lists for bariatric or metabolic surgery, which does not fully align with a population that is of direct relevance to the stakeholders i.e. people with obesity waiting for tier 3 services. In the UK, patients undergoing bariatric surgery are typically at tier 4 in the weight management pathway and have usually received tier 3 treatment prior to referral for surgery. Therefore, these patients are less representative of the population of primary interest to stakeholders. Additionally, only one study specifically targeted the stabilisation and improvement of mental health during the waiting period, revealing a significant gap in evidence.

We didn't identify any relevant economic evaluations and no study included an assessment of the cost-effectiveness of interventions, an essential factor for determining the practicality of implementing these interventions within constrained healthcare budgets.

The review did not find any studies that could be feasibly and scalably implemented within current healthcare constraints faced by obesity weight management services. The reliance of technologies (e.g. laptops for videoconferencing) in some interventions also raises equity concerns, as digitally excluding participants-particularly those without reliable internet access, digital devices, or the necessary digital literacy-could disproportionately impact vulnerable groups who are already at a disadvantage in accessing healthcare services. These barriers could limit the reach and effectiveness of such interventions, undermining their potential impact.

#### 3.3 Strengths and limitations of this Rapid Review

The methods used for this rapid review were based on an abbreviated systematic review approach, following established guidelines in an attempt to capture all relevant publications with minimal risk of bias in a timely manner. Included studies were systematically identified through an extensive search of electronic databases and grey literature.

However, the rapid review faced notable limitations. During preliminary work, no evidence directly addressing the stakeholders' original question was identified. As a result, the review's scope was broadened to capture a wider range of studies. This adjustment, while necessary to ensure potentially relevant research was identified, led to the inclusion of studies of interventions that did not directly address the initial research question. The broadened scope also resulted in the inclusion of studies that could not be feasibly or scalably implemented within current healthcare constraints experienced by stakeholders and were not specifically focused on the population of interest-individuals with obesity on waiting lists for tier 3 weight management services.

Additionally, poor reporting of results in several studies made it challenging to interpret findings accurately. Despite efforts to ensure a robust synthesis and consistency checks at each stage of the review, these reporting issues affected the overall clarity and applicability of the evidence.

#### Implications for policy and practice 3.4

Although we have little confidence in the evidence, there is more evidence of better quality (including two RCTs and two controlled studies) that suggest exercise interventions may

support the quality of life of people with obesity waiting for surgery, with no studies reporting a decline in quality of life. This evidence could be cautiously considered to inform interventions in practice but those designing interventions should be mindful of the population and setting in which they are applied.

The findings of this rapid review highlight the need for increased resource allocation to conduct and rigorously evaluate robust studies and economic evaluations investigating waiting list interventions for individuals with obesity. A particular focus should be on UK-based interventions for those awaiting weight management services. When designing interventions decision makers should consider both the feasibility to implement and scalability within the current healthcare resource constraints.

Emerging approaches in the UK show a shift towards digital health solutions to improve the availability of weight management support. This may provide an alternative option for those on waiting lists as well as people in areas where multidisciplinary weight management services are not vet provided. The National Institute for Health Excellence (NICE, 2023) Health Technology Evaluation (HTE14) of digital technologies highlights the potential of applications to deliver multidisciplinary weight-management services and prescribe and/ or monitor weight-management medicine for adults with obesity. These digital platforms could support increased capacity, or reduced demand on clinician time as well as cutting wait times. When considering digital solutions, NICE (HTE14) stresses the importance of evaluating these digital interventions to ensure they are safe, effective, and accessible. While some patients may prefer to access services remotely via digital technologies, it is imperative that services also consider and address the risk of digital exclusion, particularly for those who may need additional support. For example, those at greatest risk of digital exclusion may suffer from a visual, hearing or cognitive impairment, reduced manual dexterity, a learning disability or being unable to read English (NICE, 2023). Furthermore, considering the impact of a shift to digital solutions on those with limited access to technology or low digital literacy is crucial to prevent widening health inequalities.

### 3.5 Implications for future research

Future research should prioritise the development and evaluation of brief, scalable interventions tailored to low-resource settings, specifically designed to support patients to 'wait well' without further health deterioration while on waiting lists for tier 3 services. The current evidence base lacks practical solutions that directly address the resource constraints faced by weight management services, highlighting the need for studies that develop and test interventions which are both effective and feasible within these contexts.

Research efforts should focus on exploring digital health solutions, remote support, and minimal contact models that minimise the need for specialist equipment and extensive healthcare professional input and promote equity, including considerations for those at risk of digital exclusion. These approaches could provide scalable options that better meet the needs of healthcare providers and patients in low-resource settings. Additionally, future research should focus on interventions that address the specific needs of individuals on waiting lists for tier 3 weight management services, emphasising the development of interventions that balance effectiveness with feasibility and accessibility. It is crucial that future studies undertake rigorous evaluations, including cost-effectiveness analyses, to determine the clinical and financial viability of interventions. Understanding the economic impact of implementing these interventions will help stakeholders make informed decisions about adopting these solutions.

medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY-ND 4.0 International license .

#### 3.6 **Economic considerations\***

- There is a lack economic evidence on the impact of implementing interventions for supporting the health and well-being of people with obesity on healthcare waiting lists.
- Obesity is a considerable and prevalent public health issue in Wales that incurs significant economic cost. Obesity costs NHS Wales £73 million per annum (Welsh Government, 2011). If rates of overweight and obesity continue to rise in line with recent trends, by 2050, they will cost NHS Wales £465 million per year, with a cost to the wider society and economy of £2.4 billion (Public Health Wales, 2016).

Obesity and overweight can impact one's ability to enter and remain in work. A press release by the Health Secretary from October 2024 suggested illnesses caused by obesity cause people to take an extra four sick days a year on average (The Telegraph, 2024). Inability to access management services and support while awaiting services may contribute to such absenteeism.

\*This section has been completed by the Centre for Health Economics & Medicines Evaluation (CHEME), Bangor University

### 4. REFERENCES

Arman, N., et al. (2021). "The effects of core stabilization exercise program in obese people awaiting bariatric surgery: A randomized controlled study." Complementary Therapies in Clinical Practice 43: 101342.

Baillot, A., et al. (2024). "Acceptability and Feasibility of the Telehealth Bariatric Behavioral Intervention to Increase Physical Activity Before Bariatric Surgery: A Single-Case Experimental Study (Part I)." Obesity Surgery 34(5): 1639-1652.

Baillot, A., et al. (2017). "Feasibility and effect of in-home physical exercise training delivered via telehealth before bariatric surgery." Journal of Telemedicine & Telecare 23(5): 529-535.

Barker TH, Habibi N, Aromataris E, Stone JC, Leonardi-Bee J, Sears K, et al. The revised JBI critical appraisal tool for the assessment of risk of bias guasi-experimental studies. JBI Evid Synth. 2024;22(3):378-88.

Barker TH, Stone JC, Sears K, Klugar M, Tufanaru C, Leonardi-Bee J, Aromataris E, Munn Z. The revised JBI critical appraisal tool for the assessment of risk of bias for randomized controlled trials. JBI Evidence Synthesis. 2023;21(3):494-506

Keaver L, Xu B, Jaccard A, et al. (2020). Morbid obesity in the UK: A modelling projection study to 2035. Scand J Public Health. 2020 Jun;48(4):422-427.

Kulinski, K. and N. A. Smith (2020). "Surgical prehabilitation using mobile health coaching in patients with obesity: A pilot study." Anaesthesia & Intensive Care 48(5): 373-380.

Lau, T., et al. (2024). "Psychological Benefits of a Preoperative Educational Bridging Program for Bariatric Surgery: Does Face-to-Face versus Videoconference-based delivery make a difference?" Obesity Facts 17: 17.

Life Sciences Hub Wales (2023). Cwm Taf Morgannwg's new weight management service. Available: https://lshubwales.com/success-stories/cwm-taf-morgannwgs-new-weightmanagement-service

Marc-Hernandez, A., et al. (2019). "Impact of Exercise on Body Composition and Cardiometabolic Risk Factors in Patients Awaiting Bariatric Surgery." Obesity Surgery **29**(12): 3891-3900.

National Institute for Health and Care Excellence (2023). Digital technologies for delivering multidisciplinary weight-management services: early value assessment. Health technology evaluation, HTE14. Available: https://www.nice.org.uk/guidance/hte14

Public Health Wales (2016). Making a Difference: Investing in Sustainable Health and Wellbeing for the People of Wales, Executive Summary, Available: https://phwwhocc.co.uk/wpcontent/uploads/2020/08/Making-a-Difference-Investing-in-Sustainable-Health-and-Wellbeing-for-the-People-of-Wales.pdf

StatsWales (2023). National Survey for Wales - adult lifestyle by local authority and health board. Available: https://statswales.gov.wales/Catalogue/National-Survey-for-Wales/Population-Health/Adult-Lifestyles/adultlifestyles-by-healthboard-from-202021

The Telegraph (2024). Widening waistbands are a burden on Britain. Available: https://www.telegraph.co.uk/politics/2024/10/14/widening-waistbands-are-a-burden-onbritain/

Tokgoz, G., et al. (2022). "Effects of pre-surgical aerobic dance-based exercise on lower extremity in people with morbid obesity awaiting bariatric surgery: Randomized controlled study." Clinical Obesity 12(4): e12529.

Watkins P. (2024). The rise in obesity: some food for thought. Senedd Research web pages. Available: https://research.senedd.wales/research-articles/the-rise-in-obesity-some-food-forthought/#:~:text=The%20most%20recent%20figures%20from.are%20also%20increasing%2 0among%20children.

Welsh Government (2021). All Wales Weight Management Pathway 2021. (Adults): Core Components. Available: https://www.gov.wales/sites/default/files/publications/2021-06/allwales-weight-management-pathway-2021.pdf

Welsh Government (2011). Assessing the costs to the NHS associated with alcohol and obesity in Wales. Available: https://www.llyw.cymru/sites/default/files/statistics-andresearch/2019-08/alcoholmisuseandobesityreporten 0.pdf

Welsh NHS Confederation (2024). The Welsh NHS Confederation response to The Health and Social Care Committee's consultation into the Prevention of ill health - obesity. Available: https://www.nhsconfed.org/system/files/2024-06/wnhsc response obesity inquiry.pdf

medRxiv preprint doi: https://doi.org/10.1101/2024.11.07.24316892; this version posted November 7, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

# perpetuity. It is made available under a CC-BY-ND 4.0 International license .

### 5. RAPID REVIEW METHODS

#### **Eligibility criteria** 5.1

We searched for primary sources to answer the review question: what interventions are effective and cost-effective for supporting the health and well-being of people with obesity on healthcare waiting lists?

#### **Table 3: Eligibility Criteria**

|                                 | Inclusion criteria  | Exclusion criteria  |
|---------------------------------|---|---|
| Population                      | Adults (≥18 years) with obesity (BMI ≥ 30 kg/m <sup>2</sup> ) on a waiting list   | Children and adolescents (<18<br>years)   |
|                                 | Lower criteria by 2.5 kg/m <sup>2</sup> (i.e. BMI $\ge$ 27.5 kg/m <sup>2</sup> ) for people from black African, African-Caribbean and Asian   | Pregnant women  |
|                                 | groups.   | Studies excluding those with a BMI $\ge$ 40 kg/m <sup>2</sup> or 35 kg/m <sup>2</sup> with a co-morbidity |
| Intervention /<br>exposure      | vention /         Interventions aimed at mitigating health decline and/or         Interventions aimed at mi |   |
| Control or counter intervention | No intervention or usual care   |   |
| Outcome<br>measures             | Any outcome related to change in health status, quality of life or well-being while on a waiting list.  |   |
|                                 | Cost-effectiveness outcomes.  |   |
|                                 | Participant engagement outcomes:  |   |
|                                 | Acceptability/engagement/uptake of the waiting list intervention  |   |
|                                 | Acceptability, uptake and engagement at point of enrolment on the service/treatment that they are waiting for.  |   |
| Study design                    | Comparative studies and economic evaluations  |   |
| Countries                       | OECD countries (pre-1974)   |   |
| Language of<br>publication      | English   |   |
| Publication date                | 2010 or later   |   |
| Publication type                | Published and preprint  |   |

#### 5.2 Literature search

A search was conducted of the electronic bibliographic databases: Medline, Embase, PsycINFO, and CINAHL Plus with Full Text. The search strategy used for Medline is available in appendix 1. A search for ongoing and completed trials was conducted in Clinicaltrials.gov, and WHO International Clinical Trials Registry Platform (ICTRP). Citation tracking from sources identified during the preliminary stages was also undertaken, as well searching reference lists of potentially relevant reports previously conducted by the review team. All searches were conducted between 22/07/2024 and 23/07/2024. Terms used to cover the population and intervention elements of the eligibility criteria included those to describe obesity and phrase variations to describe waiting lists or waiting for support.

Searches were limited to studies published after 2010, the year that the All-Wales Obesity Pathway was introduced, and to those published in the English language.

#### 5.3 **Reference management**

All citations retrieved from the database searches were imported or entered manually into EndNote and duplicates removed by a single reviewer. The citations that remained were exported and imported to the systematic reviewing platform Rayyan for study selection.

#### 5.4 Study selection process

All titles and abstract were screened independently in duplicate with disagreements settled by discussion and consensus within the review team. This process was repeated for screening records at full-text.

#### 5.5 Data extraction

Data extracted was conducted by a single reviewer and was consistency checked by a second reviewer. Information extracted includes:

- Citation Study design •
- Intervention •
- Comparator
- Study aim •
- Data collection methods and dates •
- Outcomes reported •
- Sample size •
- Participants •
- Setting
- Key findings
- **Observations/Notes**

#### 5.6 Study design classification

The included studies were classified as quasi-experimental studies or RCTs. No economic evaluations were included following the study selection process.

#### 5.7 Quality appraisal

The JBI critical appraisal checklists for quasi-experimental studies (Barker et al, 2024) and RCTs (Barker et al. 2023) were used to assess the methodological guality of each included study. These checklists are not designed to assign an overall score to each study.

Quality assessment was undertaken in duplicate by two independent reviewers. Any discrepancies were discussed and resolved between reviewers. The quality assessment of individual studies can be seen in section 6.3.

#### 5.8 Synthesis

Data was synthesised narratively to provide a collective interpretation of the evidence.

#### 5.9 Assessment of body of evidence

An assessment of the overall body of evidence was made based on the relevance of the available evidence in addressing the review question, the amount and quality of the evidence, the magnitude and direction of effects, and consistency in the findings. This information is provided for the main outcomes (quality of life and mental well-being) in Table 2, and was used to classify the body of evidence for each outcome as: Very low certainty; Low certainty; Moderate certainty; and High certainty.

### 6. EVIDENCE

#### 6.1 Search results and study selection

A visual representation of the flow of studies throughout the review can be found in Figure 1. A total of 3,299 records were retrieved via the searches and, 1,953 records remained following deduplication. A total of 24 articles were screened at full text..

Seven studies were included in the rapid review. Of which, two studies were RCTs, and five were quasi-experimental, including one non-randomised controlled trial, a before and after study that used comparator groups from a previous study, two uncontrolled before and after studies, and one uncontrolled pilot before and after study.

In order to identify studies relevant to the context of weight management services waiting lists, intervention type will be categorised and method of delivery will be described. Relevant delivery methods include but will not be limited to:

- Digital
- Remote
- Self-directed
- Telephone
- Letters/postal
- Brief interventions





Figure 1. PRSIMA flow diagram

### 6.2 Data extraction

### Table 4: Summary of included studies

| Citation (Country)            | Study Details  | Participants & setting             | Key findings                           | Observations/notes                       |
|-------------------------------|--|------------------------------------|--|--|
| Arman et al (2021) <u>The</u> | Study Design:  | Sample size:                       | Functional capacity outcomes:          | Of 86 patients assessed for eligibility, |
| effects of core stabilization | Randomised controlled trial                            | 23 (21 completed)                  | The intervention group significantly   | 56 wereconsidered eligible. Of these,    |
| exercise program in obese     |  | Intervention Group: n= 10,         | improved their 6 MWT-distance          | 23 agreed to participate and were        |
| people awaiting bariatric     | Study aim:   | Control Group: n= 11               | values after eight weeks (IG: mean     | randomised. Of which one patient         |
| surgery: A randomized         | The aim of this study was to compare the               |                                    | difference = 146 ± 42.33 [CI:          | dropped out of the study due to a lack   |
| controlled study,             | effects of the core stabilisation exercise             | Participants:                      | 115.71, 176.28]). The control group    | of adherence to the program and the      |
| Complementary Therapies       | program (CSEP) and physical activity                   | Adult candidates with morbid       | 6 MWT-distance values                  | continuation of sedentary behavior.      |
| in Clinical Practice; 43,     | counselling on functional capacity, physical           | obesity (BMI ≥ 40) awaiting        | significantly worsened after eight-    | One patient failed to come to the clinic |
| p.101342.                     | fitness, physical activity, fatigue and quality        | bariatric surgery were eligible if | weeks (CG: mean difference = -         | due to a transportation problem.         |
|                               | of life (QoL) in obese people awaiting                 | they were expected to be           | 36.63 ± 50.91 [CI:-70.83,-2.43]).      |  |
| Turkey                        | bariatric surgery.                                     | operated on within 3–6 months.     | There was a statistically significant  |  |
|                               |  | Participants were referred to the  | difference in 6 MWT-distance           |  |
|                               | Type of intervention [exposure]:                       | research project by their surgical | scores between the groups after        |  |
|                               | A twice weekly, eight-week Core                        | teams. Exclusion criteria for      | the program in favour of the           |  |
|                               | Stabilisation Exercise Program consisting              | candidates were as follows;        | exercise group ( $P < 0.05$ ). Cohen's |  |
|                               | of a combination of strengthening,                     | unable to regularly attend         | d indicated on all scores of 6 MWT     |  |
|                               | endurance and balance exercises                        | supervised exercise sessions       | ranged from 0.03 to - 0.78 for the     |  |
|                               | accompanied by breathing.                              | from 6 months, have                | exercise group and - 0.05 to 0.26      |  |
|                               |  | musculoskeletal disorders or       | for the control group.                 |  |
|                               | Comparator intervention or control:                    | systemic diseases that may         |  |  |
|                               | All participants received physical activity            | prevent exercise, presence of      | Body composition outcomes:             |  |
|                               | counselling.   | uncontrolled neuropsychiatric      | The changes in all body                |  |
|                               |  | illnesses affecting cooperation    | composition scores (body weight,       |  |
|                               | Data collection methods and dates:                     | and cognitive functions, presence  | BMI, fat mass, fat free mass) were     |  |
|                               | Participants were recruited between                    | of acute pain or heart pain and    | not statistically significant in both  |  |
|                               | October 2019 and March 2020.                           | previous myocardial infarction,    | groups after the programs (p >         |  |
|                               | Assessments were conducted at the start                | subjective heart failure,          | 0.05). No between-group                |  |
|                               | and end of the study (after eight weeks).              | uncontrolled diabetes and          | differences occurred for any body      |  |
|                               |  | hypertension.                      | composition outcomes ( $p > 0.05$ ).   |  |
|                               | Outcomes (relevant):                                   |                                    |  |  |
|                               | <ul> <li>Functional capacity (6-minute walk</li> </ul> | Setting:                           | Fatigue outcomes:                      |  |
|                               | test)  | Department of Physiotherapy and    | Significant improvements in FSS        |  |
|                               | Body composition (bioimpedance                         | Rehabilitation, Faculty of Health  | and FACIT-F was found in the only      |  |
|                               | analysis)  | Science, Istanbul University-      | exercise group (FSS score: effect      |  |
|                               |  | Cerrahpasa                         | size = 0.79), no significant change    |  |

|                               | <ul> <li>Quality of life (Obesity Specific</li> </ul> |                                     | occurred in the control group.        |   |
|-------------------------------|---|-------------------------------------|---------------------------------------|---|
|                               | Quality of Life questionnaire                         |                                     | Moreover, the difference between      |   |
|                               | IOSQOL1)  |                                     | pre-and post-tests was statistically  |   |
|                               | Eatique (Eatique Severity Scale                       |                                     | significant between the exercise      |   |
|                               | and the Eulertional Assessment of                     |                                     | and control groups and $(p < 0.01)$ . |   |
|                               |   |                                     | The effect sizes were 0.79 and 0.59   |   |
|                               | Chronic liness Therapy-laugue                         |                                     | for the scales, respectively and      |   |
|                               | Scale)  |                                     | moderate for both scales.             |   |
|                               |   |                                     |                                       |   |
|                               |   |                                     | Quality of life outcomes:             |   |
|                               |   |                                     | OSQOL scores were statistically       |   |
|                               |   |                                     | significant in both groups (p <       |   |
|                               |   |                                     | 0.01). The intervention group         |   |
|                               |   |                                     | OSQOL score significantly             |   |
|                               |   |                                     | improved whereas the control          |   |
|                               |   |                                     | groups worsened. The effect sizes     |   |
|                               |   |                                     | were 0.41 and -0.48 for the           |   |
|                               |   |                                     | intervention group and control        |   |
|                               |   |                                     | group respectively.                   |   |
|                               |   |                                     | 3.0 dp . cop con c. j .               |   |
| Baillot et al (2024)          | Study Design:   | Sample size:                        | Psychosocial and quality of life      | The outcomes included in this data      |
| Acceptability and             | Quasi-experimental before and after                   | 12                                  | outcomes:                             | extraction were secondary outcomes      |
| Feasibility of the Telehealth |   |                                     | At least one minimal clinically       | designed to assess the feasibility of a |
| Bariatric Behavioural         | Study aim:  | Participants:                       | important difference for              | larger-scale intervention in the future |
| Intervention to Increase      | 1) assess the feasibility and acceptability of        | Aged >18 years: ii) scheduled to    | nsychosocial outcomes (anxiety        | and may lack appropriate statistical    |
| Physical Activity before      | the TEL Ehealth BARIatric behavioural                 | undergo a sleeve gastrectomy        | and depressive symptoms quality       | analysis                                |
| Bariatric Surgery: A Single-  | intervention (TELE-BariACTIV) trial                   | $\geq$ 12 weeks at one of the three | of life) was reached by 10/11         |   |
| Case Experimental Study       | protocol/methods and intervention, which              | participating hospitals; iii) self- | (90.9%) participants during phase     | Intervention acceptability and          |
| (Part I), Obesity Surgery,    | was designed to increase moderate-to-                 | reported ≤150 minutes of MVPA       | B1 (during the intervention) and      | engagement outcomes were reported       |
| 34(5), pp.1639-1652.          | vigorous intensity physical activity (MVPA)           | per week, and iv) had access to a   | during phase A2 (post-intervention)   | by the study: however, they were not    |
|                               | in adults awaiting MBS: and 2) estimate the           | computer with Internet and an       |                                       | comparative and therefore not           |
| Canada                        | effect of the intervention on MVPA.                   | interface with a camera.            |                                       | extracted by the review team.           |
|                               |   | Exclusion criteria were: i) having  |                                       | · · · · · · · · · · · · · · · · · · ·   |
|                               | Type of intervention [exposure]:                      | a physical contraindication to      |                                       |   |
|                               | Six weekly 45-minute PA counselling                   | physical activity: ii) already      |                                       |   |
|                               | sessions via videoconferencing designed to            | enrolled in a supervised exercise   |                                       |   |
|                               | increase MVPA   | intervention or physical activity   |                                       |   |
|                               |   | behavioural change intervention:    |                                       |   |
|                               | Participants were randomly assigned to                | iii) unable to speak and            |                                       |   |
|                               | either a 1- or 2-week baseline A1 phase:              | understand French. or iv)           |                                       |   |
|                               | subsequent phases (phases B1 and A2)                  | needing a wheelchair. cane          |                                       |   |
|                               | were of identical length for all participants.        | walker, or other assistive          |                                       |   |
|                               | The A phases were observational phases                | device(s) to move.                  |                                       |   |
|                               | The replicaces were observational phases              |                                     |                                       |   |

|                               | without intervention, and the B phases<br>were interventional with the TELE-<br>BariACTIV. Daily MVPA was assessed for<br>7 (if randomised to 1 week baseline A1) or<br>14 days (if randomized to 2-week baseline<br>A1) during the A1 phase, and then for 7<br>days in phases B1 and A2.<br><b>Data collection methods and dates:</b><br>Accelerometer, validated questionnaires,<br>and self-reported measures of data<br>collection. Recruitment took place between<br>September 2021 and July 2022. | <b>Setting:</b><br>The intervention was delivered<br>remotely. Participants were<br>recruited from three hospitals in<br>Canada, two associated tertiary<br>care centres and one associated<br>regional centre. |                                       |   |
|-------------------------------|---|---|---------------------------------------|---|
|                               | Outcomes (relevant): <ul> <li>Anxiety (GAD-7)</li> <li>Depressive symptoms (PHQ-9)</li> <li>Quality of life (RAND)</li> </ul>   |   |                                       |   |
| Baillot et al (2017)          | Study Design:   | Sample size:  | Satisfaction outcomes:                | Comparison groups were used from        |
| Feasibility and effect of in- | Quasi-experimental (with historical   | 6   | The baseline in-home telehealth       | Baillot et al (2016) study which        |
| home physical                 | comparison groups)  |   | patients' perception score was high   | compared a supervised in-person         |
| exercise training delivered   |   | Participants:   | (83.5% (80.9–91.2), and increased     | exercise intervention (PreSET) to       |
| via telehealth                | Study aim:  | Women awaiting bariatric surgery  | significantly after the intervention  | usual care.                             |
| before bariatric surgery.     | To assess the feasibility and effect of Pre-  | (aged ≥ 18 years: body mass   | (90.0% (86.8–94.1); p=0.03). The      |   |
| Journal of telemedicine       | Surgical Exercise Training (PreSET)   | index (BMI) $\geq$ 35 with  | global TelePreSET' satisfaction       | Five women for bariatric surgery        |
| and telecare 23(5)            | delivered in-home via telehealth  | comorbidities or $\geq 40 \text{ kg/m}^2$   | was $93.4\%$ (89.3–97.7) with         | refused to be contacted and 21 were     |
| nn 529-535                    | (TelePreSET) in subjects awaiting bariatric   | who were expected to be   | subscores of $97.5\%$ (91.9–98.1) for | contacted by telephone between          |
| pp.020 000.                   | surgery   | operated within 3–6 months  | quality of care provided 90.0%        | March–October 2014 Among the 21         |
| Canada                        |   | Exclusion criteria included:  | (78.0-97.0) for similarity to face-to | women 12 refused to participate (no     |
| Cunada                        | Type of intervention  | medical contraindication for  | face encounter and $100.0 \% (95.0-$  | time for exercise training n=7.         |
|                               | <b>[exposure]</b> : Twelve-weeks of endurance   | physical activity functional  | 100 0) for perception of the          | moving n=2: not interested by           |
|                               | and strength training supervised twice  | limitations >one weekly   | interaction                           | research or telehealth n=3) and nine    |
|                               | weekly using videoconferencing  | supervised exercise sessions no   |                                       | were excluded due to insufficient       |
|                               | wookly doing videocornoronoling.  | access to high speed residential  | Weight-related quality of life        | space for training (n=5) functional     |
|                               | Comparator intervention or control  | Internet and not understanding  | outcomes:                             | limitation (n=1) current involvement in |
|                               | Pre-post as well as a usual care group from   | French  | No difference was also found          | supervised exercise (n=2) and inability |
|                               | a previous study (Baillot et al 2016)   |   | between the TelePreSET and the        | to understand French (n=1)              |
|                               |   | Setting:  | usual care groups $(p>0.4)$ with a    |   |
|                               | Data collection methods and dates:  | Participants homes.   | trend for difference in the           |   |
|                               | Satisfaction and weight-related quality of  |   | confidence during exercise (7.5       |   |
|                               | life were assessed at baseline and after 12-  |   | (2.5-10.0) vs 0.0 (-15.0-5.0):        |   |
|                               | weeks (post-intervention) using validated   |   | p=0.06) and social interaction        |   |

| que<br>bet   | <ul> <li>estionnaires. Participants recruited<br/>tween March-October 2014.</li> <li>utcomes (relevant): <ul> <li>Satisfaction with telehealth</li> <li>Weight-related quality of life (The<br/>Laval questionnaire)</li> <li>Anthropometric outcomes: BMI, fat<br/>mass.</li> </ul> </li> </ul>   |   | scores (11.2 (1.0-28.6) vs 2.0 (–<br>10.2–2.0); p=0.05).<br>Anthropometric outcomes:<br>No significant change was found in<br>anthropometric, body composition<br>and blood pressure after 12 weeks<br>in each group, and no significant<br>differences between groups were<br>observed (p>0.2).  |   |
|--|--|---|---|---|
| Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Australia<br>Austra | <ul> <li><b>udy Design:</b></li> <li>Jasi-experimental before and after pilot Jdy.</li> <li><b>udy aim:</b></li> <li>test the implementation of a mobile one-based text message prehabilitation ogramme for patients with obesity valting elective surgery.</li> <li><b>rpe of intervention [exposure]:</b></li> <li>bobile phone-based automated text essage prehabilitation programme. A nk of 96 messages from the fields of trition, exercise, psychology and medical anagement, based on Australian idelines. Messages were sent four times in week for six months. Links to local analth resources also provided, e.g. rkrun, local council activities.</li> <li><b>omparator intervention or control:</b></li> <li>utcomes compared to baseline.</li> <li><b>ata collection methods and dates:</b></li> <li><b>articipant-reported including physical and thavioural measures, subjective perience, and a validated scale.</b></li> <li><b>assibility also assessed.</b> Participants cruited between October 2018 and May 19.</li> </ul> | 22 (18 completed the study)<br>Participants:<br>Adults with a BMI of 30 and<br>above, listed for a procedure with<br>a 12-month waiting time non-<br>pregnant and owning a mobile<br>phone. Exclusion criteria<br>included: participants had to own<br>a mobile phone, 30+ BMI.<br>Setting:<br>Patients were recruited from the<br>surgical waitlist within the<br>Illawarra-Shoalhaven local health<br>district. | Eight out of 18 participants (40%)<br>lost at least 2 kg by the conclusion<br>of the programme. Those who did<br>lose weight lost a mean of 6 kg<br>over six months (range 2– 12 kg),<br>with a mean weight loss of 1.7 kg<br>for the entire cohort.8 (40%)<br>reported improved BMI change.<br>Participant 5 experienced a 12 kg<br>weight loss and reported that they<br>cancelled their orthopaedic<br>procedure as a result.<br>Health-related quality of life:<br>Eleven participants (61%) reported<br>that their overall<br>'health score' as indicated by the<br>EuroQol visual analogue scale<br>improved after completing the<br>programme.<br>5 of 18 reported improved general<br>health-related quality of life. | <ul> <li>13/15 relused enrolment. Of the four non-completing patients, one was unable to participate by the end of the study due to a new health diagnosis, and three were unable to be contacted at the completion of the programme.</li> <li>2/18 participants thought that 4 messages a week were too many.</li> </ul> |

|                             | Outcomes (relevant):                                   |                                  |  |                                       |
|-----------------------------|--|----------------------------------|--|---------------------------------------|
|                             | <ul> <li>Economics (generic health quality)</li> </ul> |                                  |  |                                       |
|                             | of life measure EO-5D-3L)                              |                                  |  |                                       |
|                             |  |                                  |  |                                       |
| Lau et al (2024)            | Study Design:  | Sample size:                     | Total group:                           | The drop-out rate was $2.8\%$ (n = 10 |
| Psychological Benefits of a | Quasi-experimental before and after                    | 305                              |  | participants). There were no          |
| Preoperative Educational    |  | Face-to-face group: n = 222,     | Anxiety outcomes:                      | significant differences between       |
| Bridging Program for        | Study aim:   | online group: n = 66, both       | Anxiety symptoms significantly         | completers and dropouts at baseline.  |
| Bariatric Surgery: Does     | To assess the effects of a group-based                 | formats: n = 17                  | improved during the intervention,      |                                       |
| Face-to-Face versus         | educational programme before MBS on                    |                                  | with a mean reduction of 1.1 units     |                                       |
| Videoconference-based       | three key factors: i) patients' mental health,         | Participants:                    | (SD 4.6, z = -3.914, p < 0.001).       |                                       |
| delivery make a             | ii) the program's perceived helpfulness                | Adults with a BMI over 40, or    |  |                                       |
| difference?, Obesity Facts  | from the patients' perspective, and iii) the           | greater than 35 with             | Depression outcomes:                   |                                       |
|                             | effectiveness of delivering the program                | comorbidities awaiting bariatric | Depression symptoms significantly      |                                       |
| Germany                     | online via videoconferencing.                          | surgery. These patients have i)  | improved, with a mean reduction of     |                                       |
|                             |  | undergone conservative           | 0.9 units (SD 4.6, z = -3.771, p <     |                                       |
|                             | Type of intervention [exposure]:                       | therapies according to the       | 0.001,)                                |                                       |
|                             | Preoperative Educational Bridging                      | guidelines but still need to     |  |                                       |
|                             | Programme. The programme lasts three                   | exhaust conservative methods     | Stress outcomes:                       |                                       |
|                             | months and consists of four theoretical and            | before qualifying for MBS as the | Overall stress reduced between T0      |                                       |
|                             | three exercise sessions.                               | next-step in guideline-based     | and 11 by 4.6 (SD 15.6) points ( $z =$ |                                       |
|                             |  | therapy or ii) have a primary    | -4.976, p < 0.001). In particular, the |                                       |
|                             | Originally delivered in-person, but                    | indication for surgery, such as  | subscale "worries" showed an           |                                       |
|                             | conducted online via video conferencing                | severe weight-related diseases   | average improvement of -7.9 points     |                                       |
|                             | since the COVID-19 pandemic. The face-                 | that preclude conservative       | (SD 22.7, z = -5.761, p < 0.001). A    |                                       |
|                             | to-face sessions were attended by 12                   | treatment.                       | mean reduction of 5 points was         |                                       |
|                             | participants and each session lasted 90                | Catting                          | achieved for the subscale tension      |                                       |
|                             | minutes, while the videoconference-based               | Setting:                         | (2 - 4.237, p < 0.001) and for the     |                                       |
|                             | sessions had no more than to participants              | Hace-to-lace group. University   | subscale demands $(24.944, p)$         |                                       |
|                             |  | Online group: participants homes | < 0.001). There were no significant    |                                       |
|                             | Comparator intervention or control:                    | Chine group, participants nomes  | changes on the joy subscale.           |                                       |
|                             | Baseline, comparisons also analysed                    |                                  | Quality of life outcomes:              |                                       |
|                             | between face-to-face and online                        |                                  | There was no significant change in     |                                       |
|                             |  |                                  | the physical component between         |                                       |
|                             | Data collection methods and dates:                     |                                  | T0 and T1. However, there was a        |                                       |
|                             | Validated questionnaires, data collected               |                                  | trend toward improvement in the        |                                       |
|                             | and analysed between May 2014 and                      |                                  | mental component of quality of life.   |                                       |
|                             | August 2022.   |                                  | with a mean increase of 2.3 points     |                                       |
|                             | Ĭ  |                                  | (SD 10.8, z = 3.022, p = 0.003).       |                                       |
|                             | Outcomes (relevant):                                   |                                  |  |                                       |
|                             |  |                                  | Weight & BMI outcomes:                 |                                       |

|                            | <ul> <li>Anxiety (GAD-7)</li> </ul>            |                                      | There were almost no changes in  |                                     |
|----------------------------|--|--------------------------------------|--|-------------------------------------|
|                            | <ul> <li>Depression (PHO-9)</li> </ul>         |                                      | weight (-0.3 kg (SD 8.7, z = -0.561,                                   |                                     |
|                            |  |                                      | p = 0.575) and BMI (-0.1 units. SD                                     |                                     |
|                            | • Stress (PSQ-20)                              |                                      | 31 z = -0.277 n = 0.782) during  |                                     |
|                            | <ul> <li>Quality of life (SF-12)</li> </ul>    |                                      | the intervention   |                                     |
|                            | Weight   |                                      |  |                                     |
|                            | BMI  |                                      |  |                                     |
|                            |  |                                      | Face-to-face and   |                                     |
|                            | Satisfaction                                   |                                      | videoconference-based  |                                     |
|                            |  |                                      | Both the face-to-face and  |                                     |
|                            |  |                                      | videoconference subgroups  |                                     |
|                            |  |                                      | showed improvements over time in                                       |                                     |
|                            |  |                                      | anxiety, mental quality of life and                                    |                                     |
|                            |  |                                      | for perceived stress on the overall                                    |                                     |
|                            |  |                                      | stress scale and on the three  |                                     |
|                            |  |                                      | subscales "worries" "tension" and                                      |                                     |
|                            |  |                                      | "demands" Depressive symptoms  |                                     |
|                            |  |                                      | and quality of life remained   |                                     |
|                            |  |                                      | unchanged from T0 to T1. There   |                                     |
|                            |  |                                      | unchanged from 10 to 11. There   |                                     |
|                            |  |                                      | were no interaction effects between                                    |                                     |
|                            |  |                                      | the two subgroups indicating that                                      |                                     |
|                            |  |                                      | the mode of delivery (face-to-face                                     |                                     |
|                            |  |                                      | vs. videoconference) did not   |                                     |
|                            |  |                                      | significantly influence outcomes.                                      |                                     |
|                            |  |                                      | In addition, there were no   |                                     |
|                            |  |                                      | statistically significant differences                                  |                                     |
|                            |  |                                      | between the MCIDs achieved by  |                                     |
|                            |  |                                      | the two subgroups, supporting the                                      |                                     |
|                            |  |                                      | similar effectiveness of the two                                       |                                     |
|                            |  |                                      | delivery modes.  |                                     |
| Marc-Hernández et al       | Study Design:                                  | Sample size:                         | Body composition   | Two women of the intervention aroup |
| (2019) Impact of Exercise  | Non-randomised controlled study (Patients      | 23                                   | (bioimpedance) & BMI outcomes:   | did not finish the exercise program |
| on Body Composition and    | were assigned to the experimental or           | Intervention group: $n = 12$         | The exercise group (EG) obtained                                       | (one left the program and another   |
| Cardiometabolic Risk       | control aroun in chronological order of        | control group: $n = 11$              | significant reductions in all  | experienced physical problems)      |
| Factors in Patients        | recruitment: the first 12 patients were        |                                      | anthronometric and body  | experienced phycical presience).    |
| Awaiting Bariatric Surgery | included in the exercise aroun (EG) and the    | Particinants:                        | composition variables evaluated  |                                     |
| Obesity Surgery 29         | $\frac{1}{10000000000000000000000000000000000$ | Adults with a BMI of 40 or above     | excent in the EEM, which remained                                      |                                     |
| obesity Surgery, 23,       |  | or 35 with an obasity associated     | constant: total weight $(-7.3 \pm 4.1)$                                |                                     |
| pp.0001-0000               | Study aim:                                     | comorbidity awaiting bariatric       | kg $n < 0.01$ ) fat mass (-7.1 ± 4.7                                   |                                     |
| Spain                      | To describe the offects of an exercise         | surgeny Evolution criteria were      | (1, 1, 2, 3, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5, 5,                    |                                     |
| Spain                      | no describe the effects of all exercise        | surgery. Exclusion onlend were       | $ry, \mu > 0.01$ ), and waist<br>aircumforance (-E 2 ± 2.1 cm s =      |                                     |
|                            |  | diagona (h) any are functional       | $(-5.5 \pm 2.1 \text{ CIII}, p < 0.01)$ All these veriables evaluation |                                     |
|                            | carulometabolic fisk factors in                | limitational (a) severe functional   | U.U.T. All these variables, excluding                                  |                                     |
|                            | patients awaiting parlatric surgery.           | initiations, (c) chronic respiratory |  |                                     |
|                            |  | alseases; or (a) plans to undergo    | amerences between groups.  |                                     |

| Type of intervention [exposure]:A 12-week exercise programme which<br>combined endurance and resistance<br>training.Comparator intervention or control:<br>Control group which received advice to<br>follow an active lifestyle.                       | another exercise program during<br>the period of the study.<br><b>Setting:</b><br>Not stated. | The EG had significantly greater<br>reductions than the control group<br>(CG) in the percentage of BMI<br>reduction ( $-5.6 \pm 3.32\%$ vs $-0.57 \pm 1.95\%$ , p = 0.002), fat mass ( $-10.93 \pm 7.73$ vs $-1.75 \pm 3.05\%$ , p =<br>0.005), and visceral fat ( $-11.28 \pm 7.49\%$ vs $-2.33 \pm 5.20\%$ , p =<br>0.011). |  |
|--|---|---|--|
| Both groups received the usual care prior<br>to surgery, including psychological and<br>nutritional counselling.<br><b>Data collection methods and dates:</b><br>Physical and biochemical tests, validated<br>questionnaire, anthropometrics, Dates of |   | Waist and hip circumference<br>outcomes:<br>Waist-to-height ratio only<br>decreased significantly in the EG (p<br>= 0.007).   |  |
| data collection not stated.<br>Outcomes (relevant):  |   | Excess weight loss (EWL)<br>outcomes:<br>The %EWL was of 15.2 ± 11.2%   |  |
| <ul> <li>Body composition (bioimpedance)</li> <li>Waist and hip circumference<br/>(International Society for the<br/>Advancement of Kinanthropometry<br/>protocol)</li> <li>BMI</li> </ul>   |   | vs. $1.5 \pm 5.2\%$ in the EG and the<br>CG, respectively (p= 0.006; 95%<br>CI, 4.6–22.8; Student's t test).<br><b>Basal metabolic rate outcomes:</b><br>Significant changes in the basal   |  |
| <ul> <li>Excess weight loss</li> <li>Basal metabolic rate</li> <li>Health-related guality of life (SF-</li> </ul>  |   | any of the groups.  |  |
| 36)  |   | <b>outcomes:</b><br>the EG increased their scores on all<br>scales of the SF-36. These<br>increases were significant in 3 of<br>the 8 scales of the questionnaire<br>(physical functioning, $p = 0.002$ ,   |  |
|  |   | 95% CI, 11.7, 39.2; general health<br>perceptions, $p = 0.003$ , 95% CI,<br>8.0, 27.6; mental health, $p = 0.021$ ,<br>95% CI, 1.9, 18.8), and in one of<br>the scales, social functioning, there<br>was a large increase (d = 0.81,<br>95% CI, - 0.13, 1.68). In addition,<br>the EG also showed a significant               |  |

|                             |   |                                    | increase in the physical component       |  |
|-----------------------------|---|------------------------------------|--|--|
|                             |   |                                    |  |  |
|                             |   |                                    | summary ( $p = 0.004, 95\%$ CI, 3.1,     |  |
|                             |   |                                    | 12.3). The CG showed significant         |  |
|                             |   |                                    | increases in physical role (p =          |  |
|                             |   |                                    | 0.042, 95% CI, 1.1, 40.6), a small       |  |
|                             |   |                                    | change in bodily pain (d = 0.43,         |  |
|                             |   |                                    | 95%CI – 0.75, 1.54) and a                |  |
|                             |   |                                    | moderate change in emotional role        |  |
|                             |   |                                    | (d = 0.52, 95% Cl – 0.67, 1.63). A       |  |
|                             |   |                                    | significant difference between           |  |
|                             |   |                                    | groups was obtained in absolute          |  |
|                             |   |                                    | changes in physical functioning,         |  |
|                             |   |                                    | general health perceptions, and          |  |
|                             |   |                                    | physical component summary.              |  |
| Tokgoz et al (2022) Effects | Study Design:                               | Sample size:                       | Quality of life outcomes:                | Fifty-six patients were evaluated for  |
| of pre-surgical aerobic     | Randomised controlled trial.                | 34 (4 dropped out)                 | A statistically significant              | eligibility for the study. Thirty-four |
| dance-based exercise on     |   | Intervention group: n=13, control  | improvement was observed in all          | patients were included in the study.   |
| lower                       | Study aim:                                  | aroup: n=17.                       | comparisons of IWQOL-LITE Total          | During the study, 4 patients           |
| extremity in people with    | To investigate what are the effects of pre- |                                    | and IWQOL-LITE Physical Function         | dropped out (personal reasons n=2.     |
| morbid obesity awaiting     | surgical aerobic dance-based exercise       | Participants:                      | Scores in intervention group. It was     | transfer problems n=2).                |
| bariatric                   | programme (PSADBE) and physical activity    | Adults with morbid obesity         | statistically significant only for       | , ,                                    |
| surgery: Randomized         | counselling (PAC) programme on lower        | scheduled for laparoscopic         | comparisons of T1–T2 and T1–T3           |  |
| controlled study. Clinical  | extremity functions after surgery in people | sleeve gastrectomy within 3–6      | in the control group ( $p < .05$ ).      |  |
| Obesity 12(4) pp 12529      | with morbid obesity awaiting bariatric      | months who had not undergone       | Intergroup comparisons found             |  |
| obooldy, 12(1), pp. 12020.  | surgery (PMOABS)                            | any surgical operation at the last | statistically significant differences in |  |
| Turkey                      |   | 6 months. The exclusion criteria   | favour of the intervention group at      |  |
| lancey                      | Type of intervention [exposure]             | were: any musculoskeletal          | T1-T2 for both IWOOI -L ite total        |  |
|                             | Pre-surgical aerobic dance-based exercise   | disorder systemic illness          | (mean difference: IG= -14.46 +           |  |
|                             | for 60-minutes twice-weekly for 8-weeks     | balance problem, psychiatric or    | 16.46 CG= $1.64 + 22.08$ n =             |  |
|                             | under physiotheranist supervision in        | neurological disease acute nain    | 0.007) and IWOOI -I ITE Physical         |  |
|                             | arouns. Both groups received PAC            | or beart pain for any reason       | Eunction Scores mean difference:         |  |
|                             |   | history of myocardial infarction   | $I_{G} = -10.23 + 6.35 C_{G} = 1.70 +$   |  |
|                             | Comparator intervention or control:         | within 1 year, congestive heart    | 8.01 n = 0.001) No statistically         |  |
|                             | Devoiced activity equipoliting Individually | foilure and inshility to adopt to  | $p_{iab}$                                |  |
|                             | est for each participant, included a        | treatment                          | differences were energioup               |  |
|                             | brachura avalaining importance of abusical  | ueaunent.                          |  |  |
|                             | prochure explaining importance of physical  |                                    | 13.                                      |  |
|                             | activity, examples and recommendations,     | Cattinger                          |  |  |
|                             | participants recorded step counts and       | Setting:                           | Body composition outcomes:               |  |
|                             | activity levels for 8-weeks.                | inot stated, but aerobic dance-    | A statistically significant              |  |
|                             |   | pased exercise programme was       | Improvement was found in all             |  |
|                             | Data collection methods and dates:          | aelivered in-person in groups.     | parameters for comparisons of T1–        |  |
|                             |   |                                    | 13 and 12–13 in both groups (p <         |  |
|                             |   |                                    | .05).                                    |  |

| <ul> <li>Physical exercise tests, bioimpedance analysis, validated questionnaires and scales. Groups were evaluated before exercise (T1), after an 8-week exercise programme (T2), and at 5 months postoperative (T3). Data collection dates not stated but participants recruited between November 2019 and June 2020.</li> <li><b>Outcomes (relevant):</b> <ul> <li>Quality of life (IWQOL-LITE)</li> <li>Body composition (Bioimpedance)</li> <li>Fatigue (Fatigue Severity Scale (FSS)</li> </ul> </li> </ul> | The differences of all scores for<br>between-group comparisons of T1–<br>T2 and T2–T3 were not statistically<br>significant ( $p > .05$ ).<br><b>Fatigue outcomes:</b><br>In intergroup comparison, the<br>improvements in the FSS scores<br>for comparisons of T1–T2 and T1–<br>T3 in the intervention group and the<br>improvements in the FSS score<br>only for comparisons of T1–T3 in<br>the control group were found<br>statistically significant ( $p < .05$ ).<br>In intergroup comparison, the<br>differences |
|---|--|
|---|--|

### 6.3 Quality appraisal

#### Table 5: Quality appraisal results for quasi-experimental studies

|  |                  |                       |                        | INTERNAL VALIDITY Bias related to: Statistical Methodological |                     |  |  |   |                       |          |         |             |
|--|------------------|-----------------------|------------------------|---|---------------------|--|--|---|-----------------------|----------|---------|-------------|
|  |                  | Domain                | Temporal<br>precedence | Selection<br>and<br>allocation                                | Confounding factors | Administration<br>of intervention/<br>exposure | Assessment, F<br>detection, and r<br>measurement of the<br>outcome |   | Participant retention | validity | quality |             |
|  |                  | Question no.          | 1                      | 2   | 3                   | 4  | 5  | 6 | 7                     | 8        | 9       |             |
| Study                                  | Outcome          | Result                |                        |   |                     |  |  |   |                       |          |         |             |
| Baillot et<br>al. (2017)               | Quality of life  | Post-<br>intervention | Y                      | Y   | Y                   | U  | N  | Y | U                     | Y        | Ν       | Low quality |
| Baillot et<br>al. (2024)               | Anxiety          | Post-<br>intervention | Y                      | N   | Y                   | Y  | Ν  | Y | U                     | Y        | Ν       | Low quality |
|  | Depression       | Post-<br>intervention |                        |   |                     |  | Ν  | Y | U                     | Y        | Ν       |             |
|  | Quality of life  | Post-<br>intervention |                        |   |                     |  | Ν  | Y | U                     | Y        | Ν       |             |
| Kulinski<br>and Smith<br>(2020)        | Quality of life  | Post-<br>intervention | Y                      | N   | Y                   | Y  | N  | Y | U                     | Y        | N       | Low quality |
| Lau et al.<br>(2024)                   | Anxiety          | Post-<br>intervention | Y                      | N   | Y                   | Y  | Ν  | Y | U                     | Y        | N       | Low quality |
|  | Depression       | Post-<br>intervention |                        |   |                     |  | N  | Y | U                     | Y        | N       |             |
|  | Quality of life  | Post-<br>intervention |                        |   |                     |  | N  | Y | U                     | Y        | N       |             |
|  | Stress           | Post-<br>intervention |                        |   |                     |  | N  | Y | U                     | Y        | N       |             |
| Marc-<br>Hernández<br>et al.<br>(2019) | Quality of life  | Post-<br>intervention | Y                      | Y   | N                   | Y  | N  | Y | U                     | N        | U       | Low quality |
| Key: Y=Yes,                            | , N=No, U=Unclea | r, N/A=Not appli      | cable                  |   |                     |  |  |   |                       |          |         |             |

Is it clear in the study what is the "cause" and what is the "effect"?
 Was there a control group?

- 3. Were participants included in any comparisons similar?
- 4. Were the participants included in any comparisons receiving similar treatment/care, other than the exposure or intervention of interest?
- 5. Were there multiple measurements of the outcome, both pre and post the intervention/exposure?
- 6. Were the outcomes of participants included in any comparisons measured in the same way?

#### Table 6: Quality appraisal results for randomised controlled trials

| INTERNAL VALIDITY Bias related to: |                 |                       |                          |   |  |   |   |   |                       |                                 |   |       |                           |    |    |                  |
|------------------------------------|-----------------|-----------------------|--------------------------|---|--|---|---|---|-----------------------|---------------------------------|---|-------|---------------------------|----|----|------------------|
|                                    |                 | Domain                | Selection and allocation |   | Administration of intervention/ exposure |   | Assessment, detection, and measurement of the outcome |   | Participant retention | Statistical conclusion validity |   | usion | Methodological<br>quality |    |    |                  |
|                                    |                 | Question no.          | 1                        | 2 | 3  | 4 | 5   | 6 | 7                     | 8                               | 9 | 10    | 11                        | 12 | 13 |                  |
| Study                              | Outcome         | Result at follow up   |                          |   |  |   |   |   |                       |                                 |   |       |                           |    |    |                  |
| Arman et<br>al. (2021)             | Quality of life | Post-<br>intervention | Y                        | U | Y  | N | N   | Y | N                     | Y                               | U | Y     | U                         | Y  | Y  | Moderate quality |
| Tokgoz et<br>al. (2022)            | Quality of life | Post-<br>intervention | Y                        | N | N  | N | N   | Y | Y                     | Y                               | N | N     | N                         | N  | Y  | Low quality      |
| Key: Y=Yes,                        | N=No, U=Unc     | lear, N/A=Not ar      | plicable                 |   |  |   |   |   |                       |                                 |   |       |                           |    |    |                  |

1. Was true randomisation used for assignment of participants to treatment groups?

2. Was allocation to treatment groups concealed?

3. Were treatment groups similar at baseline?

4. Were participants blind to treatment assignment?

5. Were those delivering the treatment blind to treatment assignment?

- 6. Were treatment groups treated identically other than the intervention of interest?
- 7. Were outcome assessors blind to treatment assignment?

8. Were outcomes measured in the same way for treatment groups?

- 9. Were outcomes measured in a reliable way?
- 10. Was follow up complete and if not, were differences between groups in terms of their follow up adequately described and analysed?
- 11. Were participants analysed in the groups to which they were randomised?
- 12. Was appropriate statistical analysis used?

13. Was the trial design appropriate and deviations from the standard RCT design accounted for in the conduct and analysis of the trial?

#### 6.4 Information available on request

The following are available on request: protocol; search strategies.

### 7. ADDITIONAL INFORMATION

#### 7.1 Conflicts of interest

The authors declare they have no conflicts of interest to report.

#### 7.2 Acknowledgements

The Public Health Wales team would like to thank Enzo Battista-Dowds, Maria Cole, Megan Elliot, Praveena Pemmasani and Libby Humphries for their time, expertise and contributions during stakeholder meetings, in guiding the focus of the review and interpretation of the findings.

### 8. APPENDIX

### **APPENDIX 1:**

Ovid MEDLINE(R) ALL <1946 to July 17, 2024>

- 1 (obese or obesity).ti,ab. 394914
- 2 Obesity/ or Obesity, Abdominal/ or Obesity, Morbid/ 255657
- 3 1 or 2 449482

4 (awaiting or waitlist\* or "wait\* list\*" or backlog\* or (wait\* adj3 (support\* or interven\* or assist\* or program\* or service\* or treat\* or surg\* or refer\* or specialist\* or consult\* or clinic\* or care or appointment\* or procedure\* or therapy or therapeutic or team\* or session\* or (weight adj3 manag\*) or (obesity adj3 manag\*)))).ti,ab. 44687

5 Waiting Lists/ or Case Management/ or Patient Care Management/ or Critical Pathways/ 37247

6 "Referral and Consultation"/ and (Disease Management/ or Obesity Management/) 569

- 7 4 or 5 or 6 74724
- 8 3 and 7 1348

9 limit 8 to (english language and yr="2010 -Current") 1064

10 9 not (case reports or comment or editorial or letter).pt. 1028

12224 1 Nor 240 (1932); this version posted November 7, 2024. The copyright holder for this review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY-ND 4.0 International license .



# The Health and Care Research Wales Evidence Centre

Our dedicated team works together with Welsh Government, the NHS, social care, research institutions and the public to deliver vital research to tackle health and social care challenges facing Wales.

Funded by Welsh Government, through Health and Care Research Wales, the Evidence Centre

answers key questions to improve health and social care policy and provision across Wales.

Along with our collaborating partners, we conduct reviews of existing evidence and new research, to inform policy and practice needs, with a focus on ensuring real-world impact and public benefit that reaches everyone.

Director: Professor Adrian Edwards

Associate Directors: Dr Alison Cooper, Dr Natalie Joseph-Williams, Dr Ruth Lewis



@EvidenceWales @tystiolaethcym



healthandcareevidence@cardiff.ac.uk



www.researchwalesevidencecentre.co.uk