1	Medical Students' Knowledge, Attitudes, and Motivation towards Antimicrobial					
2	Resistance Efforts in Eastern Uganda					
3 4 5	Jonathan Babuya ^{1,4} , Daniel Waruingi ^{3,4} , Douglas Mungujakisa ^{1,4} , Osmas Ahimbisibwe ¹ , Victoria Ruth Kako ¹ , Faith Aporu ¹ , Emmanuel Mugume ¹ , Julian Nyamupachitu ³ , Kenedy Kiyimba ²					
6						
7	¹ Faculty of Health Sciences, Busitema University, Mbale, Uganda					
8 9	² Department of Pharmacology and Therapeutics, Faculty of Health Sciences, Busitema University, Mbale Uganda					
10	³ ReAct-Action on Antibiotic Resistance Africa					
11	⁴ Zihi Institute					
12						
13						
14						
15						
16						
17						
18						
19						
20						
21						
22						
23						
24						
25						

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

26

Abstract

27 Introduction

To promote holistic learning about Antimicrobial Resistance (AMR), catalyze multidisciplinary engagement, and innovative AMR interventions, it is important that learning goes beyond the classroom and students embrace different extracurricular interventions. This study aimed to determine knowledge, attitudes, and motivations influencing medical students' engagement in AMR Club Initiatives at Busitema University in Uganda.

33 Methodology

This was descriptive cross-sectional study conducted at Busitema University among undergraduate student pursuing Bachelors of Medicine and Surgery, Bachelor of Nursing, and Bachelor of Science in Anesthesia and Critical care. A semi structured pre-tested questionnaire was shared among the study participants. Bloom cut-off method was used to analyse the knowledge of the participants, chi square test used for bivariate analysis and multivariable logistic regression used for determining factors independently associated to students' engagement in AMR club activities.

41 **Results**

42 71.5% of the 193 students had sufficient knowledge (determined using Bloom's cutoff 43 categories from 60% and above as sufficient and below 60% as insufficient) on AMR with an 44 average score of 68.18% (SD= ± 16.12). 90% of participants recognized the significance of 45 incorporating AMR within their curriculum, and 87.5% appreciated the need for training AMR 46 using a One Health ApproachThe most common reasons why students engaged in 47 extracurricular efforts to address AMR such as forming AMR clubs were found to be; influence 48 from peers (n= 42), university support (n=35), and inspiration from peer mentors' work (n=35).

49 **Conclusion**

50 The students were found to have a high level of knowledge and positive attitudes towards AMR 51 but reported the need for further in-depth training. Extra-curricular engagement such as 52 participation in an AMR Club was found to positively influence students' engagement in AMR 53 interventions.

54

56 Introduction

Antimicrobial Resistance (AMR) has been declared by the World Health Organization as one 57 58 of the top 10 global health threats[1]. Despite the high mortality and burden caused by AMR, it is largely ignored and rarely prioritized relative to other global health threats[2]. In 2019, 59 60 4.95 million deaths were associated with AMR globally, and 1.27 million deaths were caused 61 directly by resistant infections[3]. In Sub-Saharan Africa, bacterial AMR was associated with 62 1.05 million deaths and attributed to 250,000 deaths[4]. In Uganda, 7,100 deaths were directly 63 caused by bacterial drug-resistant infections, and 30,700 deaths were associated with drugresistant infections[4]. 64

65 Education and awareness play a very critical role in antimicrobial use and shape other healthseeking behaviors that influence AMR[5]. The Global Action Plan on AMR developed by the 66 67 World Health Organization (WHO) lists improving awareness and understanding of AMR as its first objective[6]. It highlights the need to raise awareness and promote behavioral change 68 69 through various ways such as public communication programs implemented from a One Health 70 realm targeting human health, consumers, animal health, and agricultural practices. It also 71 makes a case for inclusion of AMR into the curricula, and the professional training of health, 72 veterinary, and agriculture sectors to promote a better understanding among professionals. The 73 Uganda National Action Plan on AMR strategic objective one proposes the promotion of public 74 awareness, training, and education through two major strategies; improving public awareness 75 and supporting education and training of human, animal, plant, and environmental health through both pre-service and in-service training. Educating them very early in their training 76 77 and practice provides them with a different perspective enabling them to challenge existing 78 norms that undermine antimicrobial stewardship and also serve as AMR champions in their 79 respective facilities. There have been several efforts in Uganda to incorporate AMR into the 80 curriculum but they are yet to be successful[7].

A study conducted across East African Universities, reported that AMR was not emphasized in the medical curricula[8]. Comparable findings were reported in Ethiopia, Ghana and Zambia among nurses and pharmacy personnel and healthcare students' in regard to knowledge of Antibiotic use, AMR, and antimicrobial stewardship programs[9]. In Uganda, while the medical students were aware of the concept of AMR, they lacked in-depth knowledge necessary for effective antimicrobial stewardship practices[10]. A research conducted amongst

medical, pharmacy and nursing interns also showcased that only a small percentage of
participants had good knowledge of AMR and rational prescription practices[11].

89 To promote behavior change, and inspire action among healthcare students, engagement should 90 extend beyond traditional classroom settings to include holistic learning that incorporates both 91 curricular and extracurricular activities[12]. From early in their training, tertiary level students 92 should be empowered to engage in AMR. Extracurricular activities such as participation in 93 clubs have been shown to enhance intrinsic motivation, and active participation in efforts to 94 address various global health challenges [13]. To develop supportive frameworks where 95 healthcare students can engage actively and empower communities on AMR, it is best to understand their perspective and drive to engage in such initiatives[14]. 96

97 Currently, there is a paucity of data from African tertiary institutions on engagement of 98 healthcare students in extracurricular activities on AMR. It is important to have evidence which 99 can be harnessed to promote effective and beneficial extracurricular engagement among 100 healthcare students. This study aimed to determine knowledge and perceptions on AMR among 101 healthcare students and the motivation that influence healthcare students' participation in AMR 102 extracurricular activities such as clubs.

103 Methods and Materials

104 Study Design

105 This study was a descriptive cross-sectional study employing quantitative data collection 106 method to assess the level of awareness and knowledge of antimicrobial resistance and factors 107 influencing healthcare students' engagement in AMR club initiatives at Busitema University.

108 Study setting

The study was conducted at Busitema university, Faculty of Health Sciences in Mbale, Eastern
Uganda between December, 2023 and February, 2024.

111 **Study population**

112 These were undergraduate medical students pursuing Bachelor of Medicine and Bachelor of

113 Surgery, Bachelor of Science in Nursing, Bachelor of Science in Anesthesia and Critical care

114 at Busitema University.

115 Eligibility criteria.

- 116 The study included undergraduate students at Busitema University, Faculty of Health Sciences
- 117 who were 18 years of age or older pursuing either Bachelor of Medicine and Bachelor of
- 118 Surgery, or Bachelor of Science in Anesthesia and Critical Care, or Bachelor of Nursing
- 119 Science. The students who were not at the Faculty due to other academic programs, such as
- 120 anesthesia rotations or in their last semester of study were excluded.

121 Study variables.

122 Dependent variable

123 The dependent variable was students' engagement in AMR related activities.

124 Independent variables.

These included; socio-demographic characteristics like the age, sex, year of study and course
of study, AMR Awareness, Knowledge levels on AMR, and Extracurricular activities
engagement.

128 Data collection tool and procedures

We used a semi structured questionnaire which was self-developed and pretested before use. 129 We uploaded the questionnaire onto KoBo Toolbox. KoBo Toolbox is an open-source software 130 131 developed by the Harvard Humanitarian Initiative with support from United Nations agencies, CISCO, and partners to support data management by researchers and humanitarian 132 133 organizations (https://www.kobotoolbox.org/). The servers are secure and encrypted with 134 strong safe guards and protection against data loss. Participants who consented to the study had 135 the questionnaire link sent to them via a WhatsApp message. The data collection tools were 136 pretested before actual data collection.

137 Sample size estimation and sampling technique.

- 138 The sample size was determined using the formula by Fisher et al. (1998): 139 $n=(Z2 \times p(1-p))/d2$
- 140 Where: n= required sample size, Z = standard normal deviation at a 95% confidence interval
- 141 (1.96), p = prevalence (0.5 due to no similar study found), d = margin of error or degree of
- 142 accuracy (0.05)

143 After calculations, n=384.16, rounded to 385 participants. Since the population was less than 144 10,000, the finite population correction formula was applied. With a population size of 500 145 students, the final sample size was determined to be 218 study participants.

146 A list of all the students at the faculty was obtained, grouped as per the year of study. Random 147 numbers were assigned to the names, they were then filled in a random selector program, that 148 generated a random list of the numbers to which the names were attached. The selected students 149 were then approached via a phone call or WhatsApp and given information about the study. 150 Those that consented to participate in the study were recruited and the link to the questionnaire 151 was sent to them. The reasons for engagement in AMR activities were first collected through 152 a pre-study survey among 20 students to populate the questionnaire since we could not find 153 any qualitative study that had been done about these reasons.

154 Data analysis and presentation

155 The data collected was meticulously cleaned and analyzed using STATA version 15.0. Categorical data was analyzed using frequencies and percentages, while numerical data was 156 157 analyzed using measures of Central Tendency and Dispersion. Pie charts were used to visually represent categorical data such as gender distribution, age groups, study programs, and year of 158 159 study while bar graphs used to illustrate comparisons, such as students' engagement in 160 extracurricular activities by gender, knowledge levels about AMR across different programs, 161 and motivations for engaging in AMR club activities. Bivariate analysis investigated factors 162 associated with students' engagement in AMR activities i.e. age groups, study program, year 163 of study, knowledge about AMR, awareness about AMR, and participation in extracurricular activities and presented in tables while multivariable analysis further examined the significant 164 165 factors identified in bivariate analysis to determine their combined impact on students' engagement in AMR activities and presented in figures including pie charts, bar graphs, and 166 167 other visual aids to succinctly display key findings.

168 **Ethical consideration.**

The study was conducted according to the Declaration of Helsinki. Ethical approval was sought and obtained from Busitema University Research Ethics committee under the reference number (BUFHS-2023-104). Written informed consent was obtained from the participants before recruitment into the study. The data collected was kept in a locked folder in the computer which was accessible only to the investigators.

174 **Results**

175 Social demographics of the participants

176 A total of 193 students participated in the study. The majority of respondents were male (58.0%), while females comprised 42.0% of the participants. Over half of the respondents 177 178 (51.3%) were below the age of 25, 40.4% were aged between 25-34, and a small proportion 179 (8.3%) were above 35. The largest group of respondents were enrolled in the Bachelor of 180 Medicine and Bachelor of Surgery program (63.2%). The Bachelor of Nursing Science program had 19.7% of respondents, and the Bachelor of Science in Anesthesia and Critical 181 182 Care program had 17.1%. The distribution across years of study was relatively even, with the highest percentage in Year 4 (24.4%) and the lowest in Year 2 (17.1%). Year 1, Year 3, and 183

- 184 Year 5 had 19.7%, 20.2%, and 18.7% of respondents, respectively(**Tabel 1**).
- **Table 1: Showing the demographic statistics of the the study participants**

	n (%)
Gender	
Female	81 (42.0%)
Male	112 (58.0%)
Age	
Below 25	99 (51.3%)
25-34	78 (40.4%)
Above 35	16 (8.3%)
Study program	
Bachelor of Medicine and Bachelor of Surgery	122 (63.2%)
Bachelor of Nursing Science	38 (19.7%)
Bachelor of Science in Anesthesia and Critical Care	33 (17.1%)
Year of study	
Year 1	38 (19.7%)
Year 2	33 (17.1%)
Year 3	39 (20.2%)
Year 4	47 (24.4%)
Year 5	36 (18.7%)

186 n=193

187 Students' engagement in extracurricular activities

Extracurricular activities, in the context of this study, refer to general institutional activities outside the classroom that students participate in. The study found a difference in extracurricular engagement between genders: 82.05% of female students participated in extracurricular activities at the university, compared to 95.5% of male students (Figure 1).

192

Figure 1: Showing students engagement in extracurricular activities; A; female, B: Male

193

194 Students' engagement in AMR club activities

AMR clubs, which are among extracurricular activities, focus specifically on interventions related to antimicrobial resistance. At Busitema University, 67.5% of female participants reported not having participated in AMR club activities, compared to 58.18% of male students(**Figure 2**).

- 199
- 200

Figure 2: Showing students engagement in AMR activities, A: Female, B: Male

201 Levels of knowledge about AMR among the students

202 Overall, 71.5% of the students had sufficient knowledge about AMR (determined using 203 Bloom's cutoff categories from 60% and above as sufficient and below 60% as insufficient). 204 The percentage of students with sufficient AMR knowledge increased with each year of study, 205 starting at 55.26% in Year 1 and rising to 91.67% in Year 5(Figure 3). Across the different 206 programs of study, Bachelor of Medicine and Bachelor of Surgery had the highest percentage, 207 76.23% of students with sufficient knowledge about AMR which was followed by Bachelor of 208 Science in Anesthesia and Critical Care, 69.7% and Bachelor of Nursing Science at 57.89% 209 (Figure 4).

- 210
- 211
- 212
- 213 Figure 3 Showing knowledge levels across different years of study
- 214 Figure 4: Showing knowledge levels of students in different study programs
- 215
- 216
- 217
- 218

219 Attitudes of students towards AMR

- A total of 8.9 % (n=17) of the study participants agreed that agreed that purchasing antibiotics without prescription from pharmacies was acceptable. Over a quarter of the participants (35.3%, n=68,) agreed to not being at risk of getting an antibiotic-resistant infection as long as they took their antibiotics correctly. Furthermore, an overwhelming majority (83.4%, n=160) agreed that prescribing broad-spectrum antimicrobials instead of narrower spectrum options contributes to antimicrobial resistance.
- Majority of the participants , 96.4% (n=186) acknowledged the critical importance of having a strong understanding of AMR in their medical careers. A substantial percentage 87.5% (n=169) agreed that poor infection control practices by healthcare professionals significantly contribute to the spread of antimicrobial resistance. Similarly, more than 80% of the students (n=164,
- 85%) believed that the excessive use of antimicrobials in livestock contributes to the problemof antimicrobial resistance.
- Regarding their practical experiences, 73.6% (n=142) of the students reported observing instances where antimicrobials were overused during their hospital rotations. Looking towards the future, over 40% (41.4%, n=80) of the participants expressed optimism that new antimicrobial agents capable of addressing resistance issues would likely be developed. These results are summarized in **Figure 5**.
- 237
- 238
- 239
- 240
- 241
- 242
- 243
- 244

²⁴⁶ Figure 5: Showing students' attitudes towards AMR: SD-Strongly disagree, D-Disagree, N- Neutral, A-Agree, SA- Strongly Agree

porpotate	<i>.</i>
It is made available under a CC-B	Y 4.0 International license .

247	
248	
249	Sources of information about AMR
250 251	Academic learning platforms were the major sources of information regarding AMR(n=152). (Figure 6).
252	
253	Figure 6: Sources of AMR information used by study participants
254	
255	
256	
257	

258 Motivations to students for engaging in AMR club activities

The largest proportion of students (n=42) were those motivated by peer to engage in AMR club activities while the least number (n=4) engaged in AMR activities to get information (*Figure* 7).

- Figure 7: Showing reasons for student engagement in AMR activities
- 263

A higher proportion of the students (59%, n=114) students engaged in extracurricular activities like clubs to create friendships and to exercise and gain physical fitness while the least number

- 266 of the students engaged in extracurricular activities due to other reasons (n=5) which included
- 267 inspiring others and getting knowledge for the future (Table 2.)

268 Table 2: Reasons for engaging in Extracurricular activities by the study particpants

	Number of students who
	selected
	77
To create friendships	114
To exercise and gain physical fitness	114
To develop leadership skills	70
To pass time	51
To contribute to well-being of society	66
To improve my CV	43
To get mentored into my future career	61
To improve job prospects after completing	
school	41
Other reasons	5

269

271 Factors associated with student participation in AMR club activities

- Age, gender and program of study were not significantly associated with student participation
- in AMR club activities independently. Third year students had 3.12 times higher odds of
- engaging in AMR club activities than second year students and the result was statistically
- 275 significant(cOR: 3.12, CI= 1.13-8.66, p=0.028).
- 276 Having sufficient knowledge about AMR had 2.21 times higher odds of engaging in AMR
- 277 club activities than having insufficient knowledge and the result was statistically significant
- 278 (cOR: 2.21, CI:1.10-4.42, p=0.026). Students who were aware about AMR had 3.24 times
- 279 higher odds of engaging in AMR club activities than those who were not aware and the result
- was statistically significant(cOR:3.24, CI:1.17-8.96, p=0.023).
- 281 Students engaging in extracurricular activities had 12.55 times higher odds of engaging in
- 282 AMR club activities than those who were engaging in extracurricular activities and the result
- was statistically significant(cOR: 12.55, CI: 1.64-96.19, p=0.015).
- However, after adjusting for confounders, student's engagement in extracurricular activities
- had 14.45 times higher odds of engaging in AMR club activities than those who did not and
- the result was statistically significant (AOR: 14.45, CI: 1.84-113.52, p=0.011)(**Table 3**)
- 287
- 288
- 289
- 290
- 291
- 292
- 293
- 294
- ____
- 295

Table 3: Showing bivariate and multivariable analysis for the factors affecting student

engagement in AMR.

Independent	Categories	Student	ts' engagement in AMR club ac	tivities	
variables		р-	Crude Odds Ratios(95%CI)	p-values	Adjusted Odds Ratios(95%CI)
		values			
Gender	Male	0.192	1.49(0.82-2.73)		
	Female	Ref	Ref		
Age(years)	Below 25	0.105	3.00(0.80-11.31)		
	25-34	0.263	2.16(0.56-8.32)		
	Above 35	Ref	Ref		
Study program	Medicine	0.745	1.15(0.51-2.59)		
	and Surgery				
	Nursing	0.509	1.39(0.52-3.67)		
	Science				
	Anesthesia	Ref	Ref		
	and Critical				
	Care				
Voor of study	Voor 1	0.255	1 82(0 65 5 12)	0.401	1 60(0 52 4 90)
rear of study	Year 2	0.233	1.82(0.03-3.12)	0.401	1.00(0.55-4.60)
	Year 2	Rej	<i>Rej</i>	Rej	<i>Rej</i>
	* Y ear 3	0.028	3.12(1.13-8.66)	0.084	2.6/(0.88-8.15)
	Year 4	0.095	2.31(0.87-6.19)	0.300	1.78(0.60-5.29)
	Year 5	0.463	1.49(0.51-4.37)	0.973	1.02(0.32-3.24)
Knowledge	Insufficient	Ref	Ref	Ref	Ref
levels	*Sufficient	0.026	2.21(1.10-4.42)	0.451	1.40(0.58-3.38)
AMR	Not aware	Ref	Ref	Ref	Ref
Awareness	*Aware	0.023	3.24(1.17-8.96)	0.191	2.34(0.65-8.41)
Extracurricular	No	Ref	Ref	Ref	Ref
activities	*Yes	0.015	12.55(1.64-96.19)	0.011	14.45(1.84-113.52)
engagement					

304 **Discussion**

305 The study findings indicate that a higher proportion of the participants (95.5% of the male 306 students, 82% of the female students) participated in extracurricular activities with students in 307 lower academic years exhibiting lower knowledge levels about antimicrobial resistance 308 compared to their fifth-year counterparts. These findings are consistent with a previous study, 309 which reported higher knowledge levels among clinical students. Additionally, a study in 310 Ghana showed a sequential increase in AMR knowledge across the years of study, suggesting 311 that students were progressively exposed to more AMR concepts as they advanced in their 312 education[10, 15]. Slightly lower levels of AMR knowledge among students have been 313 reported in various studies across Africa For instance, a study in the Democratic Republic of 314 Congo and Ethiopia found that only 55% of students had sufficient knowledge about AMR, 315 while in Zambia, 70% of students had adequate knowledge levels [9, 16, 17]. These studies 316 suggest a significant knowledge gap among medical students about AMR across the continent and attributed to a diverse factors in different countries[10]. Therefore, there is a critical need 317 318 for improved education on AMR among medical students in Uganda. Incorporating AMR 319 education into the curriculum can ensure a large number of students are well-informed through 320 a sustainable strategy.

The participants demonstrated a positive attitude towards the rational use of antibiotics, with the majority (87%, n= 168) disagreeing that it is appropriate to purchase antibiotics without a prescription. In contrast, a poor attitude was observed among participants in Sudan, where over 59.6% believed it was acceptable to buy medications from a pharmacy without a prescription[18]. Over a half of the participants demonstrated a poor understanding on the spread of drug-resistant infections

327 Less than 50% of the students (n=68, 35.3%) agreed to being at risk of getting an antibiotic-328 resistant infection as long as they took their antibiotics correctly showcasing little 329 understanding on spread of drug-resistant infections, while less than 83.1% that agreed that 330 antimicrobial resistance affects animal health and production and human health-related science 331 students had a positive feeling that inappropriate use of antimicrobials is one of the reasons for 332 the occurrence of antimicrobial resistance[19]. 83.4% (n=160) agreed that prescribing broad 333 spectrum antimicrobials when narrower spectrum antimicrobials are available increases 334 antimicrobial resistance showcasing high understanding on prescribing practices. More than 90% of the students(n=186, 96.4%) agreed that a strong knowledge of antimicrobials is 335

336 important in the medical career similar to other studies which shows that the overarching 337 recognition on the need to have strong knowledge about AMR as health professionals enabling 338 them to make better decisions for optimal health care delivery[10].

From the findings, 80% of the students (n=164, 85%) agreed that excessive use of 339 340 antimicrobials in livestock causes antimicrobial resistance similar to other studies which shows 341 appreciation on the role the one health approach in fight antimicrobial resistance by recognizing 342 other sectors' importance. About two thirds of the students (n=142, 73.6%) agreed that 343 antimicrobials were overused at the hospital where they had rotated compared to 36% from a 344 previous study[20]. This shows a better attitude towards recognizing misuse as better stewards of antimicrobials. 41.4% of the students (n=80, 41.4%) agreed that new antimicrobials will be 345 developed in the future that will keep up with the problem of resistance showcasing poor 346 347 understanding on the complexity of AMR, and the multifaceted approach required.

348 Students' engagement in extracurricular activities was associated with engagement in AMR 349 club activities even after adjusting for other factors because since AMR club was an 350 extracurricular activity alongside the others, it was easier for the students already engaged in 351 the other activities to take on AMR club activities.

352 Most students engaged in AMR club activities due to influence from peers, university support 353 and mentors while the least engaged to get information and learning. This is mainly because 354 students can easily access their peers and are more inspired and influenced by the actions of 355 people within the same bracket[21].

356 Conclusion

357 In conclusion, the study reports a sequential increase in knowledge levels on AMR across the

358 years of study amongst the study participants. The study showed that healthcare students

- 359 appreciated the need and importance of AMR training through a One Health Approach and
- 360 the importance of good infection prevention and control measures in addressing AMR.
- 361 Influence from peer, university support and inspiration by peer mentors work were found to
- 362 be the major reasons behind students' engagement in AMR related extracurricular activities.

363 **Recommendations**

364 From our findings, we recommend that, an AMR and AMS course unit should be incorporated into the preservice training of healthcare students so that there is a greater understanding of 365 366 Antimicrobial Resistance and the topic and potential mitigation efforts. Furthermore, there is

367 need to emphasize on the value of extracurricular engagement in AMR among healthcare 368 students and to leverage on peer support and mentorship to promote holistic learning and 369 effective community engagement in AMR. Lastly, we recommend a broader study 370 incorporating multiple institutions in investigating the drivers and barriers of healthcare 371 students in extracurricular AMR engagements to obtain generalizable results that can inform 372 the country's AMR policies.

373 Limitations

374375

 The study was done in one university therefore it may be difficult to generalize the findings at a country level.

376 Acknowledgements

377 The authors thank all the students who participated in the study for their feedback. We are

378 grateful the Foundation to Prevent Antibiotic Resistance for grant awarded to support this

379 project that was aimed at catalyzing grassroot engagement of tertiary level students in

380 different countries in Africa. We thank Busitema University Antimicrobial Resistance and

381 Stewardship Club and Busitema University Research and Innovation Association for

allowing us to use their structures to implement the study. We thank ReAct (Action on

- 383 Antibiotic Resistance) Africa and Zihi Institute for the technical support in implementation of
- this study.

385 Funding

A part of this study was funded by the Foundation to Prevent Antibiotic Resistance in a grant that run from 2022 – 2023 offered to Students Against Superbugs Africa (now Zihi Institute) in collaboration with ReAct Africa. The funders had no role in: the design and conduct of the study; data collection; data management, and analysis; review of the manuscript; and the decision to submit the manuscript for publication.

391 Competing interests

392 All authors have nothing to declare.

393

394

397

References

398 399	1. World Health Organisation. Antimicrobial Resistance: WHO; 2024 [cited 2024 25th October] Available from: https://www.who.int/docs/default-source/antimicrobial-
400	resistance/amr-factsheet ndf
401	2 Coque TM Canton R Perez-Cobas AE Fernandez-de-Bobadilla MD Baquero F
402	Antimicrobial Resistance in the Global Health Network: Known Unknowns and Challenges
403	for Efficient Responses in the 21st Century Microorganisms 2023:11(4):1050 Epub
404	20230417 doi: 10.3390/microorganisms11041050 PubMed PMID: 37110473: PubMed
405	Central PMCID: PMCPMC10144039
406	3 Murray CIL, Ikuta KS, Sharara F, Swetschinski L, Robles Aguilar G, Grav A, et al
407	Global burden of bacterial antimicrobial resistance in 2019 ⁻ a systematic analysis. The
408	Lancet, 2022;399(10325);629-55, doi: 10.1016/S0140-6736(21)02724-0.
409	4. Sartorius B. Grav AP. Weaver ND. Aguilar GR. Swetschinski LR. Ikuta KS. et al.
410	The burden of bacterial antimicrobial resistance in the WHO African region in 2019; a cross-
411	country systematic analysis. The Lancet Global Health. 2024:12(2):e201-e16.
412	5. Fuller W, Kapona O, Aboderin AO, Adeyemo AT, Olatunbosun OI, Gahimbare L, et
413	al. Education and Awareness on Antimicrobial Resistance in the WHO African Region: A
414	Systematic Review. Antibiotics. 2023;12(11):1613. doi: 10.3390/antibiotics12111613.
415	6. Team EE. WHO member states adopt global action plan on antimicrobial resistance.
416	Eurosurveillance. 2015;20(21):21137.
417	7. Waswa J, Kiggundu R, Joshi MP, Mpagi J, Kasujja H, Murungi M, et al. Addressing
418	gaps in AMR awareness in the public: an evidence-based policy brief to guide school
419	curriculum review in Uganda. Frontiers in Public Health. 2023;11:1287523.
420	8. Ampaire L, Muhindo A, Orikiriza P, Mwanga-Amumpaire J, Bebell L, Boum Y. A
421	review of antimicrobial resistance in East Africa. Afr J Lab Med. 2016;5(1):432. doi:
422	10.4102/ajlm.v5i1.432.
423	9. Tembo N, Mudenda S, Banda M, Chileshe M, Matafwali S. Knowledge, attitudes and
424	practices on antimicrobial resistance among pharmacy personnel and nurses at a tertiary
425	hospital in Ndola, Zambia: implications for antimicrobial stewardship programmes. JAC-
426	Antimicrobial Resistance. 2022;4(5):dlac107. doi: 10.1093/jacamr/dlac107.
427	10. Kanyike AM, Olum R, Kajjimu J, Owembabazi S, Ojilong D, Nassozi DR, et al.
428	Antimicrobial resistance and rational use of medicine: knowledge, perceptions, and training
429	of clinical health professions students in Uganda. Antimicrob Resist Infect Control.
430	2022;11(1):145. doi: 10.1186/s13756-022-01186-9.
431	11. Nabidda S, Ssennyonjo R, Atwaru J, Kanyike AM, Baryayaka S, Pangholi K, et al.
432	Antimicrobial resistance and rational prescription practices: knowledge, perceptions and
433	confidence of health profession interns in Uganda. JAC-Antimicrobial Resistance.
434	2023;5(5):diad105.
435	12. Munir S, Zaneer M. The role of extra-curricular activities in increasing student
430	engagement. Asian Association of Open Universities Journal. 2021;16(3):241-54.
43/	15. Achar Fujii Kin, Kobayasi K, Claassen Enns S, Zen Tempski P. Medical Students
430 120	Cualitative Study, Adv Med Educ Pract, 2022;12:1122, 41, doi: 10.2147/AMED.S250047
439 440	Quantative Study. Adv Wed Educ Flact. 2022,15.1155-41. doi: 10.214//AWEP.555904/.
<u>440</u> <u>44</u> 1	are the motives behind their involvement? International Journal of Psychology
<u>1</u> 11 <u>1</u> 12	2013.48(5).871.80 doi: 10.1080/00207504.2012.602703
44 <i>2</i>	$2013, \pm 0(3).071^{-}00.$ u 01. 10.1000/0020737 $\pm .2012.07273$.

- Sefah IA, Akwaboah E, Sarkodie E, Godman B, Meyer JC. Evaluation of Healthcare
 Students' Knowledge on Antibiotic Use, Antimicrobial Resistance and Antimicrobial
 Stewardship Programs and Associated Factors in a Tertiary University in Ghana: Findings
- 446 and Implications. Antibiotics. 2022;11(12):1679. doi: 10.3390/antibiotics11121679.
- 447 16. Bogale AA, Amhare AF, Chang J, Bogale HA, Betaw ST, Gebrehiwot NT, et al.
- 448 Knowledge, attitude, and practice of self-medication with antibiotics among community
- 449 residents in Addis Ababa, Ethiopia. Expert Review of Anti-infective Therapy.
- 450 2019;17(6):459-66. doi: 10.1080/14787210.2019.1620105.
- 451 17. Thriemer K, Katuala Y, Batoko B, Alworonga J-P, Devlieger H, Van Geet C, et al.
- 452 Antibiotic Prescribing in DR Congo: A Knowledge, Attitude and Practice Survey among
- 453 Medical Doctors and Students. PLoS ONE. 2013;8(2):e55495. doi:
- 454 10.1371/journal.pone.0055495.
- 455 18. Ibrahim Ahmed MA. Antibacterial Agents: Knowledge and Attitudes of Sudanese
- 456 College of Medicine Undergraduates. IJCMCR. 2023;30(1). doi:
- 457 10.46998/IJCMCR.2023.30.000729.
- 458 19. Abunna F, Gebresenbet G, Megersa B. Knowledge, attitude, and practices (KAP) of
- 459 Univer- sity students regarding antimicrobial use (AMU) and antimicrobial resistance (AMR)460 in Ethiopia. 2023.
- 461 20. Minen MT, Duquaine D, Marx MA, Weiss D. A Survey of Knowledge, Attitudes, and
- 462 Beliefs of Medical Students Concerning Antimicrobial Use and Resistance. Microbial Drug
- 463 Resistance. 2010;16(4):285-9. doi: 10.1089/mdr.2010.0009.
- 464 21. Qureshi MA, Khaskheli A, Qureshi JA, Raza SA, Yousufi SQ. Factors affecting
- students' learning performance through collaborative learning and engagement. InteractiveLearning Environments. 2023;31(4):2371-91.



Figure 1: Showing students engagement in extracurricular activities; A; female, B: Male



Figure 2: Showing students engagement in AMR activities, A: Female, B: Male



Figure 3 Showing knowledge levels across different years of study



Medicine	Nursing Science	Anestnesia and Critical	
		Care	

Figure 4: Showing knowledge levels of students in different study programs



Figure 5: Showing students' attitudes towards AMR: SD-Strongly disagree, D-Disagree, N- Neutral, A-Agree, SA- Strongly Agree





Figure 7: Showing reasons for student engagement in AMR activities