It is made available under a CC-BY-NC 4.0 International license .

Genome-wide association studies and multi-omics integrative analysis reveal novel loci and their molecular mechanisms for circulating polyunsaturated, monounsaturated, and saturated fatty acids

Yitang Sun¹, Huifang Xu¹, Kaixiong Ye^{1,2,*}

¹Department of Genetics, University of Georgia, Athens, GA, USA ²Institute of Bioinformatics, University of Georgia, Athens, GA, USA

*Correspondence: Dr. Kaixiong Ye (Kaixiong.Ye@uga.edu)

It is made available under a CC-BY-NC 4.0 International license .

1 Abstract

Background: Dietary and circulating levels of polyunsaturated fatty acids (PUFAs) have been associated with mortality and risks of various diseases. The circulating levels of fatty acids (FAs) exhibit heterogeneity across individuals due to genetic and environmental influences. Although previous genome-wide association studies (GWAS) have identified genetic loci associated with FAs, the biological mechanisms of these genetic associations remain largely unexplored.

7 Methods: Here, we conducted GWAS to identify additional genetic loci for 19 circulating fatty 8 acid (FA) traits, quantified by nuclear magnetic resonance spectroscopy, in UK Biobank 9 participants of European ancestry (N = 239,268) and five other ancestries (N = 508 - 4,663). We 10 leveraged the GWAS findings to characterize genetic correlations and colocalized regions among 11 FAs, explore sex differences, examine FA loci influenced by lipoprotein metabolism, and apply 12 statistical fine-mapping to pinpoint putative causal variants. The integrative analysis of GWAS 13 and multi-omics quantitative trait loci (QTL) reveals intermediate molecular phenotypes mediating 14 the associations between the genetic loci and FA levels.

15 Results: Altogether, we identified 215 significant loci for PUFAs-related traits in European 16 participants, 163 loci for monounsaturated fatty acids (MUFAs), and 119 loci for saturated fatty 17 acids (SFAs), including 70 novel loci for PUFAs, 61 for MUFAs, and 54 for SFAs. Of these 18 significant loci, a novel locus for total FAs, omega-6 PUFAs relative to total FAs, and total 19 MUFAs (around genes GSTT1/2/2B) overlapped with QTL signals for all six molecular 20 phenotypes examined, including gene expression, protein abundance, DNA methylation, splicing, 21 histone modification, and chromatin accessibility. Across FA traits, 65% of GWAS loci 22 overlapped with QTL signals for at least one molecular phenotype.

- 23 Conclusions: Our study identifies additional novel genetic loci and provides the first evidence
- 24 uncovering molecular mechanisms underlying FAs-relevant variants. These findings offer insights
- 25 that could facilitate patient stratification for precision nutrition.
- 26
- 27 Keywords: GWAS, polyunsaturated fatty acids, fatty acids, xQTL, multi-omics

It is made available under a CC-BY-NC 4.0 International license .

28 Introduction

29 Polyunsaturated fatty acids (PUFAs), monounsaturated fatty acids (MUFAs), and saturated fatty acids (SFAs) have been implicated in various diseases.¹⁻³ Evidence from epidemiological and 30 31 genetic studies indicates that PUFAs, especially omega-3 PUFAs, are associated with reduced 32 risks of various diseases or mortality by modifying metabolism and inflammation.⁴⁻⁷ Circulating 33 fatty acid (FA) levels are influenced by environmental and genetic factors and characterized by sex- and ancestry-specific patterns.⁸⁻¹⁰ Dietary intake, socioeconomic status, physical inactivity, 34 cigarette smoking, and alcohol consumption are common environmental factors that influence FA 35 36 levels.¹¹ Although hundreds of genetic loci associated with FAs have been identified by genomewide association studies (GWAS),¹²⁻³⁷ the biological mechanisms of these genetic associations 37 38 remain largely unexplored. Moreover, these genetic loci were mainly discovered in the European 39 (EUR) population and were focused on autosomes. Extending GWAS to non-EUR populations 40 and sex chromosomes will offer a comprehensive understanding of the genetic architecture of 41 circulating FAs.

42 In this study, we performed GWAS for 19 FA traits related to PUFAs, MUFAs, and SFAs 43 to identify more novel genetic loci. We conducted overall and sex-specific GWAS using 250,101 44 individuals of EUR ancestry and five other ancestries from the UK Biobank (UKB). Leveraging 45 the GWAS summary statistics, we characterized heritability for these FA traits and examined their 46 shared genetic basis with genetic correlation and colocalization analysis. Candidate causal variants 47 were pinpointed with statistical fine-mapping. Finally, we integrated GWAS signals with multi-48 omics data to identify the mediating molecular phenotypes and provide mechanistic insights into 49 the molecular mechanisms underlying the GWAS signals.

It is made available under a CC-BY-NC 4.0 International license .

50 Materials and Methods

51 **Participants**

52 The UKB cohort is a prospective population-based study of ~500,000 participants from 53 across the United Kingdom, aged between 37 and 73 years at recruitment from 2006 to 2010.³⁸ Of 54 the individuals with both phenotype and genotyping data, we excluded participants who have 55 withdrawn consent, mismatched information between phenotypic and genetic sex, sex 56 chromosome aneuploidy, or are outliers for heterogeneity and missing genotype rate. Genetic 57 ancestry groups have been previously defined in the Pan-ancestry genetic analysis of the UK 58 Biobank (Pan-UKB).³⁹ Together, we included up to 239,268 participants of EUR ancestry in this 59 study. We also included participants of African (AFR) (N=3,352), admixed American (AMR) 60 (N = 508), Central/South Asian (CSA) (N = 4,663), East Asian (EAS) (N = 1,445), and Middle 61 Eastern (MID) (N = 865) ancestries. Table S1 provides the overall and sex-stratified characteristics 62 of participants in each ancestry group. This study received ethical approval from the National 63 Health Service North West Centre for Research Ethics Committee, and all participants provided 64 informed consent. Data from the UKB resource were accessed under application number 48818.

65 Phenotypes – circulating fatty acids (FAs)

Plasma FAs were measured in a randomly selected subset of 274,020 participants by Nightingale Health using nuclear magnetic resonance (NMR) spectroscopy.⁴⁰ All FA levels and their ratios were normalized within each ancestry group using the rank-based inverse normal transformation in overall GWAS.⁴¹ This inverse normal transformation was applied separately for males and females to account for sex-specific differences in FA distributions. In total, 20 metabolic measurements were analyzed in the GWAS including total FAs, 5 PUFA absolute concentrations (i.e., PUFAs, omega-3 PUFAs, omega-6 PUFAs, docosahexaenoic acid (DHA), and linoleic acid

73 (LA)), their relative percentages in total FAs (i.e., PUFAs%, omega-3%, omega-6%, DHA%, and 74 LA%), the ratio of omega-3 to omega-6 PUFAs (omega-3/omega-6), the ratio of omega-6 to 75 omega-3 PUFAs (omega-6/omega-3), monounsaturated fatty acids (MUFAs), saturated fatty acids 76 (SFAs), MUFAs%, SFAs%, PUFAs/MUFAs, PUFAs/SFAs, and degree of unsaturation. Both the 77 omega-3/omega-6 ratio and omega-6/omega-3 ratio were included in our GWAS due to different 78 preferences in the field, but they revealed the same genetic loci in opposite effect directions. In 79 additional analyses, we focused on omega-3/omega-6 and did not include omega-6/omega-3, resulting in a total of 19 plasma metabolic measurements. 80

81 Genotyping and quality control

Full details about genotyping, imputation, and genotype-based quality control (QC) have been described elsewhere.⁴² Briefly, we excluded variants with imputation quality score < 0.3, minor allele frequency < 0.001, genotype missingness per variant > 0.05, and Hardy-Weinberg test *P*-value < 1×10^{-8} . We also confirmed that no individuals have genotype missingness > 0.05. For the non-pseudoautosomal region of the X chromosome (GRCh37; chrX: 2699520-154931044), males were treated as homozygotes of the reference or effect allele, and coded as 0 or 2, respectively.

89 Genome-wide association studies (GWAS)

Overall GWAS were conducted for each ancestry group and followed by sex-stratified GWAS. We carried out GWAS in participants of EUR ancestry using fastGWA from the GCTA toolbox (v.1.94.1), which controls for familial relatedness using a sparse genetic relationship matrix (GRM) with a default threshold of 0.05 based on slightly linkage disequilibrium (LD)pruned HapMap3 variants.^{43,44} Considering the relatively small sample size for other ancestry groups (AFR, AMR, CSA, EAS, and MID), the mixed-linear-model association (MLMA) method

It is made available under a CC-BY-NC 4.0 International license .

96 was used to perform GWAS using a dense GRM.⁴⁵ Additionally, sex-specific GWAS were 97 conducted for each ancestry group using fastGWA for EUR participants and MLMA for other 98 ancestry groups, excluding sex as a covariate from all models.

99 To explore the influences of potential covariates, we performed GWAS in the EUR 100 ancestry participants using 3 models. In model 1, we included age, sex, age \times sex, and the top 10 101 genetic principal components derived from the Pan-UKB as covariates. Model 2 added additional 102 covariates, including body mass index, Townsend deprivation index, smoking status, alcohol status, physical activity, and statin use.^{46,47} In model 3, we adjusted for fish oil supplementation 103 status and lipoprotein lipids including chylomicron, very-low-density lipoprotein (VLDL), 104 105 intermediate-density lipoprotein (IDL), low-density lipoprotein (LDL), high-density lipoprotein 106 (HDL) cholesterols as additional covariates. For GWAS in the non-EUR population, we applied 107 models 1 and 2.

108 Identification of novel loci

To identify novel loci, we compared our independent loci to the previously reported FA loci in the GWAS Catalog and in other relevant publications identified by literature review up until June 2024.⁴⁸ Independent loci were identified in our results or previous GWAS based on genomewide significance threshold (*P*-value $< 5 \times 10^{-8}$) and LD-based clumping ($r^2 = 0.1$, window size = 250 kb) using PLINK.^{49,50} Independent loci were merged based on proximity (±250 kb). We annotated the lead variants for their target genes and functional consequences using the Ensemble GRCh37 Variant Effect Predictor (VEP).⁵¹

116 Heritability and genetic correlation analyses

Heritability and genetic correlations were estimated with linkage disequilibrium score
 regression (LDSC) in the EUR ancestry subset.^{52,53} In-sample LD scores were derived from

It is made available under a CC-BY-NC 4.0 International license .

119 <u>s3://broad-alkesgroup-ukbb-ld/UKBB_LD/</u>.⁵⁴ As recommended, we checked for inflation using 120 the attenuation ratio in the EUR population and genomic inflation factor (λ_{GC}) in other 121 populations.⁵⁵

122 Colocalization with FA traits

We performed colocalization analysis with HyPrColoc (v.1.0) to assess whether two or more FA traits share a putative causal variant.⁵⁶ Pairwise and multi-trait colocalization analyses were conducted for each pair or all FA traits altogether. We used the default priors, including the probability of a variant being associated with a single trait (prior.1 = 1×10^{-4}) and the conditional probability of association with an additional trait (prior.c = 0.02). A posterior probability (PP) > 0.8 was considered as evidence for a shared causal variant.

129 Statistical fine-mapping of GWAS loci

To identify the most plausible causal variants for genome-wide significant loci associated with each FA trait, we used SuSiE (v.0.12.27).⁵⁷ We used the 1 Mb region surrounding the lead variant of each genetic locus for statistical fine-mapping. We utilized in-sample LD information to estimate the correlations among SNPs. The 95% credible set identified potential causal variants based on their PP. Only variants with a PP > 0.8 were reported, with higher PP values closer to 1 indicating more likely to be causal variants.

136 Gene-based and gene-set analyses

137 MAGMA v.1.08, implemented in FUMA, was used to perform gene-based and gene set 138 analyses.^{58,59} In the gene-based analysis, 19,665 SNPs were linked to protein-coding genes, and 139 Bonferroni correction was applied to establish the significance threshold (P < 0.05/19,665 = 2.54140 × 10⁻⁶). We determined the enrichment of candidate FA-associated genes in specific biological 141 pathways, cellular components, or molecular functions. A total of 15,481 gene sets (5,497 curated

It is made available under a CC-BY-NC 4.0 International license .

142 gene sets and 9,984 gene ontology terms) obtained from MsigDB were tested, with the Bonferroni-

143 corrected significance threshold set at $P < 3.23 \times 10^{-6.60}$

144 **Omics PlEiotRopic Association (OPERA) analysis**

145 To provide mechanistic interpretations of FA-associated variants, we performed an 146 integrative analysis of six types of multi-omics quantitative trait loci (QTL) summary statistics with GWAS signals using the Bayesian method, Omics PlEiotRopic Association (OPERA).⁶¹ First, 147 148 we obtained available molecular quantitative trait loci (xQTL) data, including gene expression 149 QTL (eQTL), protein QTL (pQTL), DNA methylation QTL (mQTL), splicing QTL (sQTL), 150 histone modification QTL (hQTL), and chromatin accessibility QTL (caQTL). We downloaded 151 blood *cis*-eQTLs summary statistics generated from the eQTLGen study (N = 31,684).⁶² The 152 plasma pQTL summary statistics were retrieved from the UKB Pharma Proteomics Project (UKB-PPP) using UKB participants (N = 54,219).⁶³ The peripheral blood mQTL data were generated by 153 McRae *et al.* (N = 1.980).^{64,65} We derived summary-level sQTL data from whole blood samples in 154 the Genotype Tissue Expression (GTEx) Project v.8 (N = 670).⁶⁶ The H3K27ac and H3K4me1 155 156 hQTL in monocytes were generated from the BLUEPRINT project (N = 200).⁶⁷ The caQTL data in lymphoblastoid cell line samples were estimated by Kumasaka et al. (N = 100).⁶⁸ We included 157 the molecular phenotypes with at least one xQTL with $P < 5 \times 10^{-8}$ and excluded the major 158 159 histocompatibility complex (MHC) region due to its structural complexity. After filtering 160 molecular phenotypes, we retained 15,785 genes, 2,190 proteins, 94,338 DNA methylation sites, 161 6,639 RNA splicing events, 18,152 histone marks, and 13,873 chromatin accessibility peaks. The 162 posterior probability of association (PPA) threshold of 0.9 and the multi-exposure heterogeneity 163 in dependent instruments (HEIDI) test threshold of 0.01 were used to determine if a GWAS signal 164 overlaps with the QTL signals for one or multiple molecular phenotypes.

It is made available under a CC-BY-NC 4.0 International license .

165 **Results**

166 Sample characteristics

The primary GWAS study comprised up to 239,268 individuals of EUR ancestry. GWAS were also conducted on five additional ancestry groups, including up to 3,352 AFR, 508 AMR, 4,663 CSA, 1,445 EAS, and 865 MID individuals. Among EUR participants, the average age was 57 years and females were more likely to have higher PUFA levels (Table S1). The percentage of female participants in EUR, AFR, AMR, CSA, EAS, and MID populations was 53.9%, 59.7%, 66.9%, 44.8%, 64.6%, and 41.8%, respectively. Approximately 31.6% of individuals reported regular use of fish oil supplements and the ratio of omega-3/omega-6 was around 1:8.

174 Identification of genome-wide significant loci for FAs

175 We first performed EUR ancestry GWAS for each FA trait using three models, respectively. 176 Since we are particularly interested in the genetic architecture of FA traits in the category of 177 PUFAs, we primarily use results in this category as examples throughout this article to demonstrate 178 our findings. Similar results for FA traits in the categories of MUFAs and SFAs are available in 179 supplementary materials. Figure 1A shows a Manhattan plot of GWAS for the absolute 180 concentration of total PUFAs in model 2. Table S2 reports the number of significant loci for each 181 FA from all GWAS across all models. Using model 2, we identified 122 independent genetic loci 182 for the category of PUFAs, 124 for MUFAs, and 102 for SFAs; however, only 46, 39, and 38 loci 183 respectively associated with these categories of FA traits in model 3 (Table S2). (Please elaborate 184 a bit more. Are all loci in model 3 identified in model 2? If not, how many were identified in model 2? Then, the rest is only significant in model 3.) A total of 92 loci identified in model 2 were no 185 186 longer significantly associated with total PUFAs after adjusting for fish oil intake and lipoprotein 187 lipids in model 3 (Figure 1B). We observed similar effect estimates between models 1 and 2,

whereas model 3 yielded systematically larger effect estimates than those from the other two models (Figure S1). To further explore the distinction between models 3 and 2, we conducted GWAS based on model 2 by additionally adjusting for fish oil supplementation in model 2.1 and for lipoprotein lipids in model 2.2. The differences in GWAS results between model 3 and the others were attributed to the additional adjustment for lipoprotein lipids (Table S2). For example, variants in the *LDLR* and *MTTP* genes, known for their strong correlation with lipoprotein lipids,^{69,70} were significant in model 2 but not in model 3.

195 Regarding SNP-based heritability of FA traits in the EUR population, we found that SNPs 196 explained 5–19% of the phenotypic variance for all FA traits across all three models (Figure S2A; 197 Table S3). For instance, the SNP-based heritability estimates for the absolute concentrations of 198 total PUFAs, MUFAs, and SFAs were 19%, 19%, and 16%, respectively, in model 2. The genomic 199 inflation factor (λ_{GC}) for all GWAS ranged from 1.07 to 1.29 (Table S4). The attenuation ratio 200 ranged from 0.03 to 0.18, and the LDSC intercept ranged from 1.01 to 1.12, suggesting that the 201 genome-wide elevation of association statistics was primarily due to true additive polygenic effects 202 rather than confounders such as population stratification.

203 To identify novel loci, we searched the literature and GWAS Catalog to compile known 204 loci for FAs-related traits, making it the most comprehensive collection to date (Table S5). When 205 categorizing FA traits into PUFAs-related, MUFAs-related, and SFAs-related, we identified 70, 206 61, and 54 novel loci among 215, 163, and 119 significant loci, respectively, with model 2 (Table 207 S6). The numbers of novel loci for each FA trait are provided in Table S7. Compared with all 208 previously reported FAs-related loci, the top 10 strongest novel loci identified across all FA traits 209 in model 2 were mapped to candidate genes including PEPD, SBNO1, IL1RN, INSIG2, VIM, PGS1, 210 ARID5B, NF1, FAM96A, and PEX6 (Table S8). We also compared independent loci to those

211 previously reported in GWAS for the same traits. The novel locus *VEGFA* was associated with 212 total FAs, LA%, SFAs, PUFAs/SFAs, and the degree of unsaturation in our study, and it was 213 consistently associated with MUFAs, MUFAs%, and PUFAs/MUFAs in both our and previous 214 GWAS.^{29,32,33,37} Variants in *GCKR* were found to be significantly associated with DHA% for the 215 first time, consistent with previous GWAS findings for other FA traits.

Additionally, we explored genetic associations on the X-chromosome in both overall and sex-specific GWAS. Previously, only one GWAS identified FA loci on the X-chromosome.²⁹ That GWAS reported two X-chromosomal loci associated with FA traits; however, no significant variants were identified in our overall GWAS. When evaluating males separately, one significant association with omega-3% was found on the X-chromosome.

221 Sex-specific FA GWAS

222 We then stratified the total sample by sex to perform sex-specific GWAS separately for 223 128,922 females and 110,346 males in the EUR population. Approximately 4.2% of the significant 224 loci from sex-specific GWAS were not identified in the overall GWAS (Table S7). The SNP-based 225 heritability ranged from 5% to 21% in females and 4% to 20% in males, and there was no evidence 226 of genomic inflation (Figures S2B and S2C; Table S3). In model 2, we identified 128 independent 227 loci associated with PUFAs-related traits in females and 106 loci in males, including 27 and 8 228 novel loci, respectively, which have not been reported in previous GWAS of PUFAs-related traits 229 (Table S6). GWAS results for both known and novel loci associated with each FA trait are 230 provided in Table S9 for females and Table S10 for males. Interestingly, among the three novel 231 loci associated with PUFAs compared to previous GWAS related to all FA traits, two loci mapped 232 to RNU6-1180P and KRT18P55 were significant only in females from model 2 (Figure 2). After 233 accounting for lipoprotein lipids, three additional novel loci for PUFAs, around gene UNC5CL,

RP11-328J6.1, and *SIPA1L3*, were identified in females and no novel locus in males was identified
when comparing with all previous FA GWAS in model 3 (Figure S3). In the sex-specific GWAS,
we found that approximately 43% of the genome-wide significant loci identified in models 1 and
were not identified in model 3 (Table S7). Notably, one X-chromosomal locus associated with
omega-3%, whose lead SNP is rs147828433, was identified from GWAS using model 2 in males.
Our study is the first to explore loci in sex-specific GWAS and to include the X chromosome.

240 **GWAS for FA traits in individuals of non-EUR ancestries**

241 Across all FA traits, we identified 38 significant loci in ancestry-specific GWAS, of which 242 23 are novel compared to previous studies in EUR and non-EUR populations (Table S11). The 243 genomic inflation factor (λ_{GC}) was also reported for each non-EUR ancestry, with average λ_{GC} 244 values of 0.97, 0.92, 0.99, 0.90, and 0.95 for AFR, AMR, CSA, EAS, and MID, respectively 245 (Tables S4). In the non-EUR ancestry GWAS, the most significant locus was around FADS2, 246 identified in four ancestry groups except AFR, which is a well-known locus in the EUR population 247 (Table S11). Among non-EUR populations, CSA had the largest sample size, leading to more loci 248 being identified compared to other groups, with 1-4 loci in model 1 and 0-2 in model 2 (Table 249 S2). The GWAS in CSA male participants identified a novel locus EDN1 associated with total 250 PUFAs, omega-6 PUFAs, and LA, while this locus did not reach the genome-wide significance in 251 the EUR population. In addition, variants in VPS39, which had not been reported in previous 252 GWAS, were associated with total PUFAs, omega-6 PUFAs, and LA in both total and male-only 253 CSA participants.

254 Genetic correlations among FA traits

255 We examined genetic correlations (r_g) between FA traits using EUR GWAS summary 256 statistics from three models (Table S12). Broadly, comparisons of genetic correlations between

257 models revealed similar estimates for models 1 and 2 but distinct estimates when compared to 258 model 3, as illustrated in Figure S4. Among the 171 pairs of genetic correlations, 14 pairs that had 259 significantly negative genetic correlations in models 1 and 2 became significantly positive in 260 model 3, while four pairs that were significantly positive in models 1 and 2 became significantly 261 negative in model 3. The strongest correlation was a negative correlation between MUFAs and PUFAs/MUFAs ($r_g = -0.99$) observed based on GWAS in both models 1 and 2 (Figures S5 and 262 263 S6). Notably, genetic correlations between omega-6 PUFAs and omega-6% were negative in 264 models 1 ($r_g = -0.38$) and 2 ($r_g = -0.51$), but they were positively correlated in model 3 ($r_g = 0.34$) after adjustment for lipoprotein lipids (Figure S7). Within omega-3-related traits, genetic 265 266 correlations were strong (~ 0.9) across all three models, with even stronger correlations found in 267 model 3. We also estimated pairwise phenotypic correlations using Pearson's correlation 268 coefficient (Table S13). Among significant genetic correlations, all pairs had consistent directions 269 compared to phenotypic correlations in models 1 and 2, while 83% of pairs had consistent 270 directions in model 3, suggesting that genetic effects on FA traits were altered after adding 271 lipoprotein lipids as covariates.

272 Colocalization of GWAS signals across FA traits

To assess the probability that two FA traits share the same causal variants, we conducted pairwise colocalization analyses. For the 171 pairs of FA traits, we identified 9–231 colocalized signals in model 2 and 6–114 in model 3, respectively. We found that pairs with a larger number of colocalized signals also had stronger genetic correlations (Figure S8). In model 2, potential causal variants in *DOCK7* (rs2934744), *GCKR* (rs1260326), *ZNF259* (rs964184), *FADS1* (rs174564), *CPT1A* (rs2229738), *LIPC* (rs2070895), *LIPG* (rs77960347), *LDLR* (rs73015024), *TM6SF2* (rs58542926), and *APOE* (rs7412) showed robust evidence of colocalization, indicating

280 shared genetic variants for pairs of FA traits (Table S14). After accounting for lipoprotein lipids 281 in model 3, FADS1 (rs174564), CPT1A (rs2229738), and BUD13 (chr11:116623213:TA:T) were 282 identified as colocalized and potentially causal signals for FA traits, with results consistent with 283 those from model 2 (Table S15). Next, we performed multi-trait colocalization analyses to evaluate 284 the posterior probability of a shared genetic signal across all FA traits. Overall, 168 and 159 285 colocalized signals were identified in models 2 and 3, respectively. We identified GCKR 286 (rs1260326), TRIB1 (rs28601761), FADS1 (rs174564), ZNF259 (rs964184), and APOE (rs7412) 287 as shared between multiple FA traits in model 2 (Table S16), and GCKR (rs1260326), TRIB1 288 (rs28601761), FADS1 (rs174564), and APOE (rs1065853) from model 3 as shared signals (Table 289 S17).

290 Statistical fine-mapping for candidate causal variants

291 We applied the Bayesian fine-mapping method to identify putative causal variants for FA 292 traits. We identified 477 unique potential causal variants in model 1, 422 in model 2, and 193 in 293 model 3 (Table S18). Consistent with colocalization analysis, we confirmed that the genetic 294 variants at the loci TRIB1 (rs28601761), FADS1 (rs174564), CPT1A (rs2229738), LIPC 295 (rs2070895), LIPG (rs77960347), LDLR (rs72658867), TM6SF2 (rs58542926), and APOE (rs7412) 296 are likely to be causally associated with FA traits. Notably, we found that several loci contain 297 multiple 95% credible sets, indicating the existence of multiple potential causal variants within 298 those significant loci. Among the novel loci for FA traits, we found that ZMIZ1 (rs1782652) was 299 associated with total FAs, total PUFAs, and omega-3 PUFAs, suggesting that it is a candidate 300 shared causal variant for these FAs. Additional potential causal variants for novel loci were 301 identified, including CPS1 (rs1047891), ATXN7L1 (rs118061830), ARID5B (rs77044968), SBF2

It is made available under a CC-BY-NC 4.0 International license .

302 (rs12789941), DGKZ (rs149903077), CYFIP1 (rs199854211), FAM96A (rs62023393), MIR122

303 (rs41292412), BCL2 (rs12454712), INSR (rs112630404), and PEPD (rs62102718).

304 Gene-based and gene-set analyses

305 In the genome-wide gene-based association study, 139-541 genes from model 2 and 110-306 255 genes from model 3 were found to be significantly associated with 19 FA traits after 307 Bonferroni correction (Table S19). The top gene across 19 FA traits identified in model 2 was 308 ALDH1A2 (chr15: 58245622-58790065), which overlaps with another identified gene LIPC 309 (chr15: 58702768-58861151). In the GWAS of the omega-3/omega-6 ratio using model 3, STH 310 was the top gene, which might be related to neurodegenerative diseases such as Alzheimer's 311 disease.⁷¹ Gene-tissue expression analysis revealed that the liver was the most significantly 312 enriched tissue, followed by whole blood, kidney, small intestine, spleen, nerve, and adipose tissue 313 (Tables S20 and 21). Additionally, gene set analysis was performed to investigate potential gene 314 pathways (Tables S22 and 23). We detected those genes associated with total PUFAs clustered in 315 biological processes related to lipid homeostasis, secondary alcohol metabolism, organic hydroxy 316 compound transport, regulations of plasma lipoprotein particle levels and lipid metabolism, and 317 fatty acid biosynthesis (Figure 3). Fatty acid metabolic and biosynthetic processes were 318 significantly enriched in the gene set analysis results from GWAS of total PUFAs using model 3 319 (Figure S9).

320 OPERA identifies intermediate molecular phenotypes underlying FA-associated loci

To better understand the biological mechanisms of our GWAS findings, we applied OPERA to explore the overlap of GWAS signals for FA traits and QTL signals for six molecular phenotypes, including eQTL, pQTL, mQTL, sQTL, hQTL, and caQTL. With a PPA threshold of 0.9 and a P_{HEIDI} threshold of 0.01, we identified 976 FA-associated variants that overlap with QTL

325 signals for 2,106 unique molecular phenotype measures, including the expression levels of 242 326 genes, the abundance levels of 60 proteins, the methylation levels at 1,254 DNA sites, the RNA 327 splicing events at 141 sites, the status of 233 histone marks, and the openness at 176 chromatin 328 accessibility peaks (Figure 4A and Table S24). Overlaps with xQTL signals were also observed 329 for a large number of loci after adjusting for lipoprotein lipids in GWAS (Figure 4B and Table 330 S25). Associations with DNA methylation were more frequent than with other molecular 331 phenotypes, and these findings were consistent with previous results in other complex traits.⁶¹ The 332 number of loci associated with eQTL across 19 traits ranged from 7 for the degree of unsaturation 333 to 68 for PUFAs in model 2, and from 2 to 41 in model 3. Using the pQTL data from UKB-PPP, 334 we observed that 33 loci for omega-3 PUFAs from models 2 and 3 had the highest number of 335 significant associations with proteins.

336 We found that approximately 65% of GWAS loci for 19 FA traits are shared with at least 337 one molecular phenotype. Among the well-known FA loci, the potential causal variant rs174564 338 was found to overlap with QTL signals for the expression level of MYRF (ENSG00000124920) 339 and the methylation level at cg01825921. Across FA traits such as total FAs, total PUFAs, omega-340 6 PUFAs, and total SFAs, the potential causal variant rs72658867, annotated at the LDLR gene, 341 was associated with histone modification (chr19:11105519-11214483) and chromatin accessibility 342 (peak 264429). Among the novel loci for omega-6%, rs149903077 (DGKZ) was jointly associated 343 with the nearby gene NR1H3 (ENSG0000025434), histone modification (chr11:47234446-344 47272168), and chromatin accessibility (peak 200213). It was also marginally associated with 345 methylation (cg01183595).

Notably, all six molecular phenotypes were jointly associated with total FAs, omega-6%,
and total MUFAs at the novel locus *GSTTP1* or nearby genes on chromosome 22 (Figure 5). We

- 348 observed that variant rs189660333 was associated with omega-3% jointly with five other types of
- 349 molecular phenotypes, except for histone modification. The nearby variants rs12786457 and
- 350 rs184864731 were consistently associated with these five phenotypes across GWAS related to
- 351 omega-3 PUFAs and the omega-3/omega-6 ratio. Overall, OPERA integrates six types of xQTL
- 352 data to offer mechanistic insights into genetic loci and their downstream molecular phenotypes.

It is made available under a CC-BY-NC 4.0 International license .

353 **Discussion**

354 We performed overall and sex-specific GWAS for 19 FA traits in 250,101 UKB 355 participants of EUR descent and five other ancestry groups, separately. We identified 215, 163, 356 and 119 significant loci for PUFAs-, MUFAs-, and SFAs-related traits, including 70, 61, and 54 357 novel loci, respectively, in our primary model (i.e., model 2). Adjusting for lipoprotein lipids 358 dramatically reduce the number of GWAS signals. Our analyses of genetic correlations and 359 colocalization revealed the levels of shared genetic basis and specific candidate shared variants 360 across FA traits. Using statistical fine-mapping, we identified 422 putative causal variants across 361 all FA traits. Gene-based and gene-set analyses pinpointed the liver as the most relevant tissue and 362 highlighted biological pathways underlying FA loci. In addition, we integrated GWAS signals with 363 QTL signals for six molecular phenotypes and revealed that approximately 65% of GWAS loci for 364 19 FA traits are shared with at least one molecular phenotype, offering novel mechanistic insights. 365 In EUR GWAS, we analyzed 19 plasma NMR measures of FAs from 239,268 participants 366 in the UKB. Although previous GWAS of FA traits in the UKB reported many loci, we identified approximately 90% more loci with a doubled sample size.^{29,32,33} In addition, we compared our 367 368 results with all previous GWAS of FAs-related traits to identify novel loci, including the second-369 largest published GWAS to date, which included 136,016 participants from 33 cohorts.³⁷ We 370 calculated the omega-3/omega-6 and PUFA/SFA ratios for GWAS for the first time. Although the 371 omega-6 to omega-3 ratio is commonly used in studies, the omega-3/omega-6 ratio may better 372 capture the benefits of omega-3. This is particularly relevant given the ongoing debate over the potential harmful or beneficial effects of omega-6 PUFAs.⁷² Our large GWAS of FAs showed that 373 374 the phenotypic variance explained by all SNPs ranged from 5-19%, while heritability estimates 375 from twin studies were approximately 25-62%.¹⁴ These results suggest that future studies with

It is made available under a CC-BY-NC 4.0 International license .

376 rare or structural variants from sequencing and more diverse ancestry groups could be valuable for377 identifying additional loci for FA traits.

Variants in the FADS gene cluster are well-known for modifying polyunsaturated fatty acid 378 379 desaturation, but the underlying molecular mechanisms require investigation. Using OPERA, we 380 identified the potential causal variant rs174564 in FADS1, which is associated with the MYRF gene 381 and the methylation site cg01825921. The LDLR gene variant rs72658867 was linked to histone 382 modification (19:11105519:11214483) and chromatin accessibility (264429), impacting lipid metabolism and cardiovascular health.⁷³ The variant rs149903077 in the novel DGKZ locus has 383 384 been associated with the nearby gene NR1H3, chromatin accessibility, histone modifications, and 385 DNA methylation, highlighting their combined roles in nuclear receptor regulation, lipid 386 metabolism, and immune responses.⁷⁴ The novel GSTTP1 locus, or nearby genes on chromosome 387 22, overlapped with all six molecular phenotypes and has been linked to key biological processes such as detoxification, cancer susceptibility, and cellular responses to oxidative stress.⁷⁵ 388

389 Gene set analysis findings highlight biological pathways underlying FA loci, implicating 390 lipid homeostasis, transport, and metabolism, and providing genetic support for the current understanding of fatty acid regulation.⁷⁶ Our previous study suggested that lower omega-6% may 391 392 reduce alcohol consumption, with the secondary alcohol metabolism pathway potentially 393 explaining this effect on alcohol-related behaviors.⁶ In model 3, which adjusts for lipoprotein lipid 394 levels, fatty acid metabolic and biosynthetic processes were significantly more enriched in the 395 gene set analysis of GWAS of total PUFAs. The investigation of the disparities between models 2 396 and 3 suggests that lipoprotein-related biomarkers play a critical role in FA loci. Future studies are 397 needed to thoroughly dissect these roles, and further analyses, such as Mendelian randomization,

It is made available under a CC-BY-NC 4.0 International license .

398 should properly infer results using genetic variants with or without adjustment for lipoprotein-399 related biomarkers.⁷⁸

400 We investigated the sex-specific genetic architecture of FA traits. A novel locus, 401 KRT18P55, was identified in the GWAS for females, and previous research indicates that this 402 locus is highly expressed in patients with gastric cancer.⁷⁹ Notably, only one X-chromosomal locus 403 was identified in the male-only GWAS for omega-3%, tagged by the lead variants rs147828433 404 (closest protein-coding gene: ACOT9). A study on ACOT9 in mice suggests that it regulates both fatty acid and amino acid metabolism in mitochondria.⁸⁰ For the female-only GWAS, the 405 406 heritability was higher and larger numbers of loci were identified. Our results highlight the 407 importance of tracking sex differences in genetics and may motivate future studies on gene-sex 408 interactions. We found higher plasma PUFA levels in females, consistent with previous studies, 409 which were influenced by genetic effects, sex hormones, and conversion rates in FA metabolism.⁹ 410 Randomized controlled trials show sex- and race-specific differences in the benefits of fish oil 411 supplementation and omega-3 PUFAs for preventing cardiovascular events and cognitive 412 decline.81

413 By performing ancestry-specific and sex-specific GWAS in non-EUR samples, we 414 identified additional novel loci, suggesting ancestry-specific genetic architecture of FA traits. In 415 male CSA participants, the novel locus EDN1, associated with omega-6 PUFA traits, encodes the protein endothelin 1, which has been linked to cardiovascular events and prognosis.⁸² Variants in 416 417 VPS39, associated with omega-6 PUFAs in the CSA population, have been previously identified 418 as important regulators of myoblast differentiation and muscle glucose uptake in patients with type 419 2 diabetes.⁸³ Besides these ancestry-specific signals, we also found several loci identified from 420 large EUR GWAS were transferable to other ancestry groups, including FADS2, ZNF259, MYRF,

421 *APOA5*, *GCKR*, *APOE*, and other loci. We found significantly fewer loci in the other five non-422 EUR ancestry GWAS due to small sample sizes. These results suggest that future studies with 423 larger and more diverse ancestry samples are needed to identify more loci and confirm their 424 consistency across populations.

425 To the best of our knowledge, our study is the most comprehensive GWAS on FA traits in 426 a very large cohort. First, this is the first study to integrate six types of xQTL data with FA GWAS 427 to explore the possible molecular mechanisms of FA-associated genetic variants. Second, we used 428 a large number of participants of EUR ancestry with plasma NMR measures for 19 FA traits and 429 included participants from five other ancestry groups. Third, the inclusion of the omega-3/omega-430 6 ratio in the GWAS provides better insights into the benefits of omega-3 PUFAs. Fourth, our sex-431 specific GWAS on EUR and non-EUR populations, as well as identified X-chromosomal loci, 432 uncover sex differences in FA genetic basis. Fifth, we comprehensively compared our findings to 433 all previous GWAS of FAs-related traits. Detailed information on all novel and known loci has been provided (Tables S5 and S7), which offers a good resource for researchers to further explore 434 435 FAs-related loci. Finally, we adjusted for various covariates across three models, with a particular 436 focus on genetic architecture adjusted for lipoprotein metabolism.

The study has limitations. The number of participants from non-EUR populations was low, and future studies with larger sample sizes are necessary. While we explored the molecular mechanisms of FA loci, we are eager to extend OPERA analysis to other tissues, particularly the liver, and even specific cell types, to identify cell type-specific disease mechanisms.

441 In conclusion, our findings reveal novel genetic loci and provide the first evidence of 442 molecular mechanisms underlying FA-relevant variants. These insights could aid patient

- 443 stratification for precision nutrition and the identification of novel therapeutic targets for FAs-
- 444 related diseases.

It is made available under a CC-BY-NC 4.0 International license .

445 **Data availability**

- 446 The full summary statistics for GWAS of fatty acids will be publicly available in the GWAS
- 447 Catalog database (<u>https://www.ebi.ac.uk/gwas/</u>). The blood eQTL data of eQTLGen Consortium
- 448 used in the analyses were downloaded from <u>https://molgenis26.gcc.rug.nl/downloads/eqtlgen/cis-</u>
- 449 eqtl/SMR formatted/cis-eQTL-
- 450 <u>https://zenodo.org/records/7951839/files/LBC_BSGS_meta.tar.gz?download=1SMR_20191212.</u>
- 451 tar.gz. UKBB pQTL data are available at https://metabolomips.org/ukbbpgwas/. The mQTL data 452 download available for from are 453 https://zenodo.org/records/7951839/files/LBC BSGS meta.tar.gz?download=1. Blood sOTL data 454 be downloaded can at 455 https://yanglab.westlake.edu.cn/data/SMR/GTEx V8 cis sqtl summary/sQTL Whole Blood.zi 456 The hQTL be data can assessed from <u>p</u>. http://ftp.ebi.ac.uk/pub/databases/blueprint/blueprint_Epivar/qtl_as/OTL_RESULTS/. Summary 457 458 statistics of caOTL can be found at https://zenodo.org/records/1405945/files/.

459 Acknowledgments

460 This research was approved by the University of Georgia Institutional Review Board and the UK 461 Biobank consortium (application no. 48818). This work was funded by the University of Georgia 462 Research Foundation and by the National Institute of General Medical Sciences of the National 463 Institutes of Health under Award Number R35GM143060. The content is solely the responsibility 464 of the authors and does not necessarily represent the official views of the National Institutes of 465 Health.

466 Author contributions

- 467 Y.S. and K.Y. designed the study. Y.S. and H.X. performed analyses. Y.S. and K.Y. drafted the
- 468 manuscript. All authors contributed to the review and critical revision of the manuscript.
- 469 **Declaration of interests**
- 470 The authors declare no competing interests.

It is made available under a CC-BY-NC 4.0 International license .

References

471	1.	Worthmann, A., Ridder, J., Piel, S.Y.L., Evangelakos, I., Musfeldt, M., Voss, H.,
472		O'Farrell, M., Fischer, A.W., Adak, S., Sundd, M., et al. (2024). Fatty acid synthesis
473		suppresses dietary polyunsaturated fatty acid use. Nat Commun 15, 45. 10.1038/s41467-
474		023-44364-у.
475	2.	Baum, S.J., Kris-Etherton, P.M., Willett, W.C., Lichtenstein, A.H., Rudel, L.L., Maki,
476		K.C., Whelan, J., Ramsden, C.E., and Block, R.C. (2012). Fatty acids in cardiovascular
477		health and disease: a comprehensive update. J Clin Lipidol 6, 216-234.
478		10.1016/j.jacl.2012.04.077.
479	3.	Astrup, A., Magkos, F., Bier, D.M., Brenna, J.T., de Oliveira Otto, M.C., Hill, J.O., King,
480		J.C., Mente, A., Ordovas, J.M., Volek, J.S., et al. (2020). Saturated Fats and Health: A
481		Reassessment and Proposal for Food-Based Recommendations: JACC State-of-the-Art
482		Review. J Am Coll Cardiol 76, 844-857. 10.1016/j.jacc.2020.05.077.
483	4.	Scorletti, E., and Byrne, C.D. (2013). Omega-3 fatty acids, hepatic lipid metabolism, and
484		nonalcoholic fatty liver disease. Annu Rev Nutr 33, 231-248. 10.1146/annurev-nutr-
485		071812-161230.
486	5.	Zhang, Y., Sun, Y., Yu, Q., Song, S., Brenna, J.T., Shen, Y., and Ye, K. (2024). Higher
487		ratio of plasma omega-6/omega-3 fatty acids is associated with greater risk of all-cause,
488		cancer, and cardiovascular mortality: A population-based cohort study in UK Biobank.
489		Elife 12. 10.7554/eLife.90132.
490	6.	Xu, H., Sun, Y., Francis, M., Cheng, C.F., Modulla, N.T.R., Brenna, J.T., Chiang,
491		C.W.K., and Ye, K. (2023). Shared genetic basis informs the roles of polyunsaturated
492		fatty acids in brain disorders. medRxiv. 10.1101/2023.10.03.23296500.
493	7.	Sun, Y., Chatterjee, R., Ronanki, A., and Ye, K. (2022). Circulating Polyunsaturated
494		Fatty Acids and COVID-19: A Prospective Cohort Study and Mendelian Randomization
495		Analysis. Front Med (Lausanne) 9, 923746. 10.3389/fmed.2022.923746.
496	8.	Hodson, L., Skeaff, C.M., and Fielding, B.A. (2008). Fatty acid composition of adipose
497		tissue and blood in humans and its use as a biomarker of dietary intake. Prog Lipid Res
498		47, 348-380. 10.1016/j.plipres.2008.03.003.
499	9.	Decsi, T., and Kennedy, K. (2011). Sex-specific differences in essential fatty acid
500		metabolism. Am J Clin Nutr 94, 1914S-1919S. 10.3945/ajcn.110.000893.
501	10.	Mathias, R.A., Pani, V., and Chilton, F.H. (2014). Genetic Variants in the FADS Gene:
502		Implications for Dietary Recommendations for Fatty Acid Intake. Curr Nutr Rep 3, 139-
503		148. 10.1007/s13668-014-0079-1.
504	11.	Zhuang, P., Liu, X., Li, Y., Wu, Y., Li, H., Wan, X., Zhang, L., Xu, C., Jiao, J., and
505		Zhang, Y. (2023). Circulating fatty acids, genetic risk, and incident coronary artery
506		disease: A prospective, longitudinal cohort study. Sci Adv 9, eadf9037.
507		10.1126/sciadv.adf9037.
508	12.	Tanaka, T., Shen, J., Abecasis, G.R., Kisialiou, A., Ordovas, J.M., Guralnik, J.M.,
509		Singleton, A., Bandinelli, S., Cherubini, A., Arnett, D., et al. (2009). Genome-wide
510		association study of plasma polyunsaturated fatty acids in the InCHIANTI Study. PLoS
511		Genet 5, e1000338. 10.1371/journal.pgen.1000338.
512	13.	Lemaitre, R.N., Tanaka, T., Tang, W., Manichaikul, A., Foy, M., Kabagambe, E.K.,
513		Nettleton, J.A., King, I.B., Weng, L.C., Bhattacharya, S., et al. (2011). Genetic loci
514		associated with plasma phospholipid n-3 fatty acids: a meta-analysis of genome-wide

515		association studies from the CHARGE Consortium. PLoS Genet 7, e1002193.
516		10.1371/journal.pgen.1002193.
517	14.	Kettunen, J., Tukiainen, T., Sarin, A.P., Ortega-Alonso, A., Tikkanen, E., Lyytikainen,
518		L.P., Kangas, A.J., Soininen, P., Wurtz, P., Silander, K., et al. (2012). Genome-wide
519		association study identifies multiple loci influencing human serum metabolite levels. Nat
520		Genet 44, 269-276. 10.1038/ng.1073.
521	15.	Wu, J.H., Lemaitre, R.N., Manichaikul, A., Guan, W., Tanaka, T., Foy, M., Kabagambe,
522		E.K., Djousse, L., Siscovick, D., Fretts, A.M., et al. (2013). Genome-wide association
523		study identifies novel loci associated with concentrations of four plasma phospholipid
524		fatty acids in the de novo lipogenesis pathway: results from the Cohorts for Heart and
525		Aging Research in Genomic Epidemiology (CHARGE) consortium. Circ Cardiovasc
526		Genet 6, 171-183. 10.1161/CIRCGENETICS.112.964619.
527	16.	Guan, W., Steffen, B.T., Lemaitre, R.N., Wu, J.H.Y., Tanaka, T., Manichaikul, A., Foy,
528		M., Rich, S.S., Wang, L., Nettleton, J.A., et al. (2014). Genome-wide association study of
529		plasma N6 polyunsaturated fatty acids within the cohorts for heart and aging research in
530		genomic epidemiology consortium. Circ Cardiovasc Genet 7, 321-331.
531		10.1161/CIRCGENETICS.113.000208.
532	17.	Lemaitre, R.N., King, I.B., Kabagambe, E.K., Wu, J.H., McKnight, B., Manichaikul, A.,
533		Guan, W., Sun, Q., Chasman, D.I., Foy, M., et al. (2015). Genetic loci associated with
534		circulating levels of very long-chain saturated fatty acids. J Lipid Res 56, 176-184.
535		10.1194/jlr.M052456.
536	18.	Tintle, N.L., Pottala, J.V., Lacey, S., Ramachandran, V., Westra, J., Rogers, A., Clark, J.,
537		Olthoff, B., Larson, M., Harris, W., and Shearer, G.C. (2015). A genome-wide
538		association study of saturated, mono- and polyunsaturated red blood cell fatty acids in the
539		Framingham Heart Offspring Study. Prostaglandins Leukot Essent Fatty Acids 94, 65-72.
540		10.1016/j.plefa.2014.11.007.
541	19.	Dorajoo, R., Sun, Y., Han, Y., Ke, T., Burger, A., Chang, X., Low, H.Q., Guan, W.,
542		Lemaitre, R.N., Khor, C.C., et al. (2015). A genome-wide association study of n-3 and n-
543		6 plasma fatty acids in a Singaporean Chinese population. Genes Nutr 10, 53.
544		10.1007/s12263-015-0502-2.
545	20.	Hu, Y., Li, H., Lu, L., Manichaikul, A., Zhu, J., Chen, Y.D., Sun, L., Liang, S.,
546		Siscovick, D.S., Steffen, L.M., et al. (2016). Genome-wide meta-analyses identify novel
547		loci associated with n-3 and n-6 polyunsaturated fatty acid levels in Chinese and
548		European-ancestry populations. Hum Mol Genet 25, 1215-1224. 10.1093/hmg/ddw002.
549	21.	Kettunen, J., Demirkan, A., Wurtz, P., Draisma, H.H., Haller, T., Rawal, R., Vaarhorst,
550		A., Kangas, A.J., Lyytikainen, L.P., Pirinen, M., et al. (2016). Genome-wide study for
551		circulating metabolites identifies 62 loci and reveals novel systemic effects of LPA. Nat
552		Commun 7, 11122. 10.1038/ncomms11122.
553	22.	Kalsbeek, A., Veenstra, J., Westra, J., Disselkoen, C., Koch, K., McKenzie, K.A., O'Bott,
554		J., Vander Woude, J., Fischer, K., Shearer, G.C., et al. (2018). A genome-wide
555		association study of red-blood cell fatty acids and ratios incorporating dietary covariates:
556		Framingham Heart Study Offspring Cohort. PLoS One 13, e0194882.
557		10.1371/journal.pone.0194882.
558	23.	de Oliveira Otto, M.C., Lemaitre, R.N., Sun, Q., King, I.B., Wu, J.H.Y., Manichaikul, A.,
559		Rich, S.S., Tsai, M.Y., Chen, Y.D., Fornage, M., et al. (2018). Genome-wide association

560		meta-analysis of circulating odd-numbered chain saturated fatty acids: Results from the
561		CHARGE Consortium. PLoS One 13, e0196951, 10,1371/journal.pone.0196951.
562	24.	Marklund, M., Morris, A.P., Mahajan, A., Ingelsson, E., Lindgren, C.M., Lind, L., and
563		Riserus, U. (2018). Genome-Wide Association Studies of Estimated Fatty Acid
564		Desaturase Activity in Serum and Adipose Tissue in Elderly Individuals: Associations
565		with Insulin Sensitivity. Nutrients 10. 10.3390/nu10111791.
566	25.	Locke, A.E., Steinberg, K.M., Chiang, C.W.K., Service, S.K., Havulinna, A.S., Stell, L.,
567		Pirinen, M., Abel, H.J., Chiang, C.C., Fulton, R.S., et al. (2019). Exome sequencing of
568		Finnish isolates enhances rare-variant association power. Nature 572, 323-328.
569		10.1038/s41586-019-1457-z.
570	26.	Coltell, O., Sorli, J.V., Asensio, E.M., Barragan, R., Gonzalez, J.I., Gimenez-Alba, I.M.,
571		Zanon-Moreno, V., Estruch, R., Ramirez-Sabio, J.B., Pascual, E.C., et al. (2020).
572		Genome-Wide Association Study for Serum Omega-3 and Omega-6 Polyunsaturated
573		Fatty Acids: Exploratory Analysis of the Sex-Specific Effects and Dietary Modulation in
574		Mediterranean Subjects with Metabolic Syndrome. Nutrients 12. 10.3390/nu12020310.
575	27.	Riveros-Mckay, F., Oliver-Williams, C., Karthikeyan, S., Walter, K., Kundu, K.,
576		Ouwehand, W.H., Roberts, D., Di Angelantonio, E., Soranzo, N., Danesh, J., et al.
577		(2020). The influence of rare variants in circulating metabolic biomarkers. PLoS Genet
578		16, e1008605. 10.1371/journal.pgen.1008605.
579	28.	Li-Gao, R., Hughes, D.A., van Klinken, J.B., de Mutsert, R., Rosendaal, F.R., Mook-
580		Kanamori, D.O., Timpson, N.J., and Willems van Dijk, K. (2021). Genetic Studies of
581		Metabolomics Change After a Liquid Meal Illuminate Novel Pathways for Glucose and
582		Lipid Metabolism. Diabetes 70, 2932-2946. 10.2337/db21-0397.
583	29.	Richardson, T.G., Leyden, G.M., Wang, Q., Bell, J.A., Elsworth, B., Davey Smith, G.,
584		and Holmes, M.V. (2022). Characterising metabolomic signatures of lipid-modifying
585		therapies through drug target mendelian randomisation. PLoS Biol 20, e3001547.
586		10.1371/journal.pbio.3001547.
587	30.	Yin, X., Chan, L.S., Bose, D., Jackson, A.U., VandeHaar, P., Locke, A.E., Fuchsberger,
588		C., Stringham, H.M., Welch, R., Yu, K., et al. (2022). Genome-wide association studies
589		of metabolites in Finnish men identify disease-relevant loci. Nat Commun 13, 1644.
590		10.1038/s41467-022-29143-5.
591	31.	Montasser, M.E., Aslibekyan, S., Srinivasasainagendra, V., Tiwari, H.K., Patki, A.,
592		Bagheri, M., Kind, T., Barupal, D.K., Fan, S., Perry, J., et al. (2022). An Amish founder
593		population reveals rare-population genetic determinants of the human lipidome. Commun
594		Biol 5, 334. 10.1038/s42003-022-03291-2.
595	32.	Francis, M., Sun, Y., Xu, H., Brenna, J.T., and Ye, K. (2022). Fifty-one novel and
596		replicated GWAS loci for polyunsaturated and monounsaturated fatty acids in 124,024
597		Europeans. medRxiv, 2022.2005.2027.22275343. 10.1101/2022.05.27.22275343.
598	33.	Borges, M.C., Haycock, P.C., Zheng, J., Hemani, G., Holmes, M.V., Davey Smith, G.,
599		Hingorani, A.D., and Lawlor, D.A. (2022). Role of circulating polyunsaturated fatty acids
600		on cardiovascular diseases risk: analysis using Mendelian randomization and fatty acid
601		genetic association data from over 114,000 UK Biobank participants. BMC Med 20, 210.
602		10.1186/s12916-022-02399-w.
603	34.	Liu, C., Wang, Z., Hui, Q., Chiang, Y., Chen, J., Brijkumar, J., Edwards, J.A., Ordonez,
604		C.E., Dudgeon, M.R., Sunpath, H., et al. (2022). Crosstalk between Host Genome and

605 606		Metabolome among People with HIV in South Africa. Metabolites 12.
607	25	Dawyson F. Shan Y. Godd D.A. Barnahay F. Hillary P.F. McCartney D.I.
608	55.	Adams M. Marioni P. and McIntosh A.M. (2023) Metabolomic Investigation of
600		Maior Depressive Disorder Identifies a Detentially Causal Association With
610		Delymenting of Depressive Disorder Identifies a Potentiality Causal Association with Delymenting to the Disorder Disorder Disorder (04, 620, 620)
010 611		10 1016/j historych 2022 01 027
011	20	10.1010/J.Diopsych.2023.01.027.
012	30.	Fuller, H., Iles, M.M., Moore, J.B., and Zulyniak, M.A. (2025). Metabolic drivers of
613		dysglycemia in pregnancy: ethnic-specific GWAS of 146 metabolites and 1-sample
614		Mendelian randomization analyses in a UK multi-ethnic birth conort. Front Endocrinol
615	27	(Lausanne) 14, 115/416. 10.3389/fendo.2023.115/416.
616	37.	Karjalainen, M.K., Karthikeyan, S., Oliver-Williams, C., Sliz, E., Allara, E., Fung, W.I.,
617		Surendran, P., Zhang, W., Jousilahti, P., Kristiansson, K., et al. (2024). Genome-wide
618		characterization of circulating metabolic biomarkers. Nature 628, 130-138.
619	• •	10.1038/s41586-024-07148-y.
620	38.	Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott,
621		P., Green, J., Landray, M., et al. (2015). UK biobank: an open access resource for
622		identifying the causes of a wide range of complex diseases of middle and old age. PLoS
623		Med 12, e1001779. 10.1371/journal.pmed.1001779.
624	39.	Pan-ancestry genetic analysis of the UK Biobank. <u>https://pan.ukbb.broadinstitute.org/</u> .
625	40.	Julkunen, H., Cichonska, A., Slagboom, P.E., Wurtz, P., and Nightingale Health,
626		U.K.B.I. (2021). Metabolic biomarker profiling for identification of susceptibility to
627		severe pneumonia and COVID-19 in the general population. Elife 10.
628		10.7554/eLife.63033.
629	41.	McCaw, Z.R., Lane, J.M., Saxena, R., Redline, S., and Lin, X. (2020). Operating
630		characteristics of the rank-based inverse normal transformation for quantitative trait
631		analysis in genome-wide association studies. Biometrics 76, 1262-1272.
632		10.1111/biom.13214.
633	42.	Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A.,
634		Vukcevic, D., Delaneau, O., O'Connell, J., et al. (2018). The UK Biobank resource with
635		deep phenotyping and genomic data. Nature 562, 203-209. 10.1038/s41586-018-0579-z.
636	43.	Jiang, L., Zheng, Z., Qi, T., Kemper, K.E., Wray, N.R., Visscher, P.M., and Yang, J.
637		(2019). A resource-efficient tool for mixed model association analysis of large-scale data.
638		Nat Genet 51, 1749-1755. 10.1038/s41588-019-0530-8.
639	44.	International HapMap, C., Altshuler, D.M., Gibbs, R.A., Peltonen, L., Altshuler, D.M.,
640		Gibbs, R.A., Peltonen, L., Dermitzakis, E., Schaffner, S.F., Yu, F., et al. (2010).
641		Integrating common and rare genetic variation in diverse human populations. Nature 467.
642		52-58. 10.1038/nature09298.
643	45.	Yang, J., Zaitlen, N.A., Goddard, M.E., Visscher, P.M., and Price, A.L. (2014).
644		Advantages and pitfalls in the application of mixed-model association methods. Nat
645		Genet 46, 100-106, 10,1038/ng.2876.
646	46.	Jordan, H., Roderick, P., and Martin, D. (2004). The Index of Multiple Deprivation 2000
647		and accessibility effects on health. J Enidemiol Community Health 58, 250-257
648		10.1136/jech.2003.013011
649	47	Cleland C Ferguson S Ellis G and Hunter R F (2018) Validity of the International
650	• / •	Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical

651		activity and sedentary behaviour of older adults in the United Kingdom. BMC Med Res
652		Methodol 18, 176. 10.1186/s12874-018-0642-3.
653	48.	Welter, D., MacArthur, J., Morales, J., Burdett, T., Hall, P., Junkins, H., Klemm, A.,
654		Flicek, P., Manolio, T., Hindorff, L., and Parkinson, H. (2014). The NHGRI GWAS
655		Catalog, a curated resource of SNP-trait associations. Nucleic Acids Res 42, D1001-
656		1006. 10.1093/nar/gkt1229.
657	49.	Manolio, T.A., Collins, F.S., Cox, N.J., Goldstein, D.B., Hindorff, L.A., Hunter, D.J.,
658		McCarthy, M.I., Ramos, E.M., Cardon, L.R., Chakravarti, A., et al. (2009). Finding the
659		missing heritability of complex diseases. Nature 461, 747-753, 10,1038/nature08494.
660	50.	Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A., Bender, D., Maller,
661		L. Sklar, P., de Bakker, P.L. Daly, M.L. and Sham, P.C. (2007), PLINK: a tool set for
662		whole-genome association and population-based linkage analyses. Am I Hum Genet 81
663		559-575 10 1086/519795
664	51	McLaren W Pritchard B Rios D Chen Y Flicek P and Cunningham F (2010)
665	51.	Deriving the consequences of genomic variants with the Ensembl API and SNP Effect
666		Predictor Bioinformatics 26 2069-2070 10 1093/bioinformatics/bta330
667	52	Bulik-Sullivan BK Loh PR Finucane HK Rinke S Vang L Schizonbrenia
668	52.	Working Group of the Psychiatric Genomics C Patterson N Daly M I Price A I
669		and Neale B M (2015) I D Score regression distinguishes confounding from
670		nolygenicity in genome-wide association studies. Nat Genet 47, 201-205
671		10 1038/ng 3211
672	53	Bulik-Sullivan B. Finucane H.K. Anttila V. Gusev A. Dav F.R. Loh P.R.
673	55.	ReproGen C Psychiatric Genomics C Genetic Consortium for Anorevia Nervosa of
674		the Wellcome Trust Case Control C. Duncan L. et al. (2015) An atlas of genetic
675		apprelations parces human discusses and traits. Not Const 47, 1226, 1241
676		10 1038/ng 3406
677	54	Weissbrod O. Hermezdiari F. Denner C. Cui P. Illirech I. Gazal S. Schoech A.D.
679	54.	ven de Coiin P. Deshef V. Merguez Lune C. et al. (2020) Eunetionally informed
670		fine menning and polyconic localization of complex trait heritability. Not Const 52
690		inte-mapping and polygenic localization of complex trait heritability. Nat Genet 32 ,
000 601	55	1555-1505. 10.1050/841500-020-00755-5.
(92	55.	Lon, P.K., Kichaev, G., Gazai, S., Schoech, A.P., and Price, A.L. (2018). Wixed-model
692		association for biobalik-scale datasets. Nat Genet 50, 900-908. 10.1056/841588-018-
083	56	U144-0. Falax CN Stalax IB Dream BC Sum DD Kick DDW Dumage S and Harrow
084	30.	Foley, C.N., Staley, J.R., Breen, P.G., Sun, B.B., KIrk, P.D. w., Burgess, S., and Howson,
083		J.M.M. (2021). A fast and efficient colocalization algorithm for identifying shared
080		genetic risk factors across multiple traits. Nat Commun 12 , 764. 10.1058/s41467-020-
08/	57	20883-8.
688	57.	Wang, G., Sarkar, A., Carbonetto, P., and Stephens, M. (2020). A simple new approach
689		to variable selection in regression, with application to genetic fine mapping. J K Stat Soc
690	5 0	Series B Stat Methodol $\delta 2$, $12/3-1300$. 10.1111/rssb.12388.
691	58.	Watanabe, K., Taskesen, E., van Bochoven, A., and Posthuma, D. (2017). Functional
692		mapping and annotation of genetic associations with FUMA. Nat Commun δ , 1826.
693	50	10.1038/s4146/-01/-01261-5.
694	39.	Lee, S.H., Wray, N.K., Goddard, M.E., and Visscher, P.M. (2011). Estimating missing
695		heritability for disease from genome-wide association studies. Am J Hum Genet 88, 294-
696		305. 10.1016/j.ajhg.2011.02.002.

697	60.	Liberzon, A., Subramanian, A., Pinchback, R., Thorvaldsdóttir, H., Tamayo, P., and
698		Mesirov, J.P. (2011). Molecular signatures database (MSigDB) 3.0. Bioinformatics 27,
699		1739-1740. 10.1093/bioinformatics/btr260.
700	61.	Wu, Y., Qi, T., Wray, N.R., Visscher, P.M., Zeng, J., and Yang, J. (2023). Joint analysis
701		of GWAS and multi-omics OTL summary statistics reveals a large fraction of GWAS
702		signals shared with molecular phenotypes. Cell Genom 3, 100344.
703		10.1016/j.xgen.2023.100344.
704	62.	Vosa, U., Claringbould, A., Westra, H.J., Bonder, M.J., Deelen, P., Zeng, B., Kirsten, H.,
705		Saha, A., Kreuzhuber, R., Yazar, S., et al. (2021). Large-scale cis- and trans-eOTL
706		analyses identify thousands of genetic loci and polygenic scores that regulate blood gene
707		expression. Nat Genet 53, 1300-1310. 10.1038/s41588-021-00913-z.
708	63.	Sun, B.B., Chiou, J., Traylor, M., Benner, C., Hsu, Y.H., Richardson, T.G., Surendran,
709		P., Mahajan, A., Robins, C., Vasquez-Grinnell, S.G., et al. (2023). Plasma proteomic
710		associations with genetics and health in the UK Biobank. Nature 622, 329-338.
711		10.1038/s41586-023-06592-6.
712	64.	Wu, Y., Zeng, J., Zhang, F., Zhu, Z., Qi, T., Zheng, Z., Lloyd-Jones, L.R., Marioni, R.E.,
713		Martin, N.G., Montgomery, G.W., et al. (2018). Integrative analysis of omics summary
714		data reveals putative mechanisms underlying complex traits. Nat Commun 9, 918.
715		10.1038/s41467-018-03371-0.
716	65.	McRae, A.F., Marioni, R.E., Shah, S., Yang, J., Powell, J.E., Harris, S.E., Gibson, J.,
717		Henders, A.K., Bowdler, L., Painter, J.N., et al. (2018). Identification of 55,000
718		Replicated DNA Methylation QTL. Sci Rep 8, 17605. 10.1038/s41598-018-35871-w.
719	66.	Consortium, G.T. (2020). The GTEx Consortium atlas of genetic regulatory effects
720		across human tissues. Science 369, 1318-1330. 10.1126/science.aaz1776.
721	67.	Chen, L., Ge, B., Casale, F.P., Vasquez, L., Kwan, T., Garrido-Martin, D., Watt, S., Yan,
722		Y., Kundu, K., Ecker, S., et al. (2016). Genetic Drivers of Epigenetic and Transcriptional
723		Variation in Human Immune Cells. Cell 167, 1398-1414 e1324.
724		10.1016/j.cell.2016.10.026.
725	68.	Kumasaka, N., Knights, A.J., and Gaffney, D.J. (2019). High-resolution genetic mapping
726		of putative causal interactions between regions of open chromatin. Nat Genet 51, 128-
727		137. 10.1038/s41588-018-0278-6.
728	69.	Goldstein, J.L., and Brown, M.S. (2009). The LDL receptor. Arterioscler Thromb Vasc
729		Biol 29, 431-438. 10.1161/ATVBAHA.108.179564.
730	70.	Naik, A., Kosir, R., and Rozman, D. (2013). Genomic aspects of NAFLD pathogenesis.
731		Genomics 102, 84-95. 10.1016/j.ygeno.2013.03.007.
732	71.	Gao, L., Tse, S.W., Conrad, C., and Andreadis, A. (2005). Saitohin, which is nested in
733		the tau locus and confers allele-specific susceptibility to several neurodegenerative
734		diseases, interacts with peroxiredoxin 6. J Biol Chem 280, 39268-39272.
735		10.1074/jbc.M506116200.
736	72.	Schulze, M.B., Minihane, A.M., Saleh, R.N.M., and Riserus, U. (2020). Intake and
737		metabolism of omega-3 and omega-6 polyunsaturated fatty acids: nutritional implications
738		for cardiometabolic diseases. Lancet Diabetes Endocrinol 8, 915-930. 10.1016/S2213-
739	70	8587(20)30148-0.
740	73.	Gretarsdottir, S., Helgason, H., Helgadottir, A., Sigurdsson, A., Thorleifsson, G.,
741		Magnusdottir, A., Oddsson, A., Steinthorsdottir, V., Rafnar, T., de Graaf, J., et al. (2015).
/42		A Splice Region Variant in LDLR Lowers Non-high Density Lipoprotein Cholesterol and

743		Protects against Coronary Artery Disease. PLoS Genet 11, e1005379.
744		10.1371/journal.pgen.1005379.
745	74.	Raben, D.M., and Tu-Sekine, B. (2008). Nuclear diacylglycerol kinases: regulation and
746		roles. Front Biosci 13, 590-597. 10.2741/2704.
747	75.	Sadafi, S., Choubsaz, P., Kazemeini, S.M.M., Imani, M.M., and Sadeghi, M. (2024).
748		Glutathione S-transferase theta 1 (GSTT1) deletion polymorphism and susceptibility to
749		head and neck carcinoma: a systematic review with five analyses. BMC Cancer 24, 885.
750		10.1186/s12885-024-12618-7.
751	76.	Li, H., Herrmann, T., Seessle, J., Liebisch, G., Merle, U., Stremmel, W., and Chamulitrat,
752		W. (2022). Role of fatty acid transport protein 4 in metabolic tissues: insights into obesity
753		and fatty liver disease. Biosci Rep 42. 10.1042/BSR20211854.
754	77.	Sundaram, M., and Yao, Z. (2010). Recent progress in understanding protein and lipid
755		factors affecting hepatic VLDL assembly and secretion. Nutr Metab (Lond) 7, 35.
756		10.1186/1743-7075-7-35.
757	78.	Pingault, J.B., O'Reilly, P.F., Schoeler, T., Ploubidis, G.B., Rijsdijk, F., and Dudbridge,
758		F. (2018). Using genetic data to strengthen causal inference in observational research.
759		Nat Rev Genet 19, 566-580. 10.1038/s41576-018-0020-3.
760	79.	Tayefeh-Gholami, S., Ghanbari, M., Aghazadeh, A., Rajabi, A., Saber, A., Hussen, B.M.,
761		Farsad-Akhtar, N., and Safaralizadeh, R. (2022). Prognostic Value of LncRNA
762		KRT18P55 in Patients with Intestinal Type of Gastric Cancer. J Gastrointest Cancer 53,
763		1014-1019. 10.1007/s12029-021-00744-5.
764	80.	Tillander, V., Arvidsson Nordstrom, E., Reilly, J., Strozyk, M., Van Veldhoven, P.P.,
765		Hunt, M.C., and Alexson, S.E. (2014). Acyl-CoA thioesterase 9 (ACOT9) in mouse may
766		provide a novel link between fatty acid and amino acid metabolism in mitochondria. Cell
767		Mol Life Sci 71, 933-948. 10.1007/s00018-013-1422-1.
768	81.	Welty, F.K., Daher, R., and Garelnabi, M. (2024). Fish and Omega-3 Fatty Acids: Sex
769		and Racial Differences in Cardiovascular Outcomes and Cognitive Function. Arterioscler
770		Thromb Vasc Biol 44, 89-107. 10.1161/ATVBAHA.122.318125.
771	82.	Jankowich, M., and Choudhary, G. (2020). Endothelin-1 levels and cardiovascular
772		events. Trends Cardiovasc Med 30, 1-8. 10.1016/j.tcm.2019.01.007.
773	83.	Davegardh, C., Sall, J., Benrick, A., Broholm, C., Volkov, P., Perfilyev, A., Henriksen,
774		T.I., Wu, Y., Hjort, L., Brons, C., et al. (2021). VPS39-deficiency observed in type 2
775		diabetes impairs muscle stem cell differentiation via altered autophagy and epigenetics.
776		Nat Commun 12, 2431. 10.1038/s41467-021-22068-5.

It is made available under a CC-BY-NC 4.0 International license .

Figure Captions

Main Figures

Figure 1. Manhattan plot of the genome-wide association studies for the absolute concentration of total polyunsaturated fatty acids. The association of each variant with polyunsaturated fatty acids was obtained from genome-wide association studies using models 2 (A) and 3 (B). A variant with a *P*-value below 5×10^{-8} was considered statistically significant. The nearest gene at each genome-wide significant locus is annotated, in black for known loci and in red for novel loci.

Figure 2. Miami plot of the absolute concentration of total polyunsaturated fatty acids from genome-wide association studies using model 2. The top panel shows the GWAS results in females, while the bottom panel shows the GWAS results in males. The $-\log_{10}(P$ -value) is plotted on the y-axis and chromosomal location is plotted on the x-axis. The genome-wide significance threshold (P-value $< 5 \times 10^{-8}$) is indicated by the red dashed line. The nearest gene at each genome-wide significant locus is annotated, in black for known loci and in red for novel loci.

Figure 3. Pathway analysis of polyunsaturated fatty acids from genome-wide association studies using model 2. Treemap depicting significantly enriched pathways at an adjusted *P*-value threshold of 0.05. Gene ontology terms were clustered based on semantic similarity, with terms displayed as individual rectangles. The color indicates cluster membership, and thick border lines differentiate clusters. The size of each rectangle represents the enrichment significance, and the most significantly enriched term in each cluster is highlighted in white text as the representative term.

It is made available under a CC-BY-NC 4.0 International license .

Figure 4. Number of loci for fatty acid traits associated with different molecular phenotype combinations. (A and B) The number of loci associated with 19 fatty acid traits and molecular quantitative trait loci (xQTLs) was determined based on genome-wide association studies using models 2 (A) and 3 (B). The loci numbers are based on significant OPERA association results that pass a posterior probability of association threshold of 0.9 and the multi-exposure heterogeneity in dependent instruments (HEIDI) test ($P_{\text{HEIDI}} > 0.01$). The x-axis represents the association hypotheses for different phenotype combinations, while the y-axis shows the number of loci associated with these combinations across 19 fatty acids. Each violin plot displays the distribution of loci numbers by width, with lines indicating the 25th percentile, median, and 75th percentile.

Figure 5. Prioritization of a locus near the *GSTT1/2/2B* genes for total fatty acids. The top track displays the $-\log_{10}(P$ -value) of GWAS SNPs (gray dots) for total fatty acids. Red diamonds indicate OPERA marginal PPA for gene associations using eQTL data, while blue circles show OPERA marginal PPA for associations with DNA methylation, RNA splicing, histone modification, and chromatin accessibility, respectively. The bottom track presents 14 chromatin state annotations inferred from the 127 samples of the Roadmap Epigenomics Mapping Consortium.









It is made available under a CC-BY-NC 4.0 International license .

Figure 3.

organic hydroxy compound transport organic hy compound tr	regulation of sterol transport droxy ansport	regulati plasma lip particle regula lipoprote	ion of oprotein levels ition of pla ein particle	high–densi lipoproteir particle clearance sma levels	^{ty} triglyc metabolic tr meta	eride process iglyceric bolic pro	lipid catabolic process le	
intestinal lipid absorption	positive regulation of sterol transport cholesterol import	triglyceride–rich lipoprotein particle clearance	regulation of very-low-density lipoprotein particle clearance	low–densi lipoprotei particle clearance	ty glycerolipio n catabolic process	d lipid catabol proces isop catabol	glycerolipic metabolic process renoid ic process	
secondary alcoho secondary alcohol m metabolic process	alcohol etabolic aprócess process	regulation of lipid local regulation of lipid localregulation		tive re ation lipi ation m	gulation of d metabolic process agulation of DNA letabolic process positive	recepto end	amin intracellula vitamin sport transport eptor-mediated endocytosis negative	
	lipid store	age of loc	ationC	holesterol netabolic process	derived differ	rophage I foam cell entiation		
lipid home	ostasis	triglyceride–rich lipoprotein particle remodeling		le reg lip lipa	egative Julation of poprotein se activity	sterol biosynthetic process	chromatin organization chromatin organization membrane organization	
	of regulation cat catabo process	on of cellu trig olic proces	Solic Solic ocess	atty acid osynthetic orocess	negative regulation of endocytosis	protein localization to extracellular region		



It is made available under a CC-BY-NC 4.0 International license .





Chromosome 22 Mb