

Original article

Peripheral immune profiles in individuals at genetic risk for amyotrophic lateral sclerosis and Alzheimer's disease

Laura Deecke¹, Olena Ohlei¹, David Goldeck², Jan Homann¹, Sarah Toepfer³, Ilja Demuth^{3,4}, Lars Bertram⁵, Graham Pawelec^{2,6}, Christina M. Lill^{1,7*}

1. Institute of Epidemiology and Social Medicine, University of Münster, Münster, Germany
2. Department of Immunology, University of Tübingen, Tübingen, Germany
3. Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Endocrinology and Metabolic Diseases (including Division of Lipid Metabolism), Augustenburger Platz 1, 13353 Berlin, Germany
4. Berlin Institute of Health at Charité - Universitätsmedizin Berlin, BCRT - Berlin Institute of Health Center for Regenerative Therapies, Berlin, Germany
5. Lübeck Interdisciplinary Platform for Genome Analytics (LIGA), University of Lübeck, Lübeck, Germany
6. Health Sciences North Research Institute of Canada, Sudbury, Ontario, Canada
7. Ageing and Epidemiology Unit (AGE), School of Public Health, Imperial College London, London, UK

**Corresponding Author:*

Christina M. Lill, MD, MSc, Translational Epidemiology Group, Institute of Epidemiology and Social Medicine, University of Münster, Albert-Schweitzer-Campus 1, 48149 Münster, Germany, e-mail: christina.lill@uni-muenster.de

Keywords: Immune system, immune cell, neurodegeneration, polygenic risk score, disease prediction

Abstract

The immune system plays a crucial role in the pathogenesis of neurodegenerative diseases. Here, we explored whether blood immune cell profiles are already altered in healthy individuals with a genetic predisposition to amyotrophic lateral sclerosis (ALS) or Alzheimer's disease (AD).

Using multicolor flow cytometry, we analyzed 92 immune cell phenotypes in the blood of 448 healthy participants from the Berlin Aging Study II. We calculated polygenic risk scores (PGS) using genome-wide significant SNPs from recent large genome-wide association studies on ALS and AD. Linear regression analyses were then performed of the immune cell types on the PGS in both the overall sample and a subgroup of older participants (>60 years).

While we did not find any significant associations between immune cell subtypes and ALS and AD PGS when controlling for the false discovery rate (FDR=0.05), we observed several nominally significant results ($p < 0.05$) with consistent effect directions across strata. The strongest association was observed with CD57+ CD8+ early-memory T cells and ALS risk ($p = 0.006$). Other immune cell subtypes associated with ALS risk included PD-1+ CD8+ and CD57+ CD4+ early-memory T cells, non-classical monocytes, and myeloid dendritic cells. For AD, naïve CD57+ CD8+ T cells, and mature NKG2A+ natural killer cells showed nominally significant associations.

We did not observe immune cell changes in individuals with high genetic risk for ALS or AD, suggesting immune alterations may arise later in disease progression. Additional studies are required to validate our nominally significant findings.

Introduction

The development of neurodegenerative diseases (ND) arises from a complex interplay between genetic and environmental factors. Amyotrophic lateral sclerosis (ALS) is characterized by the degeneration of both first and second motor neurons resulting in severe symptoms of muscle weakness and ultimately neuromuscular respiratory failure, making it one of the most fatal forms of ND¹. Alzheimer's Disease (AD), the most common ND, is characterized by the accumulation of amyloid-beta and tau proteins in the brain, ultimately leading to cognitive decline and dementia². In this context, increasing evidence points to a critical role of the immune system in the pathogenesis of NDs³. For instance, alterations of different T cell subsets in the central nervous system as well as in the periphery have been reported in prevalent ALS⁴ and AD⁵ patients compared to healthy controls. Recently, our group investigated possible alterations of CD8+ TEMRA cells in healthy individuals with high genetic risk for AD⁶ based on prior reports suggesting differential levels of these cells in pre-clinical AD cases⁷. Although these analyses did not confirm a role for CD8+ TEMRA cells in the early pre-clinical phase of AD, the fact that changes in immune cell composition associated with AD or ALS are likely to precede the clinical onset of these diseases by years, prompted us to investigate early immune cell alterations using a more comprehensive approach. To this end, we investigated whether healthy individuals at high genetic risk for these diseases show alterations in relevant immune cell subtypes. Specifically, we calculated polygenic risk scores (PGS) for ALS and AD and examined their associations with 92 immune cell subtypes in ~450 individuals from the Berlin Aging Study II (BASE-II).

Methods

Study participants:

A total of 448 European individuals recruited as part of BASE-II were included in this study (described in ref. ⁶). BASE-II is a multi-institutional longitudinal cohort study from the larger Berlin area in Germany that investigates determinants and mechanisms of aging (described in detail elsewhere⁸). The effective dataset investigated here consists of individuals from two age groups, comprising 309 older adults (60% females, median age: 69 years, age range:60-82 years) and 139 younger adults (59% females, median age: 29, age range: 23-35, **Supplementary Figure 1**). Individuals with neurodegenerative diseases were excluded. All individuals were immunologically healthy (no fever or immune system-related diseases), had no diagnosis of neurodegenerative diseases, and were cognitively unimpaired (previously described ref. ⁷), and quality-controlled (QC) genetic as well as immune cell data were available.

Generation and processing of immune cell and genome-wide SNP data:

Peripheral blood mononuclear cells were isolated and analyzed by multi-colour flow cytometry as described previously^{6,9}. Briefly, two different immune cell antibody panels were used to distinguish between various immune cell subtypes, i.e. subsets of T cells, B cells, natural killer cells or monocytes (**Supplementary Table 1-2**). A 3 laser BD LSRII (BD biosciences) flow cytometer and DIVA6 software were utilized for the acquisition of the data, analyzed thereafter using FlowJo version 7.5 as described previously^{6,9}. QC'ed immune cells were quantified as proportions of the major immune cell types and distributions were transformed where appropriate (based on visual inspections) using one of the following approaches: log10, root, log(100-x) or square¹⁰ (**Supplementary Table 1**). As a final step, the data were z-transformed.

The genome-wide SNP data were generated and processed as described before⁶. In short, the Affymetrix Array 6.0 was used for genotyping, followed by imputation of unmeasured genotypes using the Haplotype Reference Consortium reference panel and finalized using standard QC procedures⁶. After filtering for a minor allele frequency (MAF) threshold of <0.01 and an imputation quality threshold of <0.3, we obtained a final set of 7,512,709 SNPs. As the imputation of the established AD *APOE* risk SNP e4 (rs429358) was suboptimal in this dataset (imputation $r^2=0.64$), this SNP was directly genotyped using “Taqman” technology (Thermo Fisher Scientific, Inc.)⁶.

The calculation of the PGS for ALS was based on 11 index SNPs that showed genome-wide statistical significance in European individuals from a recent large genome-wide association study (GWAS) on ALS¹¹ (**Supplementary Table 3, Supplementary Figure 2**). By including these 11 SNPs, we relaxed the minor allele frequency threshold for one SNP (rs80265967, MAF=0.00039) to maximize the number of SNPs included in the PGS. As sensitivity analysis we re-calculated an additional PGS and subsequent association statistics excluding this SNP (**Supplementary Table 3**). PGS calculations for AD were based on 70 genome-wide significant SNPs ($\alpha=5.0E-8$) from a recent GWAS on AD¹². Of note, the beta estimate and p value of the *APOE* e4 SNP were taken from an earlier AD GWAS¹³ by the same group, because it was excluded from the summary statistics in ref. ¹² (**Supplementary Table 4**). Due to missing data of the *APOE* e4 genotype in some BASE-II participants, some individuals had to be excluded from the analyses (n=18), resulting in a maximum of 430 individuals included in the AD analyses (**Supplementary Figure 2**). The PGS was z-transformed prior to statistical analyses.

Statistical analyses:

Linear regression analyses using the z-transformed PGS as exposure and the z-transformed immune cell proportions as outcome were performed, while adjusting for sex, the age group, and the first four genetic principal components to account for potential population substructures. The analyses were conducted using the 'lm' function in R. The main analyses were performed on the full sample and the subgroup of older BASE-II participants. Additionally, exploratory analyses stratifying for sex were performed on all nominally significant results ($\alpha=0.05$). Sensitivity analyses included an adjustment for cytomegalovirus (CMV) status and utilization of a modified PGS for ALS. The threshold for the false discovery rate (FDR) was set to 5%.

Results

The linear regression analyses of PGS on 92 immune cell subtypes in the full sample and the subgroup of older individuals did not yield significant signals at an FDR of 5%, neither for ALS nor for AD (**Supplementary Figure 3**). Despite the lack of study-wide significant findings, we observed several nominally significant ($\alpha=0.05$) – and hence potentially relevant – signals:

For ALS, five cell types passed the nominal threshold in analyses of the full BASE-II sample (**Figure 1**): This included the T cell subsets CD57+ CD8+ early-memory T cells ($p=0.006$), PD-1+ CD8+ T cells ($p=0.024$), and CD57+ CD4+ early-memory T cells ($p=0.038$), as well as non-classical monocytes ($p=0.015$) and myeloid dendritic cells ($p=0.047$) with variances explained (ΔR^2) ranging from 2.1% to 1.1% (**Table 1**). Notably, CD57+ CD8+ early-memory T cells exhibited consistent effect estimates across all analyses, and importantly, showed the strongest effect estimates in men with the PGS explaining 7% of the variance in immune cell distribution in this subset ($p=0.001$). The same cell type also reached nominal significance when analyses were restricted to the older age group ($\Delta R^2=2.4%$, $p=0.012$). In addition, PD-1+ CD8+ T cells, also exhibiting consistent effect estimates across strata, showed nominally significant association not only in the full sample but also in the subset of men ($\Delta R^2=2.6%$, $p=0.041$; **Table 1**).

For AD, we did not observe any trends of association ($\alpha=0.05$) in the full BASE-II dataset. However, upon restricting our investigation to the older subsample, i.e., that closer to a potential clinical onset of AD, showed trends for an association of higher genetic AD risk with subsets of CD8+ T cells and natural killer cells (**Figure 1**). Specifically, a higher PGS was associated with increased naïve CD57+ CD8+ T cells ($p=0.015$), increased mature NKG2A+ natural killer cells ($p=0.024$), and decreased

CD8+ late effector memory T cells ($p=0.038$) with $\Delta R2$ ranging from 2.4% to 1.7%. The latter two associations are driven by the male and female subgroup, respectively (**Table 1**).

Lastly, none of the results changed appreciably in the sensitivity analyses adjusting for CMV or using the alternative PGS for ALS (**Supplementary Table 5-6**).

Discussion

In this study, we performed a comprehensive investigation as to whether alterations in peripheral blood immune cell composition are evident in healthy participants at high genetic risk for AD or ALS. While none of the associations reached study-wide significance in ~450 BASE-II participants after FDR control, we observed several nominally significant associations between genetic ALS and AD risk and various immune cell subsets. These include subsets of CD4+, CD8+ T cells, monocytes, and myeloid dendritic cells for ALS as well as subsets of CD8+ T cells and natural killer cells for AD. To our knowledge, this is the first study to analyze such a broad spectrum of immune cell subtypes in healthy individuals at high risk of ALS or AD.

Notably, overall the nominally significant results showed consistent associations across analyzed strata. The most compelling results for ALS suggest a decrease in CD57+ CD8+ early-memory T cells and in PD-1+ CD8+ T cells. Interestingly, a recent landmark study investigated both effector and memory CD8+ T cells in human monogenic ALS patients with *SETX* mutations as well as PD-1+ CD8+ T cells in mice¹⁴. The authors found a general decrease of CD8+ effector and memory T cells, which is in agreement with our observations. While they did not show data for PD-1+ CD8+ T cells in humans, they described differential proportions of these cells in a

murine ALS model. However, effect directions differed in mice vs our human data, potentially reflecting the limited comparability between mouse models and humans¹⁴.

In individuals at high genetic risk for AD, the most noteworthy immune cell alterations highlighted by our analyses relate to increased naïve CD57+ CD8+ T cells and mature NKG2A+ natural killer cells. Consistent effect directions were observed for the main and all subgroup analyses. Notably the vast majority of studies investigating immune cells analyzed prevalent ALS and AD cases. While we are not aware of specific investigations of naïve CD57+ CD8+ T cells in AD, a number of studies reported on CD8+ T cell abnormalities in AD patients¹⁵. A role for natural killer cells in the pathogenesis of AD has been reported repeatedly^{15,16}, however their exact mechanisms including mature NKG2A+ natural killer cells remain elusive at this point in time. It is noteworthy that we found differences of immune cells for individuals of AD risk only in the smaller older age but not in the overall group, a pattern which we did not observe for ALS risk. This difference may be attributable to the approximately ten year later age of onset of AD compared to ALS.

The primary strengths of this study are an extensive and thorough clinical examination of all BASE-II participants, in-depth immune cell phenotyping, and a comparatively large sample size. Potential limitations of our work include limited statistical power to detect minor effects, the missing consideration of lifestyle and environmental ND risk factors, and the lack of data in non-European individuals, as discussed in previous work from our group¹⁷. In that study, we examined alterations in immune cell types in healthy individuals with a high genetic risk for Parkinson's disease¹⁷. Together with that study, our findings suggest that changes in immune cell composition may not manifest in healthy individuals with an increased genetic risk of the investigated NDs. Given the well-established role of the immune system

response in ND pathogenesis³, this suggests that immune cell alterations become manifest at later stages of disease development. Notwithstanding, our work highlights interesting trends of alterations in several innate and adaptive immune cell subtypes, which warrant further investigation in future research.

References

1. Mead RJ, Shan N, Reiser HJ, Marshall F, Shaw PJ. Amyotrophic lateral sclerosis: a neurodegenerative disorder poised for successful therapeutic translation. *Nat Rev Drug Discov.* 2023;22(3):185-212. doi:10.1038/s41573-022-00612-2
2. Knopman DS, Amieva H, Petersen RC, et al. Alzheimer disease. *Nat Rev Dis Prim.* 2021;7(1):33. doi:10.1038/s41572-021-00269-y
3. Zhang W, Xiao D, Mao Q, Xia H. Role of neuroinflammation in neurodegeneration development. *Signal Transduct Target Ther.* 2023;8(1):267. doi:10.1038/s41392-023-01486-5
4. Béland LC, Markovinovic A, Jakovac H, et al. Immunity in amyotrophic lateral sclerosis: blurred lines between excessive inflammation and inefficient immune responses. *Brain Commun.* 2020;2(2). doi:10.1093/braincomms/fcaa124
5. Bettcher BM, Tansey MG, Dorothée G, Heneka MT. Peripheral and central immune system crosstalk in Alzheimer disease — a research prospectus. *Nat Rev Neurol.* 2021;17(11):689-701. doi:10.1038/s41582-021-00549-x
6. Deecke L, Homann J, Goldeck D, et al. No increase of CD8+ TEMRA cells in the blood of healthy adults at high genetic risk of Alzheimer's disease. *Alzheimer's Dement.* 2024;20(4):3116-3118. doi:10.1002/alz.13709
7. Gericke C, Kirabali T, Flury R, et al. Early β -amyloid accumulation in the brain is associated with peripheral T cell alterations. *Alzheimer's Dement.* Published online June 14, 2023. doi:10.1002/alz.13136
8. Bertram L, Böckenhoff A, Demuth I, et al. Cohort Profile: The Berlin Aging Study II (BASE-II)†. *Int J Epidemiol.* 2014;43(3):703-712. doi:10.1093/ije/dyt018
9. Sbierski-Kind J, Goldeck D, Buchmann N, et al. T cell phenotypes associated with insulin resistance: results from the Berlin Aging Study II. *Immun Ageing.* 2020;17(1). doi:10.1186/s12979-020-00211-y
10. Deecke L, Homann J, Goldeck D, et al. Genome-wide association study reveals different T cell distributions in peripheral blood of healthy individuals at high genetic risk of type 1 diabetes and long COVID. *medRxiv Prepr under Rev.* Published online 2024.
11. van Rheenen W, van der Spek RAA, Bakker MK, et al. Common and rare variant association analyses in amyotrophic lateral sclerosis identify 15 risk loci with distinct genetic architectures and neuron-specific biology. *Nat Genet.* 2021;53(12):1636-1648. doi:10.1038/s41588-021-00973-1
12. Bellenguez C, Küçükali F, Jansen IE, et al. New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat Genet.* 2022;54(4):412-436. doi:10.1038/s41588-022-01024-z
13. Lambert JC, Ibrahim-Verbaas CA, Harold D, et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet.* 2013;45(12):1452-1458. doi:10.1038/ng.2802
14. Campisi L, Chizari S, Ho JSY, et al. Clonally expanded CD8 T cells characterize amyotrophic lateral sclerosis-4. *Nature.* 2022;606(7916):945-952. doi:10.1038/s41586-022-04844-5
15. Morris G, Berk M, Maes M, Puri BK. Could Alzheimer's Disease Originate in the Periphery and If So How So? *Mol Neurobiol.* 2019;56(1):406-434. doi:10.1007/s12035-018-1092-y
16. Jadidi-Niaragh F, Shegarfi H, Naddafi F, Mirshafiey A. The Role of Natural Killer Cells in Alzheimer's Disease. *Scand J Immunol.* 2012;76(5):451-456. doi:10.1111/j.1365-3083.2012.02769.x
17. Deecke L, Goldeck D, Ohlei O, et al. Immune cell distributions in the blood of healthy individuals at high genetic risk of Parkinson's disease. *medRxiv Prepr Serv Heal Sci.* 2024. doi:10.1101/2024.10.31.24316489

Legends to Figures

Figure 1. Nominally significant linear regression results of neurodegenerative disease polygenic risk score with immune cell distributions in the blood of healthy BASE-II participants.

Legend. This forest plot displays beta values and 95% confidence intervals of nominal significant association results linking amyotrophic laterals sclerosis (A) and Alzheimer's disease (B) and levels immune cell subtypes in blood as well as correlations between the corresponding nominally significant cell types per disease (C).

Table 1. Immune cells showing nominally significant associations with genetic risk profiles of amyotrophic lateral sclerosis and Alzheimer’s disease

	Immune cell type	Model	Beta	ΔR^2	P
ALS	CD57+ CD8+ early-memory T cells	All	-0.135	0.018	0.006
	CD57+ CD8+ early-memory T cells	Old group	-0.155	0.024	0.012
	CD57+ CD8+ early-memory T cells	Female	-0.051	0.003	0.424
	CD57+ CD8+ early-memory T cells	Male	-0.271	0.072	0.001
	PD-1+ CD8+ T cells	All	-0.111	0.012	0.024
	PD-1+ CD8+ T cells	Old group	-0.095	0.009	0.116
	PD-1+ CD8+ T cells	Female	-0.077	0.006	0.238
	PD-1+ CD8+ T cells	Male	-0.163	0.026	0.041
	Non-classical monocytes	All	-0.141	0.021	0.015
	Non-classical monocytes	Old group	-0.117	0.015	0.129
	Non-classical monocytes	Female	-0.133	0.018	0.059
	Non-classical monocytes	Male	-0.176	0.031	0.107
	CD57+ CD4+ early-memory T cells	All	-0.106	0.011	0.038
	CD57+ CD4+ early-memory T cells	Old group	-0.118	0.014	0.057
	CD57+ CD4+ early-memory T cells	Female	-0.104	0.011	0.112
	CD57+ CD4+ early-memory T cells	Male	-0.108	0.011	0.203
	Myeloid dendritic cells	All	-0.108	0.012	0.047
	Myeloid dendritic cells	Old group	-0.045	0.002	0.507
	Myeloid dendritic cells	Female	-0.128	0.017	0.058
	Myeloid dendritic cells	Male	-0.062	0.004	0.518
AD	Naïve CD57+ CD8+ T cells	All	0.081	0.007	0.118
	Naïve CD57+ CD8+ T cells	Old group	0.156	0.024	0.015
	Naïve CD57+ CD8+ T cells	Female	0.116	0.013	0.158
	Naïve CD57+ CD8+ T cells	Male	0.206	0.039	0.053
	Mature NKG2A+ natural killer cells	All	0.081	0.007	0.138
	Mature NKG2A+ natural killer cells	Old group	0.158	0.024	0.024
	Mature NKG2A+ natural killer cells	Female	0.098	0.009	0.267
	Mature NKG2A+ natural killer cells	Male	0.291	0.079	0.018
	CD8+ late effector memory T cells	All	-0.060	0.004	0.249
	CD8+ late effector memory T cells	Old group	-0.133	0.017	0.038
	CD8+ late effector memory T cells	Female	-0.112	0.013	0.022
	CD8+ late effector memory T cells	Male	-0.071	0.005	0.491

Legend. This table visualizes the results on all immune cell analyses that show nominally significant ($\alpha=0.05$) association in a least one analysis (i.e., analyzing all or restricting to the older group). The linear regression analyses were performed across 448 healthy individuals from the Berlin Aging Study II (BASE-II) using polygenic scores (PGS) for amyotrophic laterals sclerosis (ALS) and Alzheimer’s disease (AD) as exposure and immune cell proportions as outcome. Beta = effect estimate of regression analyses of z-transformed immune cell distributions on z-transformed polygenic risk scores. ΔR^2 = variance of immune cell distribution explained by the polygenic risk score.

Author Contributions

Conceptualization: C.M.L.; methodology: L.D., C.M.L.; formal analysis or interpretation of the data: L.D., O.O., J.H., C.M.L.; data acquisition: D.G., I.D., G.P., L.B.; writing—original draft preparation: L.D., C.M.L.; writing—review and editing: D.G., O.O., J.H., I.D., L.B., G.P.; supervision, C.M.L. All authors have read and agreed to the final version of the manuscript.

Funding

The BASE-II research project (co-PIs: Lars Bertram, Ilja Demuth, Denis Gerstorf, Ulman Lindenberger, Graham Pawelec, Elisabeth Steinhagen-Thiessen, and Gert G. Wagner) has been supported by the German Federal Ministry of Education and Research (Bundesministerium für Bildung und Forschung, BMBF) under grant numbers #16SV5536K, #16SV5537, #16SV5538, #16SV5837, #01UW0808,01GL1716A, and 01GL1716B, and by the Max Planck Institute for Human Development, Berlin, Germany. Additional contributions (e.g., equipment, logistics, personnel) were made from each of the other participating sites. The responsibility for the contents of this publication lies with its authors. C.M.L. was supported by the Heisenberg program of the German Research Foundation (DFG; LI 2654/4-1) by the Cure Alzheimer's Fund, and by the CReATe Consortium. CReATe (U54 NS092091) is part of the Rare Diseases Clinical Research Network (RDCRN), an initiative of the Office of Rare Diseases Research (ORDR), NCATS and funded through collaborations between NCATS, NINDS and the ALS Association.

Data Availability

All summary statistics have been made available in the supplementary material of this manuscript. Raw and source data are available upon reasonable request. Interested researchers may contact the scientific BASE-II coordinator, Ludmila

Müller, Imueller@mpib-berlin.mpg.de. Additional information is available on the BASE-II website: <https://www.base2.mpg.de/7549/data-documentation>.

Acknowledgements

We are grateful to all BASE-II participants.


Conflict of interests

None of the authors reports any conflict of interest.




Ethical approval

All BASE-II participants provided written informed consent before participation and the study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of the Charité-Universitätsmedizin Berlin – approval number EA2/029/09.

A. Immune cell type	Model	Beta	P value
CD8+ME_CD57+	All	-0.134	0.005
CD8+ME_CD57+	Old group	-0.155	0.012
CD8+ME_CD57+	Male	-0.271	0.000
CD8+ME_CD57+	Female	-0.051	0.424
CD8+PD-1+	All	-0.111	0.024
CD8+PD-1+	Old group	-0.095	0.116
CD8+PD-1+	Male	-0.163	0.041
CD8+PD-1+	Female	-0.077	0.238
CD4+ME_CD57+	All	-0.106	0.038
CD4+ME_CD57+	Old group	-0.118	0.057
CD4+ME_CD57+	Male	-0.108	0.203
CD4+ME_CD57+	Female	-0.104	0.112
M_N	All	-0.141	0.015
M_N	Old group	-0.117	0.129
M_N	Male	-0.176	0.107
M_N	Female	-0.132	0.059
mDC	All	-0.108	0.047
mDC	Old group	-0.045	0.507
mDC	Male	-0.062	0.518
mDC	Female	-0.128	0.058



B. Immune cell type	Model	Beta	P value
CD8+naive_CD57+	All	0.081	0.118
CD8+naive_CD57+	Old group	0.156	0.015
CD8+naive_CD57+	Male	0.206	0.053
CD8+naive_CD57+	Female	0.116	0.158
NK_mature_NKG2A+	All	0.081	0.138
NK_mature_NKG2A+	Old group	0.158	0.024
NK_mature_NKG2A+	Male	0.291	0.018
NK_mature_NKG2A+	Female	0.098	0.267
CD8+LEM	All	-0.06	0.249
CD8+LEM	Old group	-0.133	0.038
CD8+LEM	Male	-0.071	0.491
CD8+LEM	Female	-0.19	0.022



medRxiv preprint doi: <https://doi.org/10.1101/2024.11.13.24316906>; this version posted December 21, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-ND 4.0 International license](https://creativecommons.org/licenses/by-nd/4.0/).