1 Prevalence and Risk Factors of Diarrhea Among Children Under Five Years in Northern

2 Kenya's Drylands: A Longitudinal Study

- 3 Bonventure Mwangi^{1*}, Valerie L. Flax², Faith Thuita^{3,4}, Joshua D. Miller⁵, Chessa Lutter²,
- 4 Dickson Amugsi¹, Estelle Sidze¹, Linda Adair⁵, Esther Anono¹, Hazel Odhiambo¹, Stephen
- 5 Ekiru¹, Gillian Chepkwony¹, Monica Ng'ang'a³, Albert Webale³, Elizabeth Kimani-Murage¹,
- 6 Calistus Wilunda¹
- 7
- 8 ¹ African Population and Health Research Center, APHRC Campus, Kitisuru, Nairobi,
- 9 Kenya.
- 10 ² RTI International, Research Triangle Park, North Carolina, United States of America.
- ³ RTI International, Nairobi, Kenya.
- ⁴ Department of Public and Global Health, University of Nairobi, Nairobi, Kenya.
- ⁵ Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC,
- 14 United States of America.
- 15
- 16 *Corresponding author
- 17 African Population and Health Research Center, P.O. Box 10787-00100, APHRC Campus,
- 18 Kitisuru, Nairobi, Kenya.
- 19 E-mail: <u>bmwangi@aphrc.org</u> (BM)

20 Abstract

21 Diarrhea is the third leading cause of malnutrition and mortality in children under five globally.

However, a few studies have examined predictors of child diarrheal disease in arid or semi-arid
regions of sub-Saharan Africa. This study aimed to assess the prevalence and risk factors of

24 diarrhea among children under five in the drylands of Northern Kenya.

25 Data are from a longitudinal, population-based study of children younger than 36 months at 26 recruitment (N = 1.211) and their caregivers in Turkana County, Kenya. Households were 27 followed from May 2021 to September 2023, with survey data collected every four months 28 across six waves. Caregivers reported on household conditions and observed episodes of 29 diarrhea among index children in the two weeks before the survey. Trends in the prevalence of 30 diarrhea were stratified by administrative zone, livelihood zone, and child sex. Risk factors of 31 child diarrhea across survey waves were identified using multivariable generalized estimating 32 equations.

33 Diarrhea prevalence decreased from 32.1% [95% confidence interval (CI): 28.3%-36.1%] at 34 Wave 1 to 8.7% (95% CI: 6.3%-11.7%) at Wave 6. Risk factors for diarrhea included 35 caregivers alcohol consumption [adjusted odds ratio (AOR) = 1.30; 95% CI: 1.04-1.62], households experiencing three (AOR = 1.78; 95% CI: 1.29-2.45) or four (AOR = 2.58; 95% 36 37 CI: 1.86-3.58) climatic, biological, economic or conflict shocks in the prior 4 months compared 38 to those experiencing less than 2 shocks, households with moderate (AOR = 1.25; 95% CI: 39 1.04-1.50) or high (AOR = 1.50; 95% CI: 1.22-1.85) water insecurity in the prior 4 weeks 40 compared to those with no-to-marginal water insecurity, and child wasting (AOR = 1.22; 95%) 41 CI: 1.05–1.41).

42 These findings suggest that multisectoral interventions that reduce alcohol consumption among 43 women, improve access to safe water services, manage malnutrition, and mitigate household 44 shocks could reduce the burden of diarrhea among child under five in this region.

45 Introduction

Diarrhea, the passage of three or more loose or liquid stools per day, is the third leading cause of death among children under five years of age globally [1]. Approximately 90% of these deaths occur in sub-Saharan Africa and South Asia [2]. Diarrhea also increases the risk of dehydration, appetite loss, malnutrition, electrolyte deficiencies, infectious disease, delayed physical growth, and cognitive impairments [1,3–5]. Malnutrition and diarrhea are interrelated, with each condition exacerbating the other, creating a cycle that severely compromises the long-term health and development of impacted children [6,7].

53

54 Previous studies in sub-Saharan Africa have identified factors associated with a higher risk of child diarrhea. These include younger child age [8–12], low maternal education [12–14], young 55 56 maternal age [12,15], maternal unemployment [12,15], poor hand hygiene [16–18], open 57 defecation [19–21], improper disposal of children's stool [22,23], and households with more 58 members [12,24,25]. Recent research also underscores the influence of behavioral, 59 sociocultural, and socioeconomic factors on the prevalence of diarrhea in rural areas [26]. 60 Given this, interventions that increase access to child-friendly sanitation infrastructure, change hand hygiene behavior and expand access to improved water services have been recommended 61 62 to reduce the burden of child diarrhea [26,27]. A longitudinal study conducted in urban settings of developing countries confirmed the importance of sanitation as a major determinant of child 63 64 diarrhea [28].

65

In sub-Saharan Africa's arid and semi-arid lands (drylands), climate change has intensified
droughts, significantly impacting livelihoods, household water, and food security [29–33].
Communities living in drylands experience food and water insecurity and predominantly rely
on nomadic or semi-nomadic pastoralist livelihoods, increasing the risk of malnutrition [34].

Recurrent household climatic shocks in the region reduce agricultural productivity, damage critical infrastructure, and negatively impact health and well-being [15,35–37]. Additionally, limited access to improved water, sanitation, and hygiene services in the area increases exposure to vectors of diarrheal disease [3,32,38,39]. Moreover., households facing health shocks find themselves impoverished due to loss of income to facilitate access to quality health care and payment of health insurance [40,41].

76

While the distinctive conditions of drylands pose heightened risks for diarrhea, limited research investigates how environmental, socioeconomic, and behavioral factors intersect to influence diarrhea risk in these regions, particularly in Northern Kenya. To address this knowledge gap, we aim to assess the prevalence and risk factors of diarrhea among children under five in the drylands of Northern Kenya.

82

83 Materials and methods

84 Study area

85 This study was carried out as part of the USAID Nawiri program [42], to inform program design and adaptation. The study used data collected in Turkana County, which is in north-86 western Kenya and has seven sub-counties, 30 wards, and 156 sub-locations [43]. This region 87 88 is characterized by low annual rainfall, high year-round temperatures, and a topographically 89 varied landscape that ranges from semi-arid to arid. Traditionally, the Turkana people are 90 nomadic pastoralists, relying on livestock and migrating in search of pasture to support their 91 livelihoods. In addition to pastoralism, some communities practice agro-pastoralism, engaging 92 in crop farming through irrigation along River Turkwel, and fishing on Lake Turkana [44].

93

94 Study design

95 Caregivers with children under 36 months were eligible to participate. A multistage sampling 96 strategy was used to ensure that findings were representative at the regional and sub-regional 97 levels. In the first stage, the population was stratified based on administrative zones. Within 98 each stratum, villages were randomly selected. Study staff identified all households in selected 99 villages to create a sampling frame for eligible households. Households were then randomly 100 selected from this frame and invited to participate. Further details on the study design are 101 available in the study protocol [45].

102

103 Data collection

Data were collected from May 2021 to September 2023 across six survey waves conducted every four months. Trained enumerators administered household and caregiver questionnaires to participating households. Study staff checked the integrity and validity of the data daily to ensure data quality.

108

109 **Dependent variable**

110 Child diarrhea in the 2 weeks before the survey (yes/no) was based on caregivers' self-report.111

112 Independent variables

Independent variables included distal, intermediate, and proximal factors, with wasting considered an additional independent variable given its bidirectional relationship with diarrhea (Figure 1). Distal factors included administrative and livelihood zones. Intermediate factors included head of household age (years), household head sex (man/woman), household wealth,

number of household members, child age (months), child sex (male/female), child nutrition
indicators, caregiver age (years), caregiver education (formal/no formal education), caregiver
marital status (in union: currently living together; not in union: separated, divorced, widowed
or single), and whether the household experienced shocks in the four months before the survey.
Proximal factors included water, sanitation, and hygiene (WASH) conditions and behaviors.
Caregivers' handwashing after visiting the toilet in the last 24 hours (yes/no) and household
practice of open defection (yes/no) were based on self-report.

124

The county was divided into four administrative zones (Central =1, North =2, South = 3 and West = 4) which comprised sub-counties: Central (Turkana Central and Loima), North (Turkana North and Kibish), South (Turkana East and Turkana South) and West (Turkana West). The livelihood zones in Turkana County were broadly categorized into four zones namely pastoral, agro-pastoral, fisher folks and urban/peri-urban. However, due to small sample sizes, the agro-pastoral, fisherfolk and urban/peri-urban zones were categorized as nonpastoral livelihood zone.

132

133 The household wealth was determined based on a factor analysis of household assets, housing materials, and access to utilities, and divided into tertiles (low, medium, and high). Household 134 135 shocks were categorized into four types: climatic (excessive rains/flooding, variable 136 rain/drought), biological (livestock/crop/human disease outbreak, crop pest invasion, weed 137 outbreak, and severe illness), economic (loss of livelihood, increased prices in 138 food/agricultural/livestock inputs, loss of land/rental property, youth unemployment, death of 139 a household member, delay in food assistance, delay in other safety net programs, and fire) and 140 conflict (theft/destruction of assets, theft of livestock, domestic violence, and community

141 conflicts) and were binary variables. Further, the shock responses were summed up, and
142 households were categorized as having experienced less than two, two, three, and four shocks.
143

144 Figure 1. A conceptual framework for diarrhea risk factors among children under five

145

146 Household water insecurity was measured using the 12-item Household Water Insecurity 147 Experiences (HWISE) Scale [46]. Caregivers were asked to report how frequently in the prior 148 4 weeks they or others in their household experienced diverse water problems (0 = "never", 1 149 = "rarely", 2 = "sometimes", 3 = "often" or "always"). Responses were summed and 150 households were categorized as experiencing no-to-marginal (scores of 0-2), low (3-11), 151 moderate (12-23), or high (24-36) water insecurity. Child wasting was assessed using weightfor-height/length Z-scores (WHZ < -2 SD), based on the 2006 World Health Organization 152 153 (WHO) child growth standards [47].

154

155 **Data analysis**

Descriptive analyses were performed to examine the weighted distribution of distal factors, 156 157 household, caregiver, and child characteristics at Wave 1. We performed univariate analysis to 158 assess the associations between the dependent variable (child diarrhea) and the independent 159 variables. The independent variables with a *p*-value < 0.20 were retained and used to build the 160 final multivariable logistic regression models. To identify salient risk factors of diarrhea, iterative modeling was used to account for the complex interrelationships between the different 161 162 independent variables of interest, adjusting for either distal, intermediate, or proximal factors 163 based on the conceptual framework in Fig. 1 [48]. Multivariable logistic regressions of child diarrhea and independent variables using generalized estimating equations with working 164 165 correlation matrices were used to account for the within-subject correlation of repeated

166 measurements across survey waves [49,50]. Only those variables with p-value < 0.05 were 167 considered factors associated with diarrhea among children under five years. Adjusted odds 168 ratios (AOR) and 95% confidence intervals (CIs) were obtained from the adjusted 169 multivariable logistic regression models and interpreted for significant covariates. All statistical analyses were performed using Stata 18 (Stata Corp, College Station, TX USA). The 170 171 survey proportion command in Stata was used to compare each period and potential factors. 172 We performed a non-parametric Cochran-Armitage test for the trend in the prevalence across 173 time using nptrend command and p-value. The xtgee command was used to perform univariate 174 and multivariable logistic regressions.

175

176 Ethical Considerations

The study was approved by the African Medical and Research Foundation Ethical and Scientific Review Committee (Amref ESRC P905/2020) and the National Commission for Science, Technology, and Innovation of Kenya. The institutional review boards at the African Population and Health Research Center and Research Triangle Institute International signed a reliance agreement. Informed consent was obtained from the legal guardians of participating children.

183

184 **Results**

A total of 1,211 households with eligible children were included in this study and their baseline descriptive statistics are summarised in Table 1. Overall, 63.9%, lived in the pastoral livelihood zone. Around two-thirds (61.9%) of the households were headed by men, while females headed a third (38.1%). 44.6% and 30.6% of households experienced three and four shocks in the 4 months before the survey respectively. Among caregivers, the average age was 33 years, 86.4%

- 190 had no formal education, 49.6% lived with 3 to 5 children, and 85.2% were in a marital union.
- 191 Almost half of the children were boys (54.8%) with 21.8% of children wasted and 32.1%
- 192 reported to have experienced diarrhea in the two weeks before the survey.
- 193

194 Table 1. Distribution of distal factors, household, caregiver, and child characteristics at

195 baseline, Turkana County (N = 1,211)

Characteristic	Weighted %
Distal factors	
Livelihood zone	
Pastoral	63.9
Non-pastoral	36.1
Administrative zone	
Central	21.0
North	25.6
South	24.1
West	29.2
Household characteristics	
Age of the household head (years) a	39.5±13.4
Sex of household head	
Man	61.9
Woman	38.1
Household size ^a	6.1±2.1
Wealth tertile	
Lowest	45.6
Middle	33.8
Highest	20.6
Shocks experienced in the last 4 months before the survey	
Less than two	4.2
Two	20.6
Three	44.6
Four	30.6
Caregiver characteristics	
Age (years) ^a	33.1±10.2
Education	
Formal education	13.6
No formal education	86.4
Marital status	
In union	85.2
Not in union	14.8
Number of living children	

Less than 3	25.9
3-5	49.6
6 or More	24.4
Current alcohol consumer	8.3
Child characteristics	
Sex	
Male	54.8
Female	45.2
Age (months) ^a	17.3±10.6
Wasting (Weight-for-height/length z score < -2SD)	
Yes	21.8
Experience of diarrhea in the 2 weeks before the survey	
Yes	32.1

^a Mean ± standard deviation

196

The overall weighted prevalence of diarrhea and selected variables in each period were 197 198 examined (Table 2). The data revealed that 32.1% of the children were reported to have 199 experienced diarrhea in Wave 1. There was a noticeable decrease from 32.1% to 27.0% in one 200 year, followed by a further steady decrease to Wave 4. Between Wave 4 to 5, there was 201 a notable increase in the prevalence by approximately 2%, followed by a decrease of 9.1% to 202 Wave 6. A similar trend was observed in the livelihood, administrative zones and child sex. A 203 statistically significant trend in the prevalence of diarrhea was observed in the overall estimates 204 and the livelihood zones. The prevalence of diarrhea in administrative zones and child age did 205 not have a statistically significant trend.

206

207

medRxiv preprint doi: https://doi.org/10.1101/2024.11.13.24317266; this version posted November 13, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in

perpetuity. It is made available under a CC-BY 4.0 International license .

Variable	Wave 1	Wave 2	Wave 3	Wave 4	Wave 5	Wave 6	P-value a
Overall	32.1 [28.3, 36.1]	30.0 [26.0, 34.4]	27.9 [24.1, 32.1]	15.7 [12.5, 19.6]	17.8 [14.2, 21.9]	8.7 [6.3, 11.7]	<0.001
Livelihood zone		-		-	-	-	
Pastoral	31.2 [26.4, 36.4]	27.8 [22.5, 33.7]	28.6 [24.0, 33.7]	12.5 [8.8, 17.5]	16.9 [12.7, 22.2]	7.0 [4.3, 11.3]	< 0.001
Non-pastoral	33.6 [27.8, 40.0]	31.9 [26.1, 38.2]	26.9 [20.9, 33.9]	22.4 [17.3, 28.4]	19.6 [13.9, 26.8]	11.9 [8.5, 16.4]	
Administrative	-	-	-	-	-	-	
zone							
Central	25.8 [20.4,	32.8 [24.9,	26.8 [20.3,	18.4 [14.7,	22.3 [18.0,	9.4 [5.7,	0.004
Central	32.1]	41.7]	34.5]	22.9]	27.2]	14.9]	
North	41.6 [33.1,	22.5 [14.5,	14.8 [8.7,	22.1 [14.9,	14.2 [7.6,	14.1 [8.1,	0.868
ivoitii	50.6]	33.2]	24.1]	31.5]	24.9]	23.5]	
South	34.8 [27.5,	37.9 [30.8,	35.3 [28.6,	17.3 [11.6,	25.3 [18.2,	8.2 [5.6,	0.335
South	42.8]	45.5]	42.6]	24.9]	34.0]	11.9]	
West	25.9 [20.4,	28.2 [21.4,	35.8 [30.6,	6.9 [3.9,	11.9 [7.2,	2.9 [1.4,	0.035
West	32.5]	36.2]	41.5]	12.3]	18.9]	6.1]	
Child sex							
Male	33.9 [28.4,	31.7 [26.1,	26.2 [21.0,	15.1 [11.1,	18.9 [14.3,	6.1 [4.0,	0.294
whate	40.0]	37.8]	32.2]	20.1]	24.4]	9.1]	
Female	29.7 [24.5,	28.2 [22.5,	29.8 [24.1,	16.5 [12.2,	16.6 [12.2,	11.5 [7.7,	
	35.7]	34.7]	36.3]	21.9]	22.0]	16.9]	

208 Table 2. Weighted prevalence of diarrhea by livelihood zone, administrative zone, and child sex across survey waves (%, [95% CI]). 209

^{*a*} *Test for the trend in the prevalence of diarrhea across time*

Note: The study data collection period; Wave 1 (May/June 2021), Wave 2 (November 2021), Wave 3 (May/June 2022), Wave 4 (October /November 2022, Wave 5 (March 2023) and Wave 6 (August/September 2023).

210

211 In bivariate analysis, the following factors were associated with diarrhea at p < 0.2: distal factors 212 (livelihood and administrative zones), intermediate factors (child age and sex, caregiver age, 213 caregiver number of living children, caregiver alcohol consumption, head of household age, household shocks), proximal factors (household water insecurity, handwashing of caregiver 214 215 after visiting toilet, household open defecation) and childhood wasting (S1 Table). After 216 adjusting for potential confounders in multivariable models (Table 3), children with caregivers who consumed alcohol were found to have higher odds of diarrhea (AOR = 1.30; 95% CI: 1.04 217 218 - 1.62) compared to those whose caregivers did not. In addition, children in households that 219 experienced three (AOR = 1.78; 95% CI: 1.29 - 2.45) or four (AOR = 2.58; 95% CI: 1.86 -220 3.58) shocks had higher odds of diarrhea compared to those in households that experienced 221 fewer than two shocks. Relative to children living in households with no-to-marginal water

222	insecurity, those in households with moderate (AOR = 1.25 ; 95% CI: $1.04 - 1.50$) or high-water
223	insecurity (AOR = 1.50; 95% CI: 1.22 - 1.85) had higher odds of diarrhea. The analysis
224	revealed that the households within the pastoral livelihood zone, children's age, number of
225	children living with the caregiver and handwashing practice by the caregiver were protective
226	factors against diarrhea.

227

228 Table 3. Factors associated with diarrhea among children under five in Turkana County,

229 Kenya.

Variable	AOR [95% CI]	P-value
Survey period (Ref: Wave 1)		
Wave 2	1.05 [0.81, 1.24]	0.566
Wave 3	0.91 [0.76, 1.08]	0.272
Wave 4	0.45 [0.37, 0.54]	< 0.001
Wave 5	0.52 [0.43, 0.63]	< 0.001
Wave 6	0.21 [0.17, 0.27]	< 0.001
Distal factors		
Livelihood zone (Ref. non-pastoral)		
Pastoral	0.87 [0.77, 0.99]	0.038
Administrative zone (Ref: Central)		
North	0.92 [0.73, 1.18]	0.523
South	1.18 [0.99, 1.42]	0.064
West	0.94 [0.78, 1.13]	0.488
Intermediate factors (adjusted for distal factors)		
Child age (months)	0.97 [0.96, 0.98]	< 0.001
Child sex (Ref: Male)		
Female	1.03 [0.90, 1.19]	0.675
Caregiver age (years)	0.99 [0.98, 1.00]	0.029
Caregiver number of living children (Ref: < 3)		
3-5	0.79 [0.67, 0.93]	0.006
6 or more	0.93 [0.74, 1.16]	0.494
Caregiver alcohol consumption (Ref: No)		
Yes	1.30 [1.04, 1.62]	0.020
Shocks experienced in the last 4 months before the survey (Ref: less than two)		
Two	1.15 [0.83, 1.59]	0.412
Three	1.78 [1.29, 2.45]	< 0.001
Four	2.58 [1.86, 3.58]	< 0.001

Proximal factors (adjusted for distal and intermediate factors)

Household water insecurity (Ref: no-to-marginal)

Low	1.20 [0.98, 1.47]	0.080	
Moderate	1.25 [1.04, 1.50]	0.020	
High	1.50 [1.22, 1.85]	< 0.001	
Caregiver wash hands after using the toilet (Ref: no)			
Yes	0.87 [0.74, 1.03]	0.090	
Household practised open defecation (Ref: No)			
Yes	1.04 [0.88, 1.22]	0.650	
Immediate factor (adjusted for distal, intermediate and proximal factors)			
Wasted (WHZ < -2SD)	1.22 [1.05, 1.41]	0.010	
Note: The study data collection period: Ways 1 (May/hung 2021)	Ways 2 (November 2021)	Nava 3	

Note: The study data collection period: Wave 1 (May/June 2021), Wave 2 (November 2021), Wave 3 (May/June 2022), Wave 4 (October /November 2022, Wave 5 (March 2023) and Wave 6 (August/September 2023).

230

Discussion 231

In this study, we found that caregiver alcohol consumption, household water insecurity, 232 233 household shocks, and child wasting analysis were associated with higher odds of diarrhea in

234 children. These findings align with prior studies and can help to inform policies and programs

235 that aim to reduce the high burden of child diarrhea in Turkana County, Kenya.

236

237 Households in the drylands of sub-Saharan Africa suffer from frequent shocks, extended periods of poverty, and vulnerability, which hinder access to food, water and primary 238 239 healthcare. Studies have shown that shocks often exacerbate existing vulnerabilities, particularly among marginalized communities that heavily depend on livestock [33,51]. 240 241 Further, research indicates that these vulnerabilities adversely affect child nutrition, a key risk 242 factor for childhood morbidity and mortality, as families struggle to maintain adequate food 243 intake and access to clean water during crises [52–54]. Our study provides further evidence on 244 the effects of household experience of shocks on the risk of childhood diarrhea in an arid land 245 in Kenya.

246

247 Pastoral livelihood in sub-Saharan Africa increases vulnerability to shocks due to recurrent 248 droughts and famines rendering many destitute and chronically dependent on food aid. Some 249 households adapt to this situation by diversifying into activities that generate cash income, such as petty trade, crafting, weaving, and basketry. Research has shown that when pastoral 250 251 households experience loss of livestock, children are particularly vulnerable to illnesses such 252 as diarrhea and respiratory infections, primarily due to poor nutrition and limited access to 253 primary healthcare [55,56]. Further, a study conducted in three pastoralist communities in 254 northern Kenya provides evidence for the importance of a contextualized approach detailing 255 how social environments that include endemic conflict produce variations in health [57]. The 256 present study has shown that pastoral livelihood is a protective factor for childhood diarrhea 257 contrary to the previous findings, opening a research question for further investigations in the 258 future.

259

Our study revealed that children of caregivers who reported alcohol consumption had significantly higher odds of experiencing diarrhea. While caregiver alcoholism has been associated with persistent malnutrition in children under five [58], there is limited research directly connecting it to diarrhea. Qualitative data from Wave 1 of this longitudinal study suggested that alcohol use compromised caregivers' abilities to provide adequate and hygienic care, likely contributing to the observed association [59].

266

Greater experiences of household water insecurity (which includes issues with water availability, accessibility, and sufficiency for diverse household uses) have been similarly positively associated with diarrhea risk among children under 5 in Cameroon [60], Ethiopia [61], and Nigeria [62]. Most studies, however, have only examined child diarrhea related to

directly observable, supply-side water indicators, such as water source and quality [63,64]. Our findings demonstrate that experiential measures of water insecurity provide complementary information to traditional water metrics and make clear that water insecurity is an important modifiable risk factor that should be targeted to improve child health and well-being.

275

Malnutrition, particularly wasting, was also significantly associated with the odds of child diarrhea. Malnutrition undermines child development by depleting nutritional reserves and compromising the immune system, thereby increasing susceptibility to diarrheal infections [65]. Diarrhea, in turn, can exacerbate malnutrition, leading to a bidirectional relationship between these two conditions that ultimately increases the risk of child mortality [66–68]. Previous studies have associated childhood wasting with severe diarrhea episodes [69,70]; our findings further support wasting as a key risk factor for diarrheal infection.

283

284 Our study had several limitations. First, caregiver-reported child diarrhea is subject to recall 285 bias. Further, we did not ask about the type of diarrhea observed (e.g., acute watery, acute bloody, or persistent), as each might have different health impacts. Systematic reviews have, 286 287 however, noted a lack of evidence supporting the validity of instruments used in pediatric 288 diarrheal disease trials [71]. Future research is needed to develop and validate better child 289 diarrhea measures so that analyses can be disaggregated by diarrhea sub-type. Second, this 290 longitudinal study was not originally designed to assess diarrhea as a primary outcome, 291 although the large sample size reduces the risk of underpowered analyses.

292

293 Conclusion

In this study of diarrhea among children under five in the drylands of Northern Kenya, we found that the prevalence of diarrhea decreased as children got older. Baseline surveys were

conducted during the dry season, which may have also contributed to the prevalence of diarrhea
observed at this time point. The findings on diarrhea prevalence disaggregated across survey
waves can serve as useful referents for future targeted interventions in this region, particularly
among non-pastoralists and households in the North zone, where the prevalence was highest.
Key risk factors for diarrhea in Turkana County included caregiver alcohol use, household
experience of shocks, household water insecurity, and child wasting.

302

A multisectoral approach is recommended to reduce the burden of diarrheal disease among children under five in this region. Such an approach should prioritize interventions that improve access to improved WASH services, address malnutrition prevention and management, and reduce the burden of household economic and climatic shocks. Lastly, there's a need for intervention to reduce alcohol consumption among women as this may contribute towards reducing the burden of diarrhea among children under five years.

309

310 Funding information

The Nawiri Longitudinal Study was made possible by the generous support of the American
people through the United States Agency for International Development (USAID) (Award
Number: 72DFFP19CA00003). The contents of this paper are the responsibility of the
authors and do not necessarily reflect the views of USAID or the United States Government.

316 **Conflict of interest**

317 All authors declare that they do not have any competing interests to declare.

318

319 Data sharing

- 320 The data underlying this study will be made available to researchers through the APHRC's
- 321 Microdata Portal (<u>https://aphrc.org/microdata-portal/</u>).
- 322

323 Author contributions

- 324 FT, DA, VLF, CKL, AW, and ES designed the study. CW, DA, BM, EA, HO, SE, and GC
- 325 implemented fieldwork. CW, ES, FT, and EK-M provided study oversight. BM, JDM, and
- 326 CW analyzed the data. BM prepared the first draft of the manuscript and incorporated
- 327 comments from co-authors. All authors contributed to the revision of the manuscript and
- 328 approved the final version.

329 **References**

330 331	1.	World Health Organization (WHO). Diarrhoeal disease. Fact Sheets. Mar 2024. Available: https://www.who.int/news-room/fact-sheets/detail/diarrhoeal-disease
332 333 334 335	2.	Troeger C, Blacker BF, Khalil IA, Rao PC, Cao S, Zimsen SRM, et al. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of diarrhoea in 195 countries: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Infect Dis. 2018;18: 1211–1228.
336 337	3.	Young H. Nutrition in Africa's drylands: a conceptual framework for addressing acute malnutrition. Feinstein International Center, Tufts University. 2020.
338 339 340 341 342	4.	Council AW. Vulnerability of arid and semi-arid regions to climate change–Impacts and adaptive strategies. Perspective Document for the 5th World Water Forum, World Water Council, Marseille, Co-Operative Programme on Water and Climate (CPWC). The Hague, IUCN, Gland and International Water Association (IWA) The Hague; 2009.
343 344 345	5.	Bulled N, Singer M, Dillingham R. The syndemics of childhood diarrhoea: a biosocial perspective on efforts to combat global inequities in diarrhoea-related morbidity and mortality. Glob Public Health. 2014;9: 841–853.
346 347 348	6.	Guerrant RL, Oriá RB, Moore SR, Oriá MOB, Lima AAM. Malnutrition as an enteric infectious disease with long-term effects on child development. Nutr Rev. 2008;66: 487–505.
349 350 351	7.	Moore SR, Lima AAM, Conaway MR, Schorling JB, Soares AM, Guerrant RL. Early childhood diarrhoea and helminthiases associate with long-term linear growth faltering. Int J Epidemiol. 2001;30: 1457–1464.
352 353 354	8.	Colombara D V, Hernández B, McNellan CR, Desai SS, Gagnier MC, Haakenstad A, et al. Diarrhea prevalence, care, and risk factors among poor children under 5 years of age in Mesoamerica. Am J Trop Med Hyg. 2016;94: 544.
355 356 357	9.	Pokharel M, Sherchand JB, Upreti HC, Katuwal A, Gauchan P. A perspective study on the etiology of diarrhea in children less than 12 years of age attending Kanti Children's Hospital. Journal of Nepal Paediatric Society. 2009;29: 10–16.
358 359 360	10.	Atnafu A, Sisay MM, Demissie GD, Tessema ZT. Geographical disparities and determinants of childhood diarrheal illness in Ethiopia: further analysis of 2016 Ethiopian Demographic and Health Survey. Trop Med Health. 2020;48: 1–12.
361 362 363 364	11.	Buchwald AG, Verani JR, Keita AM, Jahangir Hossain M, Roose A, Sow SO, et al. Etiology, presentation, and risk factors for diarrheal syndromes in 3 sub-Saharan African countries after the introduction of rotavirus vaccines from the Vaccine Impact on Diarrhea in Africa (VIDA) study. Clinical Infectious Diseases. 2023;76: S12–S22.
365 366 367	12.	Demissie GD, Yeshaw Y, Aleminew W, Akalu Y. Diarrhea and associated factors among under five children in sub-Saharan Africa: evidence from demographic and health surveys of 34 sub-Saharan countries. PLoS One. 2021;16: e0257522.

368 369 370	13.	Hajara A, Faisal M. Rabi'U. Risk factors associated with prevalence of diarrhea among under 5 years children in low socioeconomic area of urban Bangladesh. Int Res J Nat Appl Sci. 2017;4: 12.
371 372	14.	Habtu M, Nsabimana J, Mureithi C. Factors contributing to diarrheal diseases among children less than five years in Nyarugenge District, Rwanda. J Trop Dis. 2017;5: 238.
373 374 375	15.	He Z, Ghose B, Cheng Z. Diarrhea as a Disease of Poverty Among Under-Five Children in Sub-Saharan Africa: A Cross-Sectional Study. INQUIRY: The Journal of Health Care Organization, Provision, and Financing. 2023;60: 00469580231202988.
376 377 378 379	16.	Agegnehu MD, Bewket Zeleke L, Goshu YA, Ortibo YL, Mehretie Adinew Y. Diarrhea Prevention Practice and Associated Factors among Caregivers of Under-Five Children in Enemay District, Northwest Ethiopia. J Environ Public Health. 2019;2019: 5490716.
380 381 382	17.	Dagnew AB, Tewabe T, Miskir Y, Eshetu T, Kefelegn W, Zerihun K, et al. Prevalence of diarrhea and associated factors among under-five children in Bahir Dar city, Northwest Ethiopia, 2016: a cross-sectional study. BMC Infect Dis. 2019;19: 1–7.
383 384 385	18.	Solomon ET, Gari SR, Kloos H, Alemu BM. Handwashing effect on diarrheal incidence in children under 5 years old in rural eastern Ethiopia: a cluster randomized controlled trial. Trop Med Health. 2021;49: 1–11.
386 387 388 389	19.	Ayalew AM, Mekonnen WT, Abaya SW, Mekonnen ZA. Assessment of Diarrhea and Its Associated Factors in Under-Five Children among Open Defecation and Open Defecation-Free Rural Settings of Dangla District, Northwest Ethiopia. J Environ Public Health. 2018;2018: 4271915.
390 391 392 393	20.	Islam M, Rahman M, Unicomb L, Kafi MAH, Rahman M, Alam M, et al. Child defecation and feces management practices in rural Bangladesh: Associations with fecal contamination, observed hand cleanliness and child diarrhea. PLoS One. 2020;15: e0236163.
394 395 396	21.	Hailu B, Ji-Guo W, Hailu T. Water, Sanitation, and Hygiene Risk Factors on the Prevalence of Diarrhea among Under-Five Children in the Rural Community of Dangila District, Northwest Ethiopia. J Trop Med. 2021;2021: 2688500.
397 398 399	22.	Mengistie B, Berhane Y, Worku A. Prevalence of diarrhea and associated risk factors among children under-five years of age in Eastern Ethiopia: A cross-sectional study. Open J Prev Med. 2013;3: 446.
400 401 402	23.	Dessalegn M, Kumie A, Tefera W. Predictors of under-five childhood diarrhea: Mecha District, west Gojam, Ethiopia. Ethiopian journal of health development. 2011;25: 192–200.
403 404 405 406	24.	Berde A, YALÇIN S, Ozcebe L, Üner S, Karadag-Caman Ö. Determinants of childhood diarrhea among under-five year old children in Nigeria: A population-based study using the 2013 demographic and health survey data. Turkish Journal of Pediatrics. 2018;60.

407 408 409	25.	Teshale AB, Liyew AM, Tesema GA. Factors associated with zinc utilization for the management of diarrhea in under-five children in Ethiopia. BMC Public Health. 2020;20: 1–7.
410 411 412	26.	Pienaah CKA, Yoshida Y, Saaka SA, Ategeeng FN, Luginaah I. Determinants of diarrhea prevalence among children under 5 years in semi-arid Ghana. Journal of Water, Sanitation and Hygiene for Development. 2024;14: 583–594.
413 414 415 416	27.	Dimitrova A, Gershunov A, Levy MC, Benmarhnia T. Uncovering social and environmental factors that increase the burden of climate-sensitive diarrheal infections on children. Proceedings of the National Academy of Sciences. 2023;120: e2119409120.
417 418 419	28.	Genser B, Strina A, Teles CA, Prado MS, Barreto ML. Risk factors for childhood diarrhea incidence: dynamic analysis of a longitudinal study. Epidemiology. 2006;17: 658–667.
420 421 422	29.	Kurui JE, Ogendi MG, Moturi NW, Nyawanga OD. Household water handling practices in the arid and semi-arid lands in Kenya. The Relevance of Hygiene to Health in Developing Countries. IntechOpen; 2019. pp. 79–89.
423 424 425	30.	Maldonado MEM. Evaluating Drivers and Sources of Pathogens to Surface Waters in Primarily Arid and Semi-Arid Tribal Lands of the United States. Arizona State University; 2023.
426 427	31.	Zelaya S. Climate change in the African drylands: options and opportunities for adaptation and mitigation. 2009.
428 429	32.	Ketiem P, Diarra A, Soura A, Konou R. Strategies for adapting to water stress in the arid and semi-arid regions of Africa. Synthesis brief. 2015.
430 431 432 433	33.	Lind J, Sabates-Wheeler R, Kohnstamm S. Changes in the drylands of Eastern Africa: A review of evidence and data and their implications for efforts to strengthening resilience. Institute of Development Studies Available at: https://opendocs.ids.ac uk/opendocs/handle/20500. 2016;12413: 12082.
434 435 436	34.	Hobbs N, Hug J, de Pee S. High Non-affordability of Diets and Malnutrition in Africa's Drylands: Systems Analysis to Guide Action. Food Nutr Bull. 2023;44: S45–S57.
437 438 439	35.	Johnson N, Wambile A. The impacts of the Arid Lands Resource Management Project (ALRMPII) on livelihoods and vulnerability in the arid and semi-arid lands of Kenya. ILRI (aka ILCA and ILRAD); 2011.
440 441 442	36.	Mokomane M, Kasvosve I, Melo E de, Pernica JM, Goldfarb DM. The global problem of childhood diarrhoeal diseases: emerging strategies in prevention and management. Ther Adv Infect Dis. 2018;5: 29–43.
443 444	37.	Amwata DA, Nyariki DM. Climate variability, land-use, pastoral and agropastoral livelihoods in arid and semi-arid areas of Kenya. 2021.

445 446	38.	Ngugi RK, Nyariki DM. Rural livelihoods in the arid and semi-arid environments of Kenya: Sustainable alternatives and challenges. Agric Human Values. 2005;22: 65–71.
447 448 449	39.	Moreira RP, Costa AC, Gomes TF, de Oliveira Ferreira G. Climate and climate- sensitive diseases in semi-arid regions: a systematic review. Int J Public Health. 2020;65: 1749–1761.
450 451	40.	Atake E-H. Health shocks in Sub-Saharan Africa: are the poor and uninsured households more vulnerable? Health Econ Rev. 2018;8: 1–13.
452 453	41.	Dinku Y, Fielding D, Genç M. Health shocks and child time allocation decisions by households: evidence from Ethiopia. IZA J Labor Econ. 2018;7: 1–23.
454	42.	USAID Nawiri fact sheet 2023.
455 456 457	43.	Kihu SM, Gachohi JM, Ndungu EK, Gitao GC, Bebora LC, John NM, et al. Sero- epidemiology of Peste des petits ruminants virus infection in Turkana County, Kenya. BMC Vet Res. 2015;11: 1–13.
458 459	44.	Akall G. Effects of development interventions on pastoral livelihoods in Turkana County, Kenya. Pastoralism. 2021;11: 23.
460 461 462	45.	Wilunda C, Sidze E, Thuita F, Amugsi D, Abajobir A, Mutua M, et al. Examining the complex dynamics influencing acute malnutrition in Turkana and Samburu counties: study protocol. Matern Child Nutr. 2024; e13744.
463 464 465 466	46.	Young SL, Boateng GO, Jamaluddine Z, Miller JD, Frongillo EA, Neilands TB, et al. The Household Water InSecurity Experiences (HWISE) Scale: development and validation of a household water insecurity measure for low-income and middle-income countries. BMJ Glob Health. 2019;4: e001750.
467 468	47.	Organization WH. World Health Organization releases new child growth standards. World Health Organization releases new child growth standards. 2006. p. 2.
469 470	48.	Victora CG, Huttly SR, Fuchs SC, Olinto MT. The role of conceptual frameworks in epidemiological analysis: a hierarchical approach. Int J Epidemiol. 1997;26: 224–227.
471 472	49.	Liang K-Y, Zeger SL. Longitudinal data analysis using generalized linear models. Biometrika. 1986;73: 13–22.
473 474	50.	Zeger SL, Liang K-Y, Albert PS. Models for longitudinal data: a generalized estimating equation approach. Biometrics. 1988; 1049–1060.
475 476	51.	Agulonye UVP. Fragilities and shocks effects on households and communities in West Africa. 2023.
477 478	52.	Pérez-Mesa D, Marrero GA, Darias-Curvo S. Child health inequality in Sub-Saharan Africa. Econ Hum Biol. 2022;47: 101176.
479 480	53.	Bonjean CA, Brunelin S, Simonet C. Impact of climate related shocks on child's health in Burkina Faso. 2012.
481 482	54.	Cornia GA, Rosignoli S, Tiberti L. The impact of the food and Financial Crises on child mortality: The case of sub-Saharan Africa. UN; 2011.

483 484 485	55.	Fratkin E, Roth EA, Nathan MA. Pastoral sedentarization and its effects on children's diet, health, and growth among Rendille of Northern Kenya. Hum Ecol. 2004;32: 531–559.
486 487	56.	Catley A, Admassu B, Bekele G, Abebe D. Livestock mortality in pastoralist herds in Ethiopia and implications for drought response. Disasters. 2014;38: 500–516.
488 489 490	57.	Pike IL, Straight B, Oesterle M, Hilton C, Lanyasunya A. Documenting the health consequences of endemic warfare in three pastoralist communities of northern Kenya: a conceptual framework. Soc Sci Med. 2010;70: 45–52.
491 492 493 494	58.	Abaasa CN, Rukundo GZ, Ayesiga S, Atukunda SP, Campisi S, O'Hearn S, et al. Healthcare providers and caregivers' perspectives on factors responsible for persistent malnutrition of under 5 children in Buhweju district, South Western Uganda; a phenomenological qualitative study. BMC Public Health. 2021;21: 1495.
495 496 497 498 499	59.	USAID. Examining the complex dynamics influencing acute malnutrition in Turkana county: a longitudinal mixed-methods study to support community-driven activity design. Kenya; 2021. Available: https://www.advancingnutrition.org/sites/default/files/2023-04/Mercy Corps_USAID Nawiri_Turkana LS Qualitative Report.pdf
500 501 502	60.	Nounkeu C, Kamgno J, Dharod J. Assessment of the relationship between water insecurity, hygiene practices, and incidence of diarrhea among children from rural households of the Menoua Division, West Cameroon. J Public Health Afr. 2019;10.
503 504 505	61.	Aydamo AA, Gari SR, Mereta ST. The nexus between household water insecurity, mother's handwashing practices, and diarrheal diseases among under-five children. J Water Health. 2024; jwh2024026.
506 507 508	62.	Akinyemi PA, Afolabi OT, Aluko OO. The effects of seasonal variations on household water security and burden of diarrheal diseases among under 5 children in an urban community, Southwest Nigeria. BMC Public Health. 2022;22: 1354.
509 510 511 512	63.	Wolf J, Hubbard S, Brauer M, Ambelu A, Arnold BF, Bain R, et al. Effectiveness of interventions to improve drinking water, sanitation, and handwashing with soap on risk of diarrhoeal disease in children in low-income and middle-income settings: a systematic review and meta-analysis. The Lancet. 2022;400: 48–59.
513 514 515	64.	Liu Q, Liu M, Liu J. Association of drinking water services with the disease burden of diarrhea in children under five in 200 countries from 2000 to 2021. Cell Reports Sustainability. 2024.
516 517	65.	Schaible UE, Kaufmann SHE. Malnutrition and infection: complex mechanisms and global impacts. PLoS Med. 2007;4: e115.
518 519 520	66.	Poda GG, Hsu C-Y, Chao JC-J. Factors associated with malnutrition among children< 5 years old in Burkina Faso: evidence from the Demographic and Health Surveys IV 2010. International Journal for Quality in Health Care. 2017;29: 901–908.

521 522 523	67.	Ogayo M, Damaris O, Sadia M, Odipo JO, Ayodo G. Risk of Bidirectional Diarrhea- Malnutrition among Under-Five-Year-Old Children Admitted in Referral Hospitals in Western Kenya. Afr J Health Sci. 2024;36: 578–587.
524 525 526	68.	Roy SK, Buis M, Weersma R, Khatun W, Chowdhury S, Begum A, et al. Risk factors of mortality in severely-malnourished children hospitalized with diarrhoea. J Health Popul Nutr. 2011;29: 229.
527 528	69.	Shah SM, Yousafzai M, Lakhani NB, Chotani RA, Nowshad G. Prevalence and correlates of diarrhea. The Indian Journal of Pediatrics. 2003;70: 207–211.
529 530 531	70.	Tickell KD, Pavlinac PB, John-Stewart GC, Denno DM, Richardson BA, Naulikha JM, et al. Impact of childhood nutritional status on pathogen prevalence and severity of acute diarrhea. Am J Trop Med Hyg. 2017;97: 1337.
532 533 534	71.	Johnston BC, Shamseer L, da Costa BR, Tsuyuki RT, Vohra S. Measurement issues in trials of pediatric acute diarrheal diseases: a systematic review. Pediatrics. 2010;126: e222–e231.

535

536

Supplementary Tables 537

- 538 S1 Table. Bivariate logistic regression analysis of factors associated with diarrhea among
- 539 children under five years of age



Figure