1	The lived experience of rejection sensitivity in ADHD - a qualitative exploration
2	
3	Annabel Rowney-Smith ¹ , Beth Sutton ^{2,3} , Lisa Quadt ^{1,3} , Jessica A Eccles ^{1,3,*}
4	
5	1 Department of Clinical Neuroscience, Brighton & Sussex Medical School,
6	University of Sussex, Brighton, UK
7	2 School of Global Studies, University of Sussex, Brighton, UK
8	3 Sussex Partnership NHS Foundation Trust, Worthing, UK
9	
10	Short title: Rejection sensitivity in ADHD
11	
12	* Corresponding Author
13	Dr Jessica Eccles
14	Trafford Centre
15	University of Sussex
16	BN1 9RY
17	Brighton
18	United Kingdom
19	J.Eccles@bsms.ac.uk
20	

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

21

Abstract

22 Attention Deficit Hyperactivity Disorder (ADHD) is a neurodevelopmental condition. 23 commonly associated with differences in attention, and/or hyperactivity and 24 impulsiveness. A lesser known, but highly impactful characteristic of ADHD is 25 emotional dysregulation, which causes difficulties in emotional expression and 26 identification. An aspect of emotional dysregulation which remains relatively 27 unexplored in ADHD, is rejection sensitivity. In people who experience rejection 28 sensitivity, perceived rejection and/or criticism can evoke extreme dysphoria. To 29 investigate the lived experience of rejection sensitivity in ADHD individuals, five 30 undergraduates participated in focus-group interviews. The subsequent thematic 31 analysis revealed three key themes which encapsulated the experience; withdrawal, 32 masking, and bodily sensations. The participants explained how rejection sensitivity 33 elicited unpleasant bodily sensations, anxiety, and misery, and how, in turn, they used masking to camouflage these feelings. They then discussed how the use of 34 35 masking had caused them to become dissociated from themselves and to withdraw 36 from others, which often resulted in loneliness. It was apparent that rejection 37 sensitivity significantly impacts mental wellbeing, eliciting feelings such as 38 anxiousness, despair, and embarrassment. In turn, social function is significantly 39 impaired, alongside career opportunities, and daily life. Our findings indicate a need 40 for deeper understanding of ADHD traits and emotional dysregulation, which may in 41 turn lessen the presence of rejection sensitivity in ADHD individuals.

42

43 Keywords: rejection sensitivity, ADHD, emotion

44

Introduction

45 ADHD is a neurodevelopmental condition, characterised by differences in attention 46 and emotion regulation, possible hyperactivity, and impulsiveness (1). The 47 neurodiversity movement (2) advocates that neurodevelopmental conditions are 48 often inseparable from one's identity (3). Thus, to align our research with the 49 neurodiversity movement's values, the language in this paper is henceforth adapted 50 to avoid the perpetuation of ableist stances. For example, medicalised language will 51 be avoided, e.g., 'traits' instead of 'symptoms', and identity-first language will be 52 used instead of person-first language (e.g., 'ADHD person', or 'ADHDer', instead of 53 'person with ADHD').

54 Emotional dysregulation is overrepresented in ADHD and may be prevalent in up to 55 70% of ADHD adults (4). emotional dysregulation is multi-dimensional, but often 56 presents as increased and/or reduced emotional reactivity, lack of emotional 57 awareness, and intense emotional expression (5). Consequently, ADHDers may 58 experience an array of emotional dysregulation-related symptoms, including intense 59 mood swings, and reactivity to seemingly benign situations. Thus, emotional 60 dysregulation is thought to significantly contribute to psycho-social problems more 61 often faced by ADHD people, such as unemployment and interpersonal issues (6) 62 and has been found to be more negatively influential on guality of life than inattentive 63 and hyperactivity traits (7). Despite its common association with ADHD, and its 64 significant effects, emotional dysregulation remains an often-neglected trait, which is evidenced in its exclusion from diagnostic criteria (8). In recent years, anecdotal 65 66 reports have highlighted a crucial dimension of emotional dysregulation experienced 67 by ADHD people; rejection sensitivity (9).

Rejection sensitivity is defined as dysphoria induced by perceived or real rejection,
and/or criticism (9). People who experience rejection sensitivity readily detect

70 rejection, and therefore perceive rejection/criticism more frequently than others (10). 71 The ensuing dysphoric episode often involves intense feelings of extreme misery. 72 anxiety, and pain (11). The source of rejection sensitivity is disputed, however; 73 evidence suggests that childhood interpersonal trauma, such as caregiver rejection. 74 may contribute to causality (12). Although rejection sensitivity is defined as a 75 personality disposition (13), it frequently features in mental health disorders typically 76 characterised by emotional dysregulation, such as depression, Body Dysmorphic 77 Disorder, and Borderline Personality Disorder (14). 78 To date, few studies have investigated the potential association between ADHD and 79 rejection sensitivity. Moreover, these studies, for the most part, have attempted to 80 establish the *extent* of the rejection sensitivity -ADHD association, rather than 81 explore the *nature* of the association. However, although they did not directly 82 measure rejection sensitivity, one study (15) explored the experience of received 83 criticism in an ADHD population, via qualitative techniques (open-ended written 84 responses).

85 Since emotional dysregulation presents as a significant influence on guality of life for 86 ADHDers, and rejection sensitivity may be a contributing factor in this, there is an 87 obvious need for more research exploring the experience of rejection sensitivity in 88 ADHD cohorts. Motivated by this gap in existing research, our study aimed to 89 provide rich accounts of rejection sensitivity in an ADHD population, via qualitative 90 methods: focus-group interviews and subsequent thematic analysis. We hope that 91 this research will improve the understanding of neurodivergent experiences and 92 shed light on the co-occurrence between rejection sensitivity and ADHD.

93

94

<u>Methods</u>

95 Measures

To explore the lived experience of rejection sensitivity, a qualitative study design was 96 97 utilised; focus groups interviews. This approach was selected with the aim of capturing a rich, and illustrative collection of perspectives, as participants can 98 99 explore and discuss their experiences alongside like-minded peers, and with limited 100 restriction. At the start of the focus groups, informed verbal consent was gained from 101 the participants, and time was allocated for introductions. The interviews were lightly 102 structured by a topic guide which comprised one broad, open-ended guestion with 103 associated sub-guestions. The broad, over-arching guestion was aimed to frame the 104 ensuing discourse; "If you do, how do you identify with Rejection Sensitive 105 Dysphoria?"; whereas the associated sub-questions were intended to encourage 106 participants' engagement with the topic, and included questions such as, "Has 107 rejection sensitivity D interfered with your daily life?", and "Do you experience any 108 particular feelings when you are experiencing rejection sensitivity D?". To increase 109 accessibility, the focus groups were conducted via online video-call, through Zoom.

110 **Participants**

Five undergraduate students were recruited from the University of Sussex, via flyers distributed across the university campus. To participate in the study, participants had to satisfy inclusion/exclusion criteria; all participants were undergraduates with formal ADHD diagnoses and were over 18 years old. The participants were also incentivised to take part, via a £16 GiftPay voucher. Two focus group interviews took place, whereby one group contained two participants, and one group contained three. Participants were sent a copy of the consent form and an information sheet

prior to providing verbal consent, to allow adequate understanding of the studyexpectations and the research process.

120 Analysis

121 The focus group content was audio-recorded via a Dictaphone and transcribed 122 verbatim into written text. Following this, the transcribed data was analysed and 123 interpreted, informed by Braun and Clarke's (16) six-step thematic analysis 124 framework (16). To promote familiarity with the data, the focus-group transcriptions 125 were firstly read multiple times. Next, seemingly important data were identified, and 126 then organised into themes according to patterns of similarity. The potential themes 127 were then discussed and reviewed amongst the research team, whereby three 128 defined themes emerged; masking, withdrawal, and bodily sensations. Two themes 129 (problems with self-esteem and cognitions) were rejected because they overlapped 130 with the other themes, and were determined as subthemes of our three defined 131 themes.

132 Our study method was devised by University of Sussex researchers and

133 undergraduate students alike, with neurodivergent perspectives.

134 Ethical approval for this study was obtained by the Research Governance and Ethics

135 Committee at Brighton and Sussex Medical School (application no. ER/AR649/1).

136 Data was collected between 25th July 2022 and 2nd September 2022.

137

<u>Results</u>

- 138 Participants spoke in depth about their feelings, thought processes, and physical
- 139 experiences before, during, and after rejection/criticism. In commonality, they
- 140 described rejection sensitivity as a painful and overwhelming experience, which often

- 141 lasts a significant amount of time (between hours and weeks) and can resurface
- 142 many years later. Three primary themes emerged from the data; *masking*,
- 143 *withdrawal*, and *bodily sensations* (Fig. 1).

144 [Figure 1 here]

- 145 Figure 1: Three primary themes
- 146

147 **Withdrawal**

- 148 One of the most common themes identified was 'withdrawal'. The participants
- 149 explained that rejection sensitivity often caused them to withdraw from friendships,
- 150 family and romantic relationships, university life, and job opportunities. All
- 151 participants agreed that the *expectation* of rejection elicited more dysphoria than the
- rejection *itself*; "there's a general idea that I will be rejected". Thus, participants
- reported to pre-emptively withdraw to avoid potential rejection; "I just get myself out
- 154 of the situation where the rejection might come, just to avoid it".
- 155 Participants discussed how withdrawal negatively impacts their relationships with
- 156 friends and family. It was disclosed that they "rarely reach out" to people, and that
- 157 when they *do* contact people, they consistently assess for possible
- rejection/criticism. For instance, they may assume that an unanswered text message
- 159 is a rejection; "I just ended up being awake until four in the morning... like constantly
- 160 checking my phone, being like, why was this person not responding?". Therefore,
- 161 participants explained that it is easier to avoid contacting people, as this pre-
- 162 emptively lessen rejection sensitivity anxiety. In turn, their quantity and quality of
- 163 relationships is negatively impacted. One participant expressed that there were
- 164 "limited people in my life that I'm close with". Similarly, another participant shared
- 165 that they keep their "friendship circles low and full of neurodivergent people" as they

had suffered "so much rejection from allistic people". Consequently, participantsdisclosed how they often felt "lonely", "friendless", and "unlovable".

168 Withdrawal also affects participants' university work; most participants agreed that 169 the fear of rejection/criticism had caused them to delay, or not submit assignments. 170 A few participants admitted that in order to protect themselves from criticism, they 171 had purposefully submitted university work which was below their usual standard, or 172 had submitted it late (which typically incurs a 10% grade deduction). They explained 173 that this allowed them an explanation for potential rejection/criticism in the form of a 174 disappointing grade. Focus group members also discussed withdrawal from the 175 workplace. One participant had avoided applying for career opportunities "to avoid 176 rejection". Interestingly, participants explained how they often experienced rejection 177 sensitivity during the application process, as opposed to when they received the 178 rejection. This conforms with the earlier assertion, that the expectation of rejection "is 179 more damaging than the actual rejection". Participants discussed that they do not 180 feel they deserve good university grades or successful job applications, with one 181 participant employing the term; "imposter syndrome".

182 Masking

Masking is often utilised by neurodivergent people to suppress and camouflage neurodivergent traits (17). This is achieved via the employment of multiple tools (both consciously and sub-consciously), and subsequent alterations in behaviour (17). Most participants, either explicitly or implicitly, used masking to camouflage their thoughts and feelings about a perceived rejection/critique. The participants discussed their individual difficulties with distinguishing between "real" and their perceived rejection. A participant provided an example of this; they explained that

190 they could not differentiate between jokes and critiques. Thus, participants explained 191 that they often felt the need to ask friends/family for clarification and/or reassurance. 192 However, this often led to embarrassment, anxiety, invalidation, and labels such as 193 "oversensitive". To avoid this, participants mask to convey robustness, and a 194 nonchalant attitude to rejection; "a mask of toughness". This enables them to hide 195 their rejection sensitivity from others and helps them feel less like an outsider, and 196 while it helps them temporarily feel like an outsider, participants agreed that masking 197 has negative consequences.

198 Participants revealed that consistent masking causes disconnection from their 199 emotions and their authentic selves, as they struggle to distinguish between real and 200 faux feelings; "I think at some point it's not just masking towards others, but you are 201 also masking with yourself... you can't identify if you're bothered or not, because you 202 lose connection". Participants also described how masking caused them to feel 203 disconnected from social interaction; "observing this interaction that's happening to 204 you". One participant disclosed how they had benefitted from therapy, because it 205 enabled them to identify and reconnect with their emotions. The participants also 206 explained that masking perpetuates their social withdrawal. Two participants 207 described this as a "vicious circle", whereby others take advantage of their 208 nonchalant mask, and often incorrectly assume that rejection and criticism does not 209 affect them. Consequently, rejections and critiques increase, and they subsequently 210 withdraw further from social situations, and further reinforce the mask; a vicious 211 circle.

212 **Bodily Sensations**

213 Participants described how perceived rejection evokes overwhelming and 214 unpleasant bodily sensations. Interestingly, sensations were varied and unique to the individual. For example, one participant felt sensations in their heart and throat; "it's 215 216 like a pinch in your heart, like a very swift, brief, like, like your heart or like your throat 217 closing up... feels like your heart kind of drops maybe a bit". Another participant 218 explained how they felt as if "a chair (had) been taken out from beneath you". One 219 participant used a powerful metaphor to explain the flooding of heat throughout their 220 body; "you know the hand warmers, when you click, it becomes solid and warm... 221 getting warmer inside me.... everything inside me starts to burn". In contrast, a 222 participant discussed how they felt their body enter a "paralysed state". Stomach 223 sensations were the only sensation common amongst participants; "sick to your stomach", and nausea "in my gut". 224 225 Figure 1 here 226 Figure 1. Graphic depicting the key themes in the analysis of rejection sensitivity in 227 ADHD 228 229 Discussion

Our findings indicate that rejection sensitivity is a multidimensional experience;
encompassing emotional pain, negative cognitions, and unpleasant physical
sensations. Our participants explained how the effects of rejection sensitivity, like
anxiety, depression, and unpleasant bodily sensations, had led them to mask their
true feelings, and withdraw from relationships and work/university commitments. It

was apparent that rejection sensitivity is detrimental to mental health, and impairsboth daily function and long-term wellbeing.

237 The findings also indicate that rejection sensitivity could be a contributing factor in 238 the psychosocial problems often faced by ADHDers. For example, the participants 239 explained how they are less likely to apply for jobs or submit university assignments 240 because they were expecting rejection and were protecting themselves from the 241 ensuing misery of rejection sensitivity. This concern could potentially limit their job 242 prospects, and any future career aspirations. Furthermore, subsequent withdrawal 243 from relationships may restrict opportunities to find a romantic partner and friends. 244 and in extreme cases, could lead to familial estrangement. Again, this could 245 contribute to psychosocial problems, as ADHDers may find themselves without a 246 support network, and may find it hard to start families of their own. Thus, the effects 247 of rejection sensitivity are far-reaching, and could prove long-term. Previous 248 research has primarily focussed on how ADHD traits affect psychosocial functioning. 249 yet our findings suggest that rejection sensitivity may be just as important in the 250 development of psychosocial issues.

251 The participants spoke at length about how their withdrawal from friends and family 252 had left them lonely and isolated. Research has found that ADHDers are more likely 253 to experience loneliness than those without ADHD (18), and rejection sensitivity -254 masking may contribute to this. Moreover, research also indicates that masking 255 threatens the development and maintenance of deep relationships. Bradley et al. 256 found that masking made their autistic participants feel disingenuous because their 257 relationships were based on an untrue representation of themselves (19). Their 258 participants felt that their relationships with others were based on deception, and that 259 very few people truly knew who they truly were. This is echoed in our study, when

260 our participants expressed how they use masking to present themselves as tougher 261 than they are, which, in turn, made others feel as if they could continue to exploit 262 them, not knowing the devastating consequences for the participants' mental health. 263 This may further contribute to the participants' withdrawal from relationships with 264 others, as they may find their relationships unfulfilling, shallow, and based on 265 deception. A study conducted by Houghton et al. (20) corroborates this, as they 266 found that ADHDers often report their friendships as low quality (20). They also 267 found that loneliness mediates the often-co-occurrent depressive disorders in ADHD 268 (20). rejection sensitivity could form part of the driving force behind the high 269 prevalence of loneliness and depression in ADHD people, as it may cause ADHDers 270 to have superficial friendships, and withdraw from developing and maintaining 271 relationships.

272 Our participants disclosed how their extensive use of masking had caused them to 273 feel dissociated from themselves: one participant explained how they needed 274 therapy to re-identify their true self. Dissociation is often allied with mental health 275 disorders like bipolar disorder (21) and borderline personality disorder (BPD) (22), 276 which are both common ADHD-co-occurrent conditions (23, 24). It may be that 277 rejection sensitivity -masking contributes to the development of bipolar disorder 278 and/or BPD in ADHDers through its dissociative consequences. For instance, our 279 participants discussed how they felt that masking had led them astray from their true 280 identities, leading them on to question who they really are. This unstable self-image 281 and de-personalisation is also frequently seen in people with bipolar disorder and 282 BPD. Furthermore, since BPD sufferers are more likely to experience unstable 283 relationships (25), rejection sensitivity -related withdrawal from relationships may 284 also contribute to BPD symptoms. Unfortunately, due to symptoms overlap (primarily

emotional dysregulation-related), ADHD is often misdiagnosed as BPD, especially in
females (26). It may be that either rejection sensitivity -masking and withdrawal
contributes to the *development* of BPD in ADHDers, or they may also be distinct
features of ADHD, which would contribute to the *misdiagnosis* of BPD.

289 To our knowledge, the unpleasant bodily sensations felt by the participants was a 290 new finding in the context of rejection sensitivity. We found that these sensations 291 varied greatly amongst the participants; each participant experienced different 292 physical sensations in different bodily areas. For example, one participant primarily 293 felt nausea in their stomach and gut, whereas another felt that their heart and throat 294 were tightening. Additionally, we found that rejection sensitivity affects the perception 295 of bodily temperature, when one participant explained that rejection sensitivity 296 evoked the feeling of progressively intense heat throughout their body. Conversely, 297 rejection sensitivity also appeared to *inhibit* the feeling of physical sensation, when 298 one participant explained that rejection sensitivity causes them to feel paralysed. 299 These types of bodily sensations are often found in anxiety disorders, and may 300 reflect an anxiety attack. Anxiety attacks are caused by activation of the sympathetic 301 nervous system, which effectively causes the body to enter a fight-or-flight 302 physiological state (27 p. 189–190). This affects multiple bodily systems, often 303 resulting in various symptoms, such as nausea, feelings of warmth or cold, chest 304 tightness, and abdominal pain (27 p. 189–190). It may be that rejection sensitivity 305 elicits a stress response in ADHDers, which may contribute to the often co-occurrent 306 diagnosis of anxiety disorders.

Perhaps the most insightful and important notion to arise from our participants'
honesty, is the idea that rejection sensitivity could be prevented or alleviated if they
received more understanding and empathy from others. The participants expressed

310 that they consistently felt they needed reassurance from others to be able to help 311 alleviate rejection sensitivity symptoms. They explained that if they did not receive 312 the reassurance they needed, their rejection sensitivity -related anxiety would 313 "spiral", because they would repeatedly analyse the rejection sensitivity -provoking 314 situation; trying to work out if they had been rejected or not. However, they often felt 315 embarrassed to ask for reassurance, because they did not want to present as 316 annoying or as overly sensitive. This was mainly due to previous negative 317 experiences, whereby asking for reassurance was followed by judgement from 318 others. Thus, gaining reassurance was a *double-edged sword*; seeking reassurance 319 induced feelings of embarrassment and judgement, and alternatively, not seeking 320 reassurance induced anxiety. This may also contribute to withdrawal from 321 relationships, as participants may wish to avoid the consequences of either decision. 322 This finding indicates that more understanding and empathy from others may 323 mediate the severity of rejection sensitivity, whereby ADHDers would not need to 324 mask as much, and would potentially be able to socialise more without such negative 325 consequences for their mental health.

326 Although our findings provide a rich insight into the experience of rejection 327 sensitivity, they do not elucidate the aetiology of rejection sensitivity. However, it 328 may be that rejection sensitivity develops as a result of sensitisation to sustained 329 criticism. Due to lack of knowledge and understanding of trait aetiology, ADHD is 330 classically stigmatised (28). ADHD traits may contravene social norms and can 331 appear to others as if they are controllable. For example, neurotypical people are 332 typically able to control their impulses, such as not interrupting others. This, paired 333 with misconceptions regarding aetiology, lead to discreditation and prejudice (28). 334 Thus, ADHD traits are often met with criticism. Since ADHD is present from

335 childhood, ADHD people are more frequently exposed to criticism and rejection from 336 a young age than neurotypical peers (29). It could be hypothesised that rejection 337 sensitivity results from sensitisation; whereby the response to upsetting, frequent, 338 and early experiences of criticism and rejection, become larger and more intense 339 over time. This fits with the data collected from our focus-groups; the participants 340 often felt that the anticipation of rejection was more painful than the rejection itself, 341 indicating that past experience of rejection frames future response. This correlates 342 with the increasing emotional dysregulation seen from childhood ADHD into 343 adulthood ADHD (4). This sensitisation theory is corroborated by Babinski et al. (30), 344 who investigated social feedback circuitry in adolescents with both ADHD and 345 rejection sensitivity. The authors found that rejection sensitivity symptoms correlated 346 with sensitisation of attentional neural circuitry for social rejection, and with 347 desensitisation for positive social feedback (30). Therefore, ADHD people who 348 experience rejection sensitivity, may be oversensitive to rejection cues, and 349 *desensitised* to social approval cues, due to alterations in neural connections. 350 Consequently, rejection is perceived more frequently, and is unassuaged by *missed* 351 social approval cues. Since ADHD is characterised by differences in attention, it may 352 be that ADHD neuronal networks are more susceptible to this attentional imbalance 353 between rejection and approval cues. Therefore, a combination of frequent and early 354 rejection/criticism, paired with attentional wiring differences, may make ADHDers 355 more vulnerable to rejection sensitivity.

Our findings aligned with previous literature, which found significant associations
 between rejection sensitivity and ADHD, with increased aggression and social

anxiety (31, 32), higher levels of victim justice sensitivity and lower levels of

359 perpetrator sensitivity (33), also withdrawal from social situations and relationships

360 (32, 34). Although a previous study investigated the experience of criticism in people 361 with ADHD traits (29), our study was, to the best of our knowledge, the first to 362 directly explore the experience of *rejection sensitivity* in ADHD individuals. 363 Although our method was successful for permitting rich and illustrative descriptions 364 of the experience of rejection sensitivity, the sample size was small, and we did not 365 screen for other neurodivergent conditions. During the interviews, two participants 366 disclosed co-occurring Autism Spectrum Condition diagnoses, whilst another 367 disclosed a Dyslexia diagnosis. Since ADHD is frequently accompanied by other 368 conditions, participants may have had other diagnoses we were unaware of. Future 369 research may be disposed to use a larger sample, and to screen for other conditions, 370 in order to improve generalisability and validity. This may also elucidate the 371 possibility that ADHD mediates associations between rejection sensitivity and other 372 conditions which are also co-occurrent with ADHD, such as BPD. We hope that our 373 findings improve understanding of neurodivergent experiences and highlight the 374 need for further research into potential underlying mechanisms.

376 Acknowledgments

We thank all of our participants for their time and openness.

378 Ethical approval statement

- 379 All elements of this study were approved by the Brighton & Sussex Medical School
- 380 Research Governance & Ethics Committee (approval no. ER/AR649/1) on July 10th,
- 381 2022. The participants provided informed verbal consent before taking part.

382 **Declaration of conflicting interests**

383 The authors declare no conflicts of interest.

384 Funding

- 385 This study was funded partially by University of Sussex for a junior research
- 386 associate summer programme.

387 Data availability statement

- 388 All relevant data are within the manuscript. Data cannot be shared publicly because
- 389 we have no permission from the respondents of our study to share the de-identified
- 390 dataset with the general public (Ethics committee: Research Governance and Ethics
- 391 Committee at Brighton and Sussex Medical School (application no. ER/AR649/1).
- 392 Data requests can be directed to the corresponding author.

393 Figure Legends

- Figure 1. Graphic depicting the key themes in the analysis of rejection sensitivity inADHD
- 396

perpetaty.	•
It is made available under a CC-BY	4.0 International license .

397		References
398		
399	1.	Retz W, Stieglitz RD, Corbisiero S, Retz-Junginger P, Rösler M. Emotional
400		dysregulation in adult ADHD: what is the empirical evidence? Expert Review
401		of Neurotherapeutics. 2012 Oct;12(10):1241–51.
402		
403	2.	Kapp SK, Gillespie-Lynch K, Sherman LE, Hutman T. Deficit, difference, or
404		both? Autism and neurodiversity. Developmental Psychology. 2013;49(1):59-
405		71.
406		
407	3.	Bottema-Beutel K, Kapp SK, Lester JN, Sasson NJ, Hand BN. Avoiding
408		Ableist Language: Suggestions for Autism Researchers. Autism in Adulthood
409		[Internet]. 2020 Sep 2;3(1):18–29.
410		https://www.liebertpub.com/doi/10.1089/aut.2020.0014
411		
412	4.	Shaw P, Stringaris A, Nigg J, Leibenluft E. Emotion Dysregulation in Attention
413		Deficit Hyperactivity Disorder. American Journal of Psychiatry [Internet]. 2014
414		Mar;171(3):276–93. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4282137/
415		
416	5.	D'Agostino A, Covanti S, Rossi Monti M, Starcevic V. Reconsidering Emotion
417		Dysregulation. Psychiatric Quarterly. 2017 Feb 14;88(4):807–25.
418		

419	6.	Hirsch O, Chavanon M, Riechmann E, Christiansen H. Emotional
420		dysregulation is a primary symptom in adult Attention-Deficit/Hyperactivity
421		Disorder (ADHD). Journal of Affective Disorders. 2018 May;232:41–7.
422		
423	7.	Riley AW, Spiel G, Coghill D, Döpfner M, Falissard B, Lorenzo MJ, et al.
424		Factors related to Health-Related Quality of Life (HRQoL) among children with
425		ADHD in Europe at entry into treatment. European Child & Adolescent
426		Psychiatry. 2006 Dec;15(S1):i38–45.
427		
428	8.	Faraone SV, Rostain AL, Blader J, Busch B, Childress AC, Connor DF, et al.
429		Practitioner Review: Emotional dysregulation in attention-deficit/hyperactivity
430		disorder - implications for clinical recognition and intervention. Journal of Child
431		Psychology and Psychiatry. 2018 Apr 6;60(2):133–50.
432		
433	9.	Dodson W, M.D., LF-APA. New Insights Into Rejection Sensitive Dysphoria
434		[Internet]. ADDitude. 2020. https://www.additudemag.com/rejection-sensitive-
435		dysphoria-adhd-emotional-dysregulation/
436		
437	10	Romero-Canyas R, Downey G, Berenson K, Ayduk O, Kang NJ. Rejection
438		Sensitivity and the Rejection-Hostility Link in Romantic Relationships. Journal
439		of Personality. 2010 Feb;78(1):119–48.

441	11. Dodson W. RSD: How Rejection Sensitive Dysphoria Feels for Adults with
442	ADHD [Internet]. www.additudemag.com. 2023 [cited 2023 Sep 19].
443	https://www.additudemag.com/rsd-rejection-sensitive-dysphoria-experiences/
444	
445	12. Feldman S, Downey G. Rejection sensitivity as a mediator of the impact of
446	childhood exposure to family violence on adult attachment behavior.
447	Development and Psychopathology. 1994;6(1):231–47.
448	
449	13. Downey G, Feldman SI. Implications of rejection sensitivity for intimate
450	relationships. Journal of Personality and Social Psychology [Internet]. 1996
451	Jun 1;70(6):1327–43.
452	https://pubmed.ncbi.nlm.nih.gov/8667172/#:~:text=Study%203%20shows%20
453	that%20people
454	
455	14. Gao S, Assink M, Cipriani A, Lin K. Associations between rejection sensitivity
456	and mental health outcomes: A meta-analytic review. Clinical Psychology
457	Review. 2017 Nov;57:59–74.
458	
459	15. Beaton DM, Sirois F, Milne E. Experiences of criticism in adults with ADHD: A
460	qualitative study. Fernández-Alcántara M, editor. PLOS ONE [Internet]. 2022
461	Feb 18;17(2):e0263366.
462	https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0263366
463	

464	16. Braun V, Clarke V. Using thematic analysis in psychology. [Internet].
465	psycnet.apa.org. 2006. https://psycnet.apa.org/record/2006-06991-002
466	
467	17. Pearson A, Rose K. A Conceptual Analysis of Autistic Masking:
468	Understanding the Narrative of Stigma and the Illusion of Choice. Autism in
469	Adulthood [Internet]. 2021 Jan 22;3(1).
470	https://www.liebertpub.com/doi/10.1089/aut.2020.0043
471	
472	18. Stickley A, Koyanagi A, Takahashi H, Ruchkin V, Kamio Y. Attention-
473	deficit/hyperactivity disorder symptoms and loneliness among adults in the
474	general population. Research in Developmental Disabilities. 2017
475	Mar;62:115–23.
476	
477	19. Bradley L, Shaw R, Baron-Cohen S, Cassidy S. Autistic Adults' Experiences
478	of Camouflaging and Its Perceived Impact on Mental Health. Autism in
479	Adulthood [Internet]. 2021 Jun 3;3(4).
480	https://www.liebertpub.com/doi/full/10.1089/aut.2020.0071
481	
482	20. Houghton S, Lawrence D, Hunter SC, Zadow C, Kyron M, Paterson R, et al.
483	Loneliness Accounts for the Association Between Diagnosed Attention Deficit-
484	Hyperactivity Disorder and Symptoms of Depression Among Adolescents.
485	Journal of Psychopathology and Behavioral Assessment. 2020 Jan
486	31;42(2):237–47.

488	21. Hariri AG, Gulec MY, Orengul FFC, Sumbul EA, Elbay RY, Gulec H.
489	Dissociation in bipolar disorder: Relationships between clinical variables and
490	childhood trauma. Journal of Affective Disorders. 2015 Sep;184:104–10.
491	
492	22. Krause-Utz A. Dissociation, trauma, and borderline personality disorder.
493	Borderline Personality Disorder and Emotion Dysregulation. 2022 Apr 19;9(1).
494	
495	23. Matthies SD, Philipsen A. Common ground in Attention Deficit Hyperactivity
496	Disorder (ADHD) and Borderline Personality Disorder (BPD)-review of recent
497	findings. Borderline Personality Disorder and Emotion Dysregulation.
498	2014;1(1):3.
499	
500	24. Schiweck C, Arteaga-Henriquez G, Aichholzer M, Thanarajah SE, Vargas-
501	Cáceres S, Matura S, et al. Comorbidity of ADHD and adult bipolar disorder:
502	A systematic review and meta-analysis. Neuroscience & Biobehavioral
503	Reviews. 2021 Jan;124.
504	
505	25. Brüne M. Borderline Personality Disorder. Evolution, Medicine, and Public
506	Health. 2016;2016(1):52–66.
507	
500	
508	26. Littman E. ADHD and BPD: Borderline Personality Disorder's Link to ADD

510	https://www.additudemag.com/adhd-and-bpd-women-borderline-personality-
511	disorder/#:~:text=Because%20ADHD%20was%20rarely%20diagnosed
512	
513	27. Robertson D. Primer on the autonomic nervous system. 2nd ed. Amsterdam ;
514	Boston: Elsevier Academic Press; 2004.
515	
516	28. Mueller AK, Fuermaier ABM, Koerts J, Tucha L. Stigma in attention deficit
517	hyperactivity disorder. ADHD Attention Deficit and Hyperactivity Disorders
518	[Internet]. 2012 Jul 8;4(3):101–14.
519	https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3430836/
520	
521	29. Beaton DM, Sirois F, Milne E. Self-compassion and Perceived Criticism in
522	Adults with Attention Deficit Hyperactivity Disorder (ADHD). Mindfulness.
523	2020 Aug 19;
524	
525	30. Babinski DE, Kujawa A, Kessel EM, Arfer KB, Klein DN. Sensitivity to Peer
526	Feedback in Young Adolescents with Symptoms of ADHD: Examination of
527	Neurophysiological and Self-Report Measures. Journal of Abnormal Child
528	Psychology. 2018 Aug 29;47(4):605–17.
529	
530	31.KOYUNCU A, ÇELEBI F, ERTEKIN E, MEMIŞ ÇÖ, TÜKEL R. The Presence
531	of Childhood Attention Deficit/Hyperactivity Disorder May Be Associated With

532	Interpersonal Sensitivity in Patients With Social Anxiety Disorder. Journal of
533	Psychiatric Practice. 2017 Jul;23(4):254–9.
534	
535	32. Motamedi M, Bierman K, Huang-Pollock CL. Rejection Reactivity, Executive
536	Function Skills, and Social Adjustment Problems of Inattentive and
537	Hyperactive Kindergarteners. Social Development. 2015 Jul 14;25(2):322–39.
538	
539	33. Bondü R, Esser G. Justice and rejection sensitivity in children and
540	adolescents with ADHD symptoms. European Child & Adolescent Psychiatry
541	[Internet]. 2014 May 31;24(2):185–98.
542	https://link.springer.com/article/10.1007/s00787-014-0560-9
543	
544	34. Narimani M, Aghajani S, Mohsenpour Z. Comparing group participation and
545	rejection sensitivity in the students with or without learning disorders. Journal
546	of Learning Disabilities [Internet]. 2020 Nov 21 [cited 2023 Oct 4];10(1):187-
547	99. https://jld.uma.ac.ir/article_1051_en.html?lang=en

Masking

Masking: when someone acts in a different way to camouflages specific traits, often in the way which is more accepted by society (3). Participants use masking as a tool to hide their over-sensitivity.

Masking causes others to view participants as tougher than they really are. They described how this led to further rejection and criticism, creating a vicious cycle, and reinforcing the <u>'tough'</u> exterior mask.



Cons preprint doines for body has 1123 for this version posted November 18,2824. The copyright holde perpetuity. within participant's made available inder of them to feel unsure of what their true feelings really are.

Withdrawal

Participants expect rejection, causing them to withdraw - such as avoiding university work, and withdrawing from friendships, and romantic relationships.



Socially withdrawing reinforces the idea that others are correct to reject them. This leads them to question if there is something wrong with them, causing them to only keep small circles of trusted people.

Bodily Sensations

Participants explained how rejection and



Figure 1