It is made available under a CC-BY-NC 4.0 International license .

Publicly available continuously updated topic specific databases of randomised clinical trials: A scoping review

Kim Boesen¹, Lars G Hemkens^{1,2}, Perrine Janiaud¹, Julian Hirt^{1,3,4}

¹ Pragmatic Evidence Lab, Research Center for Clinical Neuroimmunology and Neuroscience (RC2NB), University Hospital Basel and University of Basel, Basel, Switzerland

² Meta-Research Innovation Center at Stanford (METRICS), Stanford University, Stanford, California, USA

³ Department of Health, Eastern Switzerland University of Applied Sciences, St.Gallen, Switzerland

⁴ Institute of Health and Nursing Science, Medical Faculty, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

Authors' ORCID and email

Kim Boesen: 0000-0001-8963-8585, boesen.kim@gmail.com

Lars G. Hemkens: 0000-0002-3444-1432, lars.hemkens@usb.ch

Perrine Janiaud: 0000-0001-7684-8014, perrine.janiaud@usb.ch

Julian Hirt: 0000-0001-6589-3936, julian.hirt@usb.ch

Corresponding author

Julian Hirt, Research Center for Clinical Neuroimmunology and Neuroscience, University Hospital Basel and University of Basel, Spitalstrasse 2, CH-4031 Basel, Switzerland, julian.hirt@usb.ch

It is made available under a CC-BY-NC 4.0 International license .

Words: 2793 Figures/Tables: 3

References: 69

It is made available under a CC-BY-NC 4.0 International license .

Abstract

Conducting systematic reviews of clinical trials is arduous and resource consuming. One potential solution is to design databases that are continuously and automatically populated with clinical trial data from harmonised and structured datasets. We aimed to map publicly available, continuously updated, topic-specific databases of randomised clinical trials (RCTs). We systematically searched PubMed, Embase, the preprint servers medRxiv, ArXiv, and Open Science Framework, and Google. We described seven features (access model, database architecture, data input sources, retrieval methods, data extraction methods, trial presentation, and export options) and narratively summarised the results. We did not register a protocol for this review. We identified 14 continuously updated clinical trial databases, seven related to COVID-19 (first active in 2020) and seven non-COVID databases (first active in 2009). All databases, except one, were publicly funded and accessible without restrictions. They mainly employed methods similar to those from static article-based systematic reviews and retrieved data from journal publications and trial registries. The COVID-19 databases and some non-COVID databases implemented semi-automated features of data import, which combined automated and manual data curation, whereas the non-COVID databases mainly relied on manual workflows. Most reported information was metadata, such as author names, years of publication, and link to publication or trial registry. Two databases included trial appraisal information (risk of bias assessments). Six databases reported aggregate group level results, but only one database provided individual participant data on request. We identified few continuously updated trial databases, and existing initiatives mainly employ methods known from static article -based reviews. The main limitation to create truly live evidence synthesis is the access and import of machinereadable and harmonised clinical trial data.

It is made available under a CC-BY-NC 4.0 International license .

Introduction

The number of conducted and published clinical trials and systematic reviews has increased tremendously over the past decades.^{1, 2} It is difficult, if not impossible, for any clinician or researcher to stay up to date. Traditional systematic reviews are time consuming and costly to produce due to the time required to manually search databases, screen titles, assess studies, and extract data.³ Systematic reviews are often outdated and potentially even misleading when published. ⁴⁻⁷ Different tools have been launched to automatise the review conduct and maintenance to reduce the costs and resources.⁸⁻¹¹ Those tools focus mainly on text mining and machine learning assistance to screen and retrieve records¹² and extract data from published reports of clinical trials.¹³

Due to the systematic reviews' limitations of staying up to date, the concept of 'living systematic reviews' has popularised. There is no clear definition of what a 'living review' is other than it is a (systematic) review being continuously updated.^{14,15} In theory it should mitigate delays and the risk of missing important evidence. In 2017, the Cochrane Collaboration published what they call "the world's first 'living systematic review". The review was published in 2017, and it has still not being updated in the Cochrane Library.^{16,17} In recent years, the COVID-19 pandemic ignited a torrent of such 'live' projects reflecting the rapidly evolving evidence, including the COVID-19 trial tracker, ¹⁸ the NMA-Covid Project, ¹⁹ and WHO's living guideline²⁰ that were known to us prior to this scoping review (S1, Appendix).

The first requirement to enable such continuously updated databases of clinical trial data is to have all trial data available in a machine-readable format and structured in a harmonised fashion.²¹ The idea of such machine-readable repository of RCTs goes 25 years back with the Global Trial Bank. It was a system envisioned to go beyond trial registration to also contain trial results reporting in a computable fashion.²²⁻²⁵ The Global Trial Bank never materialised, but along these lines the International Committee for Journal Medical Editors' clinical trial registry requirement as a pre-empt for publication came into place in 2004.²⁶ Subsequently, it also became mandatory to submit summary results on ClinicalTrials.gov (2007) and on the European Union Clinical Trial Register (2012).²⁷

Some researchers have voiced that clinical trial data should be made available in a structured dataset, like the Common Technical Document format used for commercial drug

It is made available under a CC-BY-NC 4.0 International license .

applications.²⁸ Others have argued that evidence synthesis, like systematic reviews, should rely primarily on clinical trial registries as the main data source, rather than the traditional decentralised publication-based system, to enable more timely and exhaustive evidence synthesis.²⁹ The concept of the Global Trial Bank and of continuously updated, clinical trial databases lies in the intersection of these two ideas.

Whether existing living projects and topic specific trial databases leverage advanced technologies to ensure seamless and – ideally fully automated – flow of clinical trial data into structured databases is unknown. We therefore aimed to systematically identify publicly available continuously updated topic-specific databases of clinical trials to characterise basic functionalities, infrastructure, and user interface. We used this work as a primer for the design of a continuously updated database on specific cancer immunotherapies.³⁰

Methods

We conducted a scoping review and due to its exploratory nature, we did not follow a protocol or register this review since we anticipated numerous unforeseeable challenges during mapping this field. We report this scoping review according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses statement for Scoping Reviews (PRISMA-ScR).³¹

Eligibility criteria

We included any publicly available continuously updated database retrieving RCTs (participants allocated to one of two or more groups); irrespective of retrieval method (e.g. manually, automated, or a combination), input source (e.g. databases of published literature, trial registries, or both), and we did not define a minimum update frequency threshold. We considered field or topic specific databases only, i.e. exclusive to one condition, specialty, intervention, or indication, such as a COVID-19. Databases that also retrieved other research designs, such as observational studies, were still considered eligible. We did not include broad, unspecific trial databases, such as clinical trial registries. We

applied no restrictions on language, release date, intended use (i.e., for research or clinical decision-making), or the database status (i.e. live, archived, or unknown status).

We did not include conventional, static, article-based 'living systematic reviews' without an eligible database adhering to the inclusion criteria above.

Information sources and search strategy

We employed a two-step systematic search strategy. First, we did a "cold search" of various resources to identify key words and seed references to subsequently build a systematic database search.

Step 1: Cold ('seed') search

One author (KB) searched PubMed (including using the "similar articles" function), preprint servers medRxiv and ArXiv, Open Science Framework, and Google (first five pages for each search) in January 2023 (S2, Appendix).

Step 2: Systematic search

Two authors (KB, JH) designed a three-component search strategy containing keywords for "living" (i.e. live, up-to-date, updated, continuously updated, accumulating, dynamic, digital, and interactive), keywords for "library" (i.e. database, platform, hub, repository, collection, overview, archive, body of evidence, index, bibliography, catalogue, compilation, portal, inventory, and map), and study design (i.e. RCT, trial, and clinical study). One author (JH) searched PubMed and Embase (S3 and S4, Appendix) on February 1, 2023, deduplicated search results using Citavi,³² and imported the results into the screening tool Rayyan.³³ One author (KB) screened titles, abstracts, and full texts and decided on inclusion in one step.

Data extraction and analysis

We described and narratively summarised the following seven database features. One reviewer (KB) extracted information, and one reviewer (JH) double-checked and confirmed the extractions.

Access model

How is the database content accessed? Unrestricted access (i.e. data is freely available without any constraints), controlled access (e.g. users have to submit requests and the provider grants or denies access), or a hybrid between the two.

Database architecture

What is the underlying database structure and used database management software? Simple Excel spreadsheet or relational databases,³⁴ and were proprietary data management tools used, like RedCap.³⁵

Input sources

Which data input sources were searched? Databases of published literature like PubMed or Embase, trial registries, e.g. ClinicalTrials.gov, or other sources.

Retrieval methods

How is the database populated with trials? Manually, automated e.g. by application programming interfaces (APIs), or community-based data submissions, like ClinicalTrials.gov or the nucleotide sequencing database GenBank, where researchers submit data to the database.³⁶

Data extraction

How is the trial information extracted and curated? Manually, fully automated, or a combination of machine-assisted manual curation, usually referred to as 'semi-automated'.

Trial presentation

What trial information are provided? Meta data (e.g. title, author names, publication or trial registry information), detailed trial information (e.g. study design, sample size, treatment descriptions, funding), results (outcomes and effect estimates), type of results (aggregate group level or individual participant data), and trial appraisal (e.g. risk of bias assessments, limitations in the trial design).

Export options

What are the options for downloading and reusing data? Free download of the dataset (e.g. as a CSV file), federated access ³⁷ where users can view and work with the data remotely,

e.g. in browser-based applications without downloading it to their own personal computer such as Vivli.com, ³⁸ or no option for reusing data.

Results

Search results

Our systematic database search returned 1707 hits (deduplicated), of which 15 were assessed in full-text from which one database (TrialsResultsCenter³⁹⁻⁴⁰) was included. One database (Evidence Finder⁴¹⁻⁴³) was found through other sources, nine databases (Cochrane COVID-19 study register,⁴⁴ COVID-evidence,⁴⁵ COVID TrialsTracker,¹⁸ COVID-NMA initiative,¹⁹ EPPI Centre Covid-19 Living Map of the Evidence,⁴⁶ MetaEvidence breast cancer,⁴⁷ MetaEvidence COVID,⁴⁸ MetalO,⁴⁹ and MetaPreg⁵⁰) were known to us before our systematic search, and our cold searches yielded three other eligible databases (Infectious Disease Data Observatory (IDDO),⁵¹ Worldwide Antimalarial Resistance Network (WWARN),⁵²⁻⁵⁴ and EvidenceMap⁵⁵⁻⁵⁷; S5, Appendix).

In total, we included 14 databases (Table 1), seven non-COVID databases on youth mental health (Evidence Finder), pancreatic surgery (Evidence Map), cardiology and oncology trials (TrialResultsCenter), cancer (MetaEvidence breast cancer and MetalO), pregnancy (MetaPreg), and malaria (WWARN), and seven COVID-19 databases (Cochrane COVID-19 study register, COVID-evidence, COVID TrialsTracker, COVID-NMA initiative, EPPI Centre Covid-19 Living Map of the Evidence, IDDO, and MetaEvidence COVID).

Basic database characteristics (non-COVID)

TrialResultsCenter was active from 2009 until 2017 (and is no longer updated), EvidenceFinder was released in 2010 (last update in July 2021, uncertain if it is still maintained), WWARN was released in 2011 (uncertain if it still maintained; the last trial in their Data Inventory⁵² dates from 2018 but their trial summary⁵¹ says it is current until 2022), and EvidenceMap was released in 2021 (last update 26 July 2023). For three databases (MetaEvidence breast cancer, MetalO, and MetaPreg) the start was not stated. Six of the non-COVID databases were either publicly funded or funded by a non-profit

organisation, and one (Evidence Map) was funded by a pharmaceutical company. All database websites were fully accessible with no restrictions.

The databases generally searched common sources, mainly databases of published reports including PubMed/MEDLINE, Embase, and Web of Science, using methods resembling those of regular systematic reviews with manual database searches, extraction, and data curation. EvidenceMap was built on a relational Microsoft SQL server database (Windows Form .NET management system), WWARN uses a proprietary data management tool (RedCap), and it was not described for the other five. Five databases (EvidenceMap, MetaEvidence breast cancer, MetalO, MetaPreg, and WWARN) made their data readily available for download, one (Trial Results Center) upon request, and one (Evidence Finder) did not mention options for download (Table 2).

All databases provide basic metadata information, e.g. title, author names, and link to publication/trial registry. Three databases (MetaEvidence breast cancer, MetalO, and MetaPreg) reported aggregated group level results and risk of bias assessments, one (TrialsResultsCenter) reported aggregated group level results, two (Evidence Finder and TrialsResultsCenter) reported some trial information, and one (EvidenceMap) reported trial appraisal in the shape of risk of bias assessment (Table 2; S6, Appendix).

Basic database characteristics (COVID)

The seven COVID databases were initiated shortly after the SARS-COV-2 outbreak in early 2020, and by January 2024 the last database, the Cochrane COVID-19 study register, stopped being updated. They were all publicly funded or funded by non-profit institutions and open to the public. All websites were fully accessible with no restrictions.

Three of the seven COVID databases (Cochrane COVID-19 study register, EPPI Centre Covid-19 Living Map of the Evidence, and MetaEvidence COVID) generally searched common sources, mainly databases of published reports including PubMed/MEDLINE and Embase. Six of the seven databases searched (in addition) one or more trial registries. Three databases (Cochrane COVID-19 study register, COVID-evidence, COVID TrialsTracker, EPPI Centre Covid-19 Living Map of the Evidence, and Meta-evidence) used semi-automated methods for retrieval of studies and data extraction, mainly for trial registry records, and two (COVID

NMA Initiative and IDDO) manually retrieved and extracted data only. Two (IDDO and COVID-evidence) used a proprietary management tool (RedCap and Directus), and for the others it was not described. Six databases (Cochrane COVID-19 study register, COVID-evidence, COVID TrialsTracker, EPPI Centre Covid-19 Living Map of the Evidence, MetaEvidence COVID, and IDDO) made their data available for download and one (COVID NMA initiative) made it available upon request (Table 3).

All databases reported basic metadata and also trial information (such as study type or design, sample size or expected starting date extracted from ClinicalTrials.gov), two (Covid NMA Initiative and MetaEvidence COVID) reported aggregate group level results, and one (MetaEvidence COVID) also reported trial appraisal in the shape of risk of bias assessments (Table 3; S6, Appendix).

Discussion

We identified 14 continuously updated clinical trials databases, of which 7 were COVID-19 related. The COVID projects used semi-automated methods to retrieve and extract data, primarily from clinical trial registries, whereas the non-COVID databases mainly employed manual methods from regular systematic reviews, which are time consuming and difficult to maintain. All databases presented mainly metadata whereas curated trial information, results, or trial appraisal were sparsely reported. Only the IDDO database explicitly stated to provide the option of sharing individual participant data, which can then be accessed on request.

Findings in context

Our scoping review highlights a dearth of continuously updated topic-specific trial databases. Existing databases borrow methods and technology from static article-based systematic reviews. The main issue seems that most information from published clinical trials must be retrieved, extracted, and curated from unstructured journal articles, and that data captured or exported from clinical trial registries are not easily incorporated into databases. None of the 14 platforms solved this problem, for instance by retrieving data from other sources such

It is made available under a CC-BY-NC 4.0 International license .

as standardised machine-readable datasets. We do not have knowledge of such publicly available structured datasets. The public-private FDA Critical Path Initiative ⁵⁸ has established databases of Alzheimer's, ^{59,60} multiple sclerosis ⁶¹, Parkinson's ⁶² and tuberculosis ⁶³ clinical trial data, although these are not publicly available. The prevalence of such public-private or fully private data platforms is unknown to us.

There are several reasons why existing trial registries are challenging as a data source for creating topic-specific trial databases. First of all, the various trial registries do not harmonise database schemes or data format structures beyond the WHO Trial Registration Data Set.⁶⁴ It is a high-level reporting checklist that do not specify exhaustive outcome specifications and results reporting. Secondly, based on our initial experiences with the CIEL database,³⁰ we assume that the error rates of inserting wrong data or into the wrong fields in clinical trial registries is very high. This makes the usefulness of machine-readable trial registry datasets (e.g. ClinicalTrials.gov's dataset that is accessible through an application programming interface or API), rather limited. Imported data needs a substantial amount of "data cleaning" to become useful. Thirdly, only a few registries enable data export, whereas others, such as the European Union Clinical Trial Register does not have an option for automated access to data.

One potential tool to solve the problem of extracting data from journal publications is TrialStreamer.⁶⁵ It is not a topic-specific database, but a mega-collection of PubMed indexed clinical trials (n = 852.723 by 20 Dec 2023), with annotated PICO information (population, information, comparator, outcome). Trial Streamer uses a mix of machine-learning and crowd-sourced manual extraction to retrieve the data. TrialStreamer may be useful for highlevel meta-epidemiological studies, but for clinical applications or guideline development, more granular trial descriptions are needed.

The importance of living evidence synthesis was recently acknowledged by the Wellcome Trust.⁶⁶ They will invest GBP 45 million over a five-year period to establish and accelerate the development of data infrastructure for future living evidence projects.

Limitations

Our scoping review has limitations. First, we did not prespecify our methodology and register a protocol for this work but we believe that the impact is minimal considering the scope of the review with its exploratory nature. Second, we have likely missed datasources due to the scarcity and disparity of databases, lacking common terminology, and lacking related published articles. We are confident that many 'stealth'⁶⁷ commercial solutions exist. An example of a public commercial solution is the LiveSLR program,^{68,69} which is a software combining an annotated library of publications (called LiveRef) with a machine-assisted search/screening/ranking workflow, which creates an interactive live platform used in drug applications and health technology dossiers. Unfortunately, there is little transparency when it comes to such commercial solutions, including LiveSLR, and we cannot describe such solutions in more details. Finally, we decided to not make detailed assessments of the database infrastructure, such as granular descriptions of how data was retrieved and extracted automatically from ClinicalTrials.gov. Mainly because it was not transparently reported on the websites (nor appeared from the journal publications that we found in our systematic search). Transparent and exhaustive reporting of the database infrastructure should be a priority because automation is the main methodological challenge. For example, the metaPreg platform described to use "automatic software robots" to screen databases, with no further specifications.⁵⁰ We acknowledge that there may be several reasons for vaguely describing the methodology, for example because such setups are likely updated and adjusted frequently. However, open and complete descriptions of platform infrastructure, including how automation is attempted and achieved, is the fastest way to advance this field.

Conclusions

This is the first review on the topic of continuously updated clinical trial databases. It could inform the design of future databases and outline technical and methodological challenges. We found only few topic-specific databases of clinical trials, which mainly relied on manual searches and manual data extraction. Such approaches are time consuming and inefficient to keep the databases updated. Recent COVID-19 trial databases implemented semiautomated workflows, albeit confined to clinical trial registries. A major challenge remains how to automatically extract trial data from journal publications. It seems timely to

It is made available under a CC-BY-NC 4.0 International license .

reconsider ideas such as the Global Trial Bank to centralise, not only trial registration, but also trial results reporting and data sharing. Facilitating the shift from article-based publications to a fully 'digital' machine-readable system of clinical trial data could enable true live evidence synthesis.

It is made available under a CC-BY-NC 4.0 International license .

Acknowledgements

None.

Conflicts of interest

LGH, PJ, JH designed and worked on the COVID-evidence and PragMeta database. All authors worked on the continuously updated trial database, Cancer Immunotherapy Evidence Living (CIEL).

Authors' contributions

KB, PJ, and JH made substantial contributions to the conception and design of the work; KB have drafted the work and all authors substantively revised it; all authors have approved the submitted version; all authors have agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work, even ones in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature.

Data sharing statement

All data related to this review are available in the paper and the Appendix.

Funding

The CIEL Project was funded by the Basel Cancer League (KLbB-5577-02-2022). RC2NB (Research Center for Clinical Neuroimmunology and Neuroscience Basel) is supported by Foundation Clinical Neuroimmunology and Neuroscience Basel.

It is made available under a CC-BY-NC 4.0 International license .

References

- 1. Bastian H, Glasziou P, Chalmers I. Seventy-five trials and eleven systematic reviews a day: how will we ever keep up? *PLoS Med* 2010;7: e1000326.
- 2. Andersen MZ, Zeinert Ph, Rosenberg J, Fonnes S. Comparative analysis of Cochrane and non-Cochrane reviews over three decades. *Systematic Rev* 2024;13:120.
- 3. Borah R, Brown AW, Capers PL, Kaiser KA. Analysis of the time and workers needed to conduct systematic reviews of medical interventions using data from the PROSPERO registry. *BMJ Open* 2017;7: e012545.
- 4. Tricco AC, Brehaut J, Chen MH, Moher D. Following 411 Cochrane protocols to completion: a retrospective cohort study. *PLoS ONE* 2008;e3684.
- 5. Andersen MZ, Gülen S, Fonnes S, et al. Half of Cochrane reviews were published more than 2 years after the protocol. *J Clin Epidemiol* 2020;124:85-93.
- 6. Bashir R, Surian D, Dunn AG. Time-to-update of systematic reviews relative to the availability of new evidence. *Syst Rev* 2018;7:195.
- 7. Beller EM, Chen JKH, Wang ULH, Glasziou PP. Are systematic reviews up-to-date at the time of publication? *Syst Rev* 2013;2:36.
- 8. Marshall C, Sutton A, O'Keefe H, Johnson E (Eds.). (2022). The Systematic Review Toolbox. Available from: <u>http://www.systematicreviewtools.com/</u>.
- 9. Marshall IJ, Wallace BC. Toward systematic review automation: a practical guide to using machine learning tools in research synthesis. *Syst Rev* 2019;8:163.
- 10. Khalil H, Ameen D, Zarnegar A. Tools to support the automation of systematic reviews: a scoping review. *J Clin Epidemiol* 2022;144:22-42.
- Cowie K, Rahmatullah A, Hardy N, Holub K, Kallmes K. Web-Based Software Tools for Systematic Literature Review in Medicine: Systematic Search and Feature Analysis. JMIR Med Inform 2022;10:e33219.
- 12. Harrison H, Griffin SJ, Kuhn I, Usher-Smith JA. Software tools to support title and abstract screening for systematic reviews in healthcare: an evaluation. *BMC Med Res Methodol* 2020;20:7.
- Schmidt L, Olorisade BK, McGuinness LA, Thomas J, Higgins JPT. Data extraction methods for systematic review (semi)automation: A living systematic review. *F1000Res* 2021; 10:401.

14. Living Systematic Reviews. Cochrane Community.

https://community.cochrane.org/review-production/production-resources/livingsystematic-reviews .

- 15. Elliot JH, Turner T, Clavisi O, et al. Living Systematic Reviews: An Emerging Opportunity to Narrow the Evidence-Practice Gap. *PLoS Med* 2014;11: e1001603.
- 16. Cochrane. <u>https://www.cochrane.org/news/cochrane-welcomes-global-</u> <u>commitment-evidence-synthesis</u>.
- Millard T, Synnot A, Elliott J, et al. Feasibility and acceptability of living systematic reviews: results from a mixed-methods evaluation. *Syst Rev* 2019;8:325. doi:10.1186/s13643-019-1248-5
- 18. Trials tracker. <u>https://covid19.trialstracker.net/</u>.
- 19. COVID-NMA. <u>https://covid-nma.com/</u>.
- 20. Therapeutics and COVID-19: Living guideline, 13 January 2023. <u>https://www.who.int/publications/i/item/WHO-2019-nCoV-therapeutics-2023.1</u>.
- 21. Canham S, Ohmann C. A metadata schema for data objects in clinical research. *Trials* 2016;17:557.
- 22. Sim I, Owens DK, Lavori PW, Rennels GD. Electronic trial banks: a complementary method for reporting randomized trials. *Med Decis Making* 2000;20:440-50.
- 23. Sim I, CariniS, Olasov B, Jeng S. Trial bank publishing: phase I results. *Stud Health Technol Inform* 2004;107:1476-80.
- 24. Sim I, Detmer DE. Beyond Trial Registration: A Global Trial Bank for Clinical Trial Reporting. *PLOS Medicine* 2005;2:e365.
- 25. The trial Bank Project. <u>https://grantome.com/grant/NIH/R01-LM006780-10</u>.
- 26. Angelis CD, Drazen JM, Frizelle FA, et al. Clinical trial registration: a statement from the International Committee of Medical Journal Editors. CMAJ 2004; 171:606-7.
- 27. Goldacre B, DeVito NJ, Heneghan C, et al. Compliance with requirement to report results on the EU Clinical Trials Register: cohort study and web ressource. *BMJ* 2018;362:k3218.
- 28. Wieseler B. Beyond journal publications a new format for the publication of clinical trials. *Z Evid Fortbild Qual Gesundhwes* 2017;120:3-8.
- 29. Dunn AG, Bourgeois FT. Is it time for computable evidence synthesis? *J Am Med Inform Assoc* 2020;27:972-75.

- 30. Boesen K, Hirt J, Dueblin P, Laubli H, Kassenda B, Hemkens, LG et al. (2024). Rationale and design of the Cancer Immunotherapy Evidence Living (CIEL) Library: A continuously updated clinical trial database of cancer immunotherapies. https://doi.org/10.1101/2024.04.26.24306436
- Tricco AC, Lillie E, Zarin W, O'Brien KK, Colquhoun H, Levac D et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169:467–473.
- 32. Citavi. https://www.citavi.com/en
- 33. Rayyan. <u>https://www.rayyan.ai/</u>.
- BM. What is a relational database. Not dated. https://www.ibm.com/topics/relational-databases.
- Research Electronic Data Capture REDCap. <u>https://www.project-redcap.org/</u>. Not dated.
- 36. Genbank overview. What is Genbank? <u>https://www.ncbi.nlm.nih.gov/genbank/</u>
- 37. Rujano MA, Boiten JW, Ohmann C, Canham S, Contrino S, David R et al. (2024). Sharing sensitive data in life sciences: an overview of centralized and federated approaches. *Brief Bioinform* 2024;25:bbae262.
- 38. Vivli. Center for global clinical research data. https://vivli.org/.
- 39. TrialResults-Center.org. Searchable database of clinical trials results. <u>https://www.trialresultscenter.org/evidenceTable/default_f.html</u>.
- 40. Cucherat M. Trialresults-center: A web-based clinical trial results database providing dynamic systematic reviews and meta-analysis of clinical trials in cardiology. *Journal of the American College of Cardiology* 2010;55:Supplement 9 A132, E1239. <u>https://www.embase.com/records?subaction=viewrecord&id=L70096434</u>.
- 41. Hetrick SE, Parker Ag, Callahan P, Purcell R. Evidence mapping: illustrating an emerging methodology to improve evidence-based practice in youth mental health. *J Eva Clin Prac* 2010;6:1025-30.
- 42. Silva SD, Bailey AP, Parker AG, Montague AE, Hetrick SE. Open-access evidence database of controlled trials and systematic reviews in youth mental health. *Early Interv Psychiatry* 2019;3:474-7.
- 43. Evidence Finder. <u>https://orygen.org.au/Training/Evidence-Finder</u>.
- 44. Cochrane. <u>https://covid-19.cochrane.org/</u>.

- 45. COVID-evidence. https://covid-evidence.org/.
- 46. EPPI Centre COVID-19 Living Map of the Evidence. <u>https://eppi.ioe.ac.uk/eppi-</u> vis/Review/Index.
- 47. MetaEvidence breast cancer. <u>http://www.metaevidence.org/default0.aspx</u>.
- 48. Meta Evidence COVID-19. <u>http://www.metaevidence.org/default0.aspx</u>.
- 49. MetalO. <u>http://www.metaevidence.org/default0.aspx</u>.
- 50. MetaPreg. http://www.metaevidence.org/default0.aspx.
- 51. Infectious Disease Data Observatory. A living systematic review of registered COVID-19 trials. <u>https://www.iddo.org/covid-19/live-systematic-review-trials</u>.
- 52. Worldwide antimalarial resistance network (WWARN) Malaria Data Inventory. <u>https://www.wwarn.org/tools-resources/literature-reviews/wwarn-clinical-trials-</u> <u>publication-library</u>.
- 53. WWARN. Malaria Data Inventory. Not dated. https://www.iddo.org/wwarn/accessing-data.
- 54. Takata J, Sondo P, Humphreys G, et al. The WorldWide Antimalarial Resistance Network Clinical Trials Publication Library: A Live, Open-Access Database of Plasmodium Treatment Efficacy Trials. *Am J Trop Med Hyg* 2020;103:359-68.
- 55. Probst P, Hüttner FJ, Meydan Ö, et al. Evidence map of pancreatic surgery: protocol for a living systematic review and meta-analysis. BMJ Open 2019;9:e032353.
- 56. Probst P, Hüttner FJ, Meydan Ö, et al. Evidence Map of Pancreatic Surgery-A living systematic review with meta-analyses by the International Study Group of Pancreatic Surgery (ISGPS). Surgery 2021;5:1517-24.
- 57. Evidence map. Living evidence map of pancreatic surgery. https://www.evidencemap.surgery/.
- 58. Critical Path Institute. https://c-path.org/tools-platforms/.
- 59. Hubscher E, Chennakrishnaiah S, Forsythe A. HTA102 Machine-Learning Technology Assisted Curated Reference Libraries as an Approach for Rapid Global Value Dossier Updates to Support Living Health Technology Assessment: A Case Study in Triple Refractory Multiple Myeloma (TRMM). *Value in Health* 2022;25:S316.
- 60. Critical Path Institute. https://c-path.org/programs/cpad/.

It is made available under a CC-BY-NC 4.0 International license .

- 61. LaRocca NG, Hudson LD, Rudick R, Amtmann D, Balcer L, Benedict R, et al. The MSOAC approach to developing performance outcomes to measure and monitor multiple sclerosis disability. *Multiple Sclerosis*. 2018;24:1469–84.
- 62. Critical Path Institute. https://c-path.org/program/critical-path-for-parkinsons/.
- 63. Critical Path Institute. <u>https://c-path.org/tools-platforms/tb-pacts/</u>.
- 64. WHO Trial Registration Data Set (Version 1.3.1). Not dated. <u>https://www.who.int/clinical-trials-registry-platform/network/who-data-set</u>.
- 65. TrialStreamer. https://trialstreamer.ieai.robotreviewer.net/.
- 66. Wellcome Trust. <u>https://wellcome.org/news/evidence-synthesis-infrastructure-</u> <u>collaborative</u>
- 67. Cristea IA, Cahen EM, Ioannidis JPA. Stealth research: Lack of peer-reviewed evidence from healthcare unicorns. Eur J Clin Invest 2019;49:e13072.
- 68. Cytel LiveSLR. https://www.cytel.com/live-slr.
- 69. Liu J, Verhoek A, Thorlund K, Heeg B, Sarri G, Forsythe A. HTA35 Should Health Technology Assessment (HTA) Bodies Utilize Living HTA Tools? Validation of LiveSLR® and LiveNMA[™] Tools Using ICER'S Class Review in Relapsed Refractory Multiple Myeloma (RRMM). *Value in Health* 2022; 25:S302-3.

Table 1. Included continuously updated field-specific clinical trials databases

Name	Year	Field	Funding/responsible organisation	Access
Cochrane COVID-19 study register	2020-2024	COVID	Cochrane and the Federal	Fully open
			Ministry of Education and	
			Research of Germany	
			(CEOsys project)	
COVID-e vidence	2020-2022	COVID	Swiss National Science	Fully open
			Foundation	
COVID NMA initiative	2020-2023	COVID	Centre for Research in	Fully open
			Epidemiology and	
			Statistics (CRESS) and	
			Cochrane France	
COVID Trials Tracker	2020-2022	COVID	Centre for Evidence Based	Fully open
			Medicine, Oxford	
			University	
EPPI Centre Covid-19 Living Map of	2019-2023	COVID	Evidence for Policy and	Fully open
the Evidence			Practice (EPPI) Centre and	
			the UK National Institute	
			for Health and Care	
			Research (NIHR) Policy	
			Research Programme	
			Reviews Facility	
Evidence Finder	2010 - 2021	Youth mental health	Orygen Ltd [®]	Fully open
Evidence Map	2021 – active	Pancreatic surgery	International Study Group	Fully open
-			for Pancreatic Surgery and	
			Viatris ^b	
IDDO Infectious disease Data	2020 - active	COVID	Coalition for Equitable	Fully open
Observatory Living systematic			ResearCh in Low-resource	
review			sEttings (CERCLE) ^c	
MetaEvidence breast cancer	Not stated	Breast cancer	EU Horizon 2020 Qualitop,	Fully open
			Hospices Civils de Lyon,	
			CNRS UMR5558 LBBE	
Meta Evidence COVID	2020-2022	COVID	Hospices Civils de Lyon	Fully open
MetalO	Not stated	Can cer	EU Horizon 2020 Qualitop,	Fully open
			Hospices Civils de Lyon,	
			CNRS UMR5558 LBBE	
MetaPreg	Not stated	Pregnancy	ANSM (agence Nationale	Fully open
			de Sécurité du	
			Médicament), Hospices	
			Civils de Lyon and	
			University Lyon-1 (LBBE)	
TrialResultsCenter	2009 - 2017	Cardiology and	Hospices Civils de Lyon	Fully open
		oncology		
WWARN Worldwide Antimalarial	2011 - active	Malaria	Coalition for Equitable	Fully open
Resistance Network Clinical Trials			ResearCh in Low-resource	
Publication Library			sEttings (CERCLE) ^d	

a) An Australian charity devoted to mental health issues (https://www.orygen.org.au/).

b) We could not find information on the International Study Group for Pancreatic Surgery including its funding. Viatris is a pharmaceutical company.

c) An international research network currently funded by the German Ministry of Education and Research.

d) WWARN is one of IDDO's projects, thus we assume the funding is the same.

It is made available under a CC-BY-NC 4.0 International license .

Name	Database architecture	Input sources	Retrieval methods (frequency)	Curation	Trial presentation	Export
Evidence Finder	Not described	MEDLINE, Embase, PsychINFO, Cochrane Library	Manually (yearly)	Manual	Metadata, trial information	Not mentioned
Evidence Map	Relational database (Windows Form .NET management system)	PubMed, CENTRAL, Web of Science	Manually (frequency not described)	Manual	Metadata, trial appraisal	Excel download
MetaEvidence Breast cancer	Not described *	Not described *	Not described *	Not described *	Metadata, trial information, trial appraisal, results (group level)	No option
MetalO	Not described *	Not described*	Not described*	Not described*	Metadata, trial information, trial appraisal, results (group level)	No option
MetaPreg	Not described ("Proprietary meta-analysis platform")	PubMed, Scopus, Web of Science, 17 preprint servers	Semi-automated (daily) ("using automatic software robots")	Manual	Metadata, trial information, trial appraisal, results (group level)	No option
Trial Results Center	Not described	MEDLINE, Embase, CENTRAL, ClinicalTrials.gov, ISRCTN, ICTRP	Manually (weekly)	Manual	Metadata, trial information, results (group level)	On request
WWARN	Proprietary data management system (RedCap)	PubMed, Embase, Web of Science	Manually (every 6 months)	Manual	Metadata	Exceldownload

Table 2. Database basic characteristics (non-COVID)

There is no separate method s description for these platforms. They link to the metaEvidence COVID methods.

It is made available under a CC-BY-NC 4.0 International license .

Name	Database architecture	Input sources	Retrieval methods (frequency)	Cura tion	Trial presentation	Export
Cochrane COVID- 19 study register	Not described	PubMed, Embase, Cochrane Central Register of Controlled Trials (CENTRAL), ClinicalTrials.gov, WH O International Clinical Trials Registry Platform (ICTRP), medRxiv, Retraction Watch	Semi-automated (daily to monthly)	Semi- automated	Metadata, trial information	CSV (no longer available)
COVID-evidence	Directus	ClinicalTrials.gov, ICTRP, LOVE platform	Semi-automated (weekly)	Semi- automated	Metadata, trial information	Excel and CSV download
COVID NMA initiative	Not described	ICTRP, LOVE, Cochrane COVID-19 register	Manual (monthly)	Manual	Metadata, trial information, results (group level)	On request
COVID Trials Tracker	Not described	ICTRP	Semi-automated (weekly)	Semi- automated	Metadata, trial information	Excel and CSV download
EPPI Centre Covid-19 Living Map of the Evidence	Not described	Microsoft Academic Graph (MAG) dataset, OpenAlex (since 2020; before MEDLINE and Embase)	Semi-automated (bi-weekly to monthly)	Semi- automated	Metadata, trial information	Excel, Citations (.txt), and Research Information System (RIS)
IDDO	RedCap	ICTRP	Manual (monthly)	Manual	M etadata, trial information, individual trial participant data (on request)	Excel download
Me ta Evide nce COVID	Not described	PubMed, Lit Covid, Scopus, Web of Science, ClinicalTrials.gov, WHO I CTRP, 17 preprint servers	Semi-automated (daily)	Semi- automated*	Metadata, trial information, results (group level), trial appraisal	Excel download

Table 3. Database basic characteristics (COVID specific)

* "Abstracts of studies identified in the above search were examined by a biocurator and an automatic classification algorithm. Each

abstract is analyzed by this automatic classification algorithm that determines an index of relevance of the abstract for meta-analysis. This

automatic classification algorithm is based on naïve Bayes classifier."