

Factors influencing healthcare providers' knowledge and skills retention following helping mothers survive and helping babies survive training in Tanzania: A mixed methods cross-sectional study

Beatrice Erastus Mwilike^{1*}, Martha Rimoy², Lucy Mabada², Nicodem Komba², Feddy Mwangi², Rashid Gosse¹, Joel Ambikile³

Affiliations:

1. Department of Community Health Nursing, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania
2. Tanzania Midwives Association (TAMA), Dar es Salaam, Tanzania
3. Department of Clinical Nursing, Muhimbili University of Health and Allied Sciences, Dar es Salaam, Tanzania

*Corresponding author

Beatrice Erastus Mwilike

E-mail: beatyerastus@gmail.com

Phone: [+255 712 620 924](tel:+255712620924)

1 Abstract

2 The Tanzanian Midwives Association (TAMA), in collaboration with development
3 partners, implemented a project (*50,000 Happy Birthdays*) from 2018 to 2020 to improve
4 providers' knowledge and skills in saving lives at birth. The project was implemented
5 under two training modules namely *Helping Mothers Survive* (HMS) and *Helping Babies*
6 *Survive* (HBS). Through a mixed-methods cross-sectional design, knowledge retention
7 was assessed by comparing follow-up and post-test knowledge scores among healthcare
8 providers in the selected health facilities in Tanga, Geita, and Katavi regions. Four
9 knowledge areas were evaluated using a written test and five skills areas were assessed
10 using skills observation checklists for each. Quantitative data was analyzed using IBM
11 SPSS version 25 by performing descriptive analysis, *t*-test and One-Way ANOVA with
12 the level of significance determined at $p < 0.05$. Qualitative analysis was conducted
13 through a thematic analysis approach and themes were generated to describe the factors
14 influencing retention. 210 respondents participated in the study with more than half
15 holding diploma ($n=116$; 55.2%) and working in urban area ($n=123$; 58.6%). There was
16 a statistically significant drop in mean knowledge scores for controlling bleeding after
17 birth ($t=15.404$, $p<0.001$), in helping babies breathe ($t=8.580$, $p<0.001$), and in essential
18 care for small babies ($t=19.620$, $p<0.001$). Working in a rural area had a statistical
19 significant higher drop in mean knowledge scores for managing pre-eclampsia ($F=4.491$,
20 $p=0.035$) and for helping babies breathe ($F=5.005$, $p=0.026$). Education level also showed
21 a significant difference in the mean knowledge score for managing pre-eclampsia
22 ($F=4.850$, $p=0.003$).

23 There was poor knowledge retention following receiving training on HMS and HBS. The
24 level of professional education and residential area significantly influenced knowledge

25 retention. Frequent refresher training and other relevant training among healthcare
26 workers may be helpful in knowledge retention regarding maternal and neonatal care,
27 with much emphasis on providers working in rural areas and those with lower level of
28 professional education.

29

30 **Keywords:** Knowledge, Post-training, Helping mothers survive, Helping babies survive,
31 Tanzania

32 **Introduction**

33 Despite the global, regional, and local efforts to improve maternal and neonatal services,
34 maternal and neonatal mortality have remained major public health issues for decades,
35 particularly in low- and middle-income countries (LMICs). Globally, every two minutes,
36 a woman dies from preventable causes related to pregnancy, which is statistically
37 equivalent to approximately 800 maternal deaths per day [1]. Of these, almost 95% occur
38 in LMICs [1, 2], with the limited skills of the birth attendants being one of the leading
39 factors [2, 3]. Besides, approximately 6,500 newborns (within the first 28 days of birth)
40 die every day. In 2022, 2.3 million newborns died in the first month of life, accounting
41 for about 47 percent of all under-five deaths [4], mainly attributed to a lack of quality care
42 at birth or skilled care and treatment immediately after birth and in the first days of life
43 [5].

44 According to the 2022 Tanzania Demographic Health and Malaria Indicator Survey
45 (TDHS - MIS), the maternal mortality ratio (MMR) was 104 maternal deaths per 100,000
46 live births, indicating a notable decline in MMR from 556 per 100,000 live births in
47 2015/2016 [6], and a step forward towards achieving the Standard Developmental Goal
48 (SDG) 3.1 target of reducing MMR to less than 70 deaths per 100,000 live births by 2030
49 [7]. Unlikely, the neonatal mortality rate has been stagnant over the past two decades,
50 decreasing at a slow pace from 40 deaths per 1,000 live births in 1999 to 25 deaths per
51 1,000 live births in 2015/2016 to 24 deaths per 1,000 live births in 2022 [6], indicating
52 slow progress towards achieving the SDG 3.2.2 target of ending preventable newborn
53 deaths by 2030 [7].

54 To reach the SDG 3.1 and 3.2.2 targets, deliberate analysis and corresponding
55 interventions are needed across various settings, with extensive focus on those showing

56 sluggishness in progress [6, 7]. Recently, the Tanzanian government has increased health
57 infrastructure by building new health facilities and expanding the existing ones in an
58 effort to improve accessibility to health services, including reproductive, maternal,
59 newborn, and child health (RMNCH) services [8]. Subsequently, services such as
60 caesarean sections, which were mainly provided at referral and tertiary hospitals, have
61 been decentralized to primary healthcare settings, particularly district hospitals and health
62 centers that meet the criteria. Despite these efforts, the skilled health personnel needed to
63 provide quality services in these health facilities, as per the World Health Organization
64 (WHO) recommendation, is one of the critical challenges [9, 10]. The WHO recommends
65 high-quality care for all women during the antenatal, intranatal, and postnatal periods [11]
66 and that all births be attended by skilled birth attendants (SBA) who are supplied with the
67 necessary requirements, including in-service training [11, 12]. Yet, Tanzania faces an
68 estimated half-shortage of skilled health workforce [13], and a notable shortage of SBA
69 [14]. In fact, 66% of maternal deaths and 43% of newborn deaths can be prevented in
70 births that occur with the assistance of SBAs who are supported and trained in basic life-
71 saving competencies [15, 16], and one of the strategies to achieve the SDG 3.1 target by
72 2030 is to have at least 90% of births attended by SBA [7].

73 In response to this shortage and the need for high-quality services, in-service training has
74 been implemented in multiple settings to strengthen the knowledge, skills, and
75 performance of healthcare providers in maternal and newborn care units [17–20]. These
76 trainings take into account various approaches and implementation frameworks, even
77 though the content is quite similar [17, 21]. Of these, Helping Mothers Survive (HMS)
78 and Helping Babies Survive (HBS) programs have been implemented as innovative
79 training strategies that are cost-effective while meeting the demands for more basic and

80 practical training, particularly in resource-limited settings [17, 22, 23]. It is mainly
81 focused on the management of the leading contributors to maternal and neonatal deaths,
82 including controlling bleeding after birth, active management of the third stage of labour,
83 administration of magnesium sulphate, manual removal of the placenta, helping babies
84 breathe, essential care for small babies, and feeding with a nasogastric tube [22, 23]. HMS
85 and HBS training have shown varying effectiveness in the settings where they have been
86 implemented. A review conducted in Tanzania reports a notable improvement in
87 knowledge, skills, and performance among SBA and a reduction in early neonatal deaths
88 and fresh stillbirths following in-job training on neonatal resuscitation [24]. Similarly,
89 reviews from other regions have also highlighted post-training increases in knowledge,
90 skills, and confidence [25, 26]. Even though, in terms of clinical practice, some studies
91 have reported improvement [27–30], while others have reported no significant
92 improvement [31]. It is further recommended that for better maternal and neonatal
93 outcomes, multifaceted interventions with structured follow-up interventions are rather
94 encouraged than single interventions or interventions with no follow-up [25, 26, 32].
95 Although HBS and HMS programs have shown significant impact in terms of
96 improvement of knowledge and skills [28], post-training follow-up surveys show a
97 decline in knowledge and skills [33, 34]. A training in rural Tanzania among SBA led to
98 an improvement in knowledge, skills, and confidence immediately post-training [33], but
99 a nine-month follow-up revealed a decline in knowledge and simulated skills [35].
100 Similarly, a cluster-randomised trial involving 61 facilities in Tanzania revealed a decline
101 of skills to 4.0% at a 10-month follow-up of in-facility training on helping mothers
102 survive bleeding after birth (HMS BAB) [34]. This entails the short-term retention of
103 knowledge, skills, and competencies gained in training, thus necessitating the need to find

104 out the factors that may be influencing post-training retention of knowledge and skills.
105 Previous studies in other settings have highlighted factors such as frequency of training,
106 years of clinical experience [36, 37], cadre variations [37, 38], and training modality [38].
107 However, they are not comprehensive enough to inform about knowledge and skill
108 retention in our context.

109 Therefore, as we owe it to enhance the training outcomes in terms of long-term retention
110 of knowledge and skills and subsequent improved clinical outcomes, our study aimed at
111 exploring the factors influencing healthcare providers' knowledge and skills retention
112 following helping mothers survive and helping babies survive training in Tanzania, as
113 they are not yet clear. In 2018 to 2020 the healthcare providers received training through
114 a programme known as 50,000 Happy Birthdays (50KHB). The programme aimed to
115 increase the professional capacity of midwives in three countries: Ethiopia, Rwanda and
116 Tanzania. This was a post-training follow-up survey conducted among the healthcare
117 providers who previously received HMS and HBS training mainly in four knowledge and
118 five skills areas. We, therefore, aimed to assess the factors influencing competence
119 retention following HMS and HBS in Tanzania.

120

121 **Materials and Methods**

122 **Study setting**

123 This study was conducted in six purposefully selected districts in three *50,000 Happy*
124 *Birthday* project regions of mainland Tanzania namely Geita, Tanga and Katavi. The
125 districts were Geita Town Council, Chato District Council, Tanga Town Council, Muheza
126 District Council, Mpanda District Council and Mpanda Town Council. These districts

127 were selected from regions where the 50KHB programme was conducted and impact
128 evaluated. For each of the six districts, the team randomly selected three health facilities
129 of different levels among those facilities which were part of the project. In each region,
130 facilities were selected using the following criteria:

131 (i) Facilities that have providers who are already trained on HMS and HBS modules.

132 (ii) For the three regions selected, selected a mix of facilities that represent hospital,
133 health center and dispensary.

134 (iii) Facilities representing urban, and rural contexts from each region.

135

136 **Study design**

137 Our study utilized a mixed-methods cross-sectional design to compare the current
138 situation at selected sites with endline data from the *50,000 Happy Birthdays* project.
139 Leveraging existing master trainers in each region for efficient and consistent data
140 collection, the study employed standard *HMBS* evaluation documents – multiple-choice
141 knowledge assessments and objective structured clinical evaluations (OSCEs) – as well
142 as supplemental tools like facility-level interviews, clinical observations, and desk
143 reviews to capture all relevant qualitative and quantitative information that may influence
144 results.

145 We engaged a significant and representative sample of study participants to ensure that
146 the findings can be applied to national best practices: we identified a diverse array of
147 health facilities (district hospitals, health centres, dispensaries) and maternal/newborn
148 service providers from each of the three project regions to evaluate. A subsequent,
149 centralised analysis was conducted to understand common factors associated with
150 competence retention among providers.

151 **Study Participants**

152 This study involved two different populations: healthcare providers and tutors. Healthcare
153 providers and tutors were enrolled from the following target populations:

154 **Target population 1** – Healthcare providers from selected health facilities who have been
155 trained on HMS and HBS modules.

156 **Target population 2** – Supervisors, DRCHCOs and RRCHCOs, and educators who have
157 been trained, and are proficient in the provision of maternal newborn and childcare in
158 their contexts.

159 **Target population 3** – Tutors who were trained on HMS and HBS modules in the
160 selected Nursing and Midwifery Colleges.

161

162 **Inclusion and exclusion criteria**

163 Healthcare providers (HCPs) and tutors were eligible for inclusion in the study if they
164 were currently living in Geita, Katavi and Tanga regions, were classified as being one of
165 the three target populations and had consented to participate in skills drills, focus groups
166 or interviews. Participants were excluded if they were sick or on study leave and they
167 plan to move from the region during the study period.

168

169 **Sampling Methods and Procedure**

170 **Sampling for Quantitative Research**

171 All the trained providers in the 28 purposively selected facilities and three training
172 institutions were approached and enrolled in the study if they consented. Healthcare
173 providers and tutors from selected health facilities who have been trained in HMS and
174 HBS modules were recruited. Consecutive recruitment was employed until all the trained

175 providers available were enrolled. Below in table 1 is the list of Health facilities and
 176 Training centers supported under 50KHB projects in Tanga, Geita, and Katavi regions.
 177 Overall, we were able to access a total of 210 participants from all the study sites (**Table**
 178 **1**).

Table 1. Sample of the study sites			
	Tanga	Geita	Katavi
District	Tanga Town Council	Geita Town Council	Mpanda Municipal Council
Education institution	Tanga COHAS	Geita School of Nursing	n/a
Hospital	Bombo Regional Referral Hospital	Geita Regional Referral Hospital	Katavi Referral Regional Hospital
Primary health facilities	Ngamiani HC, Mikanjuni HC, Duga dispensary	Nyankumbu HC, Kasamwa HC, Bung’wankoko dispensary	Town council clinic, Kakese dispensary, Mwamkulu dispensary
District	Muheza District Council	Chato District Council	Mpanda District council
Education institution	St. Augustine Muheza IHAS	n/a	n/a
Hospital	Muheza DH	Chato DH	n/a
Primary health facilities	Mkuzi HC, Bulwa HC, Misozwe dispensary, Mkanyageni dispensary	Bwanga HC, Buzilamoyo dispensary, Butalama dispensary, Kibehe dispensary	Mwese HC, Mishamo HC, Kalema HC, Katuma dispensary, Sibwesa dispensary, Kasekeseke dispensary

179

180

181 **Sampling for Qualitative research**

182 The sampling of providers was purposive to meet the desirable characteristics. Providers
183 were selected from diverse backgrounds to collate feedback from participants of different
184 socio-demographic characteristics. Focus groups were arranged to include groups of
185 similar characteristics. Participants included supervisors, DRCHCOs, RRCHCOs, and
186 educators who have been trained and are proficient in the provision of maternal, newborn
187 and childcare in their contexts.

188 The final sample sizes were based on theoretical saturation, which refers to the point at
189 which no new concepts emerge from the review of data drawn from a sample that is
190 diverse in pertinent characteristics and experiences. We achieved theoretical saturation
191 after having three focus groups with target populations composed of 12 participants each.
192 Also, we conducted a total of 16 in-depth interviews across all three regions.

193

194 **Data collection method and procedures**

195 Data was collected from January to March 2023, approximately two years following the
196 online survey for the 50,000 Happy Birthday project. The assessment of HCPs'
197 knowledge and skills utilized qualitative and quantitative approaches, including in-depth
198 interviews, OSCE, and questionnaires. Participants were recruited from the three regions
199 from 16 January until 31 March 2023.

200

201 **Qualitative data** from interviews and discussions were collected using an audio recorder.
202 Data were then transcribed verbatim and translated from Kiswahili into English. For
203 qualitative data collection, an interview guide will be used. Data were collected by a team

204 of main researchers. The interviews lasted for about 30 to 60 minutes and were conducted
205 in Kiswahili language.

206

207 **Quantitative data** were collected using the modified tools that were used for pre and
208 posttest during the 50,000 Happy Birthday program intervention. The questionnaires were
209 paper printed and participants were asked to fill in after providing signed informed
210 consent. The researchers and research assistants were responsible for data collection.

211 Tools were prepared to assess the knowledge and skills retention in HMS and HBS among
212 healthcare providers and tutors. The assessment focused on both knowledge and skills in
213 managing two HMS modules (Bleeding After Birth Complete (BABC) and Pre-
214 Eclampsia/Eclampsia (PEE)) and three HBS modules (Helping Babies Breathe (HBB2.0),
215 Essential Care for Every Baby (ECEB) and Essential Care for Small Babies (ECSB)).

216 The knowledge test tools comprise of socio demographic information and multiple-choice
217 questions for different topics covered during the training. The skills test tools comprise
218 of socio demographic information, instructions for conducting the skills and observation
219 checklists.

220 The knowledge test for HMS tools included a) Helping Mothers Survive: Bleeding after
221 Birth Complete Knowledge Assessment b) Helping Mothers Survive: Knowledge Test
222 Pre-Eclampsia & Eclampsia. The skills test for HMS tools included OSCE for a) Active
223 Management of Third Stage of Labour (AMTSL) b) Shock Management, and c)
224 Administering the loading dose of MgSO₄. The knowledge test for HBS tools included
225 a) Essential Care for Every Baby Knowledge Check b) Essential Care for Small Babies:
226 Knowledge check. The skills test for HBS tools included a) Helping baby breath and b)
227 Care for small babies.

228

229 **Data analysis**

230 *Quantitative data analysis*

231 Quantitative data from questionnaires were transferred to an Excel database and later to
232 Statistical Package for the Social Sciences (SPSS) version 25 for analysis. Descriptive
233 analyses were conducted using frequency, central tendencies, and other relevant methods
234 to establish follow up knowledge and skills retention. Similarly, these measures were used
235 in determining the impact of the training program on improvement of the maternal and
236 neonatal health outcomes. Data were compiled, cleaned, and analyzed for each specific
237 objective as shown in the following Table 2.

238

239 *Qualitative data analysis*

240 This study utilized deductive-inductive thematic analysis approach to identify, analyze,
241 and interpret patterns within the qualitative data. The audio recording information was
242 transcribed and transcripts in Swahili language were prepared. The researchers read
243 through the translated scripts repetitively and compare them with the original audio record
244 to check for accuracy. Thereafter, a list of meaningful information found was made and
245 codes assigned. Then, the list was classified into themes and each theme was classified
246 into sub-themes. Then, the data was translated and be presented in a text form. Validity
247 in qualitative data based on triangulation, member checking, and collaboration with other
248 investigators.

249

250 **Ethical approval**

251 The ethical approval for this study was obtained from the Muhimbili University of Health
252 and Allied Sciences Research and Ethics Committee with Ref.No.DA.282/298/01.C/.
253 Permission to conduct the study was sought from the Ministry of Health (MoH), the
254 President's Office Regional Administration and Local Government (PORALG), and from
255 regional and council/municipal authorities – with information being transmitted to facility
256 authorities. Participation in this study was voluntary and participants were required to
257 sign the written informed consent form. Time compensation was given to the participants
258 after the interviews. The research team ensured that data remains confidential, and no
259 identification will be published along with the reports of manuscripts. Interviews were
260 conducted in isolated rooms to ensure confidentiality and data will be stored in locked
261 cabinet in TAMA offices for a period of five years.

262

263 **Results**

264 **Characteristics of the study participants**

265 A total of 210 healthcare providers participated in the study, with the mean (SD) age of
266 35.7 (7.5), and most of them (n=145; 69.0%) were females. The majority (n=116; 55.2%)
267 were diploma holders, followed by certificate holders (n=76; 36.2%), and just more than
268 a half (n=110; 52.4%) were working in a regional or district referral hospital. The region
269 with the highest number of participants was Geita (n=78; 37.1%) followed by Tanga
270 (N=70: 33.3%), with Geita and Tanga municipal councils contributing to most
271 participants (n=47; 22.4% and n=46: 21.9%, respectively). Regarding the residential area
272 (not shown in table), more than half (n=123; 58.6%) of the participants were working in
273 health facilities located in the urban area (**Table 2**).

Table 2. Demographic characteristics of participants and mean follow-up knowledge and skills scores for managing mothers and babies during and after child birth (n=210)

		Knowledge Scores										Skills Scores										
		n	%	CBAB		Preeclampsia		HBB		ECSB		MRP		AMTSL		AdMgSO ₄		HBB		FNGT		
		n	%	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	\bar{x}	SD	
Mean age (SD)		35.7 (7.5)																				
Sex	Female	145	69.0	69.0	10.1	52.8	14.3	78.1	13.0	64.3	11.3	69.7	15.0	66.8	14.8	67.4	20.8	67.8	14.0	65.7	14.5	
	Male	65	31.0	68.3	11.6	55.6	14.5	78.1	11.2	63.3	12.1	7.7	18.1	67.3	13.8	64.8	20.1	67.0	15.6	67.9	14.7	
Education	Certificate/EN	76	36.2	67.0	10.7	49.9	15.3	74.7	13.4	61.9	11.4	65.1	15.7	64.7	14.6	64.2	23.1	65.9	15.8	61.8	17.0	
	Diploma (RN/AMO)	116	55.2	69.3	10.3	55.0	13.4	79.6	11.9	64.5	11.6	70.4	15.6	69.2	14.0	68.4	17.9	68.2	13.4	68.4	12.6	
	Degree (RN/MD)	16	7.6	73.5	11.1	60.9	13.6	83.5	8.3	69.1	10.0	72.2	19.8	60.4	15.7	64.1	26.6	67.3	13.7	73.1	10.9	
	Others (MA/CA)	2	1.0	72.5	9.2	55.0	7.1	78.0	7.1	65.0	7.1	55.5	7.8	75.0	0.0	72.5	7.8	91.5	6.4	70.0	7.1	
Health Facility	Dispensary	37	17.6	69.9	9.4	59.9	12.9	74.6	11.6	60.0	11.0	60.3	10.8	66.0	10.8	70.6	10.1	68.7	14.2	65.5	13.2	
	Health Centre	63	30.0	70.0	9.3	52.0	14.0	80.9	11.4	64.2	12.5	70.6	13.4	66.1	15.7	70.1	17.6	67.1	13.3	62.3	14.6	
	Hospital	110	52.4	67.8	11.5	52.5	14.7	77.7	13.0	65.1	10.8	70.0	18.1	67.8	14.9	63.3	24.0	67.3	15.3	69.0	14.5	
Region	Tanga	70	33.3	67.0	9.2	53.1	14.0	79.7	12.6	65.1	11.3	68.0	15.5	59.0	16.1	71.8	15.7	66.4	13.3	62.7	16.3	
	Geita	78	37.1	71.3	11.3	52.0	15.3	79.1	13.0	62.7	11.2	69.5	17.9	72.7	10.7	59.4	25.6	68.2	14.0	73.1	12.6	
	Katavi	62	29.5	67.7	10.5	56.3	13.5	75.0	11.2	64.2	12.0	67.8	14.4	68.6	12.7	69.8	15.3	67.9	16.4	62.2	11.7	
District/Council	Tanga City	46	21.9	68.2	9.3	51.3	13.4	81.1	12.1	66.3	10.8	72.7	15.3	60.1	17.9	72.4	15.2	65.2	14.7	60.9	14.6	
	Muheza DC	24	11.4	64.5	8.7	56.7	14.6	77.1	13.2	62.9	12.2	59.0	11.7	56.9	12.2	70.6	16.9	68.8	10.0	66.3	19.0	
	Geita MC	47	22.4	71.1	11.5	52.4	17.4	79.8	13.6	65.3	10.9	70.1	18.1	75.2	10.1	61.9	24.5	69.0	14.0	76.9	13.6	
	Chato DC	31	14.8	71.7	11.2	51.4	12.0	78.2	12.1	59.0	10.9	68.5	17.8	69.1	10.9	55.7	27.2	66.9	14.0	67.3	8.3	
	Mpanda MC	31	14.8	65.5	11.7	53.1	13.3	77.6	9.5	68.8	12.0	72.7	11.7	71.2	14.5	71.9	18.8	68.0	14.0	57.1	10.2	
	Tanganyika DC	31	14.8	70.0	8.8	59.5	13.2	72.3	12.3	59.6	10.1	62.9	15.3	66.1	10.1	67.7	10.7	67.8	18.7	67.3	10.9	

Key: CBAB; Controlling bleeding after birth, HBB; Helping babies breath, ECSB; Essential Care for Small Babies, MRP; Manual removal of placenta, AMTSL; active management of third stage of labour, AdMgSO₄; Administration of Magnesium sulphate, HBB; Helping babies breathe, FNGT; Feeding with nasogastric tube

276 **Participants' knowledge retention scores**

277 Four knowledge areas were assessed including, two for care of the mother i.e. how to
278 control bleeding after childbirth and managing pre-eclampsia, and two for the care of the
279 newborn i.e. helping babies to breathe and essential care for small babies. Participants'
280 mean follow-up knowledge (SD) score was highest for helping babies to breathe {78.1
281 (12.4)} and lowest for managing pre-eclampsia {53.6 (14.4)}. Follow-up knowledge
282 scores were compared to post-test knowledge scores (previously assessed) and the change
283 in mean (SD) scores was computed to determine knowledge retention. The results show
284 that there was a drop in mean knowledge score in all areas assessed with the highest drop
285 observed in managing pre-eclampsia {-21.0 (15.0)} and lowest in helping babies to
286 breathe {-7.7 (13.0)}.

287 A paired *t*-test was performed to assess the differences between the follow-up mean
288 knowledge scores and the post-test mean knowledge scores (**Table 3**) to determine
289 whether the observed drop in knowledge scores were significant. For all knowledge areas
290 assessed, there were statistically significant drop in knowledge scores. Specifically, there
291 was a statistically significant drop in mean knowledge scores for controlling bleeding
292 after birth ($t=15.404, p<0.001$), pre-eclampsia ($t=20.276, p<0.001$), helping babies breath
293 ($t=8.580, p<0.001$), and essential care for small babies ($t=19.620, p<0.001$).

294

295

296

297

298

299

Table 3. Mean scores differences between follow-up and post-test knowledge scores

Area assessed	Post-test		Follow-up		Difference		Paired <i>t</i> -test		
	Mean	SD	Mean	SD	Mean	SD	95%CI	<i>t</i>	p value
CBAB	81.3	7.3	68.8	10.5	-12.5	11.7	10.86703, 14.05678	15.404	<0.001
Pre-eclampsia	74.6	8.1	53.6	14.4	-21.0	15.0	18.93671, 23.01567	20.276	<0.001
HBB	85.8	7.8	78.1	12.4	-7.7	13.0	5.91977, 9.45166	8.580	<0.001
ECSB	78.6	7.3	64.0	11.5	-14.7	10.8	16.12989, 19.620	19.620	<0.001

Key: CBAB; Controlling bleeding after birth, HBB; Helping babies breath, ECSB; Essential Care for Small Babies, CI; confidence interval

300

301 An independent *t*-test was performed to compare the mean knowledge drop scores
 302 between participants working in urban and rural areas. Participants working in rural area
 303 had a statistically significant higher drop in mean knowledge scores for managing pre-
 304 eclampsia ($F=4.491, p=0.035$) and for helping babies breath ($F=5.005, p=0.026$) (**Table**
 305 **4**)

Table 4. Univariate analysis of variance of knowledge according to residential area

Measure	Urban		Rural		Independent <i>t</i> -test		
	Mean	SD	Mean	SD	F (1, 208)	Sig.	η^2
CBAB	-13.3	11.7	-11.2	11.7	1.677	0.197	0.008
Pre-eclampsia	-22.8	15.7	-18.4	13.7	4.491	0.035	0.021
HBB	-6.0	13.2	-10.0	12.4	5.005	0.026	0.023
ECSB	-13.7	10.4	-16.0	11.3	2.421	0.121	0.012

Key: CBAB; Controlling bleeding after birth, HBB; Helping babies breath, ECSB; Essential Care for Small Babies, CI; confidence interval η^2 ; F;Eta Squared,

306

307 Another statistical analysis that was performed for assessment of knowledge drop was the
 308 One-Way ANOVA to determine differences in mean knowledge scores among
 309 participants according to their levels of professional education i.e. holders of certificate,
 310 diploma, degree, and others (**Table 5**). There was a statistically significant difference only
 311 in the mean knowledge score for managing pre-eclampsia ($F=4.850, p=0.003$). When the
 312 Post Hoc Test was performed it was revealed that there were statistically significant mean
 313 knowledge differences in pre-eclampsia between certificate holders and diploma holders
 314 ($p=0.006$), and between certificate holders and degree holders ($p=0.037$). There were no
 315 significant differences in knowledge variance with other levels of professional education.

Table 5. One-Way ANOVA by level of professional education

Measure		Sum of squares	Mean square	F (3, 206)	<i>p</i> value
CBAB	Between Groups	528.591	176.197	1.287	0.280
	Within Groups	28197.604	136.882		
	Total	28726.195			
Pre-eclampsia	Between Groups	3099.223	1033.074	4.850	0.003
	Within Groups	43875.658	212.989		
	Total	46974.881			
HBB	Between Groups	474.154	158.051	0.937	0.424
	Within Groups	34745.103	168.666		
	Total	35219.257			
ECSB	Between Groups	381.232	127.077	1.086	0.356
	Within Groups	24114.083	117.059		
	Total	24495.314			

Key: CBAB; Controlling bleeding after birth, HBB; Helping babies breath, ECSB; Essential Care for Small Babies

316

317 **Participants' skills retention scores**

318 Five skills areas were assessed including, three for the care of the mother i.e. the manual
 319 removal of placenta, active management of the third stage of labour, and administration
 320 of magnesium sulphate, and two for the care of the newborn i.e. helping babies breathe
 321 and feeding with a nasal gastric tube. Participants' follow-up mean skills (SD) score was

322 highest for manual removal of placenta {68.5 (16.1)} and lowest for feeding with nasal
323 gastric tube {66.4 (14.6)}. The results showed that there was a drop in mean skills scores
324 in all areas assessed with the highest drop observed in the administration of magnesium
325 sulphate {-17.1 (18.7)} and lowest in helping babies breathe {-11.1 (11.0)}.

326

327 A paired *t*-test was performed to assess the differences between the follow-up mean skills
328 scores and the post-test skills mean scores (**Table 6**) to determine whether the changes in
329 skills scores were statistically significant. In all skills areas assessed, a statistically
330 significant drop in scores was observed. Specifically, there was a significant drop in mean
331 skill scores for manual removal of placenta ($t=13.427, p<0.001$), active management of
332 third stage of labour ($t=15.840, p<0.001$), administration of magnesium sulphate
333 ($t=13.210, p<0.001$), helping babies breathe ($t=14.609, p<0.001$), and feeding with nasal
334 gastric tube ($t=16.113, p<0.001$)

Table 6. Mean scores differences between follow-up and post-test skills scores

	Post-test		Follow-up		Difference		Paired <i>t</i> -test		
	Mean	SD	Mean	SD	Mean	SD	95%CI	<i>t</i>	p value
MRP	83.6000	7.88688	68.4905	16.07912	-15.1095	16.30723	12.89112, 17.32793	13.427	<0.001
AMTSL	83.9952	8.15255	66.9571	14.47694	-17.0381	15.58795	14.91754, 19.15865	15.840	<0.001
AdMgSO ₄	83.6268	9.49162	66.6048	20.56920	-17.0766	18.68859	14.52805, 19.62506	13.210	<0.001
HBB	78.6190	7.54747	67.5095	14.45178	-11.1095	11.02031	9.61034, 12.60871	14.609	<0.001
FNGT	78.8571	7.13027	66.4048	14.56886	-12.4524	11.19895	10.92890, 13.97586	16.113	<0.001

Key: MRP; Manual removal of placenta, AMTSL; active management of third stage of labour, AdMgSO₄; Administration of magnesium sulphate, HBB; Helping babies breathe, FNGT; Feeding with nasogastric tube, CI; confidence interval

335

336

Table 7. Univariate analysis of variance of skills according to residential area

Measure	Urban		Rural		Independent <i>t</i> -test		
	Mean	SD	Mean	SD	F (1, 208)	<i>p</i> value	η ²
MRP	-11.6	15.3	-20.2	16.5	14.917	<0.001	0.067
AMTSL	-14.8	16.1	-20.2	14.3	6.271	0.013	0.029
AdMgSO ₄	-15.4	17.5	-19.4	20.2	2.356	0.126	0.011
HBB	-12.1	11.6	-9.7	10.0	2.367	0.125	0.011
FNGT	-13.1	12.1	-11.5	9.8	1.028	0.312	0.005

Key: MRP; Manual removal of placenta, AMTSL; active management of third stage of labour, AdMgSO₄; Administration of magnesium sulphate, HBB; Helping babies breathe, FNGT; Feeding with nasogastric tube, CI; confidence interval η²; Eta Squared

337

338 An independent *t*-test (**Table 7**) was also performed to compare the mean skills drop scores between
 339 participants working in urban and rural areas. Participants working in rural area had a statistically
 340 significant higher drop in mean skills scores for manual removal of placenta ($F=14.917, p<0.001$),
 341 and for active management of the third stage of labour ($F=6.271, p=0.013$).

342 A One-Way ANOVA was performed to assess differences in skills drop scores among participants
 343 according to their levels of professional education i.e. holders of certificate, diploma, degree, and
 344 others (**Table 8**). There was a statistically significant difference within groups only in the skills for
 345 feeding with nasogastric tube ($F=2.713, p=0.046$).

346

Table 8. One Way Analysis of Variance of skills by level of professional education

Measure		Sum of squares	Mean square	F (3, 206)	<i>p</i> value
MRP	Between Groups	1207.331	402.444	1.525	0.209
	Within Groups	54371.150	263.938		
	Total	55578.481			
AMTSL	Between Groups	1337.333	445.778	1.857	0.138
	Within Groups	49446.362	240.031		
	Total	50783.695			
AdMgSO4	Between Groups	1253.140	417.713	1.199	0.311
	Within Groups	71393.636	348.262		
	Total	72646.775			
HBB	Between Groups	716.457	238.819	1.995	0.116
	Within Groups	24666.024	119.738		
	Total	25382.481			
FNGT	Between Groups	996.368	332.123	2.713	0.046
	Within Groups	25215.656	122.406		
	Total	26212.024			

Key: Key: MRP; Manual removal of placenta, AMTSL; active management of third stage of labour, AdMgSO4; Administration of magnesium sulphate, HBB; Helping babies breathe, FNGT; Feeding with nasogastric tube

347 **Factors influencing the knowledge and skills retention**

348 **1. Learning materials**

349 **a. Availability of learning materials**

350 Participants reported the availability of learning materials such as handouts, and reference books as
351 a crucial factor for knowledge and skills retention. Learning materials allow learners to review,
352 reinforce, and apply the concepts and skills they have acquired in the classroom. Handouts and
353 reference books can also serve as feedback, guidance, and self-assessment sources for learners. By
354 having access to learning materials, learners can enhance their understanding, deepen their memory,
355 and transfer their knowledge to new situations.

356 *“For example, when you are given out a handout, and revise every time, it will help you to*
357 *build experience. Although you may not be able to attend to a patient physically but it will*
358 *assist to read the references and help put your knowledge intact.” (FGD Tanga)*

359

360 **b. Topic revision in CMEs**

361 Participants highlighted the importance of topic revision during the Clinical Medical Education
362 sessions (CMEs) as one of the factors enhancing knowledge and skills retention. Topic revision is a
363 factor for knowledge retention because it helps learners consolidate their learning and recall it more
364 easily. Moreover, revising topics after a gap can strengthen memory traces and make them more
365 resistant to forgetting.

366 *“We normally have learning sessions at our facility; we have a teaching plan. Therefore, every*
367 *Wednesday we prepare a lesson. We prepare maybe this week it will be eclampsia, next week*
368 *it will be PPH, and we learn through those also we get experience.” (FGD Tanga)*

369

370 **c. Establishment of a learning corner**

371 Participants reported the establishment of a learning corner as another factor for knowledge retention
372 because it provides a space where learners can actively engage with the content and review the
373 material at their own pace. A learning corner can also help learners apply their knowledge to real-
374 world scenarios and problems, which can increase their confidence and competence. Participants

375 declared the presence of a learning corner that they use to refresh their knowledge and skills in the
376 provision of maternal and newborn care.

377 *“There are times, you may have what we call ‘learning corner’. Sometimes you may not have a*
378 *client and the labour ward is calm no any challenge. Therefore, you go to the small room where*
379 *there is a learning corner there we have manikins ‘Mama Natalie, Mama Besiye, Mama Yu’ and*
380 *other manikins. Therefore, we are able to learn from those manikins at the end of the day you can*
381 *come across a real case or real patient and you can recall what you have gained or this and we*
382 *transfer (the knowledge and skills) to the real patient” (IDI, GRRH)*

383

384 **2. Mentorship and supportive supervision**

385 Participants highlighted that they need to have mentorship and supervision by being followed up to
386 ensure they master the content that has been taught during the training. Mentorship and supportive
387 supervision were reported to help them learn from experienced mentors, receive constructive
388 feedback, and improve their skills and performance. Mentorship and supportive supervision also
389 foster a culture of learning and collaboration, where providers can share their knowledge and insights
390 with each other.

391 *“Mentorship should be done and supervision from our experts from CHMT and when they do*
392 *supervision several times, they bring you back to class and do the required process which is*
393 *needed in the trained issues” (IDI, Kasamwa)*

394

395 **3. Practice**

396 **a. Frequent practice of the procedure**

397 Participants responded that having opportunities for frequent practice of the procedure tends to
398 enhance knowledge and skills retention. Frequent practice of the procedure is a crucial factor for
399 knowledge and skills retention. Frequent practice helps them to consolidate their memory, build

400 confidence and refine their performance, and overcome challenges. Without frequent practice,
401 learners may forget the steps of the procedure, lose confidence, and make errors.

402 *“Okay, the big thing is that we should not get tired of practicing at all! When you do ...you practice*
403 *a lot it builds...builds confidence and ability in a sense that you will not forget because you have*
404 *touched it with your own hands meaning it will remain in your mind what you did...therefore*
405 *every day the weapon is to practice nothing else.” (FGD Muheza)*

406

407 **b. Attending cases following the topics that they have been trained**

408 Participants responded that attending cases following the topics that they have been trained in is one
409 factor that can influence their knowledge retention and performance. By attending cases, they can
410 reinforce their learning, gain confidence and feedback, and adapt to different contexts and scenarios.

411 *“...it helps to remain with knowledge when we meet a case several times. For example, at that*
412 *time we received training on manual removal of the placenta. Within that same week, I went and*
413 *attended the case, and I found that I was able to manage it. Therefore, when we get several cases,*
414 *we keep on learning.” (FGD Tanga)*

415

416 **4. Staff rotation/shifting**

417 Participants have further reported staff rotation or shifting as another factor that positively and
418 negatively influences knowledge and skills retention. Staff rotation or shifting is the practice of
419 assigning healthcare providers to different tasks or positions within a facility. This can have various
420 benefits and challenges for knowledge and skills retention. On the other hand, staff rotation can lead
421 to some risks for knowledge and skills retention, such as loss of hands-on skills, reduced efficiency,
422 and increased training costs (need for re-training).

423 *“...when shifted to another unit loses the skills, because if you do not practice for a long*
424 *time...as time goes on you keep on forgetting (IDI, Katavi)”*

425

426 *“First it is supposed that midwife work in the same maternity unit, and after taking training*
427 *may be of helping the baby to breathe and then you take to the surgical ward or medical ward,*
428 *there you have not helped, therefore after training the first thing is to remain in the maternity*
429 *unit” (IDI, Ilembo)*

430

431 *“Skills, to retain skills is when someone is working in the same unit meaning for example after*
432 *training, then works for at least for some time maybe six months. He/she keeps working in the*
433 *labour unit where training was conducted helps to gain the skills and retain them for a longer*
434 *period. But if after training, she/he is shifted from labour and goes to maybe OPD...when*
435 *she/he goes back the skills will be gone since has not practised them...” (IDI, Kasamwa)*

436

437 **5. Training**

438 Training is another factor for knowledge and skills retention that the participants have described.
439 Training can take various forms, such as retaining on-the-job training, teaching others, self-directed
440 learning through individual revision, receiving updates on evidence based practice etc. Training
441 through these various forms can enhance knowledge and skills retention by increasing the confidence,
442 competence, and commitment of healthcare providers.

443 *“...doing sessions every week and we do on-job training...For example, when we receive an*
444 *emergency PPH case, when you provide the care, you are with fellow providers...and we*
445 *teach each other” (IDI, Buga)*

446

447 *“In my opinion, I think this training should be conducted several times. Therefore, if this*
448 *group receives training after some time you prepare training for another group. It will help*
449 *people at work very much. You know when you receive on-the-job training again it helps a lot*
450 *and builds morale and competence.” (FGD, Tanga).*

451

452 “...because, for you to be able to have more education and stick in your head it is when you
453 teach others. When you keep on teaching others things and things keep on sticking into your
454 head therefore eventually your competence will be improved and people will receive high-
455 quality services.”(FGD, Muheza)

456

457 **6. Availability of equipment**

458 Participants also highlighted the availability of equipment as a factor influencing knowledge and
459 skills retention among healthcare providers. This is because it enables providers to apply their
460 learning in practice and improve their performance. However, the availability of equipment alone is
461 not sufficient for knowledge and skills retention. Receiving adequate training needs to go along with
462 the availability of equipment.

463 “...for example, if you have enough equipment, they also assist you to give good services with
464 the required skills. But if you have scarce equipment also it becomes a challenge to give
465 quality services and remain with the skills...but when you have enough equipment you provide
466 good service properly.” (IDI, Buga).

467

468 **Discussion**

469 This was a mixed methods study aimed to assess factors influencing healthcare providers’ knowledge
470 and skills retention following Helping Mothers Survive and Helping Babies Survive Training.
471 Regarding knowledge, there was a significant drop in the four domains assessed including how to
472 control bleeding after childbirth, managing pre-eclampsia, helping babies to breathe, and essential
473 care for small babies. Similarly, there was a significant drop in the five skills areas assessed including
474 the manual removal of placenta, active management of the third stage of labor, administration of
475 magnesium sulphate, helping babies breathe, and feeding with a nasal gastric tube. A significant drop
476 in specific knowledge and skills domains was higher among healthcare providers with lower
477 professional education and those working in rural areas. Qualitatively, reported facilitators of

478 knowledge and skills retention encompassed the availability of learning materials, frequent practice,
479 mentorship and supportive supervision, on-job training, and staff rotation/shifting.

480

481 The current study demonstrates a statistically significant drop in knowledge and skills scores among
482 healthcare providers following receiving the training on Helping Mothers Survive and Helping Babies
483 Survive, indicating a decrease in knowledge and skills retention over time. A similar situation was
484 observed in Uganda and Sudan where knowledge and skills regarding helping babies breathe among
485 midwives fell by 12 months post-training [39, 40]. On the contrary, a previous study in the country
486 which focused on one similar domain (Helping Mothers Survive Bleeding After Birth) reported that
487 knowledge and simulated basic delivery skills decayed after nine months, while confidence and
488 simulated obstetric emergency skills were largely retained [35]. Moreover, a study in Ethiopia
489 reported an overall strong retention of knowledge and skills among health workers following maternal
490 and newborn health training [41].

491 Various factors have been implicated in the maternal and neonatal care knowledge and skills retention
492 including refresher training [40]. This is supported by a multi-country longitudinal study which
493 showed that, after training, healthcare providers retain knowledge and skills for up to 12 months, the
494 effect which likely enhanced by short repeat skills-training sessions, or, ‘fire drills’ [36]. Our findings
495 from the qualitative arm emphasized the role of on-job training in enhancing essential maternal and
496 newborn care knowledge and skills retention among healthcare providers.

497 Participants working in urban areas had a statistically significant higher drop in mean knowledge for
498 managing pre-eclampsia and a statistically significant lower drop in mean knowledge for helping
499 babies breathe. On the other hand, participants working in rural area had a statistically significant
500 higher drop in mean skills for manual removal of placenta and active management of third stage of
501 labor. This indicates differences experienced in knowledge and skills retention in the two settings. As
502 reported in a previous study [42], there is scarcity of literature on professional education, training or
503 continuous professional development in maternity care in remote and rural settings that can be

504 compared with urban settings. However, the observed differences could be multifaceted and related
505 to several factors, including differences in clinical exposure and experience, resource availability, and
506 institutional practices [43, 44].

507 With regard to the level of professional education, certificate holder had a statistically significant
508 higher drop in mean knowledge for pre-eclampsia compared to holders of diploma and degree. The
509 differences in drop of mean skills was not significant. This implies that knowledge retention drop was
510 much more experienced by the holders of certificate. This is supported by a previous study in
511 Tanzania which found that professional qualification and experience in a maternity care settings are
512 significant factors influencing not only nurses' knowledge, but also skills [45]. The possible
513 explanation is that healthcare workers with lower levels of professional education may have a weaker
514 foundation in medical knowledge and clinical skills, leading to a more rapid decline in retained
515 information [46].

516 Our study found that working in a hospital had a significantly higher drop in pre-eclampsia mean
517 knowledge compared to those working in a dispensary. However, the situation was vice versa for
518 skills for manual removal of placenta where healthcare providers working in dispensaries had a higher
519 drop compared to those working in hospitals. This implies healthcare providers in hospitals and
520 dispensaries had different experiences in various aspects of knowledge and skills retention. A
521 previous multi-country study indicated that health care facility level was not a determinant of
522 knowledge and skills retention among healthcare providers [36]. Various factors affecting each level
523 of health facilities differently such as work experience, training type, training frequency, availability
524 of equipment and facilities, and healthcare providers' educational background may explain the
525 observed differences in knowledge and skills retention [46]. Therefore, this highlights that health
526 facility-specific factors should be considered to enhance healthcare providers' knowledge and skills
527 retention following training.

528 Our findings from the qualitative arm has highlighted the availability of learning materials, frequent
529 practice, mentorship and supportive supervision, on-job training, and staff rotation/shifting as

530 fractionators of knowledge and skills retention among healthcare providers. The identification of
531 these factors emphasize that these are important factors to be considered to enhance knowledge and
532 skills retention among healthcare providers. As discussed above, various studies support these findings
533 [40, 43–46]. This implies that factors influencing knowledge and skills retention are multifaceted in
534 nature and underscores the role of contextual considerations in addressing the phenomenon.
535 There are some strengths and limitations that are noteworthy in this study. Using a mixed design, the
536 study draws its strength from the triangulation of information from both quantitative and qualitative
537 arms. Findings from the qualitative arm were key in complementing the quantitative results.
538 Moreover, the mix of urban and rural settings and various health facility levels provide a good sample
539 representation of the study population. On the other hand, this study might have been influenced by
540 social desirability limitation, particularly with the qualitative arm as healthcare providers might want
541 to provide desirable responses regarding factors influencing the knowledge and skills retention.
542 However, they were assured of confidentiality and encouraged to provide honest responses to mitigate
543 social desirability. However, we believe that this study provides invaluable insights regarding factors
544 associated with knowledge and skills retention and among healthcare providers following the Helping
545 Mothers Survive and Helping Babies Survive Training.

546

547 **Conclusion**

548 This study highlights low level of knowledge and skills retention among healthcare providers
549 following the Helping Mothers Survive and Helping Babies Survive training, with organizational
550 factors being mainly associated. This underscores the need to address organizational barriers such as
551 making available the learning materials, providing mentorship and supportive supervision, and
552 conducting frequent on-job training to enhance knowledge and skills retention regarding maternal
553 and child care among healthcare providers.

554

555 **Acknowledgements**

556 We would like to thank all the HCPs who agreed to take part in the study and their willingness to
557 share their insights regarding factors influencing competence retention. We extend our sincere
558 gratitude to Laerdal Foundation for funding this study. Finally, we thank the assistant researchers
559 who worked diligently in data collection. All regional and district authorities which gave us
560 permission to conduct this study are acknowledged.

561

562 **Author Contributions**

563 BEM, MR, FM and LM conceptualized the study, and were involved in analysis; BEM, MR, LM,
564 and JA performed formal data analysis; BEM, JA, and RG were involved in writing the original draft
565 and BEM, LM, MR, FM, RG, NK, JA contributed in writing – review and editing. Both authors
566 revised and approved the final manuscript.

567

568 **References**

- 569 [1] WHO U, WBG U. Trends in maternal mortality 2000 to 2020: estimates by WHO, UNICEF,
570 UNFPA, World Bank Group and UNDESA/Population Division.
- 571 [2] Lawrence ER, Klein TJ, Beyuo TK. Maternal Mortality in Low and Middle-Income
572 Countries. *Obstet Gynecol Clin North Am* 2022; 49: 713–733.
- 573 [3] Bauserman M, Thorsten VR, Nolen TL, et al. Maternal mortality in six low and lower-middle
574 income countries from 2010 to 2018: risk factors and trends. *Reprod Health* 2020; 17: 173.
- 575 [4] UNICEF. Neonatal mortality, <https://data.unicef.org/topic/child-survival/neonatal-mortality/>
576 (2022, accessed 30 June 2024).
- 577 [5] WHO. Newborn mortality, [https://www.who.int/news-room/fact-sheets/detail/newborn-
578 mortality#:~:text=There are approximately 6500 newborn,to 2.3 million in 2022. \(2024,
579 accessed 30 June 2024\).](https://www.who.int/news-room/fact-sheets/detail/newborn-mortality#:~:text=There are approximately 6500 newborn,to 2.3 million in 2022. (2024,)
- 580 [6] TDHS - MIS. Tanzania Demographic and Health Survey and Malaria Indicator 2022,
581 <https://dhsprogram.com/pubs/pdf/PR144/PPR144.pdf> (2023, accessed 30 June 2024).

- 582 [7] United Nations. Standard Developmental Goals, <https://sdgs.un.org/goals> (2015, accessed 1
583 July 2024).
- 584 [8] Kapologwe NA, Meara JG, Kengia JT, et al. Development and upgrading of public primary
585 healthcare facilities with essential surgical services infrastructure: a strategy towards
586 achieving universal health coverage in Tanzania. *BMC Health Serv Res* 2020; 20: 218.
- 587 [9] Sirili N. Health workforce development post-1990s health sector reforms: the case of medical
588 doctors in Tanzania.
- 589 [10] UNFPA. Boosting the health workforce in Tanzania,
590 <https://tanzania.unfpa.org/en/news/boosting-health-workforce-tanzania> (2020, accessed 30
591 June 2024).
- 592 [11] WHO. New global targets to prevent maternal deaths, [https://www.who.int/news/item/05-10-
593 2021-new-global-targets-to-prevent-maternal-deaths](https://www.who.int/news/item/05-10-2021-new-global-targets-to-prevent-maternal-deaths) (2021).
- 594 [12] Walker T, Woldegiorgis M, Bhowmik J. Utilisation of skilled birth attendant in low-and
595 middle-income countries: trajectories and key sociodemographic factors. *Int J Environ Res
596 Public Health* 2021; 18: 10722.
- 597 [13] TDHS - MIS. 2015-16 Demographic and Health Survey and Malaria Indicator Survey,
598 <https://www.dhsprogram.com/pubs/pdf/SR233/SR233.pdf> (2017, accessed 29 June 2024).
- 599 [14] Shimamoto K, Gipson JD. The relationship of women's status and empowerment with
600 skilled birth attendant use in Senegal and Tanzania. *BMC Pregnancy Childbirth* 2015; 15: 1–
601 11.
- 602 [15] Organization WH. Trends in maternal mortality 2000 to 2017: estimates by WHO, UNICEF,
603 UNFPA, World Bank Group and the United Nations Population Division.
- 604 [16] JHPIEGO. Why Skilled Birth Attendants Matters,
605 <https://www.jnj.com/sites/default/files/pdf/skilled-birth-attendants-infographic.pdf> (2019,
606 accessed 30 June 2024).
- 607 [17] Ersdal HL, Singhal N, Msemu G, et al. Successful implementation of Helping Babies

- 608 Survive and Helping Mothers Survive programs—An Utstein formula for newborn and
609 maternal survival. *PLoS One* 2017; 12: e0178073.
- 610 [18] Msemu G, Massawe A, Mmbando D, et al. Newborn mortality and fresh stillbirth rates in
611 Tanzania after helping babies breathe training. *Pediatrics* 2013; 131: e353–e360.
- 612 [19] Evans CL, Kamunya R, Tibaijuka G. Using helping mothers survive to improve intrapartum
613 care. *Pediatrics* 2020; 146: S218–S222.
- 614 [20] Dol J, Campbell-Yeo M, Bulemela J, et al. Knowledge acquisition after Helping Babies
615 Survive training in rural Tanzania. *Int Health* 2019; 11: 136–142.
- 616 [21] JHPIEGO. Helping Mothers Survive, [https://hms.jhpiego.org/wp-](https://hms.jhpiego.org/wp-content/uploads/2016/08/hms-implementation-guide.pdf)
617 [content/uploads/2016/08/hms-implementation-guide.pdf](https://hms.jhpiego.org/wp-content/uploads/2016/08/hms-implementation-guide.pdf) (2016, accessed 1 July 2024).
- 618 [22] JHPIEGO. Helping Mothers Survive, <https://www.jhpiego.org/hms/> (2015, accessed 1 July
619 2024).
- 620 [23] JHPIEGO, Laerdal Global Health. Helping Mothers and Babies Survive,
621 <https://hmbs.org/about/> (2015, accessed 1 July 2024).
- 622 [24] Ljungblad LW, Sandvik SO, Lyberg A. The impact of skilled birth attendants trained on
623 newborn resuscitation in Tanzania: A literature review. *Int J Africa Nurs Sci* 2019; 11:
624 100168.
- 625 [25] Ameh C, Adegoke A, Hofman J, et al. The impact of emergency obstetric care training in
626 Somaliland, Somalia. *Int J Gynecol Obstet* 2012; 117: 283–287.
- 627 [26] Ameh CA, Mdegela M, White S, et al. The effectiveness of training in emergency obstetric
628 care: a systematic literature review. *Health Policy Plan* 2019; 34: 257–270.
- 629 [27] Al-Beity FA, Pembe A, Hirose A, et al. Effect of the competency-based Helping Mothers
630 Survive Bleeding after Birth (HMS BAB) training on maternal morbidity: a cluster-
631 randomised trial in 20 districts in Tanzania. *BMJ Glob Heal* 2019; 4: e001214.
- 632 [28] Nelissen E, Ersdal H, Mduma E, et al. Clinical performance and patient outcome after
633 simulation-based training in prevention and management of postpartum haemorrhage: an

- 634 educational intervention study in a low-resource setting. *BMC Pregnancy Childbirth* 2017;
635 17: 1–9.
- 636 [29] Evans CL, Bazant E, Atukunda I, et al. Peer-assisted learning after onsite, low-dose, high-
637 frequency training and practice on simulators to prevent and treat postpartum hemorrhage
638 and neonatal asphyxia: a pragmatic trial in 12 districts in Uganda. *PLoS One* 2018; 13:
639 e0207909.
- 640 [30] Goudar SS, Somannavar MS, Clark R, et al. Stillbirth and newborn mortality in India after
641 helping babies breathe training. *Pediatrics* 2013; 131: e344–e352.
- 642 [31] Van den Broek N, Ameh C, Madaj B, et al. Effects of emergency obstetric care training on
643 maternal and perinatal outcomes: a stepped wedge cluster randomised trial in South Africa.
644 *BMJ Glob Heal* 2019; 4: e001670.
- 645 [32] Rowe AK, De Savigny D, Lanata CF, et al. How can we achieve and maintain high-quality
646 performance of health workers in low-resource settings? *Lancet* 2005; 366: 1026–1035.
- 647 [33] Nelissen E, Ersdal H, Østergaard D, et al. Helping mothers survive bleeding after birth: An
648 evaluation of simulation-based training in a low-resource setting. *Acta Obstet Gynecol Scand*
649 2014; 93: 287–295.
- 650 [34] Alwy Al-beity F, Pembe AB, Marrone G, et al. Predictors of change of health workers’
651 knowledge and skills after the Helping Mothers Survive Bleeding after Birth (HMS BAB) in-
652 facility training in Tanzania. *PLoS One* 2020; 15: e0232983.
- 653 [35] Nelissen E, Ersdal H, Mduma E, et al. Helping Mothers Survive Bleeding After Birth:
654 retention of knowledge, skills, and confidence nine months after obstetric simulation-based
655 training. *BMC Pregnancy Childbirth* 2015; 15: 190.
- 656 [36] Ameh CA, White S, Dickinson F, et al. Retention of knowledge and skills after Emergency
657 Obstetric Care training: A multi-country longitudinal study. *PLoS One* 2018; 13: e0203606.
- 658 [37] Ameh CA, Kerr R, Madaj B, et al. Knowledge and Skills of Healthcare Providers in Sub-
659 Saharan Africa and Asia before and after Competency-Based Training in Emergency

- 660 Obstetric and Early Newborn Care. *PLoS One* 2016; 11: e0167270.
- 661 [38] Evans CL, Johnson P, Bazant E, et al. Competency-based training “Helping Mothers
662 Survive: Bleeding after Birth” for providers from central and remote facilities in three
663 countries. *Int J Gynecol Obstet* 2014; 126: 286–290.
- 664 [39] Draiko CV, Yamarat K, Panza A, et al. Knowledge, skills and competency retention among
665 health workers one year after completing helping babies breathe training in South Sudan. *Pan
666 Afr Med J* 2019; 33: 1–13.
- 667 [40] Mildenberger C, Ellis C, Lee K. Neonatal resuscitation training for midwives in Uganda:
668 Strengthening skill and knowledge retention. *Midwifery* 2017; 50: 36–41.
- 669 [41] Gobezaayehu AG, Mohammed H, Dynes MM, et al. Knowledge and Skills Retention Among
670 Frontline Health Workers: Community Maternal and Newborn Health Training in Rural
671 Ethiopia. *J Midwifery Womens Health* 2014; 59: 1–11.
- 672 [42] Ireland J, Bryers H, Van Teijlingen E, et al. Competencies and skills for remote and rural
673 maternity care: a review of the literature. *J Adv Nurs* 2007; 58: 105–115.
- 674 [43] Yang C-W, Yen Z-S, McGowan JE, et al. A systematic review of retention of adult advanced
675 life support knowledge and skills in healthcare providers. *Resuscitation* 2012; 83: 1055–
676 1060.
- 677 [44] Binkhorst M, Coopmans M, Draaisma JMT, et al. Retention of knowledge and skills in
678 pediatric basic life support amongst pediatricians. *Eur J Pediatr* 2018; 177: 1089–1099.
- 679 [45] Angelina JA, Kibusi SM, Mwampagatwa I. Factors influencing nurses’ knowledge and skills
680 in the prevention and management of postpartum haemorrhage. *Afr J Midwifery Womens
681 Health*; 13. Epub ahead of print October 2019. DOI: 10.12968/ajmw.2018.0018.
- 682 [46] Junli A, Isa SNI, Ibrahim FS. Factors of cardiopulmonary resuscitation skills retention
683 among healthcare providers: A scoping review. *Nurse Educ Pract* 2023; 69: 103617.

684

685 **Supporting Information**

686 DATA SET ZIPPED FOLDER