

Cultural Stigma, Psychological Distress and Help-Seeking: Moderating role of Self-esteem and Self-stigma

Toni Sawma¹, Souheil Hallit^{2,3,4,5}, Chloe Joy Younis¹, Jad Jaber¹, and Myriam El Khoury-
Malhame^{1*}

1 Department of Psychology and Education, School of Arts and Sciences, Lebanese American University, Beirut, Lebanon,

2 School of Medicine and Medical Sciences, Holy Spirit University of Kaslik, Lebanon,

3 Research Department, Psychiatric Hospital of the Cross, Jal Eddib, Lebanon

4 Psychology Department, College of Humanities, Effat University, Saudi Arabia

5 Applied Science Research Center, Applied Science Private University, Amman, Jordan

*Corresponding Author

Dr Myriam El Khoury-Malhame, Associate professor of psychology at the Department of Psychology and Education at the Lebanese American University

myriam.malhame@lau.edu.lb

Abstract

Background/Objective: Mental illness is a common and often stigmatized condition. Stigma around mental illness refers to the negative attitudes and beliefs society holds about individuals with mental health conditions, and can sometimes prevent those individuals from seeking adequate therapy. This study aims to explore the intricate relation between stigma, psychological distress, self-confidence and help-seeking behavior among young adults with diagnosed mental illnesses.

Method: A cross-sectional online survey was shared via digital platforms between February 2023 and August 2024. A total of 245 participants from various educational backgrounds and located in different Lebanese regions participated in the study and filled demographics data as well as assessments for stigma (Stigma Scale), self-esteem (Rosenberg Self-Esteem scale), help seeking (Attitudes Towards Seeking Professional Psychological Help -short form (ATSPPH-SF)), distress (Kessler Psychological Distress scale (K6)) and self-stigma (Self-Stigma Questionnaire (SSQ)).

Results: Results from our multivariable analysis revealed a significant association between high levels of stigma and low self-esteem scores, high levels of stigma and high psychological distress, as well as negative attitudes towards seeking professional help. We mostly found that fear of individuals with mental illnesses, prejudice and lack of enough knowledge and awareness were driving reasons for current stigma around mental health in Lebanon.

Conclusion: Stigma remains a pervasive condition leading to more mental health suffering, decreased self-esteem and lower tendencies to seek help. These inaccurate beliefs and stereotypes lead to fear, avoidance and overall discomfort, vis-à-vis people with mental distress perceived as unpredictable, impulsive and dangerous. It is important for public policy makers within collectivist culture to better address these misconceptions and promote help-seeking attitudes.

Keywords: Mental disorders; stigma; Young adults; Lebanon; self-esteem

BACKGROUND

In addition to the challenges of suffering from mental disorder(s), individuals often have to cope with the stigma that emerges from it as mental health stigma remains a widespread social problem impacting people's well-being. According to the World Health Organization, stigma is associated with shame, embarrassment and rejection that causes an individual to be avoided or dismissed from others (WHO, 2019). When a phenomenon is stigmatized, a particular individual or group is labeled based on misconceptions and false stereotypes (Corrigan & Watson, 2004). The stigma associated with a particular condition, such as mental illness, can also result in social exclusion and discrimination (Link & Phelan, 2001). Furthermore, individuals who are subjected to stigma can experience loss of purpose and ambitions, loss of friendships and relationships as well as diminished quality of life and work opportunity threats (Lai et al., 2001). There is a significant body of research on stigma and its impacts, and scholars have investigated the different forms it can take. To begin with, the term social stigma refers to the public beliefs and impressions regarding a specific person or community (Watson et al., 2003). For example, perceiving an individual with schizophrenia as dangerous and crazy is a common illustration of social stigma. This might lead to internalized negative beliefs and perceptions that one who encounters stigma may develop about oneself are referred to as self-stigma (Link & Phelan, 2006). In fact, when individuals internalize societal perceptions, isolate themselves and feel embarrassed and ashamed of their mental illness, they are self-stigmatizing themselves. Finally, structural stigma, which emphasizes the institutional limitations, challenges and prejudice that people who suffer from mental health issues must overcome, especially in institutions and corporations (Pugh et al., 2015). One example of structural stigma might be restricting a person's opportunities in the workplace because of their mental illness.

Among the most unfortunate outcomes of stigma around mental health is the threat of losing one's self-esteem due to the ongoing instilled belief of being a disappointment and achieving nothing to be proud of (Link et al., 2001). According to numerous research (Wright et al., 2000; Link et al., 2001; Hayward et al., 2002), it has been documented that stigmatization has a negative impact on the self-esteem of those who are subjected to it since it leads to losing their sense of worth. This has been systematically reported in people who suffer from mental illness as stigma around their disorders correlates with lower self-esteem scores (Link and Phelan, 2006). In other words, individuals who have been labeled as "mentally ill" are led to believe that they are part of a social group not worthy of love and acceptance, incapable of conforming with others. Their anticipation of being discriminated against even intensifies underlying feelings of shame and guilt which in turn leads them to deny their sufferings, devalue themselves, distort self-perceptions and worsen their self-esteem (Verhaeghe et al., 2008, Corrigan et al., 2006).

This decreased self-esteem due to stigma has been recently investigated in relation to inflating mental distress on one hand and hindering health seeking behaviors on the other (Quinn & Chaudoir, 2009, Gulliver et al. 2010). Global reports highlight extremely limited help-seeking behaviors in those struggling with mental distress with more than 75-80% of people with mental disorders not receiving therapy although approximately 1 in 4 people in Europe have a mental disorder (Clement et al., 2014). This is true even after massive collective traumas whereby only 10% of the people reported seeking mental support although made freely accessible by professionals after an urban disaster (El Khoury-Malhame et al., 2024). Pervasive stigma was reported as major indicator of unwillingness to access or continue mental health care services, in addition to financial considerations (Sirey et al., 2001, Hogan, 2003). This can leave those with

mental diagnoses navigate challenges like relational complications, personal stress and anxiety (Livingston and Boyd, 2010, Vogel et al., 2006), as well as harmful coping mechanisms alone (Major & O'Brien, 2005). Social exclusion in turn decreases tendencies to seek help for stigmatized individuals with mental disorders (Gray, 2002). In fact, lack of social assistance is itself connected to high levels of sadness and anxiety (Cacioppo & Hawkley, 2003), and loneliness deteriorating altogether physical and mental conditions and prospective prognosis (Hawkley & Cacioppo, 2010). As such, individuals struggling with mental disorders additionally have to deal with extra layers of social ostracization and personal anxiety from being subjected to stigma around their mental health.

Minorities and youth are disproportionately deterred by stigma (Clement et al., 2014). It could be that for this vulnerable group, stigma extends beyond societal opinions and impacts one's judgment of one's own self (Villatoro et al., 2022). This key relationship between perceived stigma and personal stigma makes individuals with mental illnesses adopt negative attitudes towards themselves (Maeshima et al., 2022). Stigma causes negative self-perceptions and lower self-esteem levels, further burdening personal suffering inflating internalized negative stigma especially in young adults and adolescents (Ilic et al., 2012, Wood et al., 2017). Furthermore, this negative loop poses a barrier to getting professional psychological help or persevering with a prolonged treatment (Clement et al., 2015).

These widespread stigmatization around mental health seem even more problematic in the global South in general and Arab countries in particular, additionally hindering help-seeking (Fekih et al, 2022). Heightened stigma is partly accounted for by cultural and religious factors in this part of

the world attributing mental health issues to supernatural causes (Ahad et al., 2023), in addition to poverty, lack of education and social shame or familial embarrassment (Jorm et al., 2012). Taken together, this sociocultural belief system can lead to increased discrimination and social exclusion of individuals with mental illnesses. In Lebanon, a small Levantine country, research on the influence of stigma on mental health and help-seeking behavior is scarce, although the country is going through major collective protracted crises. The protracted adversities have taken a toll on Lebanese youth mental health (El Khoury-Malhame et al., 2023), amidst unprecedented sociopolitical instability (Elkhoury-Malhame et al., 2024b). Lebanese with mental health issues reportedly feel stigmatized because of the cultural misconceptions around mental health, which has a detrimental effect on their self-esteem (Rayan and Fawaz, 2018). University students with mental illnesses further express emotions of shame and embarrassment, leading to their marginalization and social isolation (Wehbe, 2011). Stigma around mental health contributes to higher levels of distress among these young adults as they document significant levels of prejudice and greater feelings of despair and anxiety (Dimassi et al., 2015). Karam et al. (2019) highlights that people who suffer from mental disorders in Lebanon face major challenges when it comes to accessing mental health services and seeking psychological treatment because of the stigmatized association of seeking psychological help with signs of weakness, inadequacy and inability to deal with one's own challenges. These constant social marginalization leads to internalized negative attitudes towards seeking assistance in young adults (Hassan, 2015), as they are fear cultural shaming of being called "crazy" (Al Laham et al., 2020).

The purpose of this research is thus to provide empirical data on the influence of stigma on self-esteem and help-seeking behaviors of diagnosed young adults in Lebanon. The objective is to

better guide development of culturally and socially sensitive strategies to address the stigma associated with mental illness, and subsequently alleviate personal struggles, human suffering and socio-economic burden.

METHODS

Ethical Approval

In accordance with the Lebanese American University's research protocol, the Institutional Review Board IRB approved this study (Tracking number LAU.SAS.MM2.8/Mar/2023). Participants electronically signed informed consents before enrolling.

Study Design and Sample

The cross-sectional survey was conducted between March 2023 and August 2024. Participants responded to the survey link shared via social media platforms such as Instagram and WhatsApp. The sample consisted of 245 participants, with 147 women (60%). They ranged mostly between 20 and 29 years with 74.7% having a Bachelor's degree. Participants were located in different Lebanese governorates and also professionally diagnosed with various types of mental illnesses.

Procedure

Only participants who were above 18, living in Lebanon and diagnosed with at least one mental illness were eligible to participate. Participants were expected to complete the survey in 15 to 20 minutes. The number of participant responses that were excluded from the analysis was 14, and the cause of exclusion was due to incomplete survey responses.

Measures

In addition to demographic data, participants were asked to fill the following scales:

The Stigma Scale: King et al. (2007) developed the 28-item stigma scale which is organized into three categories: discrimination, disclosure, and potential positive elements of mental illness. It is five-point Likert scale ranging from 1 (strongly agree) to 5 (strongly disagree), with higher scores indicating greater stigma.

Some items include example of discriminatory conduct such as “I worry about telling people I receive psychological treatment”. This discrimination subscale measures the level of prejudice, exclusion, and discrimination experienced by individuals who suffer from mental illnesses. Moreover, the disclosure subscale measures the degree to which people with mental illnesses reveal their psychological struggles with statements like “People's reactions to my mental health problems make me keep myself to myself”. The final subdivision of the scale measures the extent to which people with mental illnesses are seen as possessing positive traits like creativity and insight. This potential positive element subscale includes questions such as “My mental health problems have made me more accepting of other people”. The scale shows strong internal consistency and test-retest reliability with a Cronbach's alpha of 0.87 for the discrimination subscale, 0.85 and 0.64 for the disclosure and potential positive elements subscales respectively.

The Rosenberg Self-Esteem Scale: (Rosenberg, 1965). This scale consists of 10 items measuring self-esteem. It is a 4-point Linkert scale with response options varying from strongly agree to strongly disagree. Five items are positively worded and the other five are negatively worded. Higher scores suggest higher levels of self-esteem and self-worth. The scale has a Cronbach's alpha of 0.84.

Attitudes Towards Seeking Professional Psychological Help Scale-short form (ATSPPH-SF): (Fischer and Farina, 1995). It assesses social perceptions towards seeking psychological assistance. It includes 10 items on a 4-point Likert scale with statements such as "Personal and emotional troubles, like many things, tend to work out by themselves". Responses range from 0 (disagree) to 3 (agree). Higher score indicates a more optimistic attitude and higher tendencies to ask for psychological aid. The scale had a Cronbach's alpha of 0.84 among an Arab-speaking population (Rayan et al., 2020).

Kessler Psychological Distress Scale (Kessler et al., 2003): A popular 6-item questionnaire used to measure psychological distress. It evaluates how frequently individuals have felt anxious, hopeless, or unworthy over the past 30 days. Each item is graded using a 5-point Likert scale, with the lowest score being 0 (none of the time) and 4 being the highest (all of the time), for a total final score that can range from 0 to 24. Higher scores mean higher levels of psychological distress. The K6's internal consistency and reliability have been found to be high, with alpha coefficient being 0.86 (Ferro et al., 2019).

Self-Stigma Questionnaire SSQ: The 14-item questionnaire was developed to evaluate the degree to which people with mental illnesses internalize stereotypes and misconceptions, leading to self-stigmatization. Each item is evaluated on a 7-point Likert scale, with 1 (strongest agreement) and 7 (strongest disagreement). The scale measures how much individuals have absorbed and instilled unfavorable attitudes and beliefs about themselves and their psychological suffering, such as feelings of inferiority and shame because they have mental disorders. Three aspects of self-stigma are evaluated by the SSQ: internalized stigma, perceived discrimination and stereotypes as well as social functioning, with higher scores indicating lower levels of self-stigma. It has been confirmed

to be reliable and effective across a range of populations and languages. high level of internal consistency, and Cronbach's alpha coefficients ranging from 0.75 to 0.901 (Ochoa et al., 2015).

Statistical Analysis

The SPSS software v.25 was used for the statistical analysis. The attitude help seeking score was considered normally distributed since the skewness (= .001) and kurtosis (= -.702) values varied between -1 and +1 [ref]. The Student t was used to compare two means, the ANOVA test to compare three or more means and the Pearson test to correlate two continuous variables. The moderation analysis was conducted using PROCESS MACRO (an SPSS add-on) v3.4 model 1 [ref], taking the self-esteem, stigma and self-stigma scores as moderators between psychological distress and attitude help seeking. Results were adjusted over all variables that showed a $p < .25$ in the bivariate analysis. $P < .05$ was deemed statistically significant.

RESULTS

Sociodemographic and other characteristics of the sample

Two hundred forty-five young adults with diagnosed mental disorders participated in this study.

Socio-demographics and descriptive statistics are represented in Table 1.

Variable	N (%)
Age (years)	
18-19	16 (6.5%)
20-29	220 (89.8%)
30-39	7 (2.9%)
40 and above	2 (.8%)
Education	
High school	14 (5.7%)
Bachelor	183 (75.0%)
Master	46 (18.9%)
PhD	2 (.8%)

	Mean ± SD
Stigma (mean score)	3.54 ± .81
Self-esteem	24.78 ± 2.37
Attitude help seeking (mean score)	2.26 ± .74
Psychological distress (mean score)	3.06 ± .93
Self-stigma (mean score)	2.93 ± 1.66

Bivariate analysis of factors associated with attitude help seeking

The results showed that older participants and those with higher education compared to the other categories had significantly higher help-seeking attitude (Table 2).

Higher stigma and psychological distress were significantly associated with lower help-seeking attitude, whereas having higher self-esteem and self-stigma were significantly associated with higher attitude help seeking (Table 3).

Variable	Mean ± SD	<i>t</i> / <i>F</i>	<i>df</i> / <i>df1</i> ,<i>df2</i>	<i>p</i>
Age (years)		8.88	3, 241	<.001
18-19	2.84 ± .43			
20-29	2.18 ± .73			
30-39	2.94 ± .57			
40 and above	3.55 ± .64			
Sex		-1.43	243	.155
Male	2.17 ± .67			
Female	2.31 ± .78			
Education		10.82	3, 241	<.001
High school	3.12 ± .64			
Bachelor	2.15 ± .77			
Master	2.37 ± .30			
PhD	3.40 ± .42			

Numbers in bold indicate significant *p* values.

	1	2	3	4	5
1. Attitude help seeking	1				
2. Stigma	-.60***	1			
3. Self-esteem	.48***	-.45***	1		
4. Psychological distress	-.41***	.75***	-.31***	1	
5. Self-stigma	.48***	-.85***	.48***	-.72***	1

p* <.01; *p* <.001

Moderation analysis

The moderation models were adjusted over the following covariates: age, gender and education.

The interaction self-esteem by psychological distress was significantly associated with help-seeking attitude (Beta= .08; $t = 4.07$; $p < .001$; 95% CI .04; .12) (Table 4, Model 1).

At low (Beta = -.44; $p < .001$) and moderate (Beta = -.25; $p < .001$) levels of self-esteem, higher psychological distress was significantly associated with lower help-seeking attitude (Table 5, Model 1).

Also, the interaction self-stigma by psychological distress was significantly associated with help-seeking attitude (Beta= .07; $t = 2.68$; $p = .008$; 95% CI .02; .12) (Table 4, Model 2).

At low (Beta = -.28; $p = .001$) and moderate (Beta = -.17; $p = .011$) levels of self-stigma, higher psychological distress was significantly associated with lower help-seeking attitude (Table 5, Model 2).

Table 4. Moderation analyses taking attitude help seeking as the dependent variable				
	Beta	<i>t</i>	<i>p</i>	95% CI
Model 1: Self-esteem as the moderator				
Psychological distress	-2.19	-4.49	<.001	-3.15; -1.23
Self-esteem	-.11	-1.94	.054	-.23; .002
Interaction psychological distress by self-esteem	.08	4.07	<.001	.04; .12
Model 2: Self-stigma as the moderator				
Psychological distress	-.37	-3.34	.001	-.58; -.15
Self-stigma	-.03	-.44	.658	-.18; .11
Interaction psychological distress by self-stigma	.07	2.68	.008	.02; .12

Numbers in bold indicate significant p values.

Table 5. Conditional effects of the focal predictor (psychological distress) at values of the moderator.				
	Beta	<i>t</i>	<i>p</i>	95% CI
Model 1: Self-esteem as the moderator				
Low (= 22.41)	-.44	-6.19	<.001	-.57; -.30
Moderate (= 24.78)	-.25	-5.52	<.001	-.34; -.16
High (= 27.15)	-.06	-1.10	.273	-.18; .05

Model 2: Self-stigma as the moderator				
Low (= 1.27)	-.28	-3.28	.001	-.45; -.11
Moderate (= 2.93)	-.17	-2.55	.011	-.29; -.04
High (= 4.59)	-.05	-.75	.455	-.19; .08

Numbers in bold indicate significant p values.

DISCUSSION

The impact of stigma on mental health is a major public health issue with cultural sensitivities and is best understood when including individuals with diagnosable disorders. In our study, stigma around mental illness as shown to deter young adults' mental distress and further defer them from seeking recovery in Lebanon. The relationship between stigma, psychological distress and help-seeking intentions is further complexified and influenced by both self-esteem and self-stigma, that are shown to play crucial mediating roles respectively.

Help-Seeking in relation to Stigma and Psychological Distress

Our findings first and foremost support accumulating evidence that stigma very significantly correlates with increased psychological distress, self-stigma and deteriorated self-esteem while impairing help-seeking. This is in line with recent data conducted in the same context among Lebanese individuals with mental illnesses, whereby participants reported lower levels of self-esteem and quality of life, as well as higher levels of social isolation when subjected to stigma (El Hayek et al., 2021). Our results further echo the extent to which stigma inflates feelings of humiliation and embarrassment, leading to negative self-perceptions. Heightened levels of stigmatization can increase psychological distress symptoms of anxiety, depression (Burgess et al., 2019) all while increasing self-stigma and loss of self-worth, leading to hopelessness and helplessness (Corrigan and Watson, 2002). Individuals become even less likely to seek

professional help (Vogel et al., 2006), even when mental health services are accessible and affordable (Sirey et al., 2001). Accumulating evidence support the negative impact of self-stigma as a barrier to seeking mental health services in young adults already suffering from mental illness (Schomerus & Angermeyer, 2008, Jennings et al. 2015). They indeed seem to synchronize their disbelief with their own competences as they internalize negative misbeliefs and misconceptions with the societal pervasive stigma associated with their mental illness. This reinforces marginalization and social isolation and perpetuates prevalent stigma and discrimination (Link & Phelan, 2001). It also solidifies barriers to seeking mental health recovery in occidental (Yanos et al., 2010) as well as oriental societies (Obeid et al., 2019) as people suffering from mental illnesses fear of prejudice and discrimination of seeking treatment. In the Lebanese context, self-stigma is still a prevalent challenge and mental health is still frowned upon (Dalky, 2012).

Individuals' confidence in getting the proper care might be additionally hindered by cultural assumptions that psychological disorders are personal failings rather than conditions that need to be recognized and treated (Karam et al., 2019, Krstanoska-Blazeska & Slewa-Younan 2021). Education and older age seem to play protective factors as we found that older adults in the sample and those with highest degrees had considerably more positive attitudes towards help-seeking behavior. Charara et al. (2017) had previously demonstrated that lack of education and awareness contributed to limited understanding of mental health and to subsequent stigma. Highlighting and targeting such social and personal stigmas could thus be considered as a public health priority since studies suggest that more favorable attitudes on seeking mental health professional help are linked to improved levels of self-esteem and restituted social functioning (Vogel et al., 2006).

Help-Seeking in relation to Self-Esteem

Our results suggest that individuals with low to moderate self-esteem, who are already vulnerable to psychological distress, are less likely to seek help for their mental health concerns. This negative correlation between self-esteem and help-seeking attitudes highlights the detrimental role of stigma in exacerbating mental health difficulties. Individuals with low self-esteem may internalize societal stigma, leading to feelings of shame, guilt, and inadequacy, which can further hinder their willingness to seek professional help. In fact, individuals with high levels of self-esteem and self-worth generally describe help-seeking as empowering and beneficial rather than a sign of weakness and vulnerability (Lannin et al., 2015). University students with higher self-esteem seem to be keen on promptly recognizing and addressing issues related to mental health for better outcomes and faster recovery (Yih, 2020). These results underscore the importance of addressing stigma and promoting self-esteem to encourage help-seeking behaviors among young adults with mental disorders.

Help-Seeking in relation to Self-Stigma

This study also illustrates the role self-stigma in moderating the relationship between psychological distress and help-seeking intentions. Self-stigma refers to students' internalization of cultural mental health stigmas and feel like a disgrace and disappointment to their families. Adopt negative perceptions and stereotypes, makes feelings of worthlessness, shame and guilt emerge, reinforcing unfavorable perceptions toward mental illnesses (Cheng, 2019). The interaction effect between self-stigma and psychological distress further underscores the complexity of help-seeking behaviors. Specifically, individuals with low to moderate levels of self-stigma who also experience high psychological distress are less likely to seek help. This finding suggests that self-stigma can amplify the negative impact of psychological distress on help-

seeking attitudes. As self-stigma increases, individuals may become more reluctant to acknowledge their mental health problems, fear negative judgment, and avoid seeking necessary support. Within this framework, self-esteem may serve as a protective factor, mitigating the negative effects of both self-stigma and psychological distress on help-seeking intentions. Individuals with higher self-esteem may be more resilient to the negative impacts of stigma and distress, making them more likely to seek help. These positive attitudes seem better supported with the positive assistance of healthcare providers (Abi Hana et al., 2022).

Limitations and Future Research

Increasing the sample size could benefit in more adequate generalization of the findings. Qualitative reports that specifically target the Lebanese cultural context could provide additional support to some sensitive aspect of stigma and would better encompass the range of diversity and experiences within different Lebanese communities related to gender, religious or urban versus rural settings. Conducting a longitudinal design and including therapy outcome could also track potential fluctuations in stigma, self-stigma, self-esteem, psychological distress and help-seeking tendencies across several mental conditions. Lastly, it could be useful to track the psychiatric diagnoses of participants to overcome some social desirability biases when self-reporting mental disorders.

Conclusion

In conclusion, this study sheds light on the significance of mental health stigma on psychological distress and help-seeking. Findings highlight the pivotal role of self-esteem and self-stigma in moderating the above intricate relations in a collectivist crisis enriched context. This builds on top

of recommendations from the Lancet Commission on ending stigma and discrimination in mental health (2022). If policy makers capitalize on positive stigma awareness and incentivize engagement in proactive behaviors towards destigmatizing and treating mental illness this could increase help-seeking attitudes and behaviors, with favorable outcomes on mental and physical health.

Declarations.

Human Ethics approval and Consent to participate declaration. In accordance with the Declaration of Helsinki, all participants approved and signed the informed consent as per the IRB ethics committee approval and tracking number LAU.SAS.MM2.8/Mar/2023

Availability of data and materials. The datasets used and/or analyzes during the current study are available from the corresponding author on reasonable request.

Competing interests. The authors declare that they have no competing interests

Funding. Not Applicable

Authors' contributions. M.EK and T.S designed the study and wrote the manuscript. C. Y and J.J collected data and drafted part of the main manuscript and S.H prepared the analyses and corresponding tables. All authors reviewed the manuscript.

Acknowledgements. The authors wish to acknowledge all those who made the project possible.

References

Abi Doumit, C., Haddad, C., Sacre, H., Salameh, P., Akel, M., Obeid, S., ... & Soufia, M. (2019). Knowledge, attitude and behaviors towards patients with mental illness: Results from a national Lebanese study. *PloS one*, *14*(9), e0222172.

Abi Hana, R., Arnous, M., Heim, E., Aeschlimann, A., Koschorke, M., Hamadeh, R. S., Thornicroft, G., Kohrt, B. A., Sijbrandij, M., Cuijpers, P., & El-Chammy, R. (2022). Mental health stigma at primary health care centres in Lebanon: qualitative study. *International journal of mental health systems*, *16*(1), 23. <https://doi.org/10.1186>

Ahad AA, Sanchez-Gonzalez M, Junquera P. Understanding and Addressing Mental Health Stigma Across Cultures for Improving Psychiatric Care: A Narrative Review. *Cureus*. 2023 May 26;15(5):e39549. doi: 10.7759/cureus.39549

Al Laham, D., Ali, E., Mousally, K., Nahas, N., Alameddine, A., & Venables, E. (2020). Perceptions and health-seeking behaviour for mental illness among Syrian refugees and Lebanese community members in Wadi Khaled, North Lebanon: a qualitative study. *Community Mental Health Journal*, *56*, 875-884.

Burgess, A. M., Molina, K. M., Bhandari, K., & Dibartolo, P. M. (2018). Stigma consciousness and evaluative concerns: A pathway to psychological distress for academically stigmatized students. *Journal of Social and Clinical Psychology*, *37*(4), 275-296.

Cacioppo, J. T., & Hawkley, L. C. (2003). Social isolation and health, with an emphasis on underlying mechanisms. *Perspectives in biology and medicine*, *46*(3), S39-S52.

Charara, Raghid, Mohammad Forouzanfar, Mohsen Naghavi, Maziar Moradi-Lakeh, Ashkan Afshin, Theo Vos, Farah Daoud et al. "The burden of mental disorders in the eastern Mediterranean region, 1990-2013." *PloS one* 12, no. 1 (2017): e0169575.

Cheng, C. M., Chang, C. C., Wang, J. D., Chang, K. C., Ting, S. Y., & Lin, C. Y. (2019). Negative impacts of self-stigma on the quality of life of patients in methadone maintenance treatment: The mediated roles of psychological distress and social functioning. *International journal of environmental research and public health*, 16(7), 1299.

Clement, S., Schauman, O., Graham, T., Maggioni, F., Evans-Lacko, S., Bezborodovs, N., ... & Thornicroft, G. (2015). What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychological medicine*, 45(1), 11-27.

Corrigan, W. Barr (2006) Corrigan, PW, Watson, AC, & Barr, L.(2006). The self-stigma of mental illness: Implications for self-esteem and self-efficacy. *Journal of Social and Clinical Psychology*, 25(9), 875-884.

Corrigan, P. W., & Watson, A. C. (2004). At issue: Stop the stigma: Call mental illness a brain disease. *Schizophrenia bulletin*, 30(3), 477-479.

Corrigan, P. W., & Watson, A. C. (2002). Understanding the impact of stigma on people with mental illness. *World psychiatry*, 1(1), 16.

Dalky, H. F. (2012). Perception and coping with stigma of mental illness: Arab families' perspectives. *Issues in mental health nursing*, 33(7), 486-491.

Dimassi, H., Karam, E. G., Mneimneh, Z. N., Fayyad, J. A., Karam, A. N., Nasser, S. C., & Kessler, R. C. (2015). Lifetime prevalence of mental disorders in Lebanon.

Elhai JD, Schweinle W, Anderson SM. Reliability and validity of the Attitudes Toward Seeking Professional Psychological Help Scale-Short Form. *Psychiatry Res.* 2008 Jun 30;159(3):320-9.

doi: 10.1016/j.psychres.2007.04.020. Epub 2008 Apr 22. PMID:

El Hayek S, Cherro M, El Alayli A, El Hayek R, Ghossoub E. Stigma Towards Mental Illness at an Academic Tertiary Care Center in Lebanon. *Acad Psychiatry.* 2021 Dec;45(6):725-732. doi:

10.1007/s40596-021-01511-1. Epub 2021 Jul 20. PMID: 34286455; PMCID:

PMC8294272.18433879.

El Khoury-Malhame M., Sfeir M, Hallit S, Sawma T. (2023). Factors associated with posttraumatic growth: gratitude, PTSD and distress; one year into the COVID-19 pandemic in Lebanon. *Curr Psychol.* 2023;1-10. doi:10.1007/s12144-022-04159-8

El Khoury-Malhame M., Harajli A. D., Reykowska D.,and Ohme R. (2024) Can the Phoenix Still Rise? Traumatic effect of Beirut port explosion on people’s experiences. *Psychological Trauma: Theory, Research, Practice, and Policy.* 10.1037/tra0001744. Advance online publication. <https://doi.org/10.1037/tra0001744>

El Khoury-Malhame M., Bou Malhab S., Chaaya R., Sfeir M. and El Khoury S. (2024b). Coping during socio-political Uncertainty. *Frontiers in Psychiatry.* 14: 1267603

Fekih-Romdhane, F., Jahrami, H., Stambouli, M., Alhuwailah, A., Helmy, M., Shuwiekh, H. A. M., ... & Cheour, M. (2022). Cross-cultural comparison of mental illness stigma and help-seeking attitudes: a multinational population-based study from 16 Arab countries and 10,036 individuals. *Social Psychiatry and Psychiatric Epidemiology*, 1-16.

Ferro, M. A. (2019). The psychometric properties of the Kessler Psychological Distress Scale (K6) in an epidemiological sample of Canadian youth. *The Canadian Journal of Psychiatry*, 64(9), 647-657.

Fischer E. H., Farina A. (1995). Attitudes toward seeking professional psychological help: a shortened form and considerations for research. *J. Coll. Stud. Dev.* 36, 368–373.

Gray, A. J. (2002). Stigma in psychiatry. *Journal of the royal society of medicine*, 95(2), 72-76.

Gulliver, A., Griffiths, K. M., & Christensen, H. (2010). Perceived barriers and facilitators to mental health help-seeking in young people: a systematic review. *BMC psychiatry*, 10(1), 1-9.

Hassan, F. Y. (2015). Attitudes toward help-seeking behavior in a sample of university students in Lebanon (Doctoral dissertation).

Hastings, M., Northman, L., & Tangney, J. (2002). Shame, guilt, and suicide. *Suicide science*, 67-79.

Hayward, P., Wong, G., Bright, J. A., & Lam, D. (2002). Stigma and self-esteem in manic depression: an exploratory study. *Journal of affective disorders*, 69(1-3), 61-67.

Hawkey, L. C., & Cacioppo, J. T. (2010). Loneliness matters: A theoretical and empirical review of consequences and mechanisms. *Annals of behavioral medicine*, 40(2), 218-227.

Hogan, M. F. (2003). New Freedom Commission report: The president's New Freedom Commission: recommendations to transform mental health care in America. *Psychiatric Services*, 54(11), 1467-1474.

Ilic, M., Reinecke, J., Bohner, G., Hans-Onno, R., Beblo, T., Driessen, M., ... & Corrigan, P. W. (2012). Protecting self-esteem from stigma: A test of different strategies for coping with the stigma of mental illness. *International Journal of Social Psychiatry*, 58(3), 246-257.

Jennings, K. S., Cheung, J. H., Britt, T. W., Goguen, K. N., Jeffirs, S. M., Peasley, A. L., & Lee, A. C. (2015). How are perceived stigma, self-stigma, and self-reliance related to treatment-seeking? A three-path model. *Psychiatric rehabilitation journal*, 38(2), 109.

Jorm, A. F. (2012). Mental health literacy: Empowering the community to take action for better mental health. *Am. Psychol.* 67, 231–243. doi: 10.1037/a0025957

Karam, E. G., Karam, G. E., Farhat, C., Itani, L., Fayyad, J., Karam, A. N., ... & Thornicroft, G. (2019). Determinants of treatment of mental disorders in Lebanon: barriers to treatment and changing patterns of service use. *Epidemiology and psychiatric sciences*, 28(6), 655-661.

Kessler, R. C., Andrews, G., Colpe, L. J., Hiripi, E., Mroczek, D. K., Normand, S. L., Walters, E.E., & Zaslavsky, A. M. (2002). Short screening scales to monitor population prevalences and trends in non-specific psychological distress. *Psychological medicine*, 32(6), 959-976.

King, M., Dinos, S., Shaw, J., Watson, R., Stevens, S., Passetti, F., ... & Serfaty, M. (2007). The Stigma Scale: development of a standardised measure of the stigma of mental illness. *The British Journal of Psychiatry*, 190(3), 248-254.

Krstanoska-Blazeska, K., Thomson, R., & Slewa-Younan, S. (2021). Mental illness stigma and associated factors among Arabic-speaking religious and community leaders. *International Journal of Environmental Research and Public Health*, 18(15), 7991.

Lai, Y. M., Hong, C. P. H., & Chee, C. Y. (2001). Stigma of mental illness. *Singapore medical journal*, 42(3), 111-114.

Lannin, D. G., Vogel, D. L., Brenner, R. E., & Tucker, J. R. (2015). Predicting self-esteem and intentions to seek counseling: The internalized stigma model. *The Counseling Psychologist*, 43(1), 64-93.

Link, B. G., & Phelan, J. C. (2006). Stigma and its public health implications. *The Lancet*, 367(9509), 528-529.

Link, B. G., & Phelan, J. C. (2001). Variations in the Definition of Stigma & Challenges to the Stigma Concept. *Conceptualizing stigma. annual Review of Sociology*, 27, 363-385.

Link, B. G., Struening, E. L., Neese-Todd, S., Asmussen, S., & Phelan, J. C. (2001). Stigma as a barrier to recovery: The consequences of stigma for the self-esteem of people with mental illnesses. *Psychiatric services*, 52(12), 1621-1626.

Livingston, J. D., & Boyd, J. E. (2010). Correlates and consequences of internalized stigma for people living with mental illness: A systematic review and meta-analysis. *Social science & medicine*, 71(12), 2150-2161.

Maeshima, L. S., & Parent, M. C. (2022). Mental health stigma and professional help-seeking behaviors among Asian American and Asian international students. *Journal of American college health : J of ACH*, 70(6), 1761–1767. <https://doi.org/10.1080/07448481.2020.1819820>

Major, B., & O'brien, L. T. (2005). The social psychology of stigma. *Annu. Rev. Psychol.*, 56, 393-421.

- Merhej, R. (2019). Stigma on mental illness in the Arab world: beyond the socio-cultural barriers. *International Journal of Human Rights in Healthcare*.
- Meier, A., Csiernik, R., Warner, L., & Forchuk, C. (2015). The stigma scale: A Canadian perspective. *Social Work Research*, 39(4), 213-222.
- Obeid, S., Haddad, C., Zakhour, M., Fares, K., Akel, M., Salameh, P., & Hallit, S. (2019). Correlates of self-esteem among the Lebanese population: a cross-sectional study. *Psychiatria Danubina*, 31(4), 429-439.
- Ochoa, S., Martínez-Zambrano, F., Garcia-Franco, M., Vilamala, S., Ribas, M., Arenas, O., ... & Haro, J. M. (2015). Development and validation of the Self-Stigma Questionnaire (SSQ) for people with schizophrenia and its relation to social functioning. *Comprehensive psychiatry*, 62, 93-99.
- Pugh, T., Hatzenbuehler, M., & Link, B. (2015). Structural stigma and mental illness. Commissioned Paper for Committee on the Science of Changing Behavioral Health Social Norms, Mailman School of Public, Columbia University (August 2015).
- Quinn, D. M., & Chaudoir, S. R. (2009). Living with a concealable stigmatized identity: the impact of anticipated stigma, centrality, salience, and cultural stigma on psychological distress and health. *Journal of personality and social psychology*, 97(4), 634.
- Rayan, A., & Fawaz, M. (2018). Cultural misconceptions and public stigma against mental illness among Lebanese university students. *Perspectives in psychiatric care*, 54(2), 258-265.

Rayan, A., Baker, O., & Fawaz, M. (2020b, August 11). The psychometric properties of attitudes toward seeking professional psychological help scale-short form in Jordanian University students. *Journal of Muslim Mental Health*.

Rosenberg, M. (1965). Rosenberg self-esteem scale (RSE). Acceptance and commitment therapy. *Measures package*, 61(52), 18.

Sirey, J. A., Bruce, M. L., Alexopoulos, G. S., Perlick, D. A., Raue, P., Friedman, S. J., & Meyers, B. S. (2001). Perceived stigma as a predictor of treatment discontinuation in young and older outpatients with depression. *American Journal of psychiatry*, 158(3), 479-481.

Sirey, J. A., Bruce, M. L., Alexopoulos, G. S., Perlick, D. A., Friedman, S. J., & Meyers, B. S. (2001). Stigma as a barrier to recovery: Perceived stigma and patient-rated severity of illness as predictors of antidepressant drug adherence. *Psychiatric services*, 52(12), 1615-1620.

The Lancet (2022). Can we end stigma and discrimination in mental health?. *Lancet (London, England)*, 400(10361), 1381. [https://doi.org/10.1016/S0140-6736\(22\)01937-7](https://doi.org/10.1016/S0140-6736(22)01937-7)

Tinakon, W., & Nahathai, W. (2012). A comparison of reliability and construct validity between the original and revised versions of the Rosenberg Self-Esteem Scale. *Psychiatry investigation*, 9(1), 54.

Verhaeghe, M., Bracke, P., & Bruynooghe, K. (2008). Stigmatization and self-esteem of persons in recovery from mental illness: the role of peer support. *International Journal of Social Psychiatry*, 54(3), 206-218.

Villatoro, A. P., DuPont-Reyes, M. J., Phelan, J. C., & Link, B. G. (2022). 'Me' vs. 'Them': How Mental Illness Stigma Influences Adolescent Help-Seeking Behaviors for Oneself and

Recommendations for Peers. *Stigma and health*, 7(3), 300–310.

<https://doi.org/10.1037/sah0000392>

Vogel, D. L., Wade, N. G., & Haake, S. (2006). Measuring the self-stigma associated with seeking psychological help. *Journal of counseling psychology*, 53(3), 325.

Watson, A. C., Ottati, V., & Corrigan, P. (2003). From whence comes mental illness stigma?. *International Journal of Social Psychiatry*, 49(2), 142-157.

Wehbe, H. (2011). Predictors of Attitudes towards Psychological Counseling among Lebanese College Students. *Graduate Student Journal of Psychology*, 13, 25-29.

World Health Organization. (2019). *Advocacy for mental health, disability and human rights: WHO Quality Rights guidance module*.

Wood, L., Byrne, R., Burke, E., Enache, G., & Morrison, A. P. (2017). The impact of stigma on emotional distress and recovery from psychosis: The mediatory role of internalized shame and self-esteem. *Psychiatry Research*, 255, 94-100.

Wright, E. R., Gronfein, W. P., & Owens, T. J. (2000). Deinstitutionalization, social rejection, and the self-esteem of former mental patients. *Journal of health and social behavior*, 68-90.

Yanos, P. T., Roe, D., & Lysaker, P. H. (2010). The impact of illness identity on recovery from severe mental illness. *American journal of psychiatric rehabilitation*, 13(2), 73-93.

Yih, I. (2020). *Individual and Group Differences in Self-Concept and Help-Seeking Behavior* (Doctoral dissertation, Wellesley College).