

1 **[Title]**

2 **Exploring the acceptability and adherence of the integrated MAMI care**
3 **pathway for supporting vulnerable infants and their mothers in South**
4 **Sudan: a qualitative study**

5
6 **[Authors]**

7 Hedwig Deconinck^{1,2}
8 Marlene Traore Hebie²
9 Angelina Nasira Boi³
10 Kirk Dearden²

11
12 **[Authors affiliation]**

13 ¹Institute of Health and Society, Université catholique de Louvain, Brussels, Belgium.

14 ²MOMENTUM Integrated Health Resilience (MIHR), Corus International, USA

15 ³MOMENTUM Integrated Health Resilience (MIHR), IMA World Health (IMA), Juba, South Sudan, affiliate
16 of Corus International, Washington DC, USA

17
18 **[Authors position in case needed]**

19 Hedwig Deconinck, MAMI Consultant MOMENTUM Integrated Health Resilience Institute of Health and
20 Society, UCLouvain, Belgium

21 Marlene Traore Hebie, Senior Technical Advisor, Nutrition, MOMENTUM Integrated Health Resilience,
22 USA

23 Angelina Nasira Boi, MAMI Coordinator, MOMENTUM Integrated Health Resilience, South Sudan

24 Kirk Dearden, Senior Technical Lead, Nutrition and WASH, MOMENTUM Integrated Health Resilience,
25 USA

26
27 **[Corresponding author]**

28 Kirk Dearden, kdearden@momentumihhr.org

29
30 **[Key terms]**

31 Infant health, infant nutrition, maternal and child health, care pathway, South Sudan.

32
33 **ABSTRACT [WORD COUNT 300]**

34 **Background:** One in four infants is born too small or too early, increasing the risk of poor growth,
35 development, ill health, and death. This risk is exacerbated by recurring conflict and climatic shocks. The
36 management of small and nutritionally at-risk infants under 6 months and their mothers (MAMI) care
37 pathway was integrated into primary healthcare to ensure continuous mother-infant-centered care in
38 South Sudan. This study explored the acceptability of the MAMI approach for service providers and users
39 and the factors influencing their adherence.

40 **Methods:** A qualitative study was embedded in a mixed-methods study following 521 vulnerable mother-
41 infant pairs. MAMI care included sensitization and screening in the community and health facility, in-depth
42 assessment, and risk-based care addressing physical, mental and nutritional vulnerability of infants and
43 mothers. Semi-structured interviews of 20 health workers and 30 mothers explored acceptability and
44 adherence to MAMI care provision and use.

45 **Results:** Themes of perceived acceptability of MAMI care were mapped across coherence, self-efficacy,
46 affective attitude, burden, ethicality, opportunity cost, and perceived effectiveness. Health workers were
47 most apprehensive about the additional workload and time required to implement the care pathway.
48 Mothers expressed concern over the numerous health visits far from their homes and not having
49 supportive family members. Factors promoting adherence to MAMI care provision and use were mapped
50 across capability, opportunity, and motivation. Health workers' self-confidence was boosted by the new

Exploring the acceptability and adherence of the integrated MAMI care pathway for supporting vulnerable
infants and their mothers: a qualitative study

51 knowledge and skills they acquired through training and mentorship. Mothers reported learning practical
52 ways to care for their infants independently and to witness the positive impact on their infants' thriving.

53 **Conclusion:** MAMI care was accepted for addressing the needs of a previously overlooked vulnerable
54 group. To ensure effective adoption, delivery and use, challenges must be resolved. Adapting the MAMI
55 care pathway to the local health system context will encourage adherence from both providers and users.

56 **KEY MESSAGES OF THE ARTICLE**

- 57 • **What is already known on this topic** – *The 2023 WHO updated guideline on preventing and*
58 *treating wasting in children under 5 includes care for infants under 6 months at risk of poor growth*
59 *and development and their mothers within primary healthcare. Understanding and effectively*
60 *delivering care to this previously overlooked vulnerable group is critical for national policymakers*
61 *and implementers to address persistently high levels of infant mortality.*
- 62 • **What this study adds** – *This study explored the acceptability of the MAMI care pathway—a risk-*
63 *based, mother-infant centered care approach—among service providers and users in South*
64 *Sudan, as well as the factors influencing their adherence. It offers recommendations for*
65 *effectively adapting the approach to the local health system context to promote adherence from*
66 *both providers and users.*
- 67 • **How this study might affect research, practice or policy** – *The study generated valuable*
68 *insights and learning at both national and global levels to enhance policies and practices. It also*
69 *laid the foundation for future research and intervention development, particularly in settings with*
70 *similar demographics and healthcare challenges.*

71 **MAIN TEXT [WORD COUNT 5000]**

72 Globally, an estimated one in four infants under 6 months old (u6m) is small or nutritionally at risk, leading
73 to poor growth and development, ill health and death (1, 2). Although deaths in children under 5 years old
74 have declined in South Sudan, children with wasting have a nine times higher risk of death than healthy
75 children, overburdening the health system (3).

76 National maternal, neonatal and child health and nutrition policies address vulnerability in infants u6m and
77 their mothers using various approaches. Established primary care practices for infants u6m include
78 vaccinations at birth and at 6, 10 and 14 weeks, with over 70 percent coverage against common
79 childhood illnesses (6), and health seeking for ill infants. Maternal care and support are fragmented
80 across health units, and there is a lack of psychosocial and mental health services, feeding and care
81 counselling and early childhood development (ECD) (4). Moreover, irregular and inadequately covered
82 technical and financial support hinders effective health service delivery (4, 5).

83 To address care challenges for vulnerable mother-infant pairs, Momentum Integrated Health Resilience
84 (MIHR) partnered with the South Sudan Ministry of Health (MOH) in 2022 to adapt and field test the
85 management of small and nutritionally at-risk infants u6m and their mothers (MAMI) integrated care
86 pathway (7, 8). The MAMI care pathway (henceforth MAMI care) delivered in primary care consisted of
87 risk screening in both communities and health facilities, in-depth assessment and risk classification and
88 targeted risk-based care and counselling for physical and mental health and nutrition of both infants and
89 mothers (12, 13). The exercise aimed to generate national and global learning to improve policies and
90 practices.

91 During implementation of MAMI care, challenges and outcomes were noted by the study team and
92 discussed but not always fully understood. This study explored the acceptability of MAMI care to both
93 health workers and mothers to understand what worked for whom under what circumstances and to
94 identify factors that facilitated or hindered adherence to care by service providers and users (9).

95 **METHODS**

96 **Qualitative approach and research paradigm.** The theory-driven qualitative study used Sekhon's
97 theoretical framework of acceptability (11) and Michie's Capability, Opportunity and Motivation Behavior
98 Model (COM-B) (9) to explore acceptability and understand adherence behavior of key informants in
99 semi-structured interviews.

100 **Context.** The study was conducted in February–March 2024 in five sites in Central Equatoria, Jonglei,
101 Western Bahr el-Ghazal and Western Equatoria states in South Sudan, nested in a longitudinal multi-
102 methods study of 521 vulnerable mother-infant pairs enrolled and supported between October 2022 and
103 December 2023 (15 months) in the MAMI CP until the infants reached 6 months of age. Each site had an
104 MOH-run primary healthcare center (PHCC), a referral hospital and a County Health Department (CHD)
105 for oversight.

106 **Researcher characteristics.** Two international study investigators supported a local lead interviewer, an
107 assistant and a boma (community) health worker (BHW) as translator with proficiency in English and local
108 languages.

109 **Sampling strategy and units of study.** Fifty key informants—20 MAMI care providers and 30 MAMI
110 care users—participated in semi-structured interviews. Care providers included five MAMI assistants who
111 were previously MIHR staff and supported one of the five MAMI implementation sites, five clinical officers
112 who were MOH staff and integrated MAMI care as part of their regular consultation duties and 10 BHWs
113 appointed by their communities and supported by MIHR to integrate MAMI care in the catchment areas of
114 the MAMI sites included in the study. Gurei and Pariak sites were purposively selected to represent an
115 urban and a rural context. The clinical officers were purposively selected based on their involvement in
116 MAMI care. The BHWs were randomly selected from a list of eighteen MIHR-supported BHWs. Thirty
117 mothers were randomly selected from the MAMI individual care database and stratified in three
118 adherence categories: Group 1 attended all follow-up visits until exit, group 2 returned at least once for a
119 follow-up visit, and group 3 discontinued care immediately after enrolment. All participants were informed
120 of the purpose, date and duration of the interview and received money for transport to the interview site.
121 The number of key informants was sufficient to obtain variation and reach saturation. All care providers

Exploring the acceptability and adherence of the integrated MAMI care pathway for supporting vulnerable
infants and their mothers: a qualitative study

122 had received initial training and ongoing mentoring on the MAMI care before implementation, and all
123 mothers had agreed to enroll in the care pathway after providing informed consent.

124 **Ethical approval.** This study is part of the overall MAMI implementation study protocol that received
125 ethical approval from the Institutional Review Board (IRB) of Momentum-JSI's and the Ministry of Health
126 of South Sudan in March 2022.

127 **Data collection methods, instruments and technologies.** Interviews were conducted face-to-face
128 whenever possible, and due to travel issues, some were conducted online using Microsoft Teams. Both
129 study investigators participated in all interviews online. In-person interviews with health workers and
130 mothers took place at PHCC sites. Interviews followed semi-structured interview guides and lasted on
131 average 90 minutes. Discussions were held in English, Arabic or the local language with simultaneous
132 translation.

133 **Data processing and analysis.** The study team debriefed after each interview on key points and clarified
134 inconsistencies in information. Interviews and discussions were recorded, transcribed using [otter.ai](#) and
135 summarized by theoretical constructs in Excel. Demographic and care information was retrieved from
136 interviews and study databases. Recurrent themes were coded and analyzed from care providers' and
137 users' perspectives. Two study investigators discussed coding consistency across the interview data.
138 The acceptability framework included seven constructs: coherence, self-efficacy, affective attitude,
139 burden, ethicality, opportunity cost and perceived effectiveness. Adherence behavior factors were
140 mapped using the COM-B model's capability, opportunity and motivation.

141 **Techniques to enhance trustworthiness.** Data were triangulated and iterated for consistency of
142 interpretation of discussions within and across themes, providing reliable and valid insights. Research
143 processes and decisions were recorded in detail for study transparency and reproducibility. Reflective
144 journaling recorded biases, alterations and decisions made during the study to improve confirmability and
145 identify and document recurring and deviant patterns. The structure of this manuscript follows the
146 Standards for Reporting Qualitative Research (SRQR) checklist (10).

147 RESULTS

148 Characteristics of the MAMI care providers and users

149 Five of the 20 participating health workers were MIHR-employed MAMI assistants, and five were MOH-
150 employed clinical officers (two of the five were officers-in-charge of the PHCCs). Each MAMI assistant and
151 clinical officer represented one of the five MIHR-supported MAMI implementation sites. All clinical officers
152 involved in MAMI care had clinical nurse education and four were male. Ten community-appointed BHWs
153 receiving incentives from MIHR were from urban Gurei and rural Pariak and six were male.

154 All 30 mothers were the biological mothers of the infants, and half resided in Gurei or Pariak. Only about
155 one-third had received any education. More mothers in group 1 (younger, less educated and more
156 involved in farming), delivered in the maternity unit. Key vulnerability factors related to the mother for
157 enrolment in MAMI care were primiparity, followed by lactation difficulties and mid-upper arm
158 circumference (MUAC) below (<) 230 mm. These mothers completed on average three follow-up visits,
159 stayed longer in care (24 weeks), received more support from their mothers (or mothers-in-law) or
160 husbands and advised other mothers about MAMI. Maternal mental health measured with the Patient
161 Health Questions-9 score was low across the three groups (score below (<)10). Although some mothers
162 reported knowing the BHWs in their communities, few had received explicit support from them. The most
163 common point of referral for MAMI was the vaccination unit. Most of the 13 mother-infant pairs who ended
164 care when the infant reached 6 months of age recovered, i.e., not requiring follow-on healthcare services
165 (Table 1).

166

167 **Table 1: Characteristics of mothers participating in the South Sudan study, February–March 2024**

Characteristics of mothers and infants of enrolled pairs	Group 1 Follow-up visits until 6m of age	Group 2 Few follow-up visits	Group 3 No follow-up visit	All*
All	13	8	9	30
Urban (vs rural)	8	2	5	15
MOTHER				
Mother (vs caregiver)	13	8	9	30
Mother's age at enrolment (years) (median)	20.0	26.0	25.0	200
Mother's education:				
None	10	6	6	22
Primary	2	1	2	5
Secondary	1	1	1	3
Mothers' source of income:				
Commerce	2	2	1	5
Farming	7	2	5	14
Housekeeping	4	4	3	11
Mothers' access to health services:				
Attended ANC	4	1	5	10
Delivered at maternity unit	5	2	3	10
Mother's MUAC at enrolment (mm) (median)	256	238	250	256
Mothers' reasons for being classified as at moderate risk:				
Low MUAC (<230 mm)	3	1	1	5
Lactation difficulties	3	2	2	7
Primipara	9	3	4	16
Multipara	1	2	1	4
Adolescent mothers	2	1	1	4
Maternal mental health (N=25):				
Mental health low risk (PHQ-9 score <10)	10	7	8	25
Mental health moderate or high risk (PHQ-9 score ≥10 or positive Question9)	0	0	0	0
Mothers in MAMI care shared experiences with others:				
Mothers advised mothers (one to one)	9	4	4	17
Mothers shared MAMI information in women's groups	3	0	0	3
Mothers receiving support:				
Support from mother (or in-law)	7	2	2	11
Support from family	1	1	4	6
Boma health worker (BHW) known	4	1	4	9
Support from BHW	1	0	1	2
Mothers in MAMI care shared experiences with others:				
Mothers advised mothers (one to one)	9	4	4	17
Mothers shared MAMI information in women's groups	3	0	0	3
Number of return visits after enrolment (median)	3	2	0	3
Duration of care (enrolment to last visit) (weeks) (median)	24	14	0	15
Care outcome:				
Recovered	11	NA	NA	11
Not-recovered	2	NA	NA	2
INFANT				
Male infants	5	4	5	14
Infant age at enrolment (months) (median)	1.7	0.4	1.0	1.4
Infant's MUAC at enrolment (mm) (median)	108	100	107	104
Infants' reasons for being classified as at moderate risk:				
Mother-reported low birthweight (LBW <2500 g)	0	1	0	1
Mother-reported preterm birth (<37 weeks gestation)	0	0	1	1
Morbidity	7	5	5	17
Low MUAC (Infant <7 weeks MUAC <110 mm)	5	5	5	15
Low MUAC (Infant ≥7 weeks MUAC <115 mm)	4	3	2	9
Low WAZ (<-2)	0	3	0	3
Feeding problem	13	11	6	30

* Note that for some indicators the sample size is smaller than 30 due to missing data.

ANC= antenatal care; BHW= boma health worker; MUAC= Mid-upper arm circumference; NA= not applicable; PHQ-9= mental health assessment tool.

Exploring the acceptability and adherence of the integrated MAMI care pathway for supporting vulnerable infants and their mothers: a qualitative study

168 Infants in group 1 were older on average on enrolment. Across the three groups, almost all infants <7
169 weeks had a MUAC <110 mm, and two-thirds of infants above or equal to (\geq) 7 weeks had a MUAC <115
170 mm. Key vulnerability factors related to the infant for enrollment were feeding problems, morbidity and low
171 MUAC (Table 1).

172 **Thematic analysis of acceptability**

173 Analysis of the perception of MAMI care among health workers and mothers yielded 29 and 33 themes,
174 respectively, across the seven theoretical constructs of healthcare acceptability, some of which were
175 repeated when a different construct lens was applied (Figures 1 and 2).

176 **Health workers' perceptions of care**

177 **Coherence, or understanding MAMI care and how it worked.** The coherence construct focused on
178 how health workers described MAMI care and which aspects they found important or new. All health
179 workers described MAMI care as encompassing vulnerability assessment and risk-based care for infants
180 u6m and their mothers through targeted counselling and support. They noted that since the introduction
181 of MAMI care, the vulnerability of small, sick newborns and infants under 6 months has been
182 acknowledged. Previously, a breastfed newborn was considered stable and only needed to return to
183 the PHCC for vaccination or when ill.

184 *"Before, when delivering a stable newborn, it was assumed that everything was OK, and they would only*
185 *return after 6 weeks for EPI ... but now with MAMI, following growth and development closely, this was an eye*
186 *opener." MAMI Assistant 3*

187 Some health workers mentioned the importance of community awareness to encourage mothers.

188 *"Sensitization in the community is important in supporting mothers in knowing and receiving family support to*
189 *be able to go to the HF and follow the advice." MAMI Assistant 2*

190 **Self-efficacy, or confidence in providing MAMI care.** Most health workers felt competent to assess
191 vulnerability and give mothers targeted messages about healthy behaviors.

192 *"I was able because the training was perfect, and MAMI became easy, I was able to do in my daily activity.*
193 *Previous knowledge plus new knowledge helped me..." Clinical Officer 5*

194 Some liked the tools that facilitated their work.

195 *"Assessing using the form is better because the questions are there, and the findings are recorded and cases*
196 *are identified. If referral, ... without forms, this information is written in the mother's 'book. The form is far*
197 *better and fast than being writing on any paper." Clinical Officer 4*

198 On the other hand, health workers complained about the heavy workload, lengthy consultations and lack
199 of teamwork and links with the hospital to follow up referred mother-infant pairs.

200 *"Whatever you do many times comes easy. IMCI assessment was easy because I have been doing it before.*
201 *What was challenging was time and workload and working alone." Clinical Officer 2*

202 Several felt the lack of transport to the communities or incentives hindered their ability to convince
203 mothers to be involved in MAMI care.

204 *"Mothers have a lot of expectations and challenges to address. They expect drugs or other tangible support,*
205 *and we have nothing to give." MAMI Assistant 4*

206 **Affective attitude, or interest in being involved in MAMI care.** Most health workers felt good about
207 providing care to a previously ignored but vulnerable population. Some felt they could facilitate mothers'
208 access to care across services and in turn receive recognition and respect.

209 *"I was touched by being involved with MAMI because no one talked about this very vulnerable population,*
210 *which was left out, and we provided services and linked with other services. ... Mothers were touched by the*
211 *messages. For example, expressing milk for mothers when they had to be absent, was unheard of. They*
212 *would rather give cow milk." MAMI Assistant 2*

213 Others felt frustrated when they failed to solve mothers' problems.

214 *"Mothers were comfortable to share health-related issues. It is good to talk, and if they understand they will*
215 *adhere and if not, they will not. There are a lot of misconceptions, cultural taboos and certain topics are*
216 *difficult to provide. ... Also in our tradition it is not good to enter a house where a mother has delivered,*
217 *sometimes I feel uncomfortable to assess a newborn, unless you give something good." Pariak BHW 2*

218 Nearly half of the health workers were disappointed in the lack of compensation for the extra work and
219 transport to provide MAMI care.

220 *"Discouraging is that I work but am not paid." Pariak BHW 4*

221 **Burden, or efforts to provide MAMI care.** Some health workers did not find MAMI care a burden
222 because they saw it as their professional commitment and were motivated by what they learned and the
223 technical support they received during implementation.

224 *"The collaboration and learning from each other was good. Collaboration with the MAMI Assistant helped to*
225 *start early with the MAMI cases ... The risk assessment needs a good mind set." Clinical Officer 5*

226 On the other hand, they felt MAMI care was complex and time-consuming and felt they should be paid an
227 incentive for the extra work. Although they could see the purpose of the forms and paperwork, many
228 found them a burden.

229 *"The forms were too long, and it takes longer hours and many mothers on the line were complaining. I feel like*
230 *I'm not doing enough and faster while conducting risk assessment." Clinical Officer 5*

231 **Ethicality, or whether MAMI care fits with the value system.** Most health workers recognized that
232 MAMI care saved lives and found fulfilment in aiding a vulnerable but previously neglected population.
233 Acquiring skills to serve vulnerable mother-infant pairs bolstered their confidence and engagement.
234 Positive feedback from mothers reinforced their belief in the value of their work.

235 *"Meaningful good messages that prevent death. What is different now is that MAMI also focuses on u6m."*
236 *Gurei BHW 3*

237 **Opportunity cost or sacrifices and gains from providing MAMI care.** Most health workers regretted
238 having less personal time or time for income-generating activities because of longer working hours.

239 *"Before, I was a businesswoman in the market. Since I became a BHW, I stopped. ... I work more than four*
240 *hours and have to give up income and social contact. I'm not able to cultivate. I cannot go and meet my*
241 *friends. When I reach home, it is late." Gurei BHW 2*

242 **Perceived effectiveness, or the ability of MAMI care to achieve its purpose as expected.** Most
243 health workers felt MAMI care addressed a vital need by offering specialized, risk-based care across
244 services and in the communities. About one-third appreciated that MAMI care enhanced infant health and
245 development, which in turn increased maternal trust and adherence to care practices. This improvement
246 in care adherence had tangible results.

247 *"MAMI filled a gap. Nobody paid attention to u6m before... For the infant, there was vaccination and concern*
248 *in case of illness. MAMI makes an interconnection ... a collaboration between maternity, EPI, OPD and the*
249 *movements between them." MAMI Assistant 2*

250 Mothers' perceptions of care

251 **Coherence, or understanding MAMI care and how it worked.** The coherence construct focused on
252 what mothers remembered about the MAMI Care Pathway and which messages they found important.
253 Most mothers understood that MAMI care was about identifying vulnerability factors, receiving targeted
254 counselling to improve infant health and growth and taking good care of themselves. Messages they
255 retained included exclusive and frequent breastfeeding up to 6 months, maintaining good hygiene, and
256 returning to the health facility when their infants were sick. For the mothers, counselling meant "*listening*
257 *to and following up health messages*".

258 *"MAMI is assessing infant and mother to know their health, for taking care ... In MAMI there are pictures of*
259 *breastfeeding and position, and on hygiene... My family told to give water to the infant, water is good when*
260 *the infant is thirsty. In MAMI I got the message to do exclusive breastfeeding, because breastfeeding*
261 *includes water."* Mother GUR08 Group

262 **'Self-efficacy', or confidence in applying the advice received in MAMI care.** Most mothers were
263 grateful for the advice because they wanted to take good care of their infants and managed to follow the
264 advice. Receiving the same messages across healthcare units reinforced the messages.

265 *"I was confident to follow the [new] advice of the health worker, and even when my own mother had other*
266 *ideas, but when I informed her, she accepted."* Mother PAR05 Group 1

267 Most mothers were unconcerned about the gender of the health workers. However, they did not consider
268 psychological stress or mental wellbeing as health issues, and therefore did not discuss them with health
269 workers. Family or couple conflicts were kept within the family.

270 *"Personal issues I cannot discuss ... because I feel ashamed to share because these are family issues, not*
271 *health issues, so I cannot share."* Mother PAR05 Group 1

272 A few mothers did not follow the advice because they misunderstood MAMI care, were not told clearly
273 what it was about or were not motivated. One deliberately ignored the advice to attend supplementary
274 feeding because she did not like the food given. Some mothers felt they were wasting their time when
275 they did not receive any tangible incentive in MAMI care.

276 *"... I expected to be given milk... I am selling tea. I leave my infant at home with infant formula. My job does*
277 *not allow me to take the baby... I was told to return after one month. I did not return. I was wasting my time."*
278 *Mother GUR09 Group 3*

279 **Affective attitude, or interest in being involved in MAMI care.** Mothers were grateful for the messages
280 they received and appreciated the skills they acquired to better care for their infants. Even mothers who
281 never returned or returned only a few times appreciated the messages, listened and applied the advice.

282 *"I got good messages when I needed it most, this made me very happy. ... Messages on "Take good care of*
283 *baby, take good care of yourself" helped me a lot."* Mother GUR37 Group 1

284 Many mothers mentioned that the messages were new to them, and some enthusiastically recalled their
285 MUAC classification.

286 *"My child was OK but I was thin, my size was less than 230."* Mother PAR74 Group 3

287 The novelty of attention to the mothers and the explicit link between infants and mothers made the
288 mothers feel supported.

289 *"I always feel good [about MAMI care], ... it gave important information that I did not know. ... I liked that I was*
290 *given messages which were very crucial which can really help a mother on how to take care of the baby and*
291 *you yourself as a mother."* PAR19 Group 2

292 MAMI care was especially appreciated by first-time mothers, who usually felt unskilled and were
293 dominated by older women.

294 *"My positive feeling is that I feel happy about MAMI because I got good messages and when I put them into*
295 *practice result in having good health of the baby, so I felt happy about it."* Mother PAR62 Group 3

296 Less than half of the mothers felt returning for follow-up visits to monitor their infants' growth and their
297 own wellbeing important. Some said they did not return because they were not told. But most mothers
298 who went to the PHCC did so for their infants' vaccination, a well-established healthcare service mothers
299 adhered to.

300 *"My family members were supportive to come for vaccination. When it is time for me to come for vaccination,*
301 *they can give me transport fee to use motorcycle to come."* Mother PAR74 Group 3

302 Some mothers found the number of questions and unsolicited attention they received when they came to
303 the PHCC for vaccination worrying and confusing.

304 *"I was not used to all these questions and I felt worried, as if there was something wrong with me and my*
305 *baby. ... I saw MUAC taken of older kids, but not so small as mine."* Mother GUR51 Group 2

306 Some mothers were unhappy about long lines at the PHCC, drug stockouts that meant having to buy
307 medicines from local vendors, lack of incentives such as soap or mosquito nets and the need to pay for
308 notebooks to keep consultation notes as a condition of access to the PHCCs.

309 *"I like the care and receiving drugs. I do not like the long waiting time, and there is no water to drink... If you*
310 *do not pay for the book, then you cannot see the doctor."* Mother GUR 18 Group 3

311 **Burden, or efforts to adhere to MAMI care and advice.** Most mothers felt that the advice to support
312 their infants' health made their participation in MAMI care less burdensome. Some were relieved by
313 encouragements from mothers (mothers-in-law) or husbands.

314 *"Not a burden to listen to messages and follow the advice. I believe that has helped me, changed my life, now*
315 *I know how to take care in the right way."* Mother GUR70 Group 2

316 Concerns were mostly related to distance to the PHCC, time and travel costs, having to leave other
317 children at home, missing work or household duties, being criticized by household members or inquisitive
318 neighbors and insecurity of travel in some areas. Some mothers did not understand the advice or could
319 not follow it.

320 *"I have no transport, I came on foot, I am at home alone with the children, so I cannot come as I would like."*
321 *Mother GUR94 Group 2*

322 **Ethicality, or whether MAMI care fits with the value system.** Most mothers felt able to provide better
323 care for their infants, the most valuable perceived benefit of MAMI care. Their gratitude for new
324 knowledge and skills made them want other mothers to benefit. A few said that consistent messages
325 across healthcare units made them feel secure and confident.

326 *"... I would not have known ..., the women in my home would not have told me ... I could not have taken care*
327 *of myself. I delivered in the PHCC, and I received the same messages there when I came for antenatal care*
328 *and delivery."* Mother PAR49 Group 2

329 **Opportunity cost, or sacrifices and gains from adhering to MAMI care.** Mothers felt they gained
330 knowledge and skills from MAMI that they would never have learned or obtained elsewhere.

331 *"It was possible to return [four times]. I was motivated to come, each time I received a check and*
332 *encouragements. Even neighbors started asking: "where do you go and why"? And answered, "To improve*
333 *the health of my baby". On each visit, it was great to have direct access to the doctor".* Mother GUR73 Group
334 1

335 Conversely, regardless of their adherence to follow-up visits, many mothers faced difficulties securing
336 transport or had to forgo income-generating and social activities.

337 *"Visits to the PHCC are difficult because it is far, and I have to leave things at home. ... MAMI messages were*
338 *helping me as a mother ... but MAMI consumed too much of my time... I had to give up important things such*
339 *as farming, going to watch wrestling, meeting friends."* Mother PAR19 Group 3

340 **Perceived effectiveness, or the ability of MAMI care to achieve its purpose as expected.** Mothers
341 strongly affirmed that the skills they gained made their infants healthier.

342 "MAMI helped me and my child and we are living better. My child is healthy not being sick no diarrhea. I see
343 child of my brother with lot of diarrhea." Mother PAR09 Group 1

344 A small proportion felt that participation in MAMI care facilitated access to health workers, who were more
345 attentive to their concerns. Others were pleased to resolve conflicting advice from the PHCCs and their
346 families.

347 "Some neighbors advised me to go and see a traditional healer, but I said no. On each visit, it was great to
348 have direct access to the doctor...Now my daughter is healthy, which I could not have done with advice from
349 neighbors." Mother GUR73 Group 3

350 **Thematic analysis of adherence behaviour**

351 Factors that influenced adherence to MAMI care were mapped on the COM-B framework of Michie et al.
352 (Table 2) (9). The framework proposes three conditions that interact to generate behavior that in turn
353 influences these conditions. Capability is knowledge and skills to engage in an activity, both psychological
354 (ability to engage in thought processes, capacity to plan) and physical (skill to undertake and follow
355 advice). Opportunity is factors that enable or prompt the desired behavior, both physical (afforded by the
356 environment) and social (afforded by the cultural context, supportive network). Motivation is attitudes and
357 aspirations that energize and direct behavior, both automatic (involving emotions and impulses) and
358 reflective (involving evaluations and plans) (9, 14).

359 **Table 2: Map of influencing factors of positive adherence behavior as perceived by MAMI care**
360 **providers and users participating in the South Sudan study, February–March 2024**

Components of behavior:	Care provider	Care user
Capability (psychological or physical capability): knowledge and skills to engage in the activity	<ul style="list-style-type: none"> – Knowledgeable, understand the relevance of the care pathway – Skilled to plan, provide care, use the tools for implementation, improve quality 	<ul style="list-style-type: none"> – Know what to do and why – Able to do new care on their own at home – Confidence in care and in themselves as carer – Able to explain and justify to others and oppose negative influences
Opportunity (physical or social opportunity): factors lying outside of the individual that enable or prompt the desired behavior	<ul style="list-style-type: none"> – Training, mentoring and support system – Access to tools for implementation and quality improvement – Continuity of care among established quality services (ANC, maternity, PNC, vaccination, IMCI consultations, CMAM) – Organization of care, including task sharing between colleagues 	<ul style="list-style-type: none"> – Respected vaccination schedule and service – Reinforced messages across health services – Facilitated access to health services – Supportive household and community network
Motivation (automatic or reflective motivation): attitudes and aspirations that energize and direct behavior	<ul style="list-style-type: none"> – New initiative with visibility, new learning – Attention for a neglected population – Confidence in providing care – Increased respect from mothers and communities – As a mother/father of infant emotional subject close to their heart – Eagerness for involving in research 	<ul style="list-style-type: none"> – Open to new knowledge and learning – Feeling supported, listened to, health worker showing interest – Improved health for infant and self – Stronger infant when comparing to other infants in the community

361 The findings of the thematic analysis in the COM-B framework show drivers of adherence to MAMI care
362 (Figure 3). For health workers, the key factor in adhering to care was self-confidence and the ability to
363 care for a previously neglected vulnerable population. For mothers, the key factor was seeing their infants
364 thrive.

365 Discussion

366 The study enhanced understanding of the effectiveness of MAMI care by examining its acceptability
367 among health workers and mothers of enrolled mother-infant pairs and identified factors that facilitated or
368 hindered adherence to MAMI care.

369 Health workers were trained and shaped by vertical health programs that lacked a holistic approach,
370 resulting in gaps in continuity of care. For example, they often overlooked infants u6m after a healthy
371 start, assuming that breastfeeding would proceed smoothly, common diseases would be infrequent, and
372 growth and development issues would only arise with the introduction of complementary feeding and
373 exposure to unhealthy environments.

374 Gaining the expertise and skills to serve this vulnerable group enhanced health worker's sense of
375 professional fulfilment. However, they did not report on challenges in addressing maternal physical and
376 mental health, an aspect the interview did not uncover. Cultural norms do not recognize psychological
377 and mental problems as health issues or allow open discussion.

378 Clinical health workers worried most about the additional workload for in-depth assessment, filling out
379 forms and providing risk-based counselling. The issue, however, seemed to be lack of routine
380 assessment. With good practice, this activity would be fast, as few mother-infant pairs would present risks
381 requiring intensive attention. Also, effective organizational skills could help manage crowds more
382 efficiently, distribute consultations more evenly or extend the activity window beyond 9 am to 12 pm. In
383 some sites, MAMI screening and assessment slowed when the MAMI assistants or the one trained
384 clinical officer, who were exclusively assigned these tasks, were absent. Vulnerability screening focused
385 heavily on measuring MUAC, which was easy, while key vulnerability questions received less
386 attention. This aligns with mothers' reports that community nutrition volunteers only measured MUAC of
387 children 6–59 months, overlooking other important health and nutrition issues.

388 BHWs trained and equipped to implement community-based healthcare, which included MIHR-specific
389 maternal and child health activities like MAMI, had mixed feelings about MAMI care. The extra workload
390 was difficult to fit in their voluntary work schedule, and cultural norms made access to households with
391 newborns difficult. Nonetheless, the BHI materials covered most vulnerabilities for infants u6m and their
392 mothers, except for measuring MUAC in infants u6m and the mother-infant pair-centered care approach.
393 BHWs were often too busy or did not focus on infants u6m, as the BHI targets children under 5 years for
394 vaccination, illness or malnutrition for 6–59 months. Newborns are kept inside the house, where BHWs
395 are not allowed without the father's permission. Consequently, small infants were not their primary
396 concern.

397 Mothers were eager to learn how to care for their infants and pleased when their infants thrived when
398 they followed MAMI advice. However, targeted counseling was often simple message sharing, which
399 might not be enough to change behavior. Mothers adhered to advice if it seemed feasible and had family
400 support, facilitated by community awareness. Urban mothers reported less family support and social
401 interaction than rural mothers. Mothers had difficulty discussing mental health issues, which were much
402 misunderstood. All mothers were concerned about frequent follow-up visits, especially traveling to PHCCs
403 far away with newborns or small infants. Transportation costs, mentioned as a major barrier to follow-up
404 visits, often masked more complex reasons. Mothers sometimes felt the trip was not worth the effort
405 compared to other NGO-supported programs that provided incentives. However, adherence to the
406 vaccination schedule was well respected and supported by the community, suggesting that MAMI follow-
407 up schedules could align with these appointments to improve adherence. Also, the contact with the BHW
408 seemed limited to measuring MUAC of older children, leaving young mothers in a vacuum in terms of
409 infant feeding and care. It is no surprise mothers were eager for MAMI care to continue.

410 Missed opportunities for health workers included adequate time and skills for counseling, addressing
411 maternal mental health, community engagement, consultation management, form proficiency, and task
412 sharing. For mothers, it included connections to BHWs and adequate family and community support.
413 Seized opportunities were the strengthened integrated management of neonatal and childhood illness
414 (IMNCI) approach and engaging mothers in their infants' health. Recommendations for improvement
415 included making MAMI care routinely available, improving patient flow, combining visits, and giving
416 consistent health advice. Follow-up visits should be mutually agreed upon with mothers, considering their
417 ability to return for care. In the community, BHW teams should adopt a holistic, person-centered
418 approach, decentralize health care, and strengthen individual counseling. Urban mothers would benefit
419 from health education through radio, theatre, telephone, and television.

420 This study faced limitations. First, the interviews were conducted by the MAMI coordinator, who was also
421 an employee of MIHR and responsible for overseeing MAMI implementation. This overlap in roles could
422 have affected participants' willingness to speak freely, as well as the coordinator's impartiality during the
423 interviews (interviewer bias). Second, interviews were mainly done at the MIHR office or the PHCC.
424 Health workers, still under MIHR employment, might have been less willing to be frank in their responses.
425 Mothers interviewed at their PHCCs might have felt pressured to provide responses they perceived as
426 socially acceptable or favorable. Third, interviews with mothers were conducted in their native language
427 and immediately translated by a proficient boma health worker. Despite the translators' expertise, subtle
428 nuances might have been lost, necessitating careful interpretation of quotes (translation accuracy). The
429 interview team and two study investigators addressed these limitations during interview debriefings to
430 minimize the impact of biases.

431 **Conclusion**

432 The study found the MAMI Care Pathway acceptable to both providers and users, providing care for a
433 previously overlooked vulnerable group. It highlighted the need for policy changes to prioritize
434 comprehensive, patient-centered care for vulnerable infants u6m and their mothers. To ensure effective
435 adoption, delivery and use challenges must be resolved. Successful adaptation will encourage adherence
436 from both providers and users.

437 The findings, though context-specific, may be applicable to other settings with similar demographics and
438 healthcare challenges, and provide a foundation for future research and intervention development.

439 **[Competing interest]** None declared.

440 **[Funding]** MOMENTUM Integrated Health Resilience is funded by the U.S. Agency for International
441 Development (USAID) as part of the MOMENTUM suite of awards, under USAID cooperative agreement
442 #7200AA20CA00005.

443 **[Acknowledgements]** Our heartfelt appreciation goes to Michael Onguech Costantino, MAMI Assistant,
444 and Abraham Kuech Majok, Boma Health Worker and translator, for their invaluable contributions
445 throughout the interview process. Special thanks are extended to all South Sudan Ministry of Health and
446 MIHR collaborators in Juba and at the Gurei, Hai Dinka, Pariak, Wau, and Yambio primary healthcare
447 centers.

448 **[Authors tasks]** HD and MHe designed the study, supported the interviews, managed and analyzed the
449 data, and wrote the report. KD participated in the study design and reporting. ANB conducted the
450 interviews.

451 **[Patient and Public Involvement]** Patients or the public WERE NOT involved in the design, or conduct,
452 or reporting, or dissemination plans of our research.

453 **[Figure/Illustration legends]** Figure 1: Themes of acceptability of the MAMI Care Pathway approach as
454 perceived by care providers participating in the South Sudan study, February–March 2024 (blue boxes
455 are positive or enabling factors, and orange boxes are negative or hindering factors).
456 Figure 2: Themes of acceptability of the MAMI Care Pathway approach as perceived by care users
457 participating in the South Sudan study, February–March 2024 (blue boxes are positive or enabling factors
458 and orange boxes are negative or hindering factors).
459 Figure 3: Behavior conditions that improve adherence to MAMI care provision and use as perceived by
460 MAMI care providers and users participating in the South Sudan study, February–March 2024.

461 REFERENCES

- 462 1. Lawn JE, Ohuma EO, Bradley E, Idueta LS, Hazel E, Okwaraji YB, et al. Small babies, big risks: global estimates
463 of prevalence and mortality for vulnerable newborns to accelerate change and improve counting. *Lancet*
464 [Internet]. 2023 May 1;401(10389):1707–19. Available from: [https://doi.org/10.1016/s0140-6736\(23\)00522-6](https://doi.org/10.1016/s0140-6736(23)00522-6)
- 465 2. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, De Onis M, et al. Maternal and child undernutrition
466 and overweight in low-income and middle-income countries. *Lancet* [Internet]. 2013 Aug 1;382(9890):427–51.
467 Available from: [https://doi.org/10.1016/s0140-6736\(13\)60937-x](https://doi.org/10.1016/s0140-6736(13)60937-x)
- 468 3. United Nations Office for the Coordination of Humanitarian Affairs (OCHA). South Sudan Humanitarian Needs
469 Overview 2023 [Internet]. 2022 Nov. Available from: <https://www.unocha.org/publications/report/south-sudan/south-sudan-humanitarian-needs-overview-2023-november-2022>
- 471 4. MOMENTUM Integrated Health Resilience (MIHR). MAMI South Sudan Capacity Analysis. Internal report;
472 2022a.
- 473 5. Deconinck H, Wrottesley SV, McGrath M, Nasira Boi A, Mane Luka P, Sabino S, et al. Implementing Care for
474 Vulnerable Infants Under Six Months and Their Mothers. 'Learning by Doing' Case Study Series: South Sudan.
475 Oxford, United Kingdom of Great Britain and Northern Ireland: Emergency Nutrition Network (ENN); 2024.
- 476 6. South Sudan. Gavi, the Vaccine Alliance. 2022 [cited 2024 Jun 1]. Available from:
477 <https://www.gavi.org/programmes-impact/country-hub/africa/south-sudan>
- 478 7. Campbell H, Hotchkiss R, Bradshaw N, Porteous M. Integrated care pathways. *BMJ British Medical Journal*
479 [Internet]. 1998 Jan 10;316(7125):133–7. Available from: <https://doi.org/10.1136/bmj.316.7125.133>
- 480 8. MAMI Global Network, Emergency Nutrition Network (ENN), London School of Hygiene and Tropical Medicine
481 (LSHTM). MAMI Care Pathway Package, Version 3 [Internet]. 2021. Available from:
482 <https://www.ennonline.net/mamicarepathway>
- 483 9. Michie S, Van Stralen MM, West R. The behaviour change wheel: A new method for characterising and
484 designing behaviour change interventions. *Implementation Science* [Internet]. 2011 Apr 23;6(1). Available from:
485 <https://doi.org/10.1186/1748-5908-6-42>
- 486 10. O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. Standards for Reporting Qualitative research.
487 *Academic Medicine* [Internet]. 2014 Sep 1;89(9):1245–51. Available from:
488 <https://doi.org/10.1097/acm.0000000000000388>
- 489 11. Sekhon M, Cartwright M, Francis JJ. Acceptability of healthcare interventions: an overview of reviews and
490 development of a theoretical framework. *BMC Health Services Research*. 2017;17(1). doi:10.1186/s12913-017-
491 2031-8
- 492 12. MOMENTUM Integrated Health Resilience (MIHR). Protocol of the Pilot Study on the Integrated Management of
493 Small and Nutritionally At-risk Infants Under 6 Months and Their Mothers (MAMI) Care Pathway in South Sudan.
494 2022b Mar.
- 495 13. Nasira Boi A, Traore Hebie M, Deconinck H. Introducing the Integrated Care Pathway for the Management of
496 Small and Nutritionally At-risk Infants (MAMI) in South Sudan, 2022–2024. Interim report; 2023 Dec.
- 497 14. Mayne J. The Capabilities, Opportunities and Motivation Behaviour-Based Theory of Change Model [Internet].
498 2016. Available from:
499 https://www.researchgate.net/publication/301701597_The_Capabilities_Opportunities_and_Motivation_Behaviour-Based_Theory_of_Change_Model
- 501 15. Bowen DJ, Kreuter M, Spring B, Cofta-Woerpel L, Linnan L, Weiner D, et al. How we design feasibility studies.
502 *American Journal of Preventive Medicine* [Internet]. 2009 May 1;36(5):452–7. Available from:
503 <https://doi.org/10.1016/j.amepre.2009.02.002>

Theoretical construct	Coherence	Self-efficacy	Affective attitude	Burden	Ethicality	Opportunity cost	Perceived effectiveness
Theme	Vulnerability assessment, risk-based care (n=20)	Knowledge and skills gained (n=17)	Pride in professional duty (n=17)	Professional commitment (n=17)	Vital contribution to life saving efforts (n=22)	Less time for social activities (n=13)	Critical gap in care of vulnerable infants & mothers (n=17)
	Targeted counselling and support (n=20)	Specific tools facilitated work (n=6)	Ability to help infants & mothers (n=6)	Complex and time-consuming work (n=17)	Valuable skills to enhance care (n=7)	Less time for income-generating activities (n=6)	Collaboration among services and communities (n=11)
	Wellbeing of infant & mother (n=20)	Heavy workload, lengthy consultations (n=12)	Recognition and respect (n=4)		Mothers' satisfaction (n=4)	Personal expenses used for transport (n=3)	Improved trust, care compliance of mothers (n=6)
	Community awareness (n=4)	Failure to meet mothers' expectations (n=12)	Inability to solve community problems (n=15)				Improved competence in care (n=6)
	Enhanced mother's participation (n=3)	Insufficient means for community work (n=6)	No compensation for extra work or transport (n=6)	Lack of mothers' understanding (n=4)			Improved health and growth of infant from care (n=8)

Figure 1: Themes of acceptability of the MAMI Care Pathway approach as perceived by care providers participating in the South Sudan study, February–March 2024 (blue boxes are positive or enabling factors, and orange boxes are negative or hindering factors)

Theoretical construct	Coherence	Self-efficacy	Affective attitude	Burden	Ethicality	Opportunity cost	Perceived effectiveness
Theme	Counselling (messages) (n=29)	Able to follow guidance for good infant care (n=23)	Satisfied with knowledge on good care (n=27)	Satisfaction with messages (n=23)	Competence to provide better care to infant (n=20)	Knowledge and skills gained (n=19)	Gained competence on nurturing infant (n=22)
	Vulnerability factors (n=23)	Discuss health concern with health worker (n=14)	Satisfied with improving care (n=23)	Support of mother (in law) and husband (n=5)	Acquired valuable information and new insights (n=17)	Transport costs not covered (n=8)	Resolution of conflicting advice and support (n=4)
	Linking infant & mother (n=19)	Unable to comprehend, follow the advice (n=5)	Proud of a healthy infant (n=8)	Self-reliant in providing infant care (n=8)	Eagerness for other mothers to access knowledge (n=11)	Neglect of income-generating and social activities (n=6)	Better access to healthcare and related services (n=6)
	Returning for monitoring (n=13)	Unable to discuss non-health issues (n=5)	Content with improved personal healthcare (n=6)	Ability to balance work and infant care (n=13)	Aligning family with learning (n=8)		
			Discouraged by health facility issues (n=6)	Challenges due to distance to health facility (n=17)	Consistency of messages across health services (n=2)		
			Neighbours envy and mistrust (n=4)	Attendance conflict with work & household tasks (n=8)			
			Inconsistent messages between health worker and home (n=2)	Difficulty to understand, follow the advice (n=6)			


Figure 2: Themes of acceptability of the MAMI Care Pathway approach as perceived by care users participating in the South Sudan study, February–March 2024 (blue boxes are positive or enabling factors and orange boxes are negative or hindering factors)

[Title of manuscript] Exploring the acceptability and adherence of the integrated MAMI care pathway for supporting vulnerable infants and their mothers in South Sudan: a qualitative study

**Care providers
(clinical and community health workers)**



COM-B framework



**Care users
(mothers)**



Figure 3: Behavior conditions that improve adherence to MAMI care provision and use as perceived by MAMI care providers and users participating in the South Sudan study, February–March 2024