

1 **Associations between weather and *Plasmodium vivax* malaria in an elimination setting in**
2 **Peru: a distributed lag analysis**

3 Gabriella Barratt Heitmann,^{1*} Xue Wu,² Anna T. Nguyen,¹ Astrid Altamirano-Quiroz,³ Sydney Fine,²
4 Bryan Fernandez-Camacho,³ Antony Barja,³ Renato Cava,³ Verónica Soto-Calle,³ Hugo Rodriguez,⁴
5 Gabriel Carrasco-Escobar,³ Adam Bennett,² Alejandro Llanos-Cuentas,³ Erin A. Mordecai,^{5,6} Michelle S.
6 Hsiang,^{2,7,8,9} Jade Benjamin-Chung^{1,9}

7

8 1. Department of Epidemiology and Population Health, School of Medicine, Stanford University,
9 Stanford, CA, USA

10 2. Malaria Elimination Initiative, Institute for Global Health Sciences, University of California San
11 Francisco (UCSF), San Francisco, CA, USA

12 3. Instituto de Medicina Tropical, Alexander von Humboldt, Universidad Peruana Cayetano Heredia,
13 Lima, Perú

14 4. Universidad Nacional de la Amazonía Peruana, Loreto, Perú

15 5. Department of Biology, Stanford University, Stanford, CA, USA

16 6. Woods Institute for the Environment, Stanford University, Stanford, CA, USA

17 7. Department of Epidemiology and Biostatistics, UCSF, San Francisco, CA, USA

18 8. Department of Pediatrics, UCSF, San Francisco, CA, USA

19 9. Chan Zuckerberg Biohub, San Francisco, CA, USA

20

21 *Corresponding author contact information:

22 Email: gabbyrbh@stanford.edu

23 Phone number: 609-972-2178

24 ORCID: <https://orcid.org/0000-0002-4212-0954>

25

26 **Conflicts of Interest Statement**

27 *The authors declare they have no conflicts of interest related to this work to disclose.*

28

29 **Funding Statement**

30 *Research reported in this publication was supported by grants to MH from the National Institute of*
31 *Allergy and Infectious Disease under Award Number U01AI157962.*

32 **ABSTRACT**

33 **Background**

34 *Plasmodium vivax* (*Pv*) is the predominant malaria species in countries approaching elimination.
35 In the context of climate change, understanding environmental drivers of transmission can guide
36 interventions, yet evidence is limited, particularly in Latin America.

37

38 **Objectives**

39 We estimated the association between temperature and precipitation and *Pv* malaria incidence in
40 a malaria elimination setting in Peru.

41

42 **Methods**

43 We analyzed malaria incidence data from 2021-2023 from 30 communities in Loreto, Peru with
44 hourly weather data from the ERA5 dataset and land cover data from MapBiomias. Predictors
45 included average weekly minimum and maximum temperature, high heat (>90th percentile mean
46 temperature), total weekly precipitation, and heavy rain (>90th percentile total precipitation). We
47 fit non-linear distributed lag models for continuous weather predictors and generalized additive
48 models for binary predictors and the lookback period was 2—16 weeks. Temperature models
49 adjusted for total precipitation; precipitation models adjusted for maximum temperature. We
50 performed subgroup analyses by season, community type, and distance to forest edge.

51

52 **Results**

53 The median vs. lowest values of weekly average minimum temperature was associated with 2.16
54 to 3.93-fold higher incidence 3-16 weeks later (5-week lag incidence ratio (IR) =3.93 [95% CI
55 2.18, 7.09]); for maximum temperature, the association was hump-shaped across lags, with
56 protective associations at 1-2 and 15-16 week lags and 1.07-1.66-fold higher incidence at 6-13
57 week lags. High heat (>27.5°C) was associated with 1.23 to 1.37-fold higher incidence at 5--9
58 week lags (9-week lag IR = 1.25 [1.02, 1.53]). Associations between total precipitation and
59 malaria incidence were hump-shaped across lags, with the strongest positive association at 750
60 mm of precipitation at a 9-week lag (IR=1.56; [1.27, 1.65]). Heavy rain (>186mm) was
61 associated with 1.22–1.60-fold higher incidence at 2–10 week lags (9-week lag IR=1.23 [1.02,
62 1.49]).

63

64 **Discussion**

65 Higher temperatures and precipitation were generally associated with higher malaria incidence
66 over 1–4 months.

67

68

69

70 INTRODUCTION

71 As countries approach malaria elimination, the proportion of malaria infections due to
72 *Plasmodium vivax* (*Pv*) increases.¹ *Pv* presents unique challenges for malaria elimination efforts
73 because of its ability to survive in a wide range of environments, including temperate settings, by
74 utilizing a dormant hypnozoite stage.² Relapsing infections occur weeks to months after a prior
75 infection or relapse, and serve as a significant reservoir for persistent transmission.^{3 4,5} That *Pv*
76 infections can be latent, as well asymptomatic, sub-patent, or minimally symptomatic makes
77 diagnosis and surveillance a challenge. Further, symptomatic infections are low density and often
78 missed through standard diagnostics.^{4,6,7} These characteristics greatly obfuscate *Pv* transmission
79 biology and make *Pv* harder to detect, treat, and ultimately eliminate.

80
81 Climatic and environmental factors such as temperature, precipitation, land use, and vegetation
82 are strong drivers of malaria transmission in various endemic settings.⁸ All initial *Pv* infections
83 are environmentally mediated, and understanding these drivers is critical to interrupting the long,
84 enduring cycles of *Pv* transmission described above. Increased precipitation can increase
85 standing water presence, soil moisture, and humidity, which in turn increase breeding habitat for
86 *Anopheles*,⁹ the vector for malaria transmission. However, this relationship is complicated: in
87 dry-land communities, heavy rain can flush mosquito larvae, leading to decreased mosquito
88 populations, whereas in riverine settings such as in the Amazon, flooding from heavy rains can
89 increase breeding ground and mosquito populations.⁹ Temperature has strong effects on the
90 capacity of the *Anopheles* vector to transmit malaria via the mosquito biting rate, mosquito
91 abundance, parasite incubation rate, and mosquito longevity.^{10,11} A schematic of the timeline of
92 these environmental drivers of transmission is presented in Figure 1, generated from prior
93 findings.^{6,12–16} Modeling suggest *Pf* malaria transmission by various *Anopheles* peaks at
94 25°C,^{10,17} with substantial decreases in transmission above this thermal optimum. More recent
95 work has supported this thermal optimum for both *Pf* and *Pv* transmission by *An. gambiae*.¹¹
96 Finally, weather could be a direct trigger for relapse infections for *Pv* infections¹⁸, and there is
97 other indirect evidence that relapses may be triggered by antigens in uninfected mosquito saliva
98 (biting rate is temperature-sensitive).³ However, there is limited research on environmental
99 drivers of *Pv*, especially in Latin America, including in the Amazon, which is estimated to bear
100 5% of the world's *Pv* malaria burden.⁶

101
102 Our objective was to assess the influence of rainfall and temperature on *Pv* incidence in Loreto
103 Region, Peru, a low transmission area key to the Peruvian government's target of malaria
104 elimination by 2030.¹⁹ Environmental risk factors for malaria tend to be highly localized,^{20,21} and
105 the Loreto Region has maximum temperatures between 30–35°C thought to be above the malaria
106 transmission thermal optimum, and daily temperature extremes are thought to impact malaria
107 transmission²². To our knowledge, the thermal biology of *Pv* transmission by *An. darlingi*, the
108 primary vector in this region, has not been well characterized in disease ecology studies.

109 Information about climatic drivers of *Pv* transmission is needed to inform effective tailoring and
110 targeting of interventions for malaria elimination.

111

112

113 **METHODS**

114 **Study design**

115 We analyzed malaria incidence data collected from January 2021 to December 2023 in 30
116 communities in San Juan Bautista, Punchana, Alto Nanay, Iquitos, and Belen districts in Loreto
117 region, Peru, an area covering 2,569 km² (Figure 2). This study used data collected prior to
118 baseline for the FocaL Mass Drug Administration for Vivax Malaria Elimination (FLAME) trial
119 in Peru (NCT05690841), and therefore communities included in this study were those that met
120 the eligibility criteria for FLAME. The study included surveillance data from riverine
121 communities only accessible via boat on the Itaya, Momón, and Nanay rivers and dryland
122 communities along the Iquitos-Nauta highway. The spatial distribution of these communities is
123 shown in Figure 2. According to a census conducted for the FLAME trial, the average population
124 size of each community was 240 individuals (range: 31 to 807).

125

126 Peru's malaria burden is concentrated in the Loreto Region, which includes a large swath of the
127 Peruvian Amazon.⁷ Since the 1990s, rapid urbanization and deforestation in the area surrounding
128 Iquitos, the capital of the Department of Loreto, has contributed to increased malaria burden⁷,
129 likely through increased forest edge habitat, which promotes mosquito breeding, survival, and
130 biting²³. While this region is approaching elimination, sustained, endemic transmission remains,
131 with an annual incidence rate of 17.4 cases per 1000 in 2019,¹⁹ and *Pv* malaria accounts for
132 approximately 80% of the malaria burden.⁷ Transmission typically peaks in Loreto between
133 February and July⁷ and there is a sizable asymptomatic caseload.¹³ Asymptomatic infections
134 typically go undetected and uncounted in routine malaria surveillance in Peru,¹³ since only
135 febrile individuals are tested.

136

137 From 2005-2010, there was a large effort towards malaria elimination via the Project for Malaria
138 Control in Andean Border Areas (PAMAFRO) program that improved case management,
139 including through community case workers, and deployed insecticide-treated bed nets.⁷
140 Transmission declined during the project, but the project did not achieve elimination, and
141 transmission has since risen.²⁴ Following PAMAFRO, the government of Peru adopted Plan
142 Malaria Cero (PMC) in 2017 to achieve malaria elimination. Control activities over the study
143 period involved some reactive (RCD) and active case detection (ACD), distribution of long-
144 lasting insecticide-treated bed nets, indoor residual spraying, and community-level
145 chemoprophylaxis²⁵.

146

147 **Outcome data**

148 Our primary outcome was weekly community-level vivax malaria incidence primarily measured
149 through passive case detection conducted by the Peruvian Ministry of Health (MINSA), though
150 PMC activities including ACD and RCD also occurred over the study period and were recorded
151 in surveillance data. Both physical records stored in health posts and online records compiled by
152 MINSA were used in this analysis. Febrile patients presented for care at a health post and were
153 tested for malaria via microscopy. According to PMC policy²⁵, blood smears are stained with 2%
154 Giemsa for 30 minutes. Parasite densities are calculated from the number of asexual parasites per
155 200 leukocytes (or per 500, if <10 asexual parasites/200 leukocytes), assuming a leukocyte count
156 of 8,000/ μ L. A blood smear is considered negative if examination of 100 high power fields does
157 not reveal asexual parasites. Thin smears are used for parasite species identification. Slides are
158 read by two microscopists. If there are discordant results, the results are determined by a third
159 microscopist. Only *Pv* cases were included in the present study, and the data did not distinguish
160 between primary infections versus relapse cases. Cases were matched to community census data
161 to confirm community of residence, and then matched to a community centroid. Population
162 counts per community were calculated from a census conducted from 2023, which was adjusted
163 via interpolation for missing and/or non-participatory households.

164

165 **Environmental variables**

166 We obtained temperature and precipitation data from the ERA5-Land Hourly dataset collected
167 by the Copernicus Climate Change Service²⁶. We used the air temperature at 2 meters above the
168 surface (K) band for all temperature variables and the total precipitation (m) band for
169 precipitation variables. Weather variables were matched to incidence data via community
170 centroid.

171

172 We imputed temperature or precipitation values less than 0 as missing. We aggregated hourly
173 temperature values into the weekly minimum, mean, and maximum temperature; the average of
174 each day's temperature range (maximum minus minimum) over a week; and total precipitation
175 observed each week. We defined a binary indicator for high heat events coded as 1 for weeks
176 with any days when the mean daily temperature exceeded the 90th percentile of mean daily
177 temperatures for that year (average over all years was 27.5°C) and 0 otherwise. Similarly, we
178 defined a binary indicator for heavy rain events coded as 1 for weeks with any days when the
179 total daily precipitation exceeded the 90th percentile of total daily precipitation for that year
180 (average of all years was 186mm) and 0 otherwise.

181

182 We conducted a distinct household-level analysis on distance to forest edge as a potential effect
183 modifier of the associations between temperature and rainfall and malaria incidence. Minimum
184 distance to forest edge was calculated from the MapBiomias Perú 2022 dataset.²⁷ We retained
185 forest classes (forest, dry forest, mangrove, and flooded forest) and removed all other classes.
186 These forest cells were transformed into vector point data for the distance calculation. For each
187 household, we calculated the minimum Euclidean distance to the nearest forest within a 5-

188 kilometer (km) radius around each household. There were no households for which the minimum
189 distance to forest edge was greater than 5km. Population density was calculated from the census
190 data. A population density raster was created by summing up the number of individuals in each
191 household with 1km by 1km resolution. Population density values were then extracted for each
192 household GPS point.

193

194 **Statistical analysis**

195 We published a pre-analysis plan at <https://osf.io/fgr6w>. Deviations from the pre-analysis plan
196 are listed in Table S1. All positive cases were matched to weather data. Thus, we performed a
197 complete case analysis.

198

199 We assessed associations between malaria incidence and weather variables using distributed lag
200 models. We chose a 2–16 week lookback period (i.e. an infection during the course of
201 epidemiologic week 20 looked back to epidemiologic weeks 2—18) to account for potential
202 variation in the influence of rainfall and temperature in *Pv* transmission and to account for
203 potential impacts on relapse cases (Figure 1). For continuous exposures (weekly maximum
204 temperature, weekly mean temperature, weekly mean temperature range, weekly minimum
205 temperature, and total precipitation), we fit non-linear distributed lag (DL) models²⁸ with a log
206 link and Poisson family using weekly malaria case counts per village as the dependent variable
207 with an offset for log community population size. For the distributed lag cross-basis function, we
208 specified a log function with 2 knots to allow for more flexible variation in the short term, and
209 natural splines with 2 degrees of freedom for the non-linear predictors, which yielded the lowest
210 AIC and BIC in model testing. All incidence ratios were calculated relative to reference weeks.
211 For temperature predictors, weeks with the minimum observed value during the study period
212 were used as the reference. For total precipitation, weeks with 0 mm of precipitation were used
213 as the reference. Continuous temperature variables were adjusted for the cross-basis matrix for
214 total precipitation, and total precipitation was adjusted for the cross-basis matrix for maximum
215 temperature.

216

217 For binary predictors (high heat and heavy rain) we fit generalized additive models (GAMs) with
218 a Poisson family and log link and an offset for community population size. We fit separate
219 models at each lag from 2–16 weeks. We adjusted the high heat model for total precipitation at
220 the concurrent lag modeled as a smooth term with $k = 3$. Similarly, we adjusted the heavy rain
221 model for maximum temperature at the concurrent lag modeled as a smooth term with $k = 3$.

222

223 To examine the influence of other environmental factors, we conducted subgroup analyses by:
224 distance to forest edge (above/below the median), season (rainy/dry), and community type
225 (riverine/dry land). We discretized the distance from each household to the nearest forest edge
226 into a binary variable for above or below median distance, and categorized months November-
227 April as the rainy season and all other months as the dry season based on both historical

228 precedence⁷ and data for our study. For distance to forest edge models, we aggregated incidence
229 per household-week and adjusted for 1km population density. We fit the same DL models as the
230 main effect models on subsets of the data according to the subgroup, and non-forest models were
231 offset for community size. We also assessed the association between the minimum distance to
232 forest edge as both a continuous and categorical predictor and fit GAMs with Poisson
233 distributions and log link functions.

234

235 For the association between maximum temperature and malaria incidence, we conducted two
236 sensitivity analyses with additional adjustments. First, we adjusted for both total precipitation
237 and minimum temperature at concurrent lags to account for potential collinearity between
238 minimum temperature and maximum temperature (Pearson's correlation coefficient = 0.48).
239 Second, we adjusted for minimum temperature at concurrent lags and total precipitation at 4-
240 week delayed lags, where we only considered the effect of precipitation beginning at 6 weeks
241 and continuing through 20 weeks. This time-delayed lag was included to account for the onset of
242 the rainy season about 4 weeks after the highest maximum temperatures were typically observed
243 (Figure 3).

244

245 For the association between total precipitation and malaria incidence, we conducted a sensitivity
246 analysis where we adjusted for minimum temperature instead of maximum temperature. The
247 Pearson's correlation coefficient for total precipitation and maximum temperature was -0.30, and
248 thus we kept this as the main effect adjustment covariate, while the Pearson's correlation
249 coefficient for total precipitation and minimum temperature was 0.1.

250

251 Because incidence data for the weather models was aggregated at the community level, we did
252 not test for spatial autocorrelation or adjust for spatial clustering. All hourly to daily aggregation
253 was performed using the Python API for Google Earth Engine on Google Colab servers. Weekly
254 aggregation and modeling were performed in R version 4.2.1. All distributed lag models were
255 built using the R package DLNM version 2.4.7²⁹. This study was approved by the Institutional
256 Review Board of Stanford University (72291) and by the Dirección Universitaria de Asuntos
257 Regulatorios de la Investigación de la Universidad Peruana Cayetano Heredia (211747).

258

259

260 **RESULTS**

261 During the study period from January 2021 to December 2023, the rainy season generally lasted
262 from November through April (Figure 3). Weekly cumulative precipitation ranged between 500–
263 1750 mm during the rainy season and did not exceed 500mm in the dry season which lasted
264 roughly from July through October.

265

266 Minimum temperature varied from 22–26°C, and maximum temperature varied from 28–38°C.

267 The weekly mean temperature range varied from 4–12°C. During the dry season, there was more

268 variation in temperature: minimum temperature dipped below 20°C in the early dry season, while
269 temperature peaked during the late dry season, with maximum temperature values as high as
270 38°C.

271
272 The mean threshold for high heat events across all study years was 27.5°C, and high heat events
273 generally occurred just prior to the rainy season. The mean threshold for heavy rain events across
274 all study years was 186 mm, and there were more heavy rain events during the rainy season than
275 the dry season. 2023 was an El Niño year, reflected by the low precipitation and high
276 temperature compared to 2022 and 2021. Malaria incidence did not appear to follow a distinct
277 seasonal trend.

278 279 **Temperature**

280 Associations between minimum or maximum temperature and malaria incidence are shown in
281 Figure 4. Weeks with a minimum temperature of 22.5°C (the median) were associated with 2.16–
282 3.93-fold higher malaria incidence for lags 3–16 weeks compared to weeks with the lowest
283 minimum temperature of 16.8°C, with the incidence ratio peaking at lag of 5 weeks (IR = 3.93,
284 95% CI 2.18–7.09) (Figure 4A, 4C). Malaria incidence was approximately 3- to 4-fold higher for
285 minimum weekly temperatures above 16.8°C at a 5-week lag (Figure 4B).

286
287 The effect of median maximum temperature on incidence was non-linear and hump-shaped
288 across lag time (Figure 4D). Median maximum temperature was associated with higher incidence
289 at lags of 6–13 weeks, whereas it was associated with lower incidence at the extremes of 3 weeks
290 and 14–16 weeks. Compared to weeks with the lowest maximum temperature (28.4°C), weeks
291 with a maximum temperature of 31.8°C (the median) were associated with 66% higher malaria
292 incidence (95 CI 1.35–2.04) at a 9-week lag (Figure 4D, 4F). The positive association between
293 median maximum temperature and malaria incidence grew stronger as maximum temperatures
294 increased to 32.5°C, then plateaued and confidence intervals were wide (Figure 4E).

295
296 In a sensitivity analysis adjusting for concurrent total precipitation and minimum temperature,
297 compared to weeks with the lowest maximum temperature, weeks with a maximum temperature
298 of 31.8°C (the median) were still associated with higher malaria incidence at 6–11 week lags
299 (Figure S1). There was no association at other lags. In another sensitivity analysis adjusting for
300 minimum temperature at concurrent lags and total precipitation at 4-week delayed lags, weeks
301 with a maximum temperature of 31.8°C (the median) were still associated with higher malaria
302 incidence at 6–9 week lags (Figure S2) and were not associated with incidence at other lags.

303
304 Weeks with high heat events (one or more days when the daily mean temperature exceeded
305 27.5°C) were associated with 35% lower malaria incidence at a 2-week lag and 1.23–1.37-fold
306 higher malaria incidence at lags 5–9 weeks compared to weeks with no high heat events (Figure
307 5A, Table S2).

308
309 For analyses of mean temperature, compared to weeks with the lowest value (23.1°C), weeks
310 with a mean temperature of 25.6°C (the median) were associated with 10–44% lower malaria
311 incidence for lags 2–8 weeks (Figure S3A, S3C), and the association was strongest at a 3-week
312 lag (IR = 0.66, 95% CI 0.51–0.84). The protective association was stronger at higher mean
313 temperatures at a 3-week lag (Figure S3B).

314
315 For analyses of temperature range, compared to weeks with the narrower ranges of 4.3°C, weeks
316 with a temperature range of 6.5°C (the median) were associated with 1.33- to 1.94-fold higher
317 malaria incidence for lags 6–14 weeks (Figure S4A, S4C), with the strongest association at a 9-
318 week lag (IR = 1.94, 95% CI 1.58–2.38). The positive association was stronger at wider mean
319 temperature ranges at a 9-week lag (Figure S4B).

320 321 **Precipitation**

322 The association between the effect of precipitation median precipitation on incidence was non-
323 linear and hump-shaped across lag time (Figure 6A). Compared to weeks with no rainfall, weeks
324 with 450mm of precipitation (the median) were associated with 1.13- to 1.45-fold higher malaria
325 incidence for lags 6–14 weeks, with the strongest association at 9 weeks (IR = 1.45, 95% CI
326 1.27–1.65) (Figure 6A, 6C). The association was also non-linear and hump-shaped across the
327 range of precipitation values, with the strongest associations with malaria incidence at
328 approximately 750mm of precipitation at a 9-week lag (Figure 6B). In a sensitivity analysis
329 adjusting for minimum temperature instead of maximum temperature, associations were still
330 positive but somewhat attenuated towards the null (Figure S5).

331
332 Compared to weeks without heavy rain, weeks with heavy rain events (one or more days when
333 the daily precipitation exceeded 186mm) were associated with 1.22–1.60-fold higher malaria
334 incidence at lags 2-10 weeks, with the strongest association at 6 weeks (IR = 1.60, 95% CI 1.33–
335 1.94) (Figure 5B). Heavy rain was associated with 24% lower malaria incidence at 16 weeks
336 (95% CI 0.63–0.92).

337 338 **Sub-group analyses**

339 In a sub-group analysis by season, high heat events were protective in the dry season and
340 associated with higher incidence in the rainy season at longer lags (Figure S6); heavy rain events
341 were associated with higher incidence in the rainy season at up to 11-week lags, while results
342 were generally null for the dry season (Figure S6).

343
344 We also performed a sub-group analysis by community type; 13 communities were classified as
345 dryland communities, while 16 were riverine. High heat events were associated with higher
346 incidence in dryland communities and lower incidence in riverine communities at longer lags
347 (Figure S7); heavy rain events were generally associated with higher incidence in dryland

348 communities across all lags, but lower incidence in riverine communities at longer lags (Figure
349 S7).

350

351 At most lags, the association between malaria incidence and maximum temperature or total
352 precipitation did not vary by a household's minimum distance to forest edge (Figure S8, S9).
353 Shorter minimum distance to forest edge was associated with higher malaria incidence (Figure
354 S10). For households living within 50m of forest edge, malaria incidence was 2.18-fold higher
355 (95% CI 2.18–2.19) compared to those who lived 500m or beyond from forest edge (Figure
356 S1A).

357

358 All main effect continuous predictor estimates can be found in Table S2. All main effect binary
359 predictor estimates can be found in Table S3.

360

361

362 **DISCUSSION**

363 In this study of associations between temperature and precipitation with malaria incidence in
364 Loreto Region, Peru, we found that higher minimum temperatures, maximum temperatures,
365 temperature ranges, and total precipitation were all associated with higher malaria incidence
366 compared to weeks with minimum values. Mean temperature was generally associated with
367 lower malaria incidence in the short (2–4 weeks) and medium (5–10 weeks) term. The effect
368 of these weather variables generally lasted over many weeks, with the longest duration and
369 strongest associations with minimum temperature. The effect of minimum temperature also
370 began sooner and lasted longer than that of maximum temperature or temperature range. This
371 may suggest that cooler minimum temperatures are a limiting factor for malaria transmission.
372 High heat events were associated with lower malaria incidence in the short term and higher
373 malaria incidence in the medium term. Heavy rain events were generally associated with
374 increased malaria risk in the short and medium term, and lower malaria incidence in the long
375 term (11–16 weeks). There was strong evidence that living further away from forest edges was
376 protective. Overall, findings suggest that higher temperatures, particularly minimum
377 temperatures, and periods of sustained rainfall could lead to periods of increased malaria burden.

378

379 In general, our findings suggest that weather influences malaria transmission over 1-4 month lags
380 in this region. This lasting association may be due to the relapse periodicity of *Pv* malaria. Even
381 after typical treatment, 6-month relapse infection rates in this region ranged between 27–33%³⁰,
382 but true relapse rates are likely higher because many *Pv* infections go undetected and untreated,
383 or are treated without medical supervision, outside of study settings³¹. However, the vast
384 majority of *Pv* infections are estimated to be relapse infections³², and thus it is likely that
385 increased incidence includes relapse cases – another study in this region found that PCR-
386 confirmed *Pv* prevalence was as high as 25%, most of which were attributed to asymptomatic,
387 low parasitemia infections.³³ Additionally, though the relapse period for *Pv* infections is poorly

388 understood in this region, relapses from the tropical strains of *Pv* are estimated to range
389 anywhere from 2–9 months.¹⁶ Though strong geographic and micro-geographic differences exist
390 in the Peruvian Amazon,³⁴ a study in neighboring Brazil found that the median time to the first
391 recurrent infection was 71 days, and that treatment delays the onset of the subsequent relapse
392 infection, while in Southeast Asia, where short latency *Pv* is also present, 90% of relapse cases
393 occurred within 6 weeks of the initial infection.³⁵ The extended association with weather over
394 many weeks could also reflect a high prevalence of asymptomatic and low parasitemia
395 infections^{4,5}, which could sustain new, onward transmission well past the initial meteorological
396 trigger.

397

398 **Precipitation**

399 We found that weeks with higher precipitation were significantly associated with higher malaria
400 incidence at lags of 6–14 weeks. These results are generally in accordance with other studies of
401 the effect of rainfall on *Pv* infections. A meta-analysis in Mauritania found that *Pv* incidence was
402 highest during and after the rainy season³⁶ and that decreased rainfall was significantly correlated
403 with decreased malaria burden; in temperate regions, one study in South Korea found that
404 increased precipitation was associated with higher malaria incidence at a 10-week lag,³⁷ while a
405 similar study in China also found positive associations with precipitation at lags 2–4 weeks.³⁸
406 The lasting influence of precipitation likely reflects increased breeding ground following rainfall,
407 which could impact multiple transmission cycles.

408

409 Interestingly, heavy rain was generally associated with higher malaria incidence in the rainy
410 season, but not in the dry season (Figure S8B). There were few heavy rain events in the dry
411 season, thus limiting the statistical power of this analysis; however, our finding for the rainy
412 season somewhat aligns with a hydrogeology hypothesis by Hiwas and Bretas⁹, wherein heavy
413 rain in the rainy season may flood otherwise dry areas, increasing malaria breeding ground. We
414 also found some evidence of qualitative effect modification by community type, with heavy rain
415 associated with lower incidence in riverine communities at longer lags and higher incidence at
416 longer lags in dryland communities (Figure S9B). This finding contradicts the hydrogeology
417 hypothesis⁹. In riverine communities, heavy rains may be protective if they flush and agitate
418 otherwise slow-moving rivers and stagnant pools of water and thus disrupt mosquito larva
419 breeding; on the other hand, in dryland communities, heavy rains may increase risk due to the
420 accumulation of surface water in otherwise dry areas. In both cases, the effect modification is
421 strongest at longer lags, signaling that heavy rain had long-lasting effects on the transmission
422 cycle.

423

424

425 **Temperature**

426 We found that higher minimum temperatures were strongly associated with higher malaria
427 incidence consistently across lags, while maximum temperature and high heat events were

428 associated with higher malaria incidence in the medium term and lower incidence in the short
429 and long term. Though the thermal optimum of the *Pv*–*An. darlingi* coupling has not been
430 studied directly, there is some mixed support in the literature from other malaria transmission
431 couplings: optimal *Pf* malaria transmission is thought to peak at 25°C,^{10,17} which generally aligns
432 with our findings for positive associations with higher minimum temperature, which were in the
433 range of 20–24°C. Our finding of the strongest associations with minimum temperature are
434 biologically plausible given that *An. darlingi* typically bite humans at nighttime³⁹ and daily
435 temperatures are lowest at night.

436

437 We were surprised to find that higher maximum temperatures in the range of 30–34°C and high
438 heat events (>27°C) were associated with increased incidence at medium lags and lower
439 incidence at short lags. Our maximum temperature findings were attenuated, but robust in
440 sensitivity analyses. High heat events may be correlated with other longer-term weather
441 variations, such as higher maximum temperature for prolonged periods, that affect incidence.
442 High temperatures can reduce the vector biting rate and survival rate, which may explain the
443 negative associations at shorter lags. However, high maximum temperatures may trigger *Pv*
444 relapse and increase the reservoir of active infections that contribute to onward malaria
445 transmission, explaining the positive association at medium lags. The positive association
446 between higher maximum temperature and high heat and malaria at medium lags conflicts with
447 previous disease ecology literature supporting a thermal optimum for *Anopheles-Pf* malaria
448 transmission at 25°C with an upper limit of 32.6°C,¹⁰ and validated *Pf* malaria transmission
449 declines above 28°C¹⁷ in Africa. Our result does have some support in the literature: a study in
450 neighboring regions of the Amazon with similar maximum temperatures (26.8–35.2°C) found
451 that a 1°C increase in maximum temperature was associated with higher incidence at 1- and 2-
452 month lags.⁴⁰ Generally, however, *Anopheles* life stages related to transmission (i.e., biting rate,
453 larva survival, fecundity, adult survival) are thought to decline at the median maximum
454 temperature range in our study (30–34°C),¹¹ indicating that vector activity is unlikely to be the
455 mechanism driving this positive association. Further research on the thermal biology of *Pv*–*An.*
456 *darlingi* in tropical settings are needed to elucidate the influence of temperature on malaria
457 transmission in this region. However, it is also possible that microclimates and small-scale
458 ecological and topographical variation in our study sites tempered high temperatures and their
459 associated negative impacts on transmission, since our weather data was at 11km resolution and
460 did not capture temperature variation more finely (i.e. in shaded forest areas, close to water
461 bodies)

462

463 One alternative explanation is that associations with maximum temperature primarily reflect
464 relapsing *Pv* infections, which could be triggered by temperature itself or co-infections that are
465 more common at higher temperatures. Extrinsic triggers of *Pv* relapse include co-infections that
466 result in host inflammation, subsequent primary *Pv* infections, seasonal changes in sunlight and
467 temperature, and mosquito bites and their associated immune responses^{16,41}. In our study site,

468 dengue is increasingly common,⁴² and it is well-documented that dengue transmission is
469 positively associated with maximum temperatures similar to those in our study^{43–45} and has a
470 higher thermal optimum (29°C) and upper limit (34.5°C) than malaria.¹⁰ If high heat events
471 trigger *Pv* relapse, symptoms would likely appear within 2 weeks, yet we found positive
472 associations with maximum temperature at lags of 6–11 weeks. It is possible that high
473 temperatures activate the hypnozoite and increase the reservoir of active infectious hosts,
474 potentially leading to onward transmission and multiple cycles of infection, explaining the
475 delayed positive association.

476
477 Relationships with high heat varied by season and community type. At lags of 3–4 months, high
478 heat was associated with higher malaria incidence in the dry season but lower malaria incidence
479 in the rainy season. High heat events may dry up surface water in the dry season, reducing vector
480 breeding, while they may lead to stagnation of water bodies in the rainy season, promoting
481 breeding. Similarly, we found that high heat was associated with higher malaria incidence in
482 dryland areas, but lower malaria incidence in riverine areas at lags of 3–4 months. In general,
483 the dryland communities in our study site were more peri-urban and more densely populated;
484 considering dengue transmission is generally thought to thrive in more urban, densely populated
485 communities,^{46,47} it is possible that elevated incidence reflects higher relapse rates due to co-
486 infection with diseases like dengue.

487 488 **Limitations**

489 Our study had several limitations. We were unable to distinguish between initial and relapse
490 infections in our incidence data; we would expect that weather would influence initial and
491 relapse cases over different lag periods, but our analysis was not able to investigate this.
492 However, using distributed lag models allowed us to investigate associations over a 4-month
493 period, including potential relapse cases. A follow-up study using genomic methods to
494 differentiate relapse and initial infections could shed light on how weather influences primary vs.
495 relapse infections. Our outcome data also only covered a 3-year period, preventing this analysis
496 from studying longer climatic trends in incidence in this region. The study period also included
497 two years (2021 and early 2022) where control efforts for the COVID-19 pandemic likely limited
498 malaria transmission as well. We were also unable to control for PMC activities nor
499 interventions over the study period.

500
501 Regarding the weather data, one of our study years (2023) was an El Niño year; a study
502 capturing a full El Niño-La Niña cycle could better elucidate associations with incidence along
503 these more variable climate cycles. While we selected ERA-5 Land remote sensing data for its
504 temporal coverage, its 11 km spatial resolution prevented identification of small-scale
505 microclimates, i.e. those created by forest cover and topographical features, and thus did not
506 capture small-resolution variations that likely have a large effect on mosquito breeding and
507 survival habitat⁴⁸. Additionally, publicly available surface water data did not reflect our ground

508 observations of surface water in the study site, so we did not include it in this analysis. Further
509 research that considers mediation by surface water could further shed light on the complicated
510 dynamics between precipitation and malaria incidence in this region. One such complication that
511 our study did not resolve is the concentration-dilution hypothesis⁹ for malaria transmission,
512 which posits that heavy rain in riverine communities can cause flooding and increase standing
513 water presence, leading to increased incidence, while the same heavy rain can flush dryland
514 communities and lead to decreased incidence. Our sub-group analyses by distance to forest edge,
515 community type, and season were also likely limited in statistical power; these findings could be
516 more conclusive with a longer look back period. Furthermore, our binary predictor analyses
517 modeled individual lags separately and did not account for time autocorrelation, limiting
518 comparisons between adjacent lags and with DL models. This study was also correlational, and
519 inferred relationships with individual weather variables may be confounded by collinearity and
520 correlation with other weather variables, and thus makes it difficult to tease out individual direct
521 effects.

522

523 **Conclusions**

524 In our study of malaria incidence in Loreto Region, Peru, we observed generally positive
525 associations with higher temperatures and higher rainfall for extended lag periods beginning 2–3
526 weeks after symptom onset and enduring for about 1–4 months. Our findings indicated that the
527 coupled transmission and relapse cycle of *Pv-An. darlingi* may have more complicated
528 associations with higher temperatures than other malaria parasite – vector pairings, a critical
529 finding in the face of climate change and global warming. These findings provide critical context
530 to ongoing malaria elimination efforts, since apparent successes or failures of malaria
531 interventions may be due in part to long-lasting effects of weather on initial infection and
532 relapse.

533

REFERENCES

- 534 1 Kattenberg JH, Erhart A, Truong MH, *et al.* Characterization of *Plasmodium falciparum* and
535 *Plasmodium vivax* recent exposure in an area of significantly decreased transmission intensity
536 in Central Vietnam. *Malaria Journal* 2018; **17**: 180.
- 537 2 Global technical strategy for malaria 2016–2030, 2021 update. 2021.
- 538 3 Hulden L, Hulden L. Activation of the hypnozoite: a part of *Plasmodium vivax* life cycle and
539 survival. *Malar J* 2011; **10**: 90.
- 540 4 Olliaro PL, Barnwell JW, Barry A, *et al.* Implications of *Plasmodium vivax* Biology for
541 Control, Elimination, and Research. *Am J Trop Med Hyg* 2016; **95**: 4–14.
- 542 5 da Silva-Nunes M, Moreno M, Conn JE, *et al.* Amazonian malaria: Asymptomatic human
543 reservoirs, diagnostic challenges, environmentally-driven changes in mosquito vector
544 populations, and the mandate for sustainable control strategies. *Acta Trop* 2012; **121**: 281–91.
- 545 6 Howes RE, Battle KE, Mendis KN, *et al.* Global Epidemiology of *Plasmodium vivax*. *Am J*
546 *Trop Med Hyg* 2016; **95**: 15–34.
- 547 7 Rosas-Aguirre A, Gamboa D, Manrique P, *et al.* Epidemiology of *Plasmodium vivax* Malaria
548 in Peru. *Am J Trop Med Hyg* 2016; **95**: 133–44.
- 549 8 Sadoine ML, Smargiassi A, Ridde V, Tusting LS, Zinszer K. The associations between
550 malaria, interventions, and the environment: a systematic review and meta-analysis. *Malar J*
551 2018; **17**: 73.
- 552 9 Hiwat H, Bretas G. Ecology of *Anopheles darlingi* Root with respect to vector importance: a
553 review. *Parasites Vectors* 2011; **4**: 177.
- 554 10Mordecai EA, Caldwell JM, Grossman MK, *et al.* Thermal biology of mosquito-borne
555 disease. *Ecol Lett* 2019; **22**: 1690–708.
- 556 11Villena OC, Ryan SJ, Murdock CC, Johnson LR. Temperature impacts the environmental
557 suitability for malaria transmission by *Anopheles gambiae* and *Anopheles stephensi*. *Ecology*
558 2022; **103**: e3685.
- 559 12Gething PW, Van Boeckel TP, Smith DL, *et al.* Modelling the global constraints of
560 temperature on transmission of *Plasmodium falciparum* and *P. vivax*. *Parasites Vectors* 2011;
561 **4**: 92.
- 562 13Gilman RH, Cabrera L, Lee E, *et al.* ENDEMIC MALARIA IN THE PERUVIAN AMAZON
563 REGION OF IQUITOS. *The American Journal of Tropical Medicine and Hygiene* 2003; **69**:
564 45–52.
- 565 14Stratman-Thomas WK. The Influence of Temperature on *Plasmodium Vivax*. *The American*
566 *Journal of Tropical Medicine and Hygiene* 1940; **s1-20**: 703–15.

- 567 15Suh E, Grossman MK, Waite JL, *et al.* The influence of feeding behaviour and temperature on
568 the capacity of mosquitoes to transmit malaria. *Nat Ecol Evol* 2020; **4**: 940–51.
- 569 16White NJ. Determinants of relapse periodicity in *Plasmodium vivax* malaria. *Malar J* 2011;
570 **10**: 297.
- 571 17Mordecai EA, Paaijmans KP, Johnson LR, *et al.* Optimal temperature for malaria transmission
572 is dramatically lower than previously predicted. *Ecology Letters* 2013; **16**: 22–30.
- 573 18Schäfer C, Zanghi G, Vaughan AM, Kappe SHI. *Plasmodium vivax* Latent Liver Stage
574 Infection and Relapse: Biological Insights and New Experimental Tools. *Annu Rev Microbiol*
575 2021; **75**: 87–106.
- 576 19malaria.pdf. <https://www.dge.gob.pe/portal/docs/vigilancia/sala/2022/SE42/malaria.pdf>
577 (accessed May 2, 2024).
- 578 20Bannister-Tyrrell M, Verdonck K, Hausmann-Muela S, Gryseels C, Muela Ribera J, Peeters
579 Grietens K. Defining micro-epidemiology for malaria elimination: systematic review and
580 meta-analysis. *Malar J* 2017; **16**: 164.
- 581 21Rosas-Aguirre A, Guzman-Guzman M, Gamboa D, *et al.* Micro-heterogeneity of malaria
582 transmission in the Peruvian Amazon: a baseline assessment underlying a population-based
583 cohort study. *Malar J* 2017; **16**: 312.
- 584 22Paaijmans KP, Blanford S, Bell AS, Blanford JI, Read AF, Thomas MB. Influence of climate
585 on malaria transmission depends on daily temperature variation. *Proceedings of the National*
586 *Academy of Sciences* 2010; **107**: 15135–9.
- 587 23MacDonald AJ, Mordecai EA. Amazon deforestation drives malaria transmission, and malaria
588 burden reduces forest clearing. *Proceedings of the National Academy of Sciences* 2019; **116**:
589 22212–8.
- 590 24Janko MM, Recalde-Coronel GC, Damasceno CP, *et al.* The impact of sustained malaria
591 control in the Loreto region of Peru: a retrospective, observational, spatially-varying
592 interrupted time series analysis of the PAMAFRO program. *The Lancet Regional Health –*
593 *Americas* 2023; **20**. DOI:10.1016/j.lana.2023.100477.
- 594 25189846_RM_244-2017-MINSA.PDF20180823-24725-1yk3i0y.pdf.
595 [https://cdn.www.gob.pe/uploads/document/file/190350/189846_RM_244-2017-](https://cdn.www.gob.pe/uploads/document/file/190350/189846_RM_244-2017-MINSA.PDF20180823-24725-1yk3i0y.PDF)
596 [MINSA.PDF20180823-24725-1yk3i0y.PDF](https://cdn.www.gob.pe/uploads/document/file/190350/189846_RM_244-2017-MINSA.PDF20180823-24725-1yk3i0y.PDF) (accessed Nov 21, 2024).
- 597 26Muñoz-Sabater J, Dutra E, Agustí-Panareda A, *et al.* ERA5-Land: a state-of-the-art global
598 reanalysis dataset for land applications. *Earth System Science Data* 2021; **13**: 4349–83.
- 599 27MapBiomias Peru Project. Collection 2 of the Annual Land Cover and Land Use Series for
600 Peru. [https://storage.googleapis.com/mapbiomas-](https://storage.googleapis.com/mapbiomas-public/initiatives/peru/collection_2/coverage/peru_coverage_2022.tif)
601 [public/initiatives/peru/collection_2/coverage/peru_coverage_2022.tif](https://storage.googleapis.com/mapbiomas-public/initiatives/peru/collection_2/coverage/peru_coverage_2022.tif) (accessed Oct 13, 2023).

- 602 28Gasparri A, Armstrong B, Kenward MG. Distributed lag non-linear models. *Stat Med* 2010;
603 29: 2224–34.
- 604 29Gasparri A. Distributed Lag Linear and Non-Linear Models in R: The Package dlnm. *J Stat*
605 *Softw* 2011; 43: 1–20.
- 606 30Llanos-Cuentas A, Lacerda MVG, Hien TT, *et al.* Tafenoquine versus Primaquine to Prevent
607 Relapse of Plasmodium vivax Malaria. *New England Journal of Medicine* 2019; 380: 229–41.
- 608 31Poespoprodjo JR, Burdam FH, Candrawati F, *et al.* Supervised versus unsupervised
609 primaquine radical cure for the treatment of falciparum and vivax malaria in Papua, Indonesia:
610 a cluster-randomised, controlled, open-label superiority trial. *The Lancet Infectious Diseases*
611 2022; 22: 367–76.
- 612 32Popovici J, Tebben K, Witkowski B, Serre D. Primaquine for *Plasmodium vivax* radical cure:
613 What we do not know and why it matters. *International Journal for Parasitology: Drugs and*
614 *Drug Resistance* 2021; 15: 36–42.
- 615 33Carrasco-Escobar G, Miranda-Alban J, Fernandez-Miño C, *et al.* High prevalence of very-
616 low Plasmodium falciparum and Plasmodium vivax parasitaemia carriers in the Peruvian
617 Amazon: insights into local and occupational mobility-related transmission. *Malar J* 2017; 16:
618 415.
- 619 34Chuquiyauri R, Peñataro P, Brouwer KC, *et al.* Microgeographical Differences of
620 Plasmodium vivax Relapse and Re-Infection in the Peruvian Amazon. *Am J Trop Med Hyg*
621 2013; 89: 326–38.
- 622 35Chu CS, White NJ. Management of relapsing Plasmodium vivax malaria. *Expert Rev Anti*
623 *Infect Ther* 2016; 14: 885–900.
- 624 36Moustapha IE, Moukah MO, Salem MSOA, *et al.* Malaria prevalence in Mauritania: a
625 systematic review and meta-analysis. *Malaria Journal* 2023; 22: 146.
- 626 37Kim Y-M, Park J-W, Cheong H-K. Estimated Effect of Climatic Variables on the
627 Transmission of Plasmodium vivax Malaria in the Republic of Korea. *Environ Health*
628 *Perspect* 2012; 120: 1314–9.
- 629 38Bi Y, Yu W, Hu W, *et al.* Impact of climate variability on Plasmodium vivax and Plasmodium
630 falciparum malaria in Yunnan Province, China. *Parasit Vectors* 2013; 6: 357.
- 631 39Moreno M, Saavedra MP, Bickersmith SA, *et al.* Implications for changes in Anopheles
632 darlingi biting behaviour in three communities in the peri-Iquitos region of Amazonian Peru.
633 *Malar J* 2015; 14: 290.
- 634 40Wangdi K, Wetzler E, Cox H, Marchesini P, Villegas L, Canavati S. Spatial patterns and
635 climate drivers of malaria in three border areas of Brazil, Venezuela and Guyana, 2016–2018.
636 *Scientific Reports* 2022; 12. DOI:10.1038/s41598-022-14012-4.

- 637 41 Schäfer C, Zanghi G, Vaughan AM, Kappe SHI. Plasmodium vivax Latent Liver Stage
638 Infection and Relapse: Biological Insights and New Experimental Tools. *Annual Review of*
639 *Microbiology* 2021; **75**: 87–106.
- 640 42 Munayco CV. Notas Desde el Campo: Brote de Dengue — Perú, 2023. *MMWR Morb Mortal*
641 *Wkly Rep* 2024; **73**. https://www.cdc.gov/mmwr/volumes/73/wr/mm7304a4_ensp.htm
642 (accessed Oct 27, 2024).
- 643 43 López MS, Gómez AA, Müller GV, Walker E, Robert MA, Estallo EL. Relationship between
644 Climate Variables and Dengue Incidence in Argentina. *Environmental Health Perspectives*
645 2023; **131**: 057008.
- 646 44 Fan J, Wei W, Bai Z, *et al.* A Systematic Review and Meta-Analysis of Dengue Risk with
647 Temperature Change. *International Journal of Environmental Research and Public Health*
648 2015; **12**: 1–15.
- 649 45 Hii YL, Rocklöv J, Ng N, Tang CS, Pang FY, Sauerborn R. Climate variability and increase
650 in intensity and magnitude of dengue incidence in Singapore. *Global Health Action* 2009; **2**:
651 2036.
- 652 46 Koyadun S, Butraporn P, Kittayapong P. Ecologic and Sociodemographic Risk Determinants
653 for Dengue Transmission in Urban Areas in Thailand. *Interdisciplinary Perspectives on*
654 *Infectious Diseases* 2012; **2012**: 907494.
- 655 47 Misslin R, Telle O, Daudé E, Vaguet A, Paul RE. Urban climate versus global climate
656 change—what makes the difference for dengue? *Annals of the New York Academy of Sciences*
657 2016; **1382**: 56–72.
- 658 48 Carrasco-Escobar G, Manrique E, Ruiz-Cabrejos J, *et al.* High-accuracy detection of malaria
659 vector larval habitats using drone-based multispectral imagery. *PLoS Negl Trop Dis* 2019; **13**:
660 e0007105.

661

662

663 **Figure 1. Timeline of main hypothesized mechanisms of the impact of temperature and**
664 **precipitation on *Plasmodium vivax* infection.** Time estimates are generalized and were generated from
665 previous findings.


666
667 **Figure 2. Study area map.** Dark green areas indicate forest cells. Circles with black outlines represent
668 the 30 study communities, colored by annual malaria incidence per 1000 individuals. The blue river to the
669 right of Iquitos is the Amazon River.


670
671 **Figure 3. Weather and malaria incidence trends over the study period.** All measures are aggregated
672 weekly. The blue dots in plot A mark weeks with a heavy rain event, defined as one or more days when
673 the daily precipitation total exceeded the 90th percentile, which averaged 186mm over the study period.
674 The orange dots in plot B mark weeks with a high heat event, defined as one or more days when the mean
675 temperature exceeded the 90th percentile. Shaded blue regions represent the rainy season, defined as
676 November–April. In plot B, the darker orange line tracks the maximum temperature observed each week,
677 and the lighter orange line tracks the minimum temperature observed each week.

678
679 **Figure 4. Association between temperature and malaria incidence.** The associations were fit using
680 distributed lag non-linear models. Plots A–C show the association for minimum temperature, and plots
681 D–F show the association for maximum temperature. For all plots, incidence ratios are relative to
682 reference weeks with the minimum observed predictor value. Plots A and D show the association between
683 temperature and malaria incidence at the median predictor value compared to the reference. For minimum
684 temperature, the reference is 16.8°C; for maximum temperature, the reference is 28.4°C. For minimum
685 temperature, the median is 22.5°C. For maximum temperature, the median is 31.8°C.


686
687 **Figure 5. Association between binary predictors and malaria incidence.** Incidence ratios were
688 calculated for weeks with the event relative to weeks without the event. The average high heat threshold
689 over the study period was 27.5°C. The average heavy rain threshold over the study period was >186 mm
690 of precipitation in one day. High heat estimates at each lag were adjusted for total precipitation at the
691 same lag. Heavy rain estimates at each lag were adjusted for maximum temperature at the same lag.

692
693 **Figure 6. Association between total precipitation and malaria incidence.** The association was fit using
694 a distributed lag non-linear model. For all plots, incidence ratios are relative to reference weeks with 0mm
695 of precipitation. Plot A shows the association between precipitation and malaria incidence at the median
696 predictor value, 450mm, compared to the reference.






A) Total weekly precipitation (mm)



B) Minimum and maximum daily temperature (C)



C) Malaria incidence per 1000 person-weeks

