1

1 Original Article

² Systematic Evaluation of Adiposity Indices as Predictors of Hypertension Risk in Children

³ and Adolescents

- 4 Short title: Adiposity Indices in Pediatric Hypertension
- ⁵ **Authors:** Bowen Zhu (phD)^{*1,2,3}, Shumin Zhan (M.S.)^{*1,2}, Hui Shi (M.S.)^{*4}, Xingyun Wang(phD)²,
- ⁶ Jingwen Yue (M.S.)³, Jianfang Gao (M.S.)², Tongshuai Wang (phD)^{2,3}, Rui Wang (phD)², Xirong
- ⁷ Guo (MD, phD)^{$2^{#}$}, Junfen Fu (MD, phD)^{$1^{#}$}
- ⁸ 1 Department of Endocrinology, Children's Hospital, Zhejiang University School of Medicine,
- ⁹ National Clinical Research Center for Child Health, Hangzhou, 310051, China.
- ¹⁰ 2 Hongqiao International Institute of Medicine, Tongren Hospital, Shanghai Jiao Tong University

¹¹ School of Medicine, No.1111, XianXia Road, Shanghai 200336, China.

- ¹² 3 Department of Clinical Research Center, Tongren Hospital, Shanghai Jiao Tong University School
- ¹³ of Medicine, No.1111, XianXia Road, Shanghai 200336, China.
- ¹⁴ 4 Xinhua Hospital, Shanghai Jiao Tong University School of Medicine, 1665 Kongjiang Road,
- ¹⁵ *200092, Shanghai, China.*
- ^{*}Bowen Zhu, Shumin Zhan and Hui Shi equally contributed to the manuscript.

17 **# Corresponding author**:

- ¹⁸ Junfen Fu, MD, PhD, Department of Endocrinology, The Children's Hospital of Zhejiang
- ¹⁹ University School of Medicine, National Clinical Research Center for Child Health, 3333 NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

2

²⁰ Binsheng Road, Hangzhou 310052, China. Email: <u>fjf68@zju.edu.cn</u>; ORCID:
 ²¹ 0000-0001-6405-1251.

Xirong Guo, MD, PhD, Professor of Pedology, Hongqiao International Institute of Medicine,
Tongren Hospital, Shanghai Jiao Tong University School of Medicine, No.1111, XianXia Road,
Shanghai 200336, China. Tel/Fax: 021-52039999; Email: xrguo@shsmu.edu.cn

25

26 Abstract

Objective To systemically examine the concomitant impact of adiposity indices on
hypertension in children and adolescents.

Methods A community-based cohort study by China Health and Nutrition Survey (2009-2011)
included 262 children and adolescents aged 7-17. Anthropometric and lipid profiles were
analyzed for hypertension risk using logistic and cross-lagged models.

Results Of the 227 participants (mean age, 16.5 [2.3] years), 147 (53.9%) were boys. Over a 32 two-year period, 26.4% developed hypertension, 5.3% had elevated systolic blood pressure 33 (SBP), and 25.6% had elevated diastolic blood pressure (DBP). After adjusting for covariates, 34 higher body mass index (BMI), waist circumference (WC), hip circumference (HC), triceps 35 36 skinfold thickness (TST), and body roundness index (BRI) were associated with increased 37 hypertension risk. Incorporating these measures into a BP-based model improved the AUC for hypertension prediction from 0.551 (95% CI: 0.463, 0.640) to 0.670 (95% CI: 0.592, 0.748). A 38 longitudinal relationship were observed between these adiposity indices and hypertension 39 40 symptoms.

3

41 Conclusions The study suggests anthropometry parameters are superior to blood lipid 42 indicators in predicting the occurrence of hypertension in children. Abdominal obesity, as 43 indicated by WC, HC, and BRI, accounts for a significant portion of the risk for hypertension, 44 particularly in children aged 7 to 12 years.

Keywords: Adiposity Indices, Pediatric Hypertension, Children and adolescents, Cross-lagged
Model

47

48 **1.Introduction**

Hypertension affects about 4% of children and 11% of adolescents globally [1]. The early 49 50 identification and proper intervention of high blood pressure or hypertension during childhood are extremely crucial because these conditions are related to cardiovascular morbidity (left 51 ventricular hypertrophy and high carotid intima-media thickness) and mortality in adulthood 52 53 and are commonly associated with obesity [2-4]. A comprehensive analysis of 2033 studies spanning 154 countries has revealed a significant prevalence of overweight and obesity among 54 children and adolescents[5]. The pooled estimates indicate that 8.5% are overweight and 14.8% 55 are obese, which translates to roughly one in every five young individuals facing the challenge of 56 excess weight. China's rapid economic progress has brought about major lifestyle and societal 57 transformations, resulting in a surge in metabolic disorders. This trend is likely to have a 58 substantial effect on the cardiovascular disease (CVD) burden in the country. The number of 59 60 children aged 5 to 19 years with obesity is predicted to rise to 254 million by 2030 [6]. It is believed that obesity-related primary hypertension has contributed to the rise in hypertension 61 prevalence, which has doubled from 1% to 4% over the past three decades [7-8]. 62

4

Enhancing our comprehension of the anthropometric and biochemical indicators of adiposity 63 64 profiles in the context of hypertension is essential for the prevention and management of this condition in children and adolescents. In a cohort study of youths, a high normal body weight 65 above the 60th percentile of BMI for age was associated with an increased risk of hypertension 66 [9]. The suggested thresholds of 0.50 and 0.46 for two distinct pediatric groups have 67 demonstrated robust predictive capabilities for identifying individuals with two or more 68 cardiometabolic risk factors in populations from six countries [10]. Furthermore, innovative 69 anthropometric indices such as the Body Shape Index (BSI), Body Roundness Index (BRI), 70 Conicity Index (CI), and Visceral Fat Index (VFI) have been introduced to more accurately 71 pinpoint subcutaneous and visceral fat distribution. A meta-analysis with 21108 children and 72 adolescents confirmed that original ABSI and modified ABSI do not perform as well as 73 traditional anthropometric indicators, such as BMI and WC. BMI continues to be the most 74 75 effective indicator for screening high blood pressure (HBP) in pediatric populations[11]. Rui Chen et al. reported that WHtR and BRI can be recommended to identify hypertension, 76 dyslipidemia, abdominal obesity, and clustered cardio-metabolic risk factors (CMRFs) in 77 7-17-year-old teenagers [12]. However, the majority of studies have been conducted with adult 78 79 populations, and the interplay and individual impact of biochemical measures, anthropometric data, and composite metrics on hypertension remains to be fully elucidated. Furthermore, there 80 is a scarcity of data that can differentiate between the hypertension risk in childhood associated 81 82 with initial obesity-related metrics and the predictive capacity. In this study, we aimed to systemically investigate the temporal and bidirectional relationship between anthropometric 83 and biochemistry measures of obesity and hypertension symptoms and further explore the 84

5

85 prediction ability of obesity-related measures for hypertension status using a large cohort.

86 2. Materials and Methods

87 2.1 Study cohorts

88	This survey encompassed children and adolescents aged 7 to 17 years, with two follow-up visits
89	conducted between 2009 and 2011. Valid blood pressure (BP) data, comprising three repeated
90	readings, were obtained. Participants with missing BP values during the follow-up period were
91	excluded from the analysis. The data was sourced from the China Health and Nutrition Survey
92	(CHNS), an ongoing, open cohort study. The CHNS is a comprehensive, nationwide, prospective
93	cohort study that includes a representative sample of the Chinese population. Its primary
94	objective is to assess the impact of health, nutrition, and family planning policies and programs
95	implemented by both national and local governments. Further details regarding the study
96	design and data collection methodologies of the CHNS have been documented in prior
97	publications [13]. Blood samples were collected in the 2009 year, and data information on
98	biochemical indicators was obtained. Since blood samples were collected in 2009 year, a total of
99	768 participants with blood biomarkers information were collected in 2009. After excluding
100	those who had 3-day high protein intake (n=17), elevated SBP or DBP (defined as age- and
101	sex-specific 90th percentile, n=221), glucose above 7.0 mmol/L or HbA1c above 6.5% (n=8),
102	BMI above 35 kg/m ² (n=1), missing data of systolic BP or diastolic BP (n=133), 388 participants
103	were included in the formal analysis. After that, a total of 161 participants were excluded as
104	they had lost visits during the follow-up duration (<i>Supplementary Figure 1</i>). Demographic and
105	behavioral characteristics of participants included in the final analysis (n=227) were compared

6

with those of excluded participants (*Supplementary Table 1*). All data and samples were
collected after written informed consent was obtained from parents or caregivers for children
and adolescents aged 7 to 17 years. The CHNS was granted approval by the Institutional Review
Board at the University of North Carolina at Chapel Hill, as well as by the local Institutional
Review Board of Ethics Committee.

111 **2.2 Data collection**

A standardized structured questionnaire was administered by trained health staff to collect 112 socio-demographic variables (in 2009) including age, gender, urban-rural residence, 113 soft/sugared fruit drinks, total protein intake, total fat intake, total carbohydrate intake, and 114 total energy intake. Physical examinations of waist circumference (WC), hip circumference (HC), 115 height, weight, triceps skinfold thickness (TST), upper arm circumference (UAC), and BP were 116 117 performed by trained clinical staff. All individuals maintained a regular life pattern for at least 118 three days before blood sample collection and 12 ml of blood was collected (in three 4 ml tubes) on an empty stomach. Biomarker data collected in CHNS 2009 involves the release of 26 fasting 119 blood measures on individuals aged 7 and older [14]. The estimated glomerular filtration rate 120 121 (eGFR) was calculated using the Schwartz formula [15]. The individuals were categorized into groups, namely, residence, urban, and rural. Individual dietary intake for 3 consecutive days was 122 determined for every household member. This step has been achieved by asking individuals 123 each day to report all food consumed away from home on a 24-hour recall basis, and the same 124 125 daily interview has been used to collect at-home individual consumption. Soft/sugared fruit

7

drinks information was assessed by the question including "Drink soft/sugared fruit drinks?" or
"How often drink soft/sugared fruit drinks?".

128 2.3 Adiposity Indices Measures

129	Adiposity indices measures included body mass index (BMI), waist circumference (WC), hip
130	circumference (HC), triceps skinfold thickness (TST), waist to hip circumference ratio (WHR),
131	upper arm circumference (UAC), triglyceride (TG), high-density lipoprotein cholesterol (HDL-c),
132	blood glucose, serum uric acid (SUA), conicity index (CI), body shape index (BSI), visceral fat
133	index (VFI), body roundness index (BRI). BMI was calculated as weight(kg)/height ² (m ²) and was
134	converted into age and gender-specific BMI percentiles [16]. Abdominal obesity was defined as
135	waist circumference age- and sex-specific 90th percentile, determined by the cut-off points of
136	standards for Chinese children and adolescents [17]. Waist to height ratio (WHR) was calculated
137	as waist circumference(cm)/ height(cm). The calculation of the novel anthropometric indices is
138	described below, with the relevant reference cited at the end of each formula:
139	$CI = WC/0.109^* (Wt/Ht)^{1/2} [18]$
140	$BSI = WC * Wt^{-2/3} * Ht^{-5/6} [19]$
141	VFI=WC/ [39.68 + 1.88*BMI*TG/1.03] * [1.31/HDL-C] (for male),
142	VFI=WC/(36.58+1.89*BMI)*(TG/0.81)*(1.52/HDL-C) (for female) [20]

- 143 BRI = $364.2 365.5 * (1 ((0.5 * WC/\pi)^2/(0.5 * Ht)^2))^{0.5}$ [21]
- 144 The unit of TG, HDL-c, and LDL-c was mmol/L.

8

TG was measured using the GPO-PAP (Hitachi 7600, Kyowa, Japan). HDL-c was measured using
the Enzymatic method (Hitachi 7600, Kyowa, UK). Glucose was measured using the GOD-PAP
(Hitachi 7600, Randox, UK).

148 2.4 Diagnostic Criteria and Definitions of Outcome

Systolic and diastolic blood pressure values were determined based on the average of three 149 150 separate measurements. Participants' blood pressure status, classified as either normal or elevated, was ascertained using sex- and age-specific cutoff values that are applicable to Chinese 151 152 school-age children and adolescents. In accordance with Cook's criteria, elevated blood pressure is defined as a systolic or diastolic blood pressure equal to or greater than the 90th percentile 153 for age, sex, and height [22]. The evaluation of blood pressure outcomes was conducted using 154 155 three distinct categorization methods: hypertension (defined by elevated SBP or elevated DBP) versus non-hypertension; elevated SBP versus normal SBP; and elevated DBP versus normal 156 DBP. 157

158 2.5 Statistical Analysis

Data were presented as mean ± standard deviation (SD) and median with interquartile range (IQR) for continuous variables or number (percentage), for categorical variables. Data on demographics, anthropometric and biochemistry measures were compared using the chi-square test, and Fisher's exact test for categorical variables. The Wilcoxon rank-sum test was performed for continuous variables. Multivariable logistic regression models were employed to evaluate the longitudinal associations of adiposity indices with incident hypertension, elevated SBP, and elevated DBP, respectively. Multivariable models were sequentially adjusted for (1) SBP, and/or

9

166 DBP; (2) plus age, and gender; and (3) plus baseline residence, eGFR, soft fruit drinks, and total 167 carbohydrate intake. The areas under the receiver operator characteristics curve (AUC) were 168 calculated to assess the prediction ability of adiposity indices for hypertension, elevated SBP, and elevated DBP. Continuous net reclassification improvement (NRI), and absolute integrated 169 170 discrimination improvement (IDI) were used to assess whether adding the obesity-related measures could improve risk discrimination and reclassification for hypertension, elevated SBP, 171 and elevated DBP prediction above the baseline SBP, /or DBP. The cross-lagged panel models 172 were applied to analyze the temporal relationship between obesity-related measures (as 173 continuous variables) and SBP, and/or DBP levels (as categorical variables). The lagged effect 174 175 between the observed variables, the correlation between the selected adiposity indices (BMI, CI, WC, HC, TST and UAC, BRI) at each phase, and the residual variance of the observed variables 176 (SBP, and/or DBP levels) were estimated in the cross-lagged panel analysis, which elucidates the 177 178 temporal relationship between correlated variables and indicates the precursor variable. Several sensitivity analyses were performed to test the robustness of our findings. First, we 179 180 treated obesity-related measures, SBP, and DBP as continuous variables, to determine the linear associations between adiposity indices and BP. Prespecified subgroup analyses were conducted 181 based on age (7-12 years and 13-17 years), and gender (boys and girls) in secondary analyses. P 182 < 0.05 (two-sided) was considered statistically significant. All analyses were conducted with SAS 183 software (v.9.4, SAS Institute Inc., Cary, NC, USA). 184

185 **3. Result**

3.1 Baseline Characteristics of Pediatric and Adolescent Participants Across Various
 Systolic and Diastolic Blood Pressure Groups

10

188 Among the 227 participants included in the present analysis (64.8% of whom were boys), the 189 mean age was 11.7 (SD, ± 2.3) years. During a median of 2 years of follow-up, 60 cases of incident 190 hypertension (26.4%), 12 cases of incident elevated SBP (5.3%), and 58 cases of elevated DBP (25.6%) were identified. Children diagnosed with hypertension exhibited significantly higher 191 192 measurements for WC, HC, BMI, TST, and BRI. Additionally, a higher proportion of these children were girls and were classified as overweight. Compared with the normal SBP group, the children 193 and adolescents diagnosed with elevated SBP had a higher level of waist circumference, hip 194 circumferences, BMI, TST, BRI, ferritin, total protein, as well as more proportions of overweight. 195 Compared with the normal DBP group, the children and adolescents diagnosed with elevated 196 197 DBP had a higher level of WC, HC, BMI, TST, low level of total carbohydrate intake, as well as more proportions of girls, overweight (*Table 1*). 198

199 **3.2 Correlations Between Body Adiposity Measures and the Onset of Hypertension**

The risk ratios (RRs) and 95% confidence intervals (CIs) of incident hypertension for each adiposity index (per SD) are shown in Figure 1. After adjusting for covariates, BMI, WC, HC, TST, and BRI were associated with a higher risk of hypertension. When defining the outcome as elevated SBP (SBP > P_{90}), BMI, WC, HC, TST, UAC, SUA, CI, and BRI were associated with a higher risk of elevated SBP. While, BMI, WC, HC, TST, TG, and BRI were associated with a higher risk of elevated DBP (*Figure 1*).

3.3 The Incremental Predictive Value of Adiposity Indices in Assessing the Risk of Hypertension, High Systolic Blood Pressure, and High Diastolic Blood Pressure

A total of seven adiposity indices were identified, including BMI, WC, HC, TST, UAC, BRI, and CI.

209 The inclusion of HC and BRI was found to enhance the net reclassification ability and the overall

11

210 predicted probability for hypertension. Similarly, incorporating WC was shown to improve the 211 net reclassification ability and the overall predicted probability for elevated SBP. Adding the WC, HC, and BRI could improve the net reclassification ability and overall predicted probability for 212 elevated DBP. Furthermore, by incorporating BMI, WC, and HC into the basic model—which 213 considers continuous SBP and DBP—the area under the curve (AUC) for hypertension 214 prediction was significantly enhanced, increasing from 0.551 (95% CI: 0.463, 0.640) to 0.670 215 (95% CI: 0.592, 0.748). For SBP prediction, the AUC remained high at 0.873, with a CI ranging 216 from 0.771 to 0.974. Similarly, for elevated DBP prediction, the AUC improved from 0.531 (95% 217 CI: 0.440, 0.621) to 0.649 (95% CI: 0.570, 0.729) (Table 2). 218 219

3.4 The Temporal Dynamics of the Association Between Body Adiposity Measures and Elevated Systolic/Diastolic Blood Pressure

The fully cross-lagged model was used when compared to both the adiposity indices measures 221 222 to BP levels (defined as elevated SBP or elevated DBP) and the BP levels to the obesity-related measures (Figure 2). Of note, there existed no directional associations between the most 223 obesity-related measures and any hypertension symptoms. Baseline adiposity indices including 224 WC, HC, TST, UAC, BRI, and CI were unidirectional longitudinally associated with hypertension 225 symptoms (elevated SBP or elevated DBP). Cross-lagged path estimates showed that only BMI 226 was directional and longitudinally associated with hypertension symptoms. Baseline SBP (in the 227 2009 year) was longitudinally negatively associated with later BMI (in 2011 year), while 228 229 baseline DBP (in 2009 year) was significantly positively associated with BMI (in 2011 year).

230 **3.5 Sensitivity and Subgroup Analysis**

12

231 In the sensitivity analysis, the linear analysis revealed significant longitudinal associations 232 between various adiposity indices including BMI, WC, HC, TST, UAC, BSI, and BRI, with a higher risk of SBP and DBP, treated as continuous variables (Supplementary Figure 2). In subgroup 233 analysis, the significant associations of BMI, WC, HC, TST, and BRI with risk of hypertension only 234 235 existed in the younger age group (7 \sim 12 years) after accounting for multiple variables. In the subgroup classified by gender, the significant associations of BMI with the risk of hypertension 236 only existed in girls. Whereas, the associations of UAC, and high level of TG with risk of 237 hypertension only existed in boys (Supplementary Table 2 and Supplementary Table 3). 238

239 4. Discussion

240 In this cohort study of the resident Chinese pediatric and adolescent population, we identified a significant correlation between elevated BMI, WC, HC, TST, and BRI with the prevalence of 241 hypertension. This association remained consistent, regardless of initial systolic and diastolic 242 243 blood pressure measurements, demographic variables, lifestyle factors, and eGFR. In contrast, we observed no significant link between hypertension and UAC, triglycerides (TG), HDL-c, blood 244 glucose levels, serum uric acid (SUA), CI, BSI, and VFI. The relationships of these adiposity 245 indices with both elevated SBP and DBP mirrored their associations with hypertension, with the 246 exception that an increased UAC and CI were linked to a higher probability of elevated SBP. It is 247 noteworthy that among the five adiposity indices, only BMI demonstrated a consistent, 248 directional longitudinal association with the symptoms of hypertension. 249

Previous research has consistently highlighted the relationship between adiposity indices and
pediatric hypertension [23,24]. Our study adds to the existing literature by providing evidence
that an increased WC is positively correlated with elevated blood pressure in pediatric

13

253 populations. The accumulation of abdominal fat is known to secrete various adipokines and 254 cytokines that can lead to endothelial dysfunction and increased peripheral vascular resistance, 255 thereby contributing to the development of hypertension [25-27]. Moreover, our study reveals that an expanded HC is associated with a higher degree of fat deposition in the buttocks and 256 thighs, consequently increasing the likelihood of hypertension in the pediatric population. TST 257 could serve as a valuable anthropometric marker for assessing cardiovascular risk in children. 258 This association may be attributed to the secretion of adipose-derived hormones and 259 inflammatory cytokines that can lead to endothelial dysfunction and heightened vascular 260 resistance [28]. The areas under the ROC curves for WC and HC in predicting pediatric 261 262 hypertension were higher than those for BMI. These findings suggested that body fat distribution, rather than overall obesity, maybe a more sensitive predictor of cardiovascular 263 health in the pediatric population [29]. 264

The areas under the receiver operating characteristic (ROC) curves for WC and HC in the prediction of pediatric hypertension were notably higher compared to those for BMI. These results suggest that the distribution of body fat, rather than overall adiposity, may be a more sensitive predictor of cardiovascular health in the pediatric population. BRI, which quantifies visceral fat and total body fat percentages, was significantly associated with elevated blood pressure in children [30].

In prior research, the BRI has been identified as a more effective predictor than other anthropometric measures for assessing the risk of a range of clinical outcomes in adults. These include conditions such as cardiometabolic disorders, renal impairment, and malignancies [31-35]. There is an emerging consensus that visceral fat poses a significantly greater health risk

14

275 compared to subcutaneous fat, as it is more strongly associated with an increased risk of various 276 diseases in adults [36-37]. While the concept that BRI can estimate total and regional fat 277 percentages seems plausible and potentially superior in adults, the evidence linking BRI to diseases in children is currently limited. In the study, the area under the ROC curves for BRI in 278 279 predicting pediatric hypertension was found to be lower compared to WC and HC. This suggests that BRI's predictive power for pediatric hypertension may not be as effective as it is in adults. 280 Theoretically, assuming the shape of the body as an ellipse with the long axis height and the 281 short axis WC, BRI can be calculated as the eccentricity of this ellipse via human modeling. 282

Acknowledging the inherent disparities in body contours between the pediatric and adult demographics, it is crucial to appreciate that indices like the BRI may not be directly comparable across these groups. The efficacy of BRI as a predictor for hypertension could be subject to

variation owing to the unique growth trajectories, body composition differences, and the 286 287 influence of developmental milestones that are characteristic of the pediatric population. Therefore, the application of BRI in pediatric assessments should be approached with caution, 288 and further research is needed to establish its validity and reliability in predicting hypertension 289 specifically within the pediatric demographic. For practical reasons, there remains a need for a 290 simple and effective children-specific indicator to better reflect visceral obesity. Our study 291 revealed that UAC and CI were significantly correlated with systolic blood pressure but not with 292 293 diastolic blood pressure. This distinction could be due to the different physiological mechanisms 294 underlying the two types of blood pressure. Systolic blood pressure is more influenced by arterial stiffness, cardiac output, and abnormal changes in brain cortical structure, whereas 295

diastolic blood pressure is more affected by peripheral vascular resistance [38,39]. The fact that

15

297 our measurements are more strongly associated with SBP may indicate a closer link with factors298 affecting arterial stiffness, and brain cortical structure.

299 Our study's findings suggest that in the pediatric population, there is no significant association between blood lipid levels (triglycerides, LDL-c, HDL-c, total cholesterol, SUA) and the 300 development of hypertension. This observation challenges the widely held belief that 301 dyslipidemia is a consistent predictor of cardiovascular risk, including hypertension in adults. 302 The absence of a correlation in our pediatric cohort could be attributed to several factors. First, 303 the pathophysiology of hypertension in children may differ fundamentally from that in adults, 304 where lipid metabolism plays a more established role [40]. Children are in a dynamic phase of 305 306 growth and development, with a rapidly changing metabolic profile that might not yet be fully aligned with the traditional risk factors seen in adulthood [41]. 307

Additionally, the dietary and lifestyle factors that significantly influence lipid profiles in adults 308 309 may not have had sufficient time to exert their effects in children. As children's dietary habits and physical activity levels evolve, their lipid profiles may not have stabilized to a point where 310 they can significantly influence blood pressure. The association between adiposity indices and 311 hypertension in children appears to be age-dependent, with our study indicating a significant 312 correlation within the 7 to 12 years age group, which was not observed in the 13 to 18 years age 313 group. This age-specific association may reflect the dynamic changes in growth, development, 314 and metabolic patterns that occur during childhood and adolescence. The physiological changes 315 316 during adolescence, including growth spurts, hormonal fluctuations, and the maturation of various organ systems, could influence the relationship between adiposity indices and blood 317 pressure in complex ways that are not yet fully understood. The distribution of body fat and its 318

medRxiv preprint doi: https://doi.org/10.1101/2024.11.26.24318031; this version posted November 28, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license .

16

metabolic impact may be different in adolescents compared to adults, potentially mitigating the
effects of excess adiposity on blood pressure regulation.

321 Strengths and Limitations

A significant advantage of our study is its community-based approach, which successfully 322 enrolled a substantial sample of children and adolescents aged 7 to 17 in China. This age group 323 has been notably underrepresented in research pertaining to pediatric hypertension. 324 Limitations include the possibility of residual confounding inherent to the observational design, 325 including the possibility of differential distribution of unmeasured or incompletely measured 326 confounders such as birth weight and gestational age. We can also not exclude limitations in the 327 generalizability of the results in this CHNS cohort with access to health care and the existence of 328 withdrawal bias. Besides, different criteria for recognizing overweight and obesity in children 329 may influence the accuracy of the estimation. 330

331 5. Conclusions and Implications

Our study identified that several adiposity indices—namely BMI, WC, HC, TST, and BRI— were correlated with a heightened risk of hypertension among Chinese children and adolescents. Notably, only BMI demonstrated a consistent, directional relationship with the longitudinal presentation of hypertension symptoms. Moreover, the pronounced associations between these adiposity indices and the risk of hypertension were predominantly observed in the younger age group, ranging from 7 to 12 years old.

338 Figure Legends

Figure 1 Associations between anthropometric indices/metabolic measures and hypertension
diagnosis (Abbreviations: BMI, body mass index; BRI, body roundness index; CI, conicity index;

17

341	DBP, diastolic blood pressure; HC, hip circumference; SBP, systolic blood pressure; TST, triceps
342	skinfold thickness; UAC, upper arm circumference; WC, waist circumference; WHR, waist-to-hip
343	ratio. * 0.01 <p≤0.05; **="" ***="" 0.001<p≤0.01;="" p<0.001)<="" td=""></p≤0.05;>
344	Figure 2 Estimates from the cross-lagged model for the associations between anthropometry
345	parameters and incident hypertension classified by elevated SBP and elevated DBP
346	(Abbreviations: BMI, body mass index; BRI, body roundness index; CI, conicity index; DBP,
347	diastolic blood pressure; HC, hip circumference; SBP, systolic blood pressure; TST, triceps
348	skinfold thickness; UAC, upper arm circumference; WC, waist circumference. All estimates were
349	standardized. * 0.01 <p≤0.05; **="" ***="" 0.001<p≤0.01;="" p<0.001).<="" td=""></p≤0.05;>
350	Reference
351	1. Song P, Zhang Y, Yu J, Zha M, Zhu Y, Rahimi K, et al. Global Prevalence of Hypertension in

Children: A Systematic Review and Meta-analysis. JAMA Pediatr. 2019;173(12):1154-1163.

doi: 10.1001/jamapediatrics.2019.3310.

Flynn JT, Kaelber DC, Baker-Smith CM, Blowey D, Carroll AE, Daniels SR, et al.;
 SUBCOMMITTEE ON SCREENING AND MANAGEMENT OF HIGH BLOOD PRESSURE IN
 CHILDREN. Clinical Practice Guideline for Screening and Management of High Blood
 Pressure in Children and Adolescents. Pediatrics. 2017;140(3):e20171904. doi:
 10.1542/peds.2017-1904.

- 359 3. Yang L, Magnussen CG, Yang L, Bovet P, Xi B. Elevated blood pressure in childhood or
 adolescence and cardiovascular outcomes in adulthood: a systematic review. Hypertension.
- 361 2020;75(4):948-955. doi:10.1161/ HYPERTENSIONAHA.119.14168

362	4.	Chung J, Robinson C, Sheffield L, Paramanathan P, Yu A, Ewusie J, et al. Prevalence of
363		pediatric masked hypertension and risk of subclinical cardiovascular outcomes: a
364		systematic review and meta-analysis. Hypertension. 2023;80(11):2280-2292.
365		doi:10.1161/HYPERTENSIONAHA.123.20967.
366	5.	Zhang X, Liu J, Ni Y, Yi C, Fang Y, Ning Q, et al. Global Prevalence of Overweight and Obesity
367		in Children and Adolescents: A Systematic Review and Meta-Analysis. JAMA Pediatr. 2024
368		Aug 1;178(8):800-813. doi: 10.1001/jamapediatrics.2024.1576. PMID: 38856986; PMCID:
369		PMC11165417.
370	6.	Lobstein T, Brinsden H. On behalf of World Obesity Federation. Atlas of childhood obesity.
371		World Obesity Federation; 2019.
372	7.	Flynn J. The changing face of pediatric hypertension in the era of the childhood obesity
373		epidemic. Pediatr Nephrol. 2013;28(7):1059-1066. doi:10.1007/s00467-012-2344-0
374	8.	Muntner P, He J, Cutler JA, Wildman RP, Whelton PK. Trends in blood pressure among
375		children and adolescents.JAMA. 2004;291(17): 2107-2113. doi:10.1001/jama.291.17.2107.
376	9.	Koebnick C, Sidell MA, Li X, Woolford SJ, Kuizon BD, Kunani P. Association of High Normal
377		Body Weight in Youths With Risk of Hypertension. JAMA Netw Open. 2023; 6(3):e231987.
378		doi: 10.1001/ jamanetworkopen.2023.1987. PMID: 3691711MC10015306.
379	10.	Zong X, Kelishadi R, Hong YM, Schwandt P, Matsha TE, Mill JG, et al. Establishing
380		international optimal cut-offs of waist-to-height ratio for predicting cardiometabolic risk in
381		children and adolescents aged 6-18 years. BMC Med. 2023; 21(1):442. doi:

19

382 10.1186/s12916-023-03169-y.

383 11. Ge W, Yi L, Xiao C, Xiao Y, Liu J, Liang F, Yin J, Hu J. Effectiveness of a body shape index in
384 predicting pediatric high blood pressure. Pediatr Res. 2022;92(3):871-879. doi:
385 10.1038/s41390-021-01844-5.

- 12. Chen R, Ji L, Chen Y, Meng L. Weight-to-height ratio and body roundness index are superior
 indicators to assess cardio-metabolic risks in Chinese children and adolescents: compared
 with body mass index and a body shape index. Transl Pediatr. 2022;11(3):318-329. doi:
- 389 10.21037/tp-21-479.
- Popkin BM, Du S, Zhai F, Zhang B. Cohort Profile: The China Health and Nutrition Survey monitoring and Understanding Socio-Economic and Health Change in China, 1989-2011. Int
 J Epidemiol (2010) 39(6):1435–40. doi: 10.1093/ije/dyp322.
- 14. Yan S, Li J, Li S, Zhang B, Du S, Gordon-Larsen P, et al. The Expanding Burden of
 Cardiometabolic Risk in China: The China Health and Nutrition Survey. Obes Rev (2012)
 13(9):810–21. doi: 10.1111/j.1467-789X.2012.01016.x.
- 396 15. Schwartz GJ, Muñoz A, Schneider MF, Mak RH, Kaskel F, Warady BA, et al. New equations to
 397 estimate GFR in children with CKD. J Am Soc Nephrol. 2009; 20(3):629-37. doi:
 398 10.1681/ASN.2008030287.
- 399 16. WS/T611-2018; High Waist Circumference Screenning Threshold among Children and
 400 Adolescents Aged 7–17 Years. National Health and Family Planning Commission of People's
 401 Republic of China: Beijing, China, 2018.

20

- 402 17. Endocrinology, Genetics and Metabolism Group, Chinese Society of Pediatrics. Definition
 403 and prevention and treatment of metabolic syndrome in children and adolescents in China.
 404 Chinese Journal of Pediatrics, 2012, 50: 420-422.
- 405 18. Valdez R. A simple model-based index of abdominal adiposity. J Clin Epidemiol.
 406 1991;44(9):955-6. doi: 10.1016/0895-4356(91)90059-i.
- 407 19. Krakauer NY, Krakauer JC. A new body shape index predicts mortality hazard independently
 408 of body mass index. PLoS One. 2012; 7(7): e39504. doi: 10.1371/journal.pone.0039504.
- 409 20. Huang J, Zhou C, Li Y, Zhu S, Liu A, Shao X, et al. Visceral adiposity index,
 410 hypertriglyceridemic waist phenotype and chronic kidney disease in a southern Chinese
 411 population: a cross-sectional study. Int Urol Nephrol. 2015; 47(8): 1387-96. doi:
 412 10.1007/s11255-015-1040-y.
- 21. Thomas DM, Bredlau C, Bosy-Westphal A, Mueller M, Shen W, Gallagher D, et al.
 Relationships between body roundness with body fat and visceral adipose tissue emerging
 from a new geometrical model. Obesity (Silver Spring). 2013; 21(11): 2264-71. doi:
 10.1002/oby.20408.

Writing group of Chinese Guidelines for Management of Hypertension; Chinese
Hypertension League; Chinese Society of Cardiology; Chinese Medical Doctor Association
Hypertension Committee; Hypertension Branch of China International Exchange and
Promotive Association for Medical and Health Care; Hypertension Branch of Chinese
Geriatric Medical Association. 2018 Chinese Guidelines for Prevention and Treatment of
Hypertension. Chin. J. Cardiovasc. Med. 2018, 24, 24–56.

- 423 23. Ferreira S. Obesity and hypertension in children: a worldwide problem. Rev Port Cardiol
 424 (Engl Ed). 2018;37(5): 433-434. doi:10.1016/j.repc.2018.04.006.
- 425 24. Koebnick C, Sidell MA, Li X, Woolford SJ, Kuizon BD, Kunani P. Association of High Normal
- Body Weight in Youths With Risk of Hypertension. JAMA Netw Open. 2023;6(3):e231987.
- 427 doi: 10.1001/jamanetworkopen.2023.1987.
- 25. Cefalo CMA, Riccio A, Fiorentino TV, Rubino M, Mannino GC, Succurro E, et al. Endothelial
 dysfunction is associated with reduced myocardial mechano-energetic efficiency in
 drug-naïve hypertensive individuals. Intern Emerg Med. 2023;18(8):2223-2230. doi:
 10.1007/s11739-023-03402-9.
- 432 26. Markopoulou P, Koutroumpa A, Mantzou A, Margeli A, Papanikolaou E, Siahanidou T.
 433 Association of Plasma Irisin Levels with Circulating Endothelial Microparticles (EMPs) and
 434 Endothelial Progenitor Cells (EPCs) in Children Born Prematurely. Metabolites. 2023;
 435 13(1):120. doi: 10.3390/metabo13010120.
- 436 27. Markopoulou P, Papanikolaou E, Loukopoulou S, Galina P, Mantzou A, Siahanidou T.
 437 Increased circulating endothelial progenitor cells (EPCs) in prepubertal children born
 438 prematurely: a possible link between prematurity and cardiovascular risk. Pediatr Res.
 439 2021;90(1):156-165. doi: 10.1038/s41390-020-01190-y.
- 28. Ouyang F, Wells JC, Zhang GH, Du K, Wang X, Shen L, et al. Maternal Prenatal Factors and
 Child Adiposity in Associations with Cardiometabolic Risk Factors in Term-Born Chinese
 Children at the Age of 2 Years. Nutrients. 2023;15(15):3342. doi: 10.3390/nu15153342.

443	29.	Kim MS, Kim WJ, Khera AV, Kim JY, Yon DK, Lee SW, et al. Association between adiposity and
444		cardiovascular outcomes: an umbrella review and meta-analysis of observational and
445		Mendelian randomization studies. Eur Heart J. 2021; 42(34):3388-3403. doi:
446		10.1093/eurheartj/ehab454.
447	30.	$Thom as {\tt DM, BredlauC, Bosy-WestphalA, et al. Relationships between body roundness with body for the standard strength of the streng$
448		atand visceral adipose tissue emerging from a newgeometricalmodel.Obesity(Silver Spring).
449		2013;21(11):2264-2271. doi:10.1002/oby.20408.
450	31.	Cai X, Song S, Hu J, Zhu Q, Yang W, Hong J, et al. Body roundness index improves the
451		predictive value of cardiovasculardiseaserisk in hypertensive patients with obstructive
452		sleep apnea: a cohort study. Clin Exp Hypertens. 2023; 45(1): 2259132.
453		doi:10.1080/10641963.2023.2259132.
454	32.	Rico-Martín S, Calderón-García JF, Sánchez-Rey P, Franco-Antonio C, Martínez Alvarez M,
455		Sánchez Muñoz, et al. Effectiveness of body roundness index in predicting metabolic
456		syndrome: a systematic review and meta-analysis. Obes Rev. 2020; 21(7):e13023.
457		doi:10.1111/obr.13023.
458	33.	Wu L, Pu H, Zhang M, Hu H, Wan Q. Non-linear relationship between the body roundness
459		index and incident type 2 diabetes in Japan: a secondary retrospective analysis. J Transl Med

- 460 2022; 20(1):110. doi:10.1186/s12967022-03321-x.
- 34. Zhang Y, Gao W, Ren R, Liu Y, Li B, Wang A. et al. Body roundness index is related to the low
 estimated glomerular filtration rate in Chinese population: a cross-sectional study. Front
 Endocrinol (Lausanne). 2023;14:1148662. doi:10.3389/fendo. 2023.1148662.

464	35.	Gao	W,	Jin	L ,]	Li D,	Zhang	g Y,	Zhao	W,	Zhao	Y, 6	et al.	The	associatio	n between	the	body
465		rour	ndne	ess i	nde	ex an	d the r	isk	of colo	rec	tal car	cer	: a cr	oss-se	ectional stu	dy. Lipids I	Healt	h Dis.

- 466 2023; 22(1):53. doi:10.1186/s12944-023-01814-2.
- 467 36. Gruzdeva O, Borodkina D, Uchasova E, Dyleva Y, Barbarash O. Localization of fat depots and
- 468 cardiovascular risk. Lipids Health Dis. 2018;17(1): 218. doi:10.1186/ s12944-018-0856-8
- 469 37. Koenen M, Hill MA, Cohen P, Sowers JR. Obesity, adipose tissue and vascular dysfunction.
 470 Circ Res. 2021;128 (7):951-968. doi:10.1161/CIRCRESAHA.121. 318093.
- 471 38. Jeemon P, Harikrishnan S. Systolic blood pressure and cardiovascular health. Nat Med.
- 472 2022;28(10):2003-2004. doi: 10.1038/s41591-022-02005-9.
- 473 39. Fatani N, Dixon DL, Van Tassell BW, Fanikos J, Buckley LF. Systolic Blood Pressure Time in
- 474 Target Range and Cardiovascular Outcomes in Patients With Hypertension. J Am Coll
- 475 Cardiol. 2021;77(10):1290-1299. doi: 10.1016/j.jacc.2021.01.014.
- 476 40. Wicklow B, Retnakaran R. Gestational Diabetes Mellitus and Its Implications across the Life
- 477 Span. Diabetes Metab J. 2023;47(3):333-344. doi: 10.4093/dmj.2022.0348.
- 478 41. Zamojska J, Niewiadomska-Jarosik K, Kierzkowska B, Gruca M, Wosiak A, Smolewska E.
- Lipid Profile in Children Born Small for Gestational Age. Nutrients. 2023; 15(22):4781. doi:
- 480 10.3390/nu15224781.

		Non-		Systolic BP> P	Systolic BP≤P	Р	Diastolic BP> P	Diastolic BP≤P	Р
	Hypertension	hypertension	P value	90	90	value	90	90	value
Participants (n)	60	167		12	215		58	169	
Demographics									
Age (years)	12.0 (2.4)	11.6 (2.3)	0.301	12.7 (2.7)	11.7 (2.3)	0.151	12 (2.5)	11.7 (2.2)	0.380
Boys (%)	28 (46.7)	119 (71.3)	< 0.001	9 (75.0)	138 (64.2)	0.651	26 (44.8)	121 (71.6)	< 0.001
Rural (%)	43 (71.7)	132 (79.0)	0.244	7 (58.3)	168 (78.1)	0.217	41 (70.7)	134 (79.3)	0.177
Anthropometry parameters									
Waist circumference (cm)	68.9 (11.4)	63.8 (9.7)	0.001	79.7 (13.7)	64.4 (9.5)	0.003	68.3 (11.1)	64.1 (9.9)	0.009
Hip circumference (cm)	82.0 (10.5)	76.8 (11.1)	0.002	90.1 (11.1)	77.5 (10.8)	< 0.001	81.5 (10.2)	77 (11.3)	0.009
WHR	0.84 (0.10)	0.83 (0.08)	0.307	0.84 (0.09)	0.83 (0.09)	0.730	0.84 (0.10)	0.83 (0.08)	0.495
BMI (kg/m ²)	18.2 (4.0)	16.7 (3.0)	0.008	20.9 (5.0)	16.9 (3.1)	< 0.001	18.0 (4.0)	16.8 (3.0)	0.028
Overweight (> P ₇₅)	10 (16.7)	15 (9.0)	0.103	5 (41.7)	20 (9.3)	< 0.001	9 (15.5)	16 (9.5)	0.204
Systolic BP (mm Hg)	95.5 (13.4)	93.3 (10.7)	0.250	109.0 (8.7)	93.1 (11.0)	< 0.001	94.8 (13.0)	93.6 (11.0)	0.496
Diastolic BP (mm Hg)	61.5 (8.0)	61 (6.7)	0.647	68.1 (5.3)	60.7 (7.0)	< 0.001	61.1 (7.9)	61.1 (6.8)	0.980
Triceps Skinfold Thickness	1 28 (0.02, 1.72)	1.07 (0.72, 1.52)	0.009	1.05 (0.07.2.7)	1 10 (0 90 1 57)	0.015	1.22 (0.02, 1.70)	1 1 (0 72 1 57)	0.020
(TST, cm)	1.28 (0.93-1.73)	1.07 (0.73-1.53)	0.008	1.95 (0.97-3.7)	1.10 (0.80-1.57)	0.015	1.22 (0.93-1.70)	1.1 (0.73-1.57)	0.030
Upper Arm Circumference			0.005		21 (4.2)	0.001	22 (1.2)	21 (4.5)	0.000
(cm)	22 (4.5)	21 (4.4)	0.095	26 (4.8)	21 (4.3)	<0.001	22 (4.3)	21 (4.5)	0.283
Blood tests									
Albumin, g/L	48.6 (3.1)	48.2 (3.2)	0.456	49.8 (3.6)	48.2 (3.1)	0.095	48.5 (2.9)	48.2 (3.3)	0.539

Table 1 Baseline characteristics of the individuals stratified by Hypertension, Systolic BP and Diastolic BP in the CHNS dataset (n=262)

Alanine Transaminase, U/L	13.0 (10.5-15.0)	14.0 (11.0-18.0)	0.286	13.5 (11.5-25.0)	14.0 (10.0-17.0)	0.271	13.0 (10.0-15.0)	14.0 (11.0-18.0)	0.183
Apolipoprotein B, mg/dL	0.7 (0.5-0.8)	0.6 (0.5-0.7)	0.142	0.7 (0.5-0.9)	0.6 (0.5-0.7)	0.212	0.6 (0.5-0.8)	0.6 (0.5-0.7)	0.287
Apolipoprotein A-1, mg/dL	0.96 (0.85-1.09)	0.98 (0.87-1.09)	0.558	1.04 (0.96-1.10)	0.97 (0.86-1.09)	0.196	0.95 (0.84-1.06)	0.98 (0.87-1.09)	0.339
Ferritin, ng/ml	43.4 (28.6-65.1)	43.9 (28.4-58.0)	0.591	64.3 (45.6-78.0)	42.8 (28.4-58.1)	0.048	42.7 (28.6-63.0)	43.9 (28.8-58.2)	0.866
Serum magnesium, mmol/L	0.94 (0.09)	0.95 (0.07)	0.424	0.93 (0.12)	0.95 (0.07)	0.579	0.95 (0.09)	0.95 (0.07)	0.569
Tana famin	287.5	295.0	0.072	299.0	293.0	0.971	287.5	295.0	0.000
Transferrin, mg/gl	(264.5-329.0)	(265.0-325.0)	0.972	(269.0-321.5)	(265.0-325.0)	0.871	(265.0-330.0)	(265.0-325.0)	0.900
Soluble Transferrin receptor,	15(1220)	15(1210)	0.654	15(1220)	15(1210)	0.077	15(1210)	15(1210)	0.675
mg/L	1.5 (1.3-2.0)	1.5 (1.3-1.9)	0.654	1.5 (1.3-2.0)	1.5 (1.3-1.9)	0.977	1.5 (1.3-1.9)	1.5 (1.3-1.9)	0.675
Total protein, g/L	76.6 (5.3)	76.2 (4.8)	0.570	79.1 (5.2)	76.1 (4.9)	0.040	76.5 (5.3)	76.2 (4.8)	0.738
Serum uric acid, mg/dL	5.2 (1.5)	5.1 (1.3)	0.801	5.7 (1.6)	5.1 (1.3)	0.108	5.2 (1.5)	5.1 (1.3)	0.937
Total cholesterol, mmol/L	4.0 (0.7)	3.8 (0.6)	0.035	4.0 (0.8)	3.8 (0.6)	0.298	4 (0.6)	3.8 (0.6)	0.068
HDL cholesterol, mmol/L	1.4 (1.2-1.7)	1.4 (1.2-1.6)	0.771	1.4 (1.3-1.6)	1.4 (1.2-1.6)	0.678	1.3 (1.2-1.6)	1.4 (1.2-1.6)	0.613
LDL cholesterol, mmol/L	2.2 (1.8-2.6)	2.2 (1.7-2.6)	0.249	2.2 (1.7-2.8)	2.2 (1.8-2.6)	0.708	2.2 (1.8-2.6)	2.2 (1.8-2.6)	0.344
Triglycerides, mg/dL	0.8 (0.5-1.3)	0.7 (0.5-1.1)	0.745	0.8 (0.5-1.3)	0.7 (0.5-1.2)	0.759	0.8 (0.5-1.2)	0.7 (0.5-1.1)	0.939
eGFR, ml/min/l.73m ²	82.4 (9.2)	81.3 (11.6)	0.481	81.6 (6.8)	81.6 (11.1)	0.991	82.6 (9.2)	81.3 (11.6)	0.427
HbA1c, %	5.3 (0.4)	5.3 (0.4)	0.907	5.2 (0.5)	5.3 (0.4)	0.768	5.3 (0.4)	5.3 (0.4)	0.862
Blood Glucose, mmol/L	4.8 (0.5)	4.9 (0.5)	0.729	5.1 (0.5)	4.8 (0.5)	0.075	4.8 (0.5)	4.9 (0.5)	0.442
Creatinine, mg/dL	67.0 (62.0-75.0)	67.0 (61.0-73.0)	0.651	70.0 (63.5-76.0)	67.0 (61.0-73.0)	0.177	66.5 (62.0-74.0)	67.0 (61.0-73.0)	0.917
Anthropometric indices									
CI	1.15 (1.07-1.23)	1.13 (1.08-1.19)	0.427	1.20 (1.11-1.31)	1.13 (1.08-1.19)	0.058	1.14 (1.07-1.25)	1.13 (1.08-1.19)	0.559
BSI	0.08 (0.08-0.09)	0.08 (0.08-0.08)	0.432	0.08 (0.08-0.09)	0.08 (0.08-0.08)	0.399	0.08 (0.07-0.09)	0.08 (0.08-0.08)	0.449
VFI	0.92 (0.48-1.44)	0.70 (0.43-1.15)	0.139	0.77 (0.51-1.42)	0.74 (0.43-1.25)	0.650	0.92 (0.48-1.43)	0.7 (0.43-1.16)	0.186
BRI	2.26 (1.67-3.31)	1.95 (1.53-2.4)	0.025	3.55 (1.68-5.17)	1.97 (1.53-2.47)	0.021	2.19 (1.59-3.20)	1.95 (1.53-2.42)	0.061
Health-related behavior									

Soft/Sugared fruit drinks	54 (90.0)	146 (88.0)	0.670	12 (100.0)	188 (87.9)	0.369	52 (89.7)	148 (88.1)	0.748
Total protein intake (g/day)	49.8 (40.5-60.7)	50.2 (39.4-67.2)	0.530	50.8 (42.7-69.4)	50.1 (39.2-64.6)	0.338	49.8 (40.5-60.5)	50.2 (39.7-67.2)	0.433
$T_{a,b,c}$	55 1 (24 7 72 2)	52 2 (2(0 71 2)	0.920	1710.0	1645.4	0.205	1621.7	1670.9	0.020
Total lat intake (g/day)	55.1 (54.7-72.2)	55.5 (30.9-71.5)	0.850	(1395.0-2078.9)	(1324.6-1937.6)	0.205	(1240.7-1928.7)	(1357.8-1947.7)	0.950
Total carbohydrate intake	218.9	235.7	0.076	218.9	234.3	0.542	218.9	235.7	0.045
(g/day)	(174.8-260.4)	(198.8-276.9)	0.076	(149.8-327.4)	(192.3-270.2)	0.563	(174.8-259.2)	(199.5-276.9)	0.045
Total energy intake	1621.7	1670.9	0.015			0.641	51.0 (24.5.52.2)	52 4 (2 4 0 50 2)	0.010
(kcal/day)	(1240.7-1933.9)	(1353.3-1947.7)	0.317	61.8 (51.0-67.2)	52.4 (35.6-71.7)	0.641	51.0 (34.7-72.2)	53.4 (36.9-70.2)	0.218

Abbreviation: BMI, body mass index; BP, blood pressure; BSI, body shape index; BRI, body roundness index; CI, conicity index; eGFR, estimated glomerular filtration rate; HDL, high-density lipoprotein; HbA1c: Hemoglobin A1C; LDL, low-density lipoprotein; VFI, visceral fat index; WHR, waist to hip circumference ratio. Data are presented as No. (%), mean \pm SD; *P values were calculated by using T test or Wilcoxon test for continuous variables and χ 2 test or fisher exact test for categorical variables.

Table 2 Performance of Risk Prediction Models for 2-Year Risk of Incident Hypertension by Including selected Anthropometry parameters and metabolic Measures

	AUC (95% CI)	Difference (95% CI)	NRI (95% CI)	Absolute IDI (95% CI)
2-Year diagnosis of Hypertension				
Basic model ^a	0.551 (0.463, 0.640)	Reference	Reference	Reference
Basic model+BMI	0.624 (0.542, 0.707)	0.073 (-0.014, 0.161)	0.165 (-0.120, 0.451)	0.036 (0.005, 0.066)*
Basic model+Waist circumference	0.641 (0.559, 0.724)	0.090 (-0.012, 0.192)	0.244 (-0.044, 0.531)	0.044 (0.013, 0.075)**
Basic model+Hip circumference	0.646 (0.563, 0.729)	0.095 (-0.001, 0.190)	$0.468\ {(0.189,\ 0.748)}^{*}$	0.040 (0.013, 0.066)**
Basic model+TST	0.625 (0.542, 0.707)	0.073 (-0.029, 0.176)	$0.148 \left(0.076, 0.658 ight)^{*}$	0.027 (-0.009, 0.063)
Basic model+BRI	0.605 (0.518, 0.692)	0.054 (-0.048, 0.156)	0.360 (0.069, 0.651)*	0.041 (0.007, 0.074)*
Basic model+BRI, Hip circumference	0.655 (0.574, 0.735)	$0.103\ {(0.003,\ 0.204)}^{*}$	0.527 (0.241, 0.812)*	0.056 (0.020, 0.092)**
Basic model+BMI, Waist circumference, Hip circumference	0.670 (0.592, 0.748)	0.119 (0.021, 0.217)*	0.455 (0.168, 0.742)**	0.051 (0.018, 0.084)**
2-Year diagnosis of Elevated Systolic BP				
Basic model ^b	0.873 (0.771, 0.974)	Reference	Reference	Reference
Basic model+BMI	0.889 (0.795, 0.982)	0.016 (-0.026, 0.058)	0.243 (-0.324, 0.818)	0.051 (0.034, 0.137)
Basic model+Waist circumference	0.896 (0.797, 0.995)	0.023 (-0.036, 0.082)	0.700 (0.153, 1.249)*	0.113 (0.001, 0.225)*
Basic model+Hip circumference	0.885 (0.783, 0.987)	0.013 (-0.029, 0.054)	0.209 (-0.365, 0.782)	0.037 (-0.02, 0.094)
Basic model+TST	0.885 (0.779, 0.991)	0.013 (-0.048, 0.073)	0.413 (-0.160,0.986)	0.144 (-0.031, 0.318)
Basic model+Upper Arm circumference	0.888 (0.801, 0.975)	0.015 (-0.026-0.057)	$0.784~(0.277,~1.290)^{*}$	0.048 (-0.018, 0.114)
Basic model+CI	0.888 (0.777, 0.999)	0.017 (-0.030, 0.063)	0.340 (-0.240, 0.919)	0.038 (-0.030, 0.105)

*

0.158 (-0.011, 0.328)	0.711 (0.142, 1.280)*	0.014 (-0.049, 0.077)	0.886 (0.766, 1.000)	Basic model+BRI
0.157 (-0.01, 0.325)	0.636 (0.066, 1.207)*	0.021 (-0.039, 0.081)	0.893 (0.776, 1.000)	Basic model+BRI, Hip circumference
0.113 (0.001, 0.224)*	0.710 (0.163, 1.258)*	0.024 (-0.034, 0.083)	0.897 (0.797, 0.996)	Basic model+BMI, Waist circumference, Hip circumference
				2-Year diagnosis of Elevated Diastolic BP
Reference	Reference	Reference	0.531 (0.440, 0.621)	Basic model ^c
0.044 (0.015, 0.072)**	0.200 (-0.089, 0.490)	0.108 (-0.027, 0.242)	0.605 (0.523, 0.687)	Basic model+BMI
0.037 (0.009, 0.065)**	$0.333\ {(0.039, 0.628)}^{*}$	0.125 (-0.012, 0.263)	0.623 (0.537, 0.709)	Basic model+Waist circumference
0.043 (0.014, 0.072)**	0.456 (0.175, 0.738)**	$0.134\ (0.008,\ 0.259)^{*}$	0.631 (0.547, 0.714)	Basic model+Hip circumference
0.013 (-0.021-0.046)	$0.322\ {(0.027, 0.618)}^{*}$	0.100 (-0.042-0.242)	0.597 (0.511-0.683)	Basic model+TST
0.052 (0.017, 0.087)**	$0.349 \left(0.054, 0.643\right)^{*}$	0.078 (-0.072, 0.228)	0.575 (0.480, 0.670)	Basic model+BRI
0.037 (0.009, 0.065)**	0.577 (0.292, 0.861)**	0.141 (0.006, 0.277)*	0.639 (0.556, 0.721)	Basic model+BRI, Hip circumference
0.044 (0.014, 0.073)**	0.349 (0.054, 0.643)**	0.152 (0.022, 0.282)*	0.649 (0.570, 0.729)	Basic model+BMI, Waist circumference, Hip circumference

2 Abbreviations: AUC, area under curve; BMI, body mass index; BRI, body roundness index; CI, conicity inex; eGFR, estimated glomerular filtration rate; TST, triceps skinfold thickness. a Basic

3 model: baseline age, gender, systolic BP, diastolic BP, and eGFR. b Basic model: baseline age, gender, systolic BP, and eGFR; c Basic model: baseline age, gender, diastolic BP.

4 0.01<P≤0.05; ** 0.001<P≤0.01; *** P≤0.001

29

Hyperte	nsion diagnosis	OR (95% CI)	Elevated Systolic BP diagnosis	OR (95% CI)	Elevated Diastolic BP diagnosis	OR (95% CI)
BMI, per SD increase, kg/m2		1.55 (1.13-2.11) 1.51 (1.09-2.10) 1.62 (1.15-2.29)		1.76 (0.99-3.12) 1.78 (1.00-3.14) 2.32 (1.20-4.49)	⊨∎−1 ⊨⊜−1 ⊢≜−1	1.49 (1.10-2.01) 1.44 (1.05-1.99) 1.53 (1.09-2.16)
Waist circumference, per SD increase, cm		1.64 (1.19-2.26) 1.69 (1.20-2.38) 1.80 (1.25-2.60)		2.51 (1.33-4.71) 2.96 (1.47-5.97) 4.32 (1.75-10.66)		1.56 (1.14-2.14) 1.59 (1.13-2.24) 1.68 (1.16-2.41)
Hip circumference, per SD increase, cm		1.70 (1.18-2.44) 1.67 (1.15-2.43) 1.76 (1.15-2.68)		2.01 (0.91-4.46) 2.29 (1.04-5.04) → 2.73 (1.08-6.92)		1.62 (1.14-2.30) 1.67 (1.12-2.48) 1.65 (1.09-2.49)
WHR, per SD increase		1.01 (0.88-1.15) 0.99 (0.87-1.12) 1.00 (0.88-1.14)	2	1.01 (0.88-1.15) 1.16 (0.97-1.38) 1.18 (0.99-1.41)		0.99 (0.87-1.13) 0.98 (0.86-1.11) 0.99 (0.87-1.12)
TST, per SD increase, cm	⊧∎-ĭ ₽₫	1.69 (1.22-2.35) 1.64 (1.17-2.29) 1.72 (1.20-2.46)		2.06 (1.19-3.57) 2.06 (1.18-3.58) 2.57 (1.30-5.06)	⊨∎⊣ ⊨●⊣ ⊨≜⊸1	1.51 (1.11-2.05) 1.47 (1.07-2.00) 1.49 (1.07-2.08)
Upper Arm Circumference per SD increase, cm		1.23 (0.90-1.67) 1.24 (0.91-1.71) 1.35 (0.96-1.89)		1.82 (1.08-3.06) 1.83 (1.10-3.03) 2.22 (1.29-3.82)	• ■ -+ •©-1 • ☆ -1	1.18 (0.88-1.59) 1.20 (0.88-1.62) 1.28 (0.92-1.78)
Triglycerides, per SD increase, mmol/L		1.08 (0.81-1.43) 1.04 (0.75-1.42) 1.04 (0.76-1.41)		1.03 (0.5-2.11) 1.00 (0.47-2.15) 1.21 (0.63-2.32)		1.51 (1.11-2.05) 1.47 (1.07-2.00) 1.49 (1.07-2.08)
HDL cholesterol, per SD increase, mmol/L		0.97 (0.72-1.31) 0.97 (0.71-1.33) 0.98 (0.71-1.35)		0.97 (0.55-1.72) 0.90 (0.49-1.67) 0.85 (0.43-1.71)		0.97 (0.72-1.31) 0.98 (0.71-1.35) 0.99 (0.71-1.37)
Glucose, per SD increase, mmol/L		0.92 (0.68-1.24) 0.98 (0.71-1.35) 0.98 (0.70-1.37)		1.18 (0.64-2.17) 1.35 (0.71-2.57) 1.56 (0.68-3.56)	3 -1 14-1	1.06 (0.79-1.41) 1.01 (0.72-1.40) 1.00 (0.73-1.38)
SUA, per SD increase, mg/dL		1.00 (0.99-1.01) 1.00 (0.99-1.01) 1.00 (0.99-1.01)	•	1.00 (0.99-1.01) 1.00 (0.99-1.01) 1.01 (1.00-1.02)		1.00 (1.00-1.01) 1.00 (1.00-1.01) 1.00 (0.99-1.01)
CI, per SD increase		1.10 (0.81-1.48) 1.09 (0.80-1.48) 1.09 (0.79-1.49)		1.86 (0.98-3.54) 1.92 (1.02-3.63) 2.01 (1.04-3.90)	₩ - 1	1.08 (0.80-1.45) 1.07 (0.79-1.45) 1.06 (0.77-1.46)
BSI, per SD increase		1.18 (0.87-1.60) 1.21 (0.89-1.66) 1.20 (0.87-1.67)		1.84 (0.95-3.56) 1.91 (0.96-3.79) 2.06 (0.95-4.46)	+ <mark>■</mark> -1 +©-1 + <u>©</u> -1	1.16 (0.86-1.57) 1.20 (0.88-1.64) 1.18 (0.85-1.65)
VFI, per SD increase	₽ ₽ ₽	1.20 (0.90-1.60) 1.09 (0.81-1.48) 1.07 (0.79-1.44)		1.14 (0.62-2.11) 1.11 (0.58-2.12) 1.2 (0.65-2.22)	9 ⊒- 1 1 0 -1 12 1 -1	1.20 (0.90-1.60) 1.08 (0.80-1.48) 1.05 (0.78-1.43)
BRI, per SD increase		2.42 (1.13-5.16) 2.42 (1.09-5.38) 2.74 (1.18-6.38)		6.57 (1.39-31.09) 7.76 (1.38-43.78) 21.59 (2.40-194.62)		2.22 (1.04-4.75) 2.23 (1.00-4.95) 2.47 (1.06-5.74)
ł	2 4	ll 6 8				<mark>⊢−−−−</mark> 6 8
Adjusted for Adjusted for Adjusted for	baseline Systolic BP, all variables in model all variables in model	and/or Diastolic BP (mode 1 plus age, and gender (mo 2 plus baseline residence, o	l 1) odel 2) eGFR, soft fruit drinks, and total car	bohydrate intake (model 3)		



7 Figure 1 Associations between anthropometric indices/metabolic measures and hypertension 8 diagnosis (Abbreviations: BMI, body mass index; BRI, body roundness index; CI, conicity index; TST, triceps skinfold thickness; WHR, waist-to-hip ratio. * 0.01<P≤0.05; ** 0.001<P≤0.01; *** 9 P<0.001) 10





- 14 elevated SBP and elevated DBP (Abbreviations: BMI, body mass index; BRI, body roundness index; CI, conicity index; eGFR, estimated glomerular
- 15 filtration rate; TST, triceps skinfold thickness. All estimates were standardized. * 0.01<P≤0.05; ** 0.001<P≤0.01; *** P<0.001).

12

31

16 Acknowledgements

The authors thank the CHNS team for their hard work and unselfsh sharing of survey data. This 17 18 research uses data from China Health and Nutrition Survey (CHNS). We are grateful to research 19 grant funding from the National Institute for Health (NIH), the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) for R01 HD30880, National Institute 20 on Aging (NIA) for R01 AG065357, National Institute of Diabetes and Digestive and Kidney 21 Diseases (NIDDK) for R01DK104371 and R01HL108427, the NIH Fogarty grant D43TW009077 22 since 1989, and the China-Japan Friendship Hospital, Ministry of Health for support for CHNS 23 2009, Chinese National Human Genome Center at Shanghai since 2009, and Beijing Municipal 24 25 Center for Disease Prevention and Control since 2011.

26 Abbreviation

AUC, Area Under Curve; BMI, Body Mass Index; BP, Blood Pressure; BRI, Body Roundness Index; 27 28 BSI, Body Shape Index; CHNS, China Health and Nutrition Survey; CI, Conicity Inex; CMRFs, Clustered Cardio-metabolic Risk Factors; CVD, Cardiovascular Disease; eGFR, estimated 29 glomerular filtration rate; HC, Hip Circumference; HDL-C, High-density Lipoprotein Cholesterol; 30 IDI, Integrated Discrimination Improvement; IQR, Interquartile Range; NRI, Net Reclassification 31 Improvement; SD, Standard Deviation; SUA, Serum Uric Acid; TG, Triglycerides; TST, Triceps 32 Skinfold Thickness; VFI, Visceral Fat Index; WC, Waist Circumference; WHR, Waist to Hip 33 **Circumference Ratio.** 34

<sup>Funding: This work was supported by the National Key Research and Development Program of
China (grant numbers 2021YFC2701900, 2021YFC2701901, 2021YFC2701903).</sup>

32

37	Author contribution: BZ, SZ, and JF contributed to the conception or design of the work. BZ, HS
38	and XW contributed to the acquisition, analysis, or interpretation of data for the work. BZ and
39	RW drafted the manuscript. JY and XG critically revised the manuscript. BZ and TW contribute to
40	the analysis or interpretation of the work. All gave final approval and agreed to be accountable
41	for all aspects of work ensuring integrity and accuracy.
42	Ethical approval: The China Health and Nutrition Survey (CHNS) was granted approval by the
43	Institutional Review Board at the University of North Carolina at Chapel Hill, as well as by the
44	local Institutional Review Board of Ethics Committee. As this study is based on secondary data
45	analysis of an existing dataset, we did not require additional ethical approval from our
46	institution. Original data generated and analyzed during this study are included in this
47	published article or in the data repositories listed in References.
48	Disclosure summary: No financial or non-financial benefits have been received or will be received
49	from any party related directly or indirectly to the subject of this article. The authors have no conflict
50	of interest to declare.
51	Data Availability: Original data generated and analyzed during this study are included in this
52	published article or in the data repositories listed in References.