

## **Long Sleep Duration, Cognitive Performance, and the Moderating Role of Depression: A Cross-Sectional Analysis in the Framingham Heart Study**

**Authors:** Vanessa M. Young<sup>1,2,3</sup>, Rebecca Bernal<sup>1</sup>, Andree-Ann Baril<sup>4,5,6</sup>, Joy Zeynoun<sup>1</sup>, Crystal Wiedner<sup>1</sup>, Carlos Gaona<sup>1</sup>, Alexa Beiser<sup>4,7,8</sup>, Antonio L. Teixeira<sup>1,11</sup>, Arash Salardini<sup>1,11</sup>, Matthew P. Pase<sup>4,9</sup>, Jayandra Jung Himali<sup>1,4,7,8,10</sup>, Sudha Seshadri<sup>1,4,7,11</sup>

### **Affiliations:**

<sup>1</sup>Glenn Biggs Institute for Alzheimer's & Neurodegenerative Diseases, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

<sup>2</sup>Graduate School of Biomedical Sciences, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

<sup>3</sup>School of Social and Behavioral Sciences, Arizona State University, Phoenix, Arizona, USA

<sup>4</sup> Framingham Heart Study, Framingham, Massachusetts, USA

<sup>5</sup>Research Center of the CIUSSS-NIM, Hôpital du Sacré-Coeur de Montréal, Montreal, Quebec, Canada was

<sup>6</sup> Department of Medicine, University of Montreal, Quebec, Canada CC

<sup>7</sup>Department of Neurology, Boston University Chobanian & Avedisian School of Medicine, Boston, Massachusetts, USA

<sup>8</sup>Department of Biostatistics, Boston University School of Public Health, Boston, Massachusetts, USA

<sup>9</sup>Turner Institute for Brain and Mental Health, Monash University, Melbourne, Victoria, Australia

<sup>10</sup> Department of Population Health Sciences, University of Texas Health Science Center, San Antonio, Texas, USA

<sup>11</sup> Department of Neurology, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

## **ABSTRACT**

**INTRODUCTION:** We investigated whether depression modified the associations between sleep duration and cognitive performance.

**METHODS:** Multivariable linear regression models examined the associations between sleep duration and cognition in 1,853 dementia- and stroke-free participants from the Framingham Heart Study. Participants were categorized in four groups: no depressive symptoms, no antidepressants; depressive symptoms without antidepressants use; antidepressant use without depressive symptoms; both depressive symptoms and antidepressant use.

**RESULTS:** Long sleep was associated with reduced overall cognitive function. Strong associations between sleep duration and cognitive performance were found in individuals with depressive symptoms, regardless of antidepressant use. Weaker but significant effects were observed in those without depressive symptoms. No significant associations were observed in participants using antidepressants without depressive symptoms.

**DISCUSSION:** These findings provide new evidence that sleep duration may be a modifiable risk factor for cognitive decline, particularly in individuals with depressive symptoms. Future research should elucidate underlying mechanisms and temporal relationships.

**Keywords:** *Sleep duration, Cognitive function, Depression*

**Corresponding author:**

*Vanessa M. Young, [Youngv1@uthscsa.edu](mailto:Youngv1@uthscsa.edu)*

*Glenn Biggs Institute, UT Health San Antonio, 7703 Floyd Curl Drive, San Antonio, TX  
78299, USA*

## **RESEARCH IN CONTEXT**

**Systematic review:** We conducted a systematic search on PubMed and Google Scholar for peer-reviewed articles using keyword combinations related to sleep, cognition, and depression. Existing evidence reveals mixed findings on the relationship between sleep duration and cognition, with limited studies examining the role of depression on this association.

**Interpretation:** We observed that only long sleep duration ( $\geq 9$ h) was associated with poorer global cognition, executive function, visuospatial memory, and verbal learning/memory. Depression moderated this association, showing stronger negative effects of long sleep on cognition in individuals with depressive symptoms, regardless of antidepressant use. This suggests that long sleep duration may serve as an early indicator or risk factor for cognitive decline in those with depressive symptoms.

**Future directions:** Longitudinal studies with objective and subjective sleep assessments across diverse populations are needed to clarify how depression and its treatment influence the relationship between sleep duration and cognitive decline and support clinical strategies.

**HIGHLIGHTS (3-5 points, no more than 85 characters with space)**

1. Sleeping  $\geq$  9hours/night was associated with worse cognitive performance.
2. This association was stronger among those with depression.
3. Long sleepers were more likely to report symptoms of depression.
4. Sleep may be a modifiable risk for cognitive decline in people with depression.

**Acknowledgments:** We thank the research team at Boston University and all Framingham Heart Study participants who volunteered their time and made this scientific effort possible.

**Funding:**

The Framingham Heart Study was supported by grants from the National Heart, Lung, and Blood Institute (contract No. N01-HC-25195, No. HHSN268201500001I, and No. 75N92019D00031), the National Institute on Aging (R01 AG054076, R01 AG049607, U01 AG052409, R01 AG059421, RF1 AG063507, RF1 AG066524, U01 AG058589), and the National Institute of Neurological Disorders and Stroke (R01 NS017950 and UH2 NS100605).

This work was made possible by grants from the National Institute of Health (P30 AG066546), the National Institute of Aging (AG0623531), and the Alzheimer's Association.

Dr. Baril is supported by the Sleep Research Society Foundation, the Fonds de recherche du Québec en Santé, the Canadian Institutes of Health Research, and the Alzheimer Society of Canada.

Dr. Pase's salary is supported by a National Health and Medical Research Council of Australia Investigator Grant (GTN2009264).

Dr. Himali is supported by an endowment from the William Castella family as William Castella Distinguished University Chair for Alzheimer's Disease Research, and Dr. Seshadri by an endowment from the Barker Foundation as the Robert R Barker

Distinguished University Professor of Neurology, Psychiatry and Cellular and Integrative Physiology. Drs. Seshadri and Himali receive support from The Bill and Rebecca Reed Endowment for Precision Therapies and Palliative Care.

**Conflict of Interests:**

Dr. Seshadri reports consulting for Eisai and Biogen. Dr. Baril reports speaking fees from Eisai. Dr. Salardini reports speaking fees from Lilly.

# 1 BACKGROUND

Cognitive decline and dementia, particularly Alzheimer's disease (AD), are pressing public health concerns, exacerbated by increasing life expectancy worldwide.<sup>1</sup> Cognitive impairment interferes with activities of daily living, threatens independence and quality of life, and leads to considerable healthcare costs and caregiver burden.<sup>2-4</sup> Despite recent advancements in treatment, the efficacy of the current anti-amyloid disease-modifying treatments is modest, and their use is associated with significant adverse effects.<sup>5</sup> Therefore, it is important to continue to investigate possible prevention strategies based on modifiable risk factors and their potential interaction.<sup>6</sup>

There is an increasing recognition of the significance of sleep as a vital physiological process for brain health.<sup>7,8</sup> Disturbances in sleep duration and patterns are present in both normal and pathological aging,<sup>9-11</sup> and they contribute to an increased risk of cognitive deficits.<sup>12</sup> The Global Council on Brain Health recommends 7 to 8 hours of nightly sleep for adults to preserve brain health.<sup>13</sup> Several studies have suggested that both excessive and insufficient sleep relative to the prescribed duration are linked to impairments in cognitive domains, including memory, attention, and executive functioning.<sup>18-21</sup> The exact nature of these relationships varies depending on the study and the characteristic of the sample. The largest such study, a pooled cohort study of more than 28,000 people in mid to late-life, found that sleep durations of less than four or more than ten hours was associated with faster cognitive decline; suggesting an inverted U curve relationship between sleep duration and cognition.<sup>19</sup> In contrast, a cross-sectional study of healthy middle-aged subjects<sup>20</sup> and a longitudinal<sup>21</sup> study of older adults reported an association solely between short sleep duration (< 6 hours/night) and diminished



cognition.<sup>22</sup> Other cross-sectional,<sup>23</sup> longitudinal<sup>24</sup> and prospective<sup>18</sup> studies have only found prolonged sleep ( $\geq 9$  hours/night) correlated with cognitive deterioration. These inconsistencies in evidence hinder our understanding of the precise nature of the association between sleep duration and cognition.

Depression, a leading cause of disability globally and a modifiable risk factor for cognitive decline<sup>6</sup>, often co-occurs with sleep disorders.<sup>25</sup> The association between sleep disorders and depression is well-established, with about 90% of people with depression reporting problems with sleep.<sup>22</sup> Sleep disorders are believed to precede depression rather than the contrary,<sup>26,27</sup> with insomnia being the most frequent sleep disorder associated with depressive symptoms.<sup>28</sup> Depression has also been associated with an increased risk of cognitive deficits.<sup>29</sup> The cognitive effects of depression appear to persist even after remission of depressive symptoms,<sup>30</sup> contributing to a heightened vulnerability to dementia later in life.<sup>31</sup> Depressive feelings frequently occur in dementia patients and may act as early markers of the condition.<sup>32</sup>

The complex relationship among sleep duration, cognitive function, and depression remains inadequately investigated. Further investigation into these often comorbid modifiable risk factors may lead to the development of more personalized prevention strategies. To address this knowledge gap, we used data from the Framingham Heart Study to cross-sectionally examine the associations of short, average, and long self-reported nighttime sleep duration with cognitive performance among dementia- and stroke-free-adults. We further explored whether depressive symptoms modified these associations. We hypothesized that (1) both long and short self-reported nighttime sleep durations would be associated with poorer cognitive performance, and (2) depressive

symptoms would moderate these associations such that the relationship between sleep duration and cognitive performance would be stronger in individuals with depressive symptoms regardless of antidepressant usage, relative to those without depressive symptoms.

## 2 METHODS

### 2.1 Sample

The Framingham Heart Study (FHS) is a multi-generational, longitudinal study started in 1948. We included 1,853 participants from the FHS Third Generation, Omni 2, and New Offspring Spouse Cohorts. The study included dementia- and stroke-free participants who completed neuropsychological testing during their second examination cycle between 2008 and 2011. Figure 1 illustrates the sample selection. This study was conducted in accordance with the Declaration of Helsinki. The Institutional Review Board at Boston Medical Center approved the study protocol. All participants signed the informed consent.

### 2.2 Sleep Assessment

Habitual self-reported sleep duration was assessed during in-clinic visits, which occurred on average approximately  $1.7 \pm 1.0$  years before the neuropsychological assessment in these cohorts. The participants responded to a prompt that asked them to list the number of hours they typically sleep. We classified sleep duration into three categories: short sleep duration (less than 6 hours), average sleep duration (between 6 and 9 hours) for the reference group, and long sleep duration (more than 9 hours).<sup>33</sup>

### 2.3 Cognitive Assessments

Trained neuropsychological raters assessed cognitive function using a comprehensive and standardized test battery.<sup>34</sup> In this study, we included the following tests: Trail Making

Test Parts A and B<sup>35</sup>, Visual Reproduction and Logical Memory Tests (sum of immediate and delayed recall scores) from the Wechsler Memory Scale—Revised<sup>36</sup>, and the Similarities Test from the Wechsler Adult Intelligence Scale.<sup>37</sup> The tests assessed attention, processing speed, and executive function; visuospatial memory; verbal learning and memory; and verbal reasoning, respectively. Trail Making Test Parts A and B were log-transformed and reversed, so that a higher score indicated better performance for all cognitive tests. A global cognitive score was derived from a principal component analysis,<sup>38</sup> combining weighted loadings of the Similarities Test, Visual Reproduction, Logical Memory Test, and Trail Making Test Part B.

## 2.4 Measures of Depression

We used the 20-item Center for Epidemiologic Studies Depression Scale (CES-D)<sup>39</sup> to assess depressive symptoms. We defined depression as either a score of  $\geq 16$  on the CES-D or the use of antidepressant medications (anatomical therapeutic chemical [ATC] code N06A).<sup>40</sup> We then categorized the participants into four groups: (1) controls – those not using antidepressants and without depressive symptoms (CES-D < 16); (2) those using antidepressants and without depressive symptoms (CES-D < 16); (3) those not using antidepressants with depressive symptoms (CES-D  $\geq 16$ ); and (4) those using antidepressants and with depressive symptoms (CES-D  $\geq 16$ ). The four-category classification affords clinical nuances in the relationship between depressive symptoms, treatment status, and cognitive outcomes that may be masked when using a binary classification (yes versus no) for overall prevalence estimates.

## 2.5 Covariates

The following covariates, potentially associated with sleep, depression, or cognitive function, were incorporated into two separate statistical models:

- Age (years) and age squared (years<sup>2</sup>) at the time of neuropsychological testing.
- Education: high school or less, some college, or college degree.
- Sex: female or male.
- Cohort: Third Generation, Omni 2, and New Offspring Spouse Cohorts from FHS.
- Time interval between sleep duration questionnaires and neuropsychological testing.
- Hypertension was characterized according to the Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure, as stage I hypertension or worse, defined by a systolic blood pressure (SBP) of  $\geq 140$  mmHg, a diastolic blood pressure (DBP) of  $\geq 90$  mmHg, or the utilization of antihypertensive medication.<sup>41</sup>
- EDTA plasma aliquots were analyzed daily at the FHS laboratory to measure:
  - (1) total cholesterol level (mg/dL), (2) high-density lipoprotein level (mg/dL)<sup>42</sup>, and (3) triglycerides (mg/dL).
- C-reactive protein (CRP) levels were categorized as low  $< 1$  mg/L, average 1-3 mg/L, and high  $> 3$  mg/L, according to the American Heart Association guidelines.<sup>43</sup>
- Body mass index (BMI) calculated using weight in kilograms and divided by the square of height in meters (m<sup>2</sup>).
- *Apolipoprotein*  $\epsilon$  4 allele carrier status (*APOE*  $\epsilon$  4 status): those with at least one  $\epsilon$  4 allele (as determined by PCR) vs those with none.

## 2.6 Statistical Analyses

We used SAS V.9.4 (SAS Institute) to perform statistical analyses and considered statistical significance at  $p < 0.10$  for test of interactions and  $p < 0.05$  for all other tests. We treated self-reported sleep duration (short, average, and long sleep duration) and depression group (control, antidepressants usage/CES-D  $< 16$ , no antidepressants usage/CES-D  $\geq 16$ , and antidepressants usage CES-D  $\geq 16$ ), as categorical variables.

First, we used multivariable linear regression models to examine the associations between the three sleep categories and global cognition. Secondary analysis assessed the associations between sleep duration categories and individual cognitive tasks. Model 1 was adjusted for age and age squared at neuropsychological testing, education, sex, time between sleep duration questionnaires and neuropsychological testing, and cohort. In Model 2, we additionally adjusted for hypertension, total cholesterol to HDL ratio, triglycerides, CRP level, BMI, and *APOE*  $\epsilon 4$  status.

We explored effect modification by depression group for the associations between sleep duration and cognitive performance by adding sleep group by depression group interaction terms in Model 1. When a statistically significant interaction was observed, we stratified the association between sleep duration and cognitive performance by depression group.

## 2.7 Sensitivity Analysis

In the primary models between sleep duration and cognitive performance, we conducted a sensitivity analysis limiting the sample to middle aged and older adults (cut-off  $\geq 45$  years, following the age groups defined by the National Library of Medicine<sup>44</sup>) to examine

whether the sleep duration-cognition association and potential effect modification by depression group remained consistent when excluding younger adults from the analysis.

## 3 RESULTS

### 3.1 Sample Characteristics

The sample consisted of 1,853 adults with an average age of 49.8 years (age range 27–85). The sample was 42.7% male, 95.3% white, and 57.2% college educated. Most of the participants reported average sleep durations of 6-9 hours (68.9%), followed by short sleep ( $\leq 6$  hours; 24.3%) and long sleep ( $\geq 9$  hours; 6.8%). Depression, defined as CES-D  $\geq 16$  or the use of antidepressant medications, was present in 448 (24.2%) participants: 315 (17.0%) reported using antidepressants, while 198 (11.0) had a CES-D score of 16 or higher. The group with long sleep duration tended to have a higher proportion of depression ( $n = 55$ , 44.0%) compared to the short ( $n = 106$ , 23.5%) and average ( $n = 287$ , 22.5%) sleep groups. **Table 1** shows the overall sample's characteristics, and **Table 2** shows the performance on domain-specific cognitive tests categorized by sleep duration.

When examining sleep duration across the four depression categories (see **Table 3**), those using antidepressants and with depressive symptoms (CES-D  $\geq 16$ ) reported the highest proportion of long sleep ( $n = 11$ ; 16.9%), followed by those using antidepressants but without depressive symptoms ( $n = 32$ ; 12.8%). In contrast, the control group (no antidepressants usage and without depressive symptoms) had the lowest proportion of long sleepers ( $n = 70$ ; 5.0%).

## 3.2 Association between Sleep and Cognition

### 3.2.1 Long Sleep Duration was Associated with Global Cognition

**Table 5** presents the association between short and long sleep durations and global cognitive performance, compared to average sleep duration (> 6 or < 9 hours vs. 6-9 hours). Long sleep duration ( $\geq 9$  hours) had a statistically significant association with poorer global cognition ( $\beta \pm SE = -0.25 \pm 0.07$ ,  $p < 0.001$ ), as measured by the global cognitive score, relative to those with average sleep duration. The association between short sleep duration ( $\leq 6$  hours) and global cognition did not reach statistical significance ( $-0.03 \pm 0.04$ ,  $p = 0.465$ ). In Model 2, the pattern of associations between long sleep duration and global cognition remained consistent with Model 1.

### 3.2.2 Long Sleep Duration was Associated with Task-specific Cognitive Domains

Long sleep duration ( $\geq 9$  hours) was associated with significantly worse performance on Trail Making Test Part B ( $-0.09 \pm 0.03$ ,  $p = 0.009$ ), Visual Reproduction ( $-1.80 \pm 0.42$ ,  $p < 0.001$ ), and Logical Memory ( $-1.50 \pm 0.60$ ,  $p = 0.013$ ) compared to the average sleep duration group. Long sleep was not associated with Trail Making Test Part A or the Similarities Test. Short sleep was not associated with any cognitive tasks. The results remained consistent in the fully adjusted model (see **Table 5**).

### 3.2.3 Sleep Duration Interacts with Depression in its Association with Cognition

We tested statistical interactions to determine whether depression groups modified the relationships between sleep duration categories and cognitive performance. We observed significant interactions indicating effects on global cognitive score ( $p = 0.028$ ), Trail Making Test Part B ( $p = 0.012$ ), Visual Reproduction ( $p = 0.064$ ), and Similarities test ( $p = 0.027$ ).

As shown in **Table 6**, in the control group (no antidepressants, CES-D < 16), long sleep was associated with poorer global cognition ( $-0.18 \pm 0.09$ ,  $p = 0.044$ ) and visual reproduction ( $-1.30 \pm 0.55$ ,  $p = 0.019$ ), while short sleep showed no significant associations with any cognitive measure. In those with depressive symptoms (CES-D  $\geq 16$ ) not using antidepressants, long sleep was associated with poorer performance across multiple domains: global cognition ( $-0.60 \pm 0.26$ ,  $p = 0.024$ ), Trails Part B ( $-0.29 \pm 0.13$ ,  $p = 0.023$ ), and visual reproduction ( $-3.70 \pm 1.70$ ,  $p = 0.031$ ). Short sleep was not significantly associated with cognitive measures in this group. Among those using antidepressants with depressive symptoms (CES-D  $\geq 16$ ), long sleep was associated with poorer global cognition ( $-0.74 \pm 0.30$ ,  $p = 0.017$ ) and visual reproduction ( $-4.30 \pm 1.89$ ,  $p = 0.027$ ). We observed no significant associations between short sleep and cognitive performance in individual using antidepressants but with depressive symptoms. Finally, among participants using antidepressants but without depressive symptoms (CES-D < 16), neither short nor long sleep duration was significantly associated with cognitive performance.

### 3.2.4 Sensitivity Analysis

In a sensitivity analysis including middle age and older adults (cut-off  $\geq 45$  years; see **Supplementary Tables 1 and 2** for sample characteristics and cognitive characteristics by age group), the pattern of associations between sleep duration and cognitive performance was largely consistent with the full sample results as shown in **Supplementary Table 3**. Significant interactions indicating effect modification by depression group were observed in the associations between sleep duration and global cognitive score ( $p = 0.009$ ), Trails Part B ( $p = 0.038$ ), Visual Reproduction ( $p = 0.050$ ), and Similarities Test ( $p = 0.006$ ). The most pronounced effects were seen in those using



antidepressants and with depressive symptoms (CES-D  $\geq$  16). Relative to average sleep duration, long sleep duration was associated with significantly worse performance in all four cognitive measures: global cognition ( $-1.10 \pm 0.35$ ,  $p = 0.004$ ), Trails Part B ( $-0.36 \pm 0.15$ ,  $p = 0.023$ ), Visual Reproduction ( $-6.59 \pm 2.24$ ,  $p = 0.005$ ), and Similarities Test ( $-3.80 \pm 1.69$ ,  $p = 0.030$ ). In contrast, for the control group, only Visual Reproduction showed a significant negative association with long sleep duration ( $-1.43 \pm 0.68$ ,  $p = 0.037$ ). No significant associations were observed among those using antidepressants and no depressive symptoms (CES-D  $<$  16) and those not using antidepressants with depressive symptoms (CES-D  $\geq$  16). See Supplementary **Table 4**.

## 4 DISCUSSION

In this study, we investigated the association between self-reported sleep duration and cognition in the FHS cohorts and examined whether this association was modified by depression. Relative to average sleep duration, we found that long but not short sleep duration was associated with poorer performance in global cognition, executive function, visuospatial memory, and verbal learning/memory performance. Further, we found the strongest negative associations among individuals with depressive symptoms, regardless of antidepressant usage status (yes versus no).

Our initial hypothesis posited that both long and short sleep durations correlate with poorer cognitive function, as the balance of evidence points to a U-shaped association between sleep duration and cognition.<sup>14-18</sup> Our findings partially support this hypothesis, corroborating the association between long sleep duration, worse global cognition, and poorer performance in multiple cognitive domains.<sup>14-18</sup> Contrary to our expectations, we did not find a statistically significant relationship between short sleep duration and

cognitive performance, with findings remaining consistent in the sensitivity analysis, including middle-aged and older adults ( $\geq 45$  years) only. These results diverge from some studies that have identified short sleep as a risk factor for cognitive impairment,<sup>14–18</sup> while aligning with others who reported a significant association between long sleep duration and poorer cognition.<sup>23,24</sup>

When exploring the role of depression in the relationship between sleep duration and cognition, we found that, in individuals with depressive symptoms ( $\text{CES-D} \geq 16$ ), associations between long sleep and cognition were more pronounced relative to controls. Specifically, in those with no prescribed antidepressants but with depressive symptoms ( $\text{CES-D} \geq 16$ ), long sleep was significantly associated with poorer performance in global cognition, executive function, and visuospatial memory. In those using antidepressants and with depressive symptoms ( $\text{CES-D} \geq 16$ ), long sleep was associated with worse global cognitive score and visuospatial memory. Conversely, no associations were found in those using antidepressants and potentially remitted depression ( $\text{CES-D} < 16$ ), suggesting that effective treatment may help normalize sleep patterns and, as a consequence, improve cognitive function in some individuals with depression.<sup>45</sup> These findings contrast with a recent study by Wang et al.,<sup>46</sup> who found that long sleep duration was associated with better cognitive scores (TICS-10, word recall, figure drawing) among those with depressive symptoms ( $\text{CES-D} 10 \geq 10$ ) among 12,589 participants aged 45 and older from the China Health and Retirement Longitudinal Study (CHARLS). While both studies highlight the complex relationship between sleep, cognition, and depression, several factors could account for the divergent results. Differences in study populations, with Wang et al. concentrating on a Chinese cohort and our research predominantly

involving White individuals residing in the U.S., suggest possible environmental, cultural, and/or ethnic differences in sleep patterns and the manifestation of depression symptoms.<sup>47</sup> In addition, the cognitive assessments varied substantially, with our study using a comprehensive neuropsychological battery, potentially capturing subtle cognitive deficits associated with long sleep duration that could have been missed by the shorter screening measures used by Wang and colleagues.<sup>46</sup> Further research is required to shed light on these relationships across diverse populations.

The reciprocal association between sleep disturbances and depression may create a cycle in which disrupted sleep exacerbates depressed symptoms, which in turn influences sleep quality.<sup>48</sup> Sleep disturbance may increase vulnerability to depression by altering neural sensitivity to inflammation<sup>49</sup> and disrupting neurotransmitter systems involved in mood regulation, such as serotonin and dopamine.<sup>50</sup> Conversely, depressive symptoms can lead to sleep disruptions due to factors like worrying and difficulty coping with stress.<sup>51</sup> Chronic stress and associated physiological changes, such as inflammation and dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis, can further exacerbate depression and cognitive impairment.<sup>52,53</sup> In those with depression, the HPA axis is usually hyperactive, potentially leading to fragmented sleep.<sup>54,55</sup> The poorer cognitive performance observed in long sleepers may partly be attributed to lighter, fragmented sleep associated with depression characterized by frequent awakenings and less time spent in restorative, consolidated sleep stages. Slow-wave sleep (SWS) is crucial in cognitive function and brain maintenance.<sup>56-58</sup> During SWS, the brain engages in a clearance process in which the glymphatic system eliminates metabolic waste products such as beta-amyloid and tau proteins.<sup>56-58</sup> Fragmented sleep can suppress or reduce the

duration of SWS,<sup>59</sup> impeding this clearance mechanisms and potentially facilitating the buildup of neurotoxic proteins leading to cognitive deterioration. Future research with using objective measures such as polysomnography and actigraphy can provide insights into sleep architecture and fragmentation and help clarify the mechanisms which link long sleep duration to cognition.

Studies have suggested that long sleep duration may be an early and potentially reversible biological marker of neurodegeneration and a risk factor for dementia onset later in life.<sup>60,61</sup> In our cross-sectional analysis, we observed that long sleep duration was associated with lower performance in specific cognitive domains, particularly executive function and visuospatial memory. These associations were present across our full age range, including adults younger than 45 years, and may reflect broader relationships between sleep duration and cognition rather than neurodegenerative processes specifically. Longitudinal studies integrating multimodal biomarker assessments of sleep, neuropsychiatric symptoms, and AD pathology are needed to better understand these relationships and to identify opportunities for therapeutic intervention.

Finally, significant associations between long sleep duration and poorer cognitive performance were also found in the control group (not using antidepressants and without depressive symptoms). The control group had a much larger sample size compared to the other three depression groups, which increases statistical power and the likelihood of detecting even small effects. The effect sizes in the control group, however, were smaller than in the groups with depressive symptoms. Hence, while the associations were statistically significant, they may not be clinically significant. Nonetheless, these associations highlight how the relationship between sleep duration and cognitive function

is not exclusive to those with depressive symptoms. Underlying mechanisms operating independently of depression, such as sleep architecture, circadian rhythms<sup>62</sup>, or subclinical depressive symptoms<sup>63</sup>, may influence the relationship sleep duration and cognition. Earlier identification of changes in sleep duration and cognitive abilities could facilitate timely prevention and intervention, particularly against accelerated cognitive decline. A multidisciplinary care model involving sleep specialists, neuropsychologists, and mental health professionals may be valuable in enhancing early detection and intervention strategies. Further research is needed to determine which specific sleep interventions, such as cognitive behavioral therapy for insomnia (CBT-I), sleep hygiene education, or light therapy, might be most effective and feasible to implement in clinical practice for people with depression and to determine whether existing interventions are effective or require adaptations to better address the unique sleep challenges faced by individuals with depression.

#### 4.1 Strengths and Limitations

We used a comprehensive battery of validated neuropsychological tests to objectively assess cognitive performance across multiple domains. This granular cognitive profiling allowed us to determine specific domains that may be preferentially associated with long sleep duration. The large, community-based sample of adults provided sufficient statistical power to detect associations between sleep exposures and various cognition outcomes. Our categorization of depression, considering both symptom severity and antidepressants usage status, allowed for a more nuanced examination of how depressive symptoms and treatment status might modify the sleep-cognition relationship, contributing valuable insights to the existing body of research. These findings highlight the need for targeted longitudinal and interventional research that incorporate objective sleep

measures to further investigate sleep and depression as co-occurrent modifiable risk factors for cognitive decline.

Our study has some limitations. Its cross-sectional design prevented us from determining causality of the relationships we observed between sleep and cognition. We utilized self-reported measures to assess sleep duration, which can be subject to recall biases, including overestimation of actual time asleep versus time spent in bed. Additionally, some antidepressant medications can prolong sleep time and alter sleep patterns and architecture, potentially contributing to the poorer cognitive performance observed in individuals with long sleep duration and active depression.<sup>31</sup> Finally, the sample primarily consisted of White participants, which limits the generalizability of the findings to more diverse populations. Longitudinal studies including diverse and ethnic backgrounds may enhance the applicability of findings to broader communities.

## 5 CONCLUSION

We found that long but not short sleep duration was associated with poorer global cognition and specific cognitive abilities like memory, visuospatial skills, and executive functions. These associations were notably stronger in people with depressive symptoms, regardless of antidepressants usage. While our study corroborates the view of the multifactorial and complex nature of cognitive function, future longitudinal studies including large-scale, multi-modal approaches—are needed to further elucidate the temporal relationship between sleep disturbances and cognitive changes. These research efforts can inform the adaptation of existing interventions to preserve cognitive health and independence in aging populations, particularly among those with depressive symptoms.

## References

1. Langa KM. Cognitive aging, dementia, and the future of an aging population. In: National Academies of Sciences, Engineering, and Medicine. *Future Directions for the Demography of Aging: Proceedings of a Workshop*. Washington, DC: National Academies Press; 2018. Accessed August 11, 2024. <https://www.ncbi.nlm.nih.gov/books/NBK513075/>
2. 2024 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2024;20(5):3708-382. doi:10.1002/alz.13809
3. Wollesen B, Schott N, Klotzbier T, et al. Cognitive, physical and emotional determinants of activities of daily living in nursing home residents—a cross-sectional study within the PROCARE-project. *Eur Rev Aging Phys Act*. 2023; 20:17. doi:10.1186/s11556-023-00327-2
4. Kawaharada R, Sugimoto T, Matsuda N, Tsuboi Y, Sakurai T, Ono R. Impact of loss of independence in basic activities of daily living on caregiver burden in patients with Alzheimer's disease: A retrospective cohort study. *Geriatr Gerontol Int*. 2019;19(12):1243–7. doi:10.1111/ggi.13803
5. Avgerinos KI, Manolopoulos A, Ferrucci L, Kapogiannis D. Critical assessment of anti-amyloid- $\beta$  monoclonal antibodies effects in Alzheimer's disease: a systematic review and meta-analysis highlighting target engagement and clinical meaningfulness. *Sci Rep*. 2024;14(1):25741. doi:10.1038/s41598-024-75204-8
6. Livingston G, Huntley J, Liu KY, et al. Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission. *The Lancet*. 2024; 404(10452):572–628. doi:10.1016/S0140-6736(24)01296-0
7. Feinberg I, Carlson VR. Sleep variables as a function of age in man. *Arch Gen Psychiatry* 1968;18(2):239–50. doi:10.1001/archpsyc.1968.01740020111014
8. Dzierzewski JM, Dautovich N, Ravyts S. Sleep and cognition in the older adult. *Sleep Med Clin*. 2018;13(1):93–106. doi:10.1016/j.jsmc.2017.09.009
9. Ju YES, Lucey BP, Holtzman DM. Sleep and Alzheimer's disease pathology--a bidirectional relationship. *Nat Rev Neurol*. 2014;10(2):115–9. doi:10.1038/nrneurol.2013.269
10. Hafycz JM, Naidoo NN. Sleep, aging, and cellular health: Aged-related changes in sleep and protein homeostasis converge in neurodegenerative diseases. *Front Aging Neurosci*. 2019;11. doi:10.3389/fnagi.2019.00140
11. Tsiknia AA, Parada H, Banks SJ, Reas ET. Sleep quality and sleep duration predict brain microstructure among community-dwelling older adults. *Neurobiol Aging*. 2023; 125:90–7. doi:10.1016/j.neurobiolaging.2023.02.001
12. Sabia S, Fayosse A, Dumurgier J, et al. Association of sleep duration in middle and old age with incidence of dementia. *Nat Commun*. 2021 Apr 20; 12(1):2289. doi:10.1038/s41467-021-22354-2

13. Global Council on Brain Health. The brain-sleep connection: GCBH recommendations on sleep and brain health. Washington, DC: *Global Council on Brain Health*; 2017. doi:10.26419/pia.00014.001
14. Xu L, Jiang CQ, Lam TH, et al. Short or long sleep duration is associated with memory impairment in older Chinese: the Guangzhou Biobank Cohort Study. *Sleep*. 2011;34(5):575–80. doi:10.1093/sleep/34.5.575
15. Wang X, Chen Y, Yue B, Li S, Liu Q, Li Q, et al. Association of changes in self-reported sleep duration with mild cognitive impairment in the elderly: a longitudinal study. *Aging*. 2021;13(11):14816–28. doi:10.18632/aging.203149
16. Lima NC, Kirov R, de Almondes KM. Impairment of executive functions due to sleep alterations: An integrative review on the use of P300. *Front Neurosci*. 2022;16:906492. doi:10.3389/fnins.2022.906492
17. Krause AJ, Simon EB, Mander BA, et al. The sleep-deprived human brain. *Nat Rev Neurosci*. 2017;18(7):404–18. doi:10.1038/nrn.2017.55
18. Sindi S, Pérez LM, Vetrano DL, et al. Sleep disturbances and the speed of multimorbidity development in old age: results from a longitudinal population-based study. *BMC Med*. 2020;18(1):382. doi:10.1186/s12916-020-01846-w
19. Ma Y, Liang L, Zheng F, Shi L, Zhong B, Xie W. Association between sleep duration and cognitive decline. *JAMA Netw Open*. 2020;3(9):e2013573. doi:10.1001/jamanetworkopen.2020.13573
20. Lo JC, Loh KK, Zheng H, Sim SKY, Chee MWL. Sleep duration and age-related changes in brain structure and cognitive performance. *Sleep*. 2014;37(7):1171–8. doi:10.5665/sleep.3832
21. Keil SA, Schindler AG, Wang MX, et al. Longitudinal sleep patterns and cognitive impairment in older adults. *JAMA Network Open*. 2023;6(12):e2346006. doi:10.1001/jamanetworkopen.2023.46006
22. Stephens ER, Sarangi A, Gude J. Short sleep duration and dementia: a narrative review. *Proc (Bayl Univ Med Cent)*. 2022;35(3):328–31. doi:10.1080/08998280.2022.2026123
23. Faubel R, López-García E, Guallar-Castillón P, Graciani A, Banegas JR, Rodríguez-Artalejo F. Usual sleep duration and cognitive function in older adults in Spain. *J Sleep Res*. 2009;18(4):427–35. doi:10.1111/j.1365-2869.2009.00759.x
24. Behrens A, Anderberg P, Berglund JS. Sleep disturbance predicts worse cognitive performance in subsequent years: A longitudinal population-based cohort study. *Arch Gerontol Geriatr*. 2023; 106:104899. doi: 10.1016/j.archger.2022.104899
25. Consensus Conference Panel, Watson NF, Badr MS, et al. Joint Consensus Statement of the American Academy of Sleep Medicine and Sleep Research Society on the Recommended Amount of Sleep for a Healthy Adult: Methodology and Discussion. *Sleep*. 2015;38(8):1161–1183. doi:10.5665/sleep.4886



26. Jaussent I, Empana JP, Ancelin ML, et al. Insomnia, daytime sleepiness and cardio-cerebrovascular diseases in the elderly: a 6-year prospective study. *PLoS One*. 2013;8(2):e56048. doi: 10.1371/journal.pone.0056048
27. Fang H, Tu S, Sheng J, Shao A. Depression in sleep disturbance: A review on a bidirectional relationship, mechanisms and treatment. *J Cell Mol Med*. 2019;23(4):2324-2332. doi:10.1111/jcmm.14170
28. Alvaro PK, Roberts RM, Harris JK. A systematic review assessing bidirectionality between sleep disturbances, anxiety, and depression. *Sleep*. 2013;36(7):1059-1068. doi:10.5665/sleep.2810
29. Perini G, Cotta Ramusino M, Sinforiani E, Bernini S, Petrachi R, Costa A. Cognitive impairment in depression: recent advances and novel treatments. *Neuropsychiatr Dis Treat*. 2019;15:1249-1258. doi:10.2147/NDT.S199746
30. Zuckerman H, Pan Z, Park C, et al. Recognition and treatment of cognitive dysfunction in major depressive disorder. *Front Psychiatry*. 2018;9:655. doi:10.3389/fpsy.2018.00655
31. Ly M, Karim HT, Becker JT, et al. Late-life depression and increased risk of dementia: a longitudinal cohort study. *Transl Psychiatry*. 2021;11(1):147. doi:10.1038/s41398-021-01269-y
32. Byers AL, Yaffe K. Depression and risk of developing dementia. *Nat Rev Neurol*. 2011;7(6):323-331. doi:10.1038/nrneurol.2011.60
33. Baril AA, Kojis DJ, Himali JJ, et al. Association of sleep duration and change over time with imaging biomarkers of cerebrovascular, amyloid, tau, and neurodegenerative pathology. *Neurology*. 2024;102(1):e207807. doi:10.1212/WNL.0000000000207807
34. Au R, Seshadri S, Wolf PA, et al. New norms for a new generation: cognitive performance in the framingham offspring cohort. *Exp Aging Res*. 2004;30(4):333-358. doi:10.1080/03610730490484380
35. Tombaugh TN. Trail Making Test A and B: normative data stratified by age and education. *Arch Clin Neuropsychol*. 2004;19(2):203-214. doi:10.1016/S0887-6177(03)00039-8
36. Elwood RW. The Wechsler Memory Scale-Revised: psychometric characteristics and clinical application. *Neuropsychol Rev*. 1991;2(2):179-201. doi:10.1007/BF01109053
37. Drozdick LW, Wahlstrom D, Zhu J, Weiss LG. The Wechsler Adult Intelligence Scale—Fourth Edition and the Wechsler Memory Scale—Fourth Edition. In: Flanagan DP, Harrison PL, eds. *Contemporary Intellectual Assessment: Theories, Tests, and Issues*. 3rd ed. New York, NY: The Guilford Press; 2012:197-223
38. Pase MP, Harrison S, Misialek JR, et al. Sleep architecture, obstructive sleep apnea, and cognitive function in adults. *JAMA Netw Open*. 2023;6(7):e2325152. doi:10.1001/jamanetworkopen.2023.25152

39. Radloff LS. The CES-D Scale: A self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977;1(3):385–401. doi: [10.1177/014662167700100306](https://doi.org/10.1177/014662167700100306)
40. ATC/DDD Index. Norwegian Institute of Public Health. Accessed July 20, 2024. [https://atcddd.fhi.no/atc\\_ddd\\_index/?code=N06A](https://atcddd.fhi.no/atc_ddd_index/?code=N06A)
41. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7). *National Heart, Lung, and Blood Institute; NIH.* Accessed June 1, 2024. <https://www.nhlbi.nih.gov/health-topics/seventh-report-of-joint-national-committee-on-prevention-detection-evaluation-and-treatment-high-blood-pressure>
42. Framingham Heart Study. Manual of procedures: section 9, phlebotomy. 2018. Accessed July 20, 2024. <https://www.framinghamheartstudy.org/files/2019/05/Gen-3-Exam-3-Onsite-Sec-9.pdf>
43. Pearson TA, Mensah GA, Alexander RW, et al. Markers of inflammation and cardiovascular disease: application to clinical and public health practice: A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation.* 2003;107(3):499-511. doi:10.1161/01.cir.0000052939.59093.45
44. National Library of Medicine. Middle age. 2017. Accessed November 3, 2024. <https://meshb-prev.nlm.nih.gov/record/ui?ui=D008875>
45. Thase ME. Depression and sleep: pathophysiology and treatment. *Dialogues Clin Neurosci.* 2006;8(2):217-226. doi:10.31887/DCNS.2006.8.2/mthase
46. Wang L, He S, Yan N, Pan R, Niu Y, Li J. Mediating role of depressive symptoms on the relationship between sleep duration and cognitive function. *Sci Rep.* 2023;13(1):4067. Published 2023 Mar 11. doi:10.1038/s41598-023-31357-6
47. Estafanous A, Sedky K. Culture, ethnicity, and sleep patterns and disorders. In: Gogineni RR, Pumariega AJ, Kallivayalil RA, et al., eds. *The WASP Textbook on Social Psychiatry: Historical, Developmental, Cultural, and Clinical Perspectives.* Oxford University Press; 2023:0. doi:10.1093/med/9780197521359.003.0031
48. Peakman G, Karunatilake N, Seynaeve M, et al. Clinical factors associated with progression to dementia in people with late-life depression: a cohort study of patients in secondary care. *BMJ Open.* 2020;10(5):e035147. doi:10.1136/bmjopen-2019-035147
49. Irwin MR, Opp MR. Sleep Health: Reciprocal regulation of sleep and innate immunity. *Neuropsych.* 2017;42(1):129-155. doi:10.1038/npp.2016.148
50. Weicker H, Strüder HK. Influence of exercise on serotonergic neuromodulation in the brain. *Amino Acids.* 2001;20(1):35-47. doi:10.1007/s007260170064
51. Zhai L, Zhang H, Zhang D. Sleep duration and depression among adults: A meta-analysis of prospective studies. *Depress Anxiety.* 2015;32(9):664–70. doi:10.1002/da.22386

52. Job E, Kirschbaum C, Steptoe A. Persistent depressive symptoms, HPA-axis hyperactivity, and inflammation: the role of cognitive-affective and somatic symptoms. *Mol Psychiatry*. 2020;25(5):1130-1140. doi:10.1038/s41380-019-0501-6
53. Hassamal S. Chronic stress, neuroinflammation, and depression: an overview of pathophysiological mechanisms and emerging anti-inflammatories. *Front Psychiatry*. 2023;14:1130989. Published 2023 May 11. doi:10.3389/fpsyt.2023.1130989
54. Nicolaidis NC, Vgontzas AN, Kritikou I, Chrousos G. HPA axis and sleep. In: Feingold KR, Anawalt B, Blackman MR, et al., eds. *Endotext*. South Dartmouth, MA: MDText.com, Inc.; 2000. Accessed July 20, 2024. <http://www.ncbi.nlm.nih.gov/books/NBK279071/>
55. Hirotsu C, Tufik S, Andersen ML. Interactions between sleep, stress, and metabolism: From physiological to pathological conditions. *Sleep Sci*. 2015;8(3):143-152. doi:10.1016/j.slsci.2015.09.002
56. Voumvourakis KI, Sideri E, Papadimitropoulos GN, et al. The Dynamic Relationship between the Glymphatic System, Aging, Memory, and Sleep. *Biomedicines*. 2023;11(8):2092. doi:10.3390/biomedicines11082092
57. Wafford KA. Aberrant waste disposal in neurodegeneration: why improved sleep could be the solution. *Cereb Circ - Cogn Behav*. 2021;2:100025. doi:10.1016/j.cccb.2021.100025
58. Li J, Vitiello MV, Gooneratne NS. Sleep in Normal Aging. *Sleep Med Clin*. 2022;17(2):161-171. doi:10.1016/j.jsmc.2022.02.007
59. Kroeger D, Vetrivelan R. To sleep or not to sleep - Effects on memory in normal aging and disease. *Aging Brain*. 2023;3:100068. Published 2023 Jan 30. doi:10.1016/j.nbas.2023.100068
60. Westwood AJ, Beiser A, Jain N, et al. Prolonged sleep duration as a marker of early neurodegeneration predicting incident dementia. *Neurology*. 2017;88(12):1172-1179. doi:10.1212/WNL.0000000000003732
61. Burke S, Grudzien A, Li T, et al. Correlations between sleep disturbance and brain structures associated with neurodegeneration in the National Alzheimer's Coordinating Center Uniform Data Set. *J Clin Neurosci*. 2022;106:204-212. doi:10.1016/j.jocn.2022.07.012
62. Leahy S, Xiao Q, Yeung CHC, Figueiro MG. Associations between circadian alignment and cognitive functioning in a nationally representative sample of older adults. *Sci Rep*. 2024;14(1):13509. Published 2024 Jun 12. doi:10.1038/s41598-024-64309-9
63. Hsieh CJ, Chiou JM, Chen TF, Chen YC, Chen JH. Association of subclinical depressive symptoms and sleep with cognition in the community-dwelling older adults. *J Formos Med Assoc*. 2023;122(8):790-799. doi:10.1016/j.jfma.2023.03.005





**Table 1.** Sample characteristics

	<b>Whole Sample</b>	<b>Short Sleep (≤6 hours)</b>	<b>Average Sleep (&gt;6-&lt;9 hours)</b>	<b>Long Sleep (≥ 9 hours)</b>
N, n (%)	1853	451 (24.3)	1277 (68.9)	125 (6.8)
Age at NP, years	49.84 (9.2)	50.57 (8.2)	49.57 (9.3)	49.92 (11.1)
Male, n (%)	791 (42.7)	240 (53.2)	511 (40.0)	40 (32.0)
Ethnicity, n (%)				
White	1760 (95.3)	422 (93.6)	1218 (95.8)	121 (96.8)
Black	19 (1.0)	12 (2.7)	7 (0.6)	0
Hispanic	33 (1.8)	7 (1.6)	22 (1.7)	4 (3.2)
Other	35 (1.9)	10 (2.2)	24 (1.9)	1 (0.8)
Education, n (%)				
Up to High School	262 (14.1)	81 (18.0)	162 (12.7)	19 (15.2)
Some College	532 (28.7)	153 (33.9)	338 (26.5)	41 (32.8)
College degree	1059 (57.2)	217 (48.1)	777 (60.9)	65 (52.0)
Time interval <sup>a</sup> , years	1.69 (1.0)	1.61 (0.9)	1.73 (1.0)	1.59 (1.0)
APOE ε4 carriers, n (%)	384 (21.8)	89 (20.7)	273 (22.5)	22 (18.8)
Total Cholesterol, mg/dL	185.98 (35.0)	185.28 (35.0)	186.05 (35.2)	187.71 (33.7)
HDL, mg/dL	60.53 (17.9)	58.26 (17.9)	61.01 (17.7)	63.67 (19.7)
Systolic BP, mm/Hg	116 (14.1)	117 (13.6)	116 (14.1)	116 (16.9)
HTN treatment, n (%)	390 (21.1)	101 (22.5)	259 (20.4)	30 (24.2)
Stage 1 Hypertension, n (%)	473 (25.7)	121 (27.0)	318 (25.0)	34 (27.4)
Triglycerides, mg/dL	112.8 (85.8)	116.89 (95.2)	111.03 (83.3)	116.16 (75.0)
CRP, mg/dL	2.83 (4.5)	2.72 (3.9)	2.73 (4.4)	4.29 (7.2)
Body Mass Index	28.19 (5.8)	29.24 (5.7)	27.83 (5.7)	28.15 (6.4)
Depression status, n (%)				
Antidepressants usage	315 (17.0)	57 (12.6)	215 (16.8)	43 (34.4)
Depression, CES-D ≤16	198 (10.7)	65 (14.4)	110 (8.6)	23 (18.4)
Control <sup>b</sup>	1405 (75.8)	345 (76.5)	990 (77.5)	70 (56.0)
Antidepressants usage/CES-D <16 <sup>c</sup>	250 (13.5)	41 (9.1)	177 (13.9)	32 (25.6)
No Antidepressants usage/CES-D ≥ 16 <sup>d</sup>	133 (7.2)	49 (10.9)	72 (5.6)	12 (9.6)
Antidepressants usage/CES-D ≥16 <sup>e</sup>	65 (3.5)	16 (3.6)	38 (3.0)	11 (8.8)

Abbreviations: NP= Neuropsychological assessment, BP = Blood Pressure; HDL = high-density lipoprotein cholesterol; HTN = hypertension; CRP = C reactive protein. All values represent mean (SD) unless otherwise indicated by n (%).

NOTE. <sup>a</sup> time interval between self-reported sleep duration and cognitive tests; <sup>b</sup> Control = no antidepressants usage and no depressive symptoms (CES-D <16); <sup>c</sup> antidepressant usage without depressive symptoms (CES-D <16); <sup>d</sup> no antidepressant usage with depressive symptoms (CES-D ≥ 16); <sup>e</sup> antidepressant usage and with depressive symptoms (CES-D ≥ 16).



**Table 2.** Sample cognitive characteristics

	<b>Whole Sample</b>	<b>Short Sleep (≤6 hours)</b>	<b>Average Sleep (&gt;6-&lt;9 hours)</b>	<b>Long Sleep (≥ 9 hours)</b>
N, n (%)	1853	451 (24.3)	1277 (68.9)	125 (6.8)
Global Cognition, <sup>a</sup> median [Q1, Q3]	0.45 [-0.12,0.98]	0.32 [-0.19,0.91]	0.49 [-0.06,1.01]	0.15 [-0.49,0.88]
Trails Part A, min, median [Q1, Q3]	0.40 [0.33,0.48]	0.40 [0.33,0.48]	0.40 [0.33,0.48]	0.40 [0.32,0.55]
Trails Part B, min, median [Q1, Q3]	0.97 [0.77,1.22]	1.00 [0.77,1.30]	0.95 [0.75,1.20]	0.97 [0.80,1.33]
Visual Reproduction <sup>b</sup> , n correct	17.99 (4.89)	17.56 (5.09)	18.31 (4.70)	16.22 (5.56)
Logical Memories <sup>b</sup> , n correct	24.46 (6.74)	24.14 (6.59)	24.71 (6.74)	23.12 (7.12)
Similarities Test, n correct	17.20 (3.15)	17.08 (3.14)	17.27 (3.17)	16.95 (3.01)

Abbreviations: Min = minutes.

NOTE. <sup>a</sup> Weighted score units; <sup>b</sup> Sum of immediate and delayed recall scores. All values represent mean (SD) unless otherwise indicated by median [Q1,Q3].



**Table 3.** Sample characteristics by depression group

	Control <sup>a</sup>	Antidepressants usage/CES-D <16 <sup>b</sup>	No Antidepressants usage/CES-D ≥ 16 <sup>c</sup>	Antidepressants usage/ CES-D ≥16 <sup>d</sup>
N, n (%)	1405 (75.8)	250 (13.5)	133 (7.2)	65 (3.5)
Age at NP, years	49.87 (9.3)	50.38 (8.7)	48.51 (8.8)	49.74 (8.8)
Male, n (%)	656 (46.7)	59 (23.6)	54 (40.6)	22 (33.8)
Ethnicity, n (%)				
White	1330 (95.0)	243 (97.2)	124 (93.2)	63 (98.4)
Black	14 (1.0)	2 (0.0)	3 (2.3)	0
Hispanic	26 (1.9)	3 (1.2)	3 (2.3)	1 (1.6)
Other	30 (2.1)	2 (0.8)	3 (2.3)	0
Education, n (%)				
Up to High School	179 (12.7)	38 (15.2)	28 (21.1)	17 (26.2)
Some College	390 (27.8)	78 (31.2)	40 (30.1)	24 (36.9)
College degree	836 (59.5)	134 (53.6)	65 (48.9)	24 (36.9)
Time interval <sup>e</sup> , years	1.74 (1.0)	1.57 (0.9)	1.47 (0.9)	1.59 (1.1)
APOE ε4 carriers, n (%)	293 (21.9)	54 (22.6)	24 (18.9)	13 (21.7)
Total Cholesterol, mg/dL	184.91 (35.0)	191.38 (33.4)	185.77 (36.6)	188.69 (32.4)
HDL, mg/dL	60.14 (17.9)	63.06 (18.1)	60.25 (18.6)	59.69 (15.8)
Systolic BP, mm/Hg	116 (14.1)	115.8 (13.3)	115 (15.0)	117 (15.2)
HTN treatment, n (%)	287 (20.5)	54 (21.7)	33 (24.8)	16 (24.6)
Stage 1 Hypertension, n (%)	347 (24.8)	62 (24.9)	43 (32.6)	21 (32.3)
Triglycerides, mg/dL	111.5 (84.3)	113.7 (64.6)	110.6 (70.7)	141.6 (171.7)
CRP, mg/dL	2.72 (4.6)	4.28 (5.3)	2.79 (3.6)	4.28 (5.3)
Body Mass Index	27.89 (5.4)	28.61 (6.2)	29.80 (7.7)	29.91 (7.1)
Short Sleep (≤6 hours)	345 (24.6)	41 (16.4)	49 (36.8)	16 (24.6)
Average Sleep (>6-<9 hours)	990 (70.5)	177 (70.8)	72 (54.1)	38 (58.5)
Long Sleep (≥ 9 hours)	70 (5.0)	32 (12.8)	12 (9.0)	11 (16.9)

Abbreviations: CES-D = Center for Epidemiologic Studies Depression Scale, NP= Neuropsychological assessment, BP = Blood Pressure; HDL = high-density lipoprotein cholesterol; HTN = hypertension; CRP = C reactive protein. All values represent mean (SD), unless otherwise indicated by n (%).

NOTE. <sup>a</sup> Control = no antidepressant usage and no depressive symptoms (CES-D <16); <sup>b</sup> antidepressant usage without depressive symptoms (CES-D <16); <sup>c</sup> no antidepressant usage with depressive symptoms (CES-D ≥ 16); <sup>d</sup> antidepressant usage and with depressive symptoms (CES-D ≥ 16).

<sup>e</sup> time interval between self-reported sleep duration and cognitive tests



**Table 4.** Cognitive characteristics by depression group

	<b>Control <sup>a</sup></b>	<b>Antidepressants usage /CES-D &lt;16 <sup>b</sup></b>	<b>No Antidepressants usage /CES-D ≥ 16 <sup>c</sup></b>	<b>Antidepressants usage /CES-D ≥16 <sup>d</sup></b>
N, n (%)	1405 (75.82)	250 (13.49)	133 (7.18)	65 (3.51)
Global Cognition <sup>e</sup> , median [Q1, Q3]	0.47 [-0.09, 0.99]	0.41 [-0.19, 0.91]	0.42 [-0.33, 0.96]	0.12 [-0.38, 0.75]
Trails Part A, min, median [Q1, Q3]	0.40 [0.33, 0.48]	0.40 [0.33, 0.48]	0.42 [0.33, 0.50]	0.42 [0.32, 0.52]
Trails Part B, min, median [Q1, Q3]	0.95 [0.75, 1.22]	0.98 [0.78, 1.20]	0.95 [0.75, 1.27]	1.12 [0.88, 1.45]
Visual Reproduction <sup>f</sup> , n correct	18.21 (4.81)	17.44 (4.64)	17.70 (5.53)	15.88 (5.56)
Logical Memories <sup>f</sup> , n correct	24.44 (6.70)	24.99 (7.03)	24.17 (6.53)	23.46 (7.04)
Similarities Test, n correct	17.26 (3.10)	17.12 (2.97)	16.87 (3.42)	16.94 (4.10)

Abbreviations: CES-D = Center for Epidemiologic Studies Depression Scale, Min = minutes.

NOTE. <sup>a</sup> Control = no antidepressants usage and no depressive symptoms (CES-D <16); <sup>b</sup> antidepressant usage without depressive symptoms (CES-D <16); <sup>c</sup> no antidepressant usage with depressive symptoms (CES-D ≥ 16); <sup>d</sup> antidepressant usage and with depressive symptoms (CES-D ≥ 16).

<sup>e</sup> weighted score units; <sup>f</sup> Sum of immediate and delayed recall scores combined). All values represent mean (SD) unless otherwise indicated by median [Q1, Q3]

**Table 5.** Association between short and long sleep duration and cognitive scores.

Cognition	Model	Average (>6-<9 hours)	Sleep Duration Categories			
			Short Sleep (≤6 hours)		Long Sleep (≥ 9 hours)	
			β±SE	p	β±SE	p
Global Cognition	1	REF	-0.03±0.04	0.465	<b>-0.25±0.07</b>	<b>&lt;0.001</b>
	2	REF	-0.04±0.04	0.280	<b>-0.27±0.07</b>	<b>&lt;0.001</b>
Trails Part A	1	REF	-0.001±0.02	0.959	-0.03±0.03	0.205
	2	REF	-0.01±0.02	0.550	-0.03±0.03	0.279
Trails Part B	1	REF	-0.02±0.02	0.235	<b>-0.09±0.03</b>	<b>0.009</b>
	2	REF	-0.04±0.02	0.067	<b>-0.07±0.03</b>	<b>0.041</b>
Visual Reproduction	1	REF	-0.41±0.25	0.104	<b>-1.80±0.42</b>	<b>&lt;0.001</b>
	2	REF	-0.45±0.26	0.081	<b>-1.76±0.44</b>	<b>&lt;0.001</b>
Logical Memory	1	REF	0.25±0.35	0.480	<b>-1.50±0.60</b>	<b>0.013</b>
	2	REF	0.22±0.36	0.538	<b>-2.16±0.62</b>	<b>&lt;0.001</b>
Similarities Test	1	REF	0.01±0.17	0.954	-0.06±0.28	0.839
	2	REF	0.002±0.17	0.990	-0.11±0.29	0.701

NOTE. Model 1 was adjusted for age and age squared at neuropsychological testing, education, sex, time between sleep duration questionnaires and neuropsychological testing, and cohort. Model 2 was further adjusted for hypertension, total cholesterol level to HDL ratio, triglycerides, C reactive protein level, body mass index, and APOE ε4 status.

**Table 6.** Effect modification by depression group on the association between self-reported sleep duration and cognitive scores.

		Global Cognition		Trails Part B		Visual Reproduction		Similarities	
Sleep Duration Interaction		p=0.028		p= 0.012		p=0.064		p=0.027	
		β±SE	p	β±SE	p	β±SE	p	β±SE	p
<b>Control<sup>a</sup></b>	Average Sleep	REF		REF		REF		REF	
	Short Sleep	-0.03±0.04	0.522	-0.03±0.02	0.235	-0.31±0.28	0.270	-0.05±0.19	0.776
	Long Sleep	<b>-0.18±0.09</b>	<b>0.044</b>	-0.04±0.04	0.379	<b>-1.30±0.55</b>	<b>0.019</b>	0.24±0.37	0.512
<b>Antidepressants usage and CES-D &lt;16<sup>b</sup></b>	Average Sleep	REF		REF		REF		REF	
	Short Sleep	-0.06±0.13	0.632	-0.07±0.06	0.273	-0.76±0.74	0.304	-0.13±0.50	0.790
	Long Sleep	-0.07±0.14	0.621	-0.03±0.07	0.632	-0.46±0.83	0.581	0.18±0.55	0.748
<b>No Antidepressants usage and CES-D ≥ 16<sup>c</sup></b>	Average Sleep	REF		REF		REF		REF	
	Short Sleep	-0.12±0.15	0.453	-0.05±0.08	0.500	-1.23±1.00	0.220	0.42±0.60	0.490
	Long Sleep	<b>-0.60±0.26</b>	<b>0.024</b>	<b>-0.29±0.13</b>	<b>0.023</b>	<b>-3.70±1.70</b>	<b>0.031</b>	-1.15±1.03	0.267
<b>Antidepressants usage and CES-D ≥16<sup>d</sup></b>	Average Sleep	REF		REF		REF		REF	
	Short Sleep	0.23±0.26	0.384	0.17±0.12	0.164	0.04±1.65	0.979	1.04±1.17	0.375
	Long Sleep	<b>-0.74±0.30</b>	<b>0.017</b>	-0.25±0.13	0.067	<b>-4.30±1.89</b>	<b>0.027</b>	-1.95±1.33	0.149

NOTE. Sleep categories: short ≤6h; average 7-8h; long ≥9h.

<sup>a</sup> Control = no antidepressants usage and no depressive symptoms (CES-D <16); <sup>b</sup> antidepressant usage without depressive symptoms (CES-D <16); <sup>c</sup> no antidepressant usage with depressive symptoms (CES-D ≥ 16); <sup>d</sup> antidepressant usage and with depressive symptoms (CES-D ≥ 16).

Model 1 was adjusted for age and age squared at neuropsychological testing, education, sex, time between sleep duration questionnaires and neuropsychological testing and cohort. Model 2 was further adjusted for hypertension, total cholesterol level to HDL ratio; triglycerides, C reactive protein level, body mass index, and APOE ε4 status. Significant findings at  $p < 0.10$  for interaction and at  $p < 0.05$  for the stratified model are shown in bold.

No significant interaction was found with Trails Part A ( $p=0.114$ ), and Logical Memory ( $p=0.624$ ).