The oral microbiome and all-cause mortality in the US population

Emily Vogtmann, PhD, MPH^{1*}, Yukiko Yano, PhD, MPH^{1*}, Jianxin Shi, PhD¹, Yunhu Wan,

PhD^{1, 2}, Vaishnavi Purandare, MS¹, Jody McLean, MPH³, Shilan Li, PhD⁴, Rob Knight, PhD⁵,

Lisa Kahle, BA⁶, Autumn G. Hullings, PhD, MPH⁷, Xing Hua, PhD^{1, 2}, Barry I. Graubard, PhD¹,

Maura L. Gillison, MD, PhD⁸, J. Gregory Caporaso, PhD⁹, Nicholas A. Bokulich, PhD¹⁰, Martin

J. Blaser, MD¹¹, Neal D. Freedman, PhD, MPH¹², Anil K. Chaturvedi, PhD^{1*}, Christian C.

Abnet, PhD, MPH^{1*}

- Division of Cancer Epidemiology and Genetics, National Cancer Institute, Bethesda, MD, USA
- Frederick National Laboratory for Cancer Research/Leidos Biomedical Research Laboratory, Inc, Frederick, MD, USA
- National Center for Health Statistics, Centers for Disease Control and Prevention, Hyattsville, MD, USA
- 4) Johns Hopkins University, Baltimore, MD, USA
- Center for Microbiome Innovation, University of California San Diego, La Jolla, CA, USA
- 6) Information Management Services, Inc, Calverton, MD, USA
- 7) Department of Nutrition, University of North Carolina, Chapel Hill, NC, USA
- Department of Thoracic and Head and Neck Medical Oncology, MD Anderson Cancer Center, Houston, TX, USA

 Center for Applied Microbiome Science, Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, AZ, USA

 Laboratory of Food Systems Biotechnology, Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland

- 11) Center for Advanced Biotechnology and Medicine, Rutgers, Piscataway, NJ, USA
- 12) Division of Cancer Control and Population Sciences, National Cancer Institute, Bethesda,

MD, USA

*These authors contributed equally

Acknowledgements

We would like to acknowledge Gail Ackermann, Amnon Amir, James Gaffney, Grant Gogul, Greg Humphrey, Tara Schwartz, and Weihong Xiao for their role in project management and data production. This study was supported in part by a grant from the National Institute for Dental and Craniofacial Research (R01DE023175), a grant from the US Food and Drug Administration's Center for Tobacco Products, a grant from the National Cancer Institute Informatics Technology for Cancer Research (1U24CA248454-01), the Intramural Research Program of the National Cancer Institute, and Federal funds from the National Cancer Institute, National Institutes of Health, under NCI Contract number 75N910D00024. We also acknowledge the research contributions of the Cancer Genomics Research Laboratory for their expertise, execution, and support of this research in the bioinformatics analysis of generated data. The content of this publication are those of the authors and does not necessarily reflect the views or policies of the Department of Health and Human Services or the official position of the

Centers for Disease Control and Prevention. The mention of trade names, commercial products, or organizations does not imply endorsement by the US Government.

Data sharing statement

Unlinked public-use National Health and Nutrition Examination Survey (NHANES) data files

are available on the NHANES website (https://www.cdc.gov/nchs/nhanes). The linked mortality

(National Death Index) files are available in a public-use version, available online

(https://www.cdc.gov/nchs/data-linkage/mortality.htm), and a restricted-use file available for

analysis at the National Center for Health Statistics Research Data Center

(https://www.cdc.gov/rdc/index.htm). The restricted-use mortality file was used for this analysis.

Correspondence

Christian C. Abnet, PhD, MPH Director, Metabolic Epidemiology Branch Division of Cancer Epidemiology & Genetics, National Cancer Institute 9609 Medical Center Drive, 6E410 Rockville, MD 20852 Email: abnetc@mail.nih.gov

Key points

Question: Does the human oral microbiome impact an individual's risk of mortality? Findings: In this prospective study including 7,721 individuals of which 426 died over followup, specific measures of the oral microbiome were associated with all-cause mortality.

Meaning: The microbes living in the oral cavity may play an important role in human health.

Abstract (350 words max)

Importance: Poor oral health, including periodontal disease, is associated with oral microbiome changes and increased mortality risk. However, no large studies have evaluated whether the oral microbiome is directly associated with mortality.

Objective: To evaluate whether measures of the oral microbiome is prospectively associated with all-cause mortality.

Design: A cross-sectional survey with samples collected from 2009-2012 and mortality linkage to the restricted-use National Death Index (NDI) through 2019.

Setting: The National Health and Nutrition Examination Survey (NHANES) 2009-2012, a multistage probability sample of the US population.

Participants: NHANES participants 20- to 69-years-old who were eligible for linkage to the NDI and provided oral rinse specimens (N=7,721, representing approximately 194 million individuals).

Exposure: Oral microbiome ascertained by sequencing the V4 region of the 16S rRNA gene of extracted DNA from oral rinse specimens. Alpha diversity, beta diversity, and genus-level data were generated using DADA2 and QIIME.

Main outcome and measure: All-cause mortality.

Results: After an average of 8.8 years, a total of 426 participants died. Using Cox proportional hazards regression and after controlling for multiple comparisons where appropriate, continuous alpha diversity was inversely associated with all-cause mortality, but only the association for the Shannon-Weiner index was significant with full adjustment for major risk factors (hazard ratio [HR] per standard deviation [SD]=0.85; 95% confidence interval [CI]=0.74-0.98). The principal coordinate analysis (PCoA) vector 2 from the Bray-Curtis dissimilarity matrix (HR per SD=0.83;

95% CI=0.73-0.93) and PCoA1 from weighted UniFrac (HR per SD=0.86; 95% CI=0.75-0.98) were significantly associated with all-cause mortality after full adjustment. Few associations were observed at the genus-level after Bonferroni correction, but an increase in 1 SD of the relative abundance of *Granulicatella* and *Lactobacillus* were associated with a 17% (95% CI=1.11-1.24) and 11% (95% CI=1.06-1.16) increase in mortality risk, respectively. Compared to participants with no detectable *Bacteroides*, participants in the highest tertile of *Bacteroides* had decreased mortality risk (HR=0.54; 95% CI=0.40-0.74).

Conclusions and relevance: Some measures of the oral microbiome were associated with allcause mortality in this representative population cohort. These results suggest that oral bacterial communities may be important contributors to health and disease.

Introduction

The oral microbiome serves multiple functions in human health, locally and systemically. Specific oral microbes are closely linked to periodontal disease,¹ and many others are associated with oral health conditions such as caries.² The oral microbiome also has been associated with cardiovascular disease³ and multiple site-specific cancers including lung cancer.⁴ These microorganisms also play a central role in the metabolism of ethanol to its carcinogenic intermediate acetaldehyde.⁵ The oral microbiome may be related to mortality, as previous studies have identified associations between oral health conditions and mortality,^{6,7} but no large studies of the oral microbiome including genus-level taxonomy have been conducted to directly investigate mortality associations.

We analyzed associations between the oral microbiome from participants aged 20-69 years in the 2009-2012 cycles of the National Health and Nutrition Examination Survey (NHANES)⁸ and all-cause mortality from linkage to the National Death Index (NDI) while controlling for numerous potential confounders.

Methods

NHANES is a continuous survey conducted by the National Center for Health Statistics (NCHS) to characterize the health and nutritional status of the US population. Participants are interviewed about demographics, diet, and other health-related data and then visit a mobile examination center (MEC) for biospecimen collection and health examinations.⁹ NDI mortality data through December 31, 2019 were available for eligible adult participants from the NCHS restricted-use linked mortality file.¹⁰

From 2009-2012, oral rinse samples were collected from participants aged 14-69 years which were used for microbiome analyses. After DNA extraction,¹¹ the V4 region of the 16S rRNA gene was PCR amplified and sequenced. Sequencing data were processed using DADA2 and QIIME and taxonomy was assigned using the SILVA v123 database. After removal of the amplicon sequencing variant (ASV) SV1032, a human pseudogene, alpha and beta diversity metrics were generated at an even sampling depth of 10,000 reads per sample.¹²

As fully described in the companion manuscript (MEDRXIV/2024/318415), we incorporated many potential confounders in our analysis, including continuous age and spline variables for non-linear age relationships, sex, self-identified race and ethnicity categories grouped by NHANES standard practice, education, marital status, and the income-to-poverty ratio. Detailed categories of self-reported lifetime cigarette smoking history and alcohol consumption were used. Periodontal disease status, tooth counts, and body mass index categories were calculated based on health examinations. We included self-reported use of prescription medications, including antibiotics, anti-lipidemic drugs, inhaled respiratory drugs, and drugs for treatment of gastroesophageal reflux disease. Finally, we determined each participant's diabetes and hypertension status using self-report, laboratory and examination data.

Analyses were restricted to MEC participants, aged 20-69 years with microbiome and NDI data (N=7,721, representing approximately 194 million people). We presented the characteristics of the population by mortality status and estimated hazard ratios (HR) and 95% confidence intervals (CI) for these characteristics from multivariable Cox proportional hazards models with time since the MEC visit as the underlying time metric. We estimated HRs for the association between the alpha diversity metrics (observed ASVs, Faith's phylogenetic diversity [PD], Shannon-Weiner index, and Simpson index) and principal coordinate analysis (PCoA)

vectors from the beta diversity matrices (unweighted UniFrac, weighted UniFrac, and Bray-Curtis). We estimated HRs for genus-level presence, relative abundance, centered log-ratio (CLR) transformed abundances, and a combined variable incorporating both the presence and tertiles of relative abundance. Bonferroni-adjusted p-values cutpoints were determined for genuslevel models. We used SAS and SUDAAN and incorporated the complex survey design, including post-stratification weights.

Results

After an average of 8.8 years of follow-up, a total of 426 participants died, including 135 deaths from malignant neoplasms and 91 from heart diseases. Participants who died were more likely to be older, male, experiencing poverty, categorized with Class 2 or 3 obesity, and have periodontal disease or edentulism (Table 1).

All continuous alpha diversity metrics were inversely associated with all-cause mortality after minimal adjustment (e.g., $HR_{Observed ASVs}$ per SD=0.82; 95% CI=0.68-0.98), but associations were attenuated after full adjustment (Figure 1 and Supplemental Table 1). In fully adjusted models, only the Shannon-Weiner index association remained statistically significant (HR per SD=0.85; 95% CI=0.74-0.98). The associations for quartiles of alpha diversity were inverse, but only the association for the highest quartile for the Simpson index reached statistical significance (HR_{Q4} =0.62; 95% CI=0.39-0.99; Supplemental Table 1). For beta diversity, PCoA2 from the Bray-Curtis dissimilarity (HR per SD=0.83; 95% CI=0.73-0.93) and PCoA1 from weighted UniFrac (HR per SD=0.86; 95% CI=0.75-0.98) matrices were statistically significantly associated with all-cause mortality after full adjustment (Figure 1 and Supplemental Table 1).

After Bonferroni correction, none of the genus-level presence associations were statistically significant in fully adjusted models, although some strong associations were noted (Figure 1 and Supplemental Table 2). For relative abundance, each SD increase in *Granulicatella* or *Lactobacillus* was associated with 1.17 (95% CI=1.11-1.24) and 1.11 (95% CI=1.06-1.16) times the mortality risk, respectively, which remained statistically significant after Bonferroni correction (Figure 1 and Supplemental Table 3). However, none of the CLR-transformed abundances were statistically significant in fully adjusted models, although the association with *Lactobacillus* was similar to the relative abundance analysis (HR 1.13; 95% CI: 0.99-1.28; Figure 1 and Supplemental Table 4). In the combined presence and relative abundance analysis, compared to participants with no *Bacteroides* detected, participants in the highest tertile had reduced mortality risk (HR=0.54; 95% CI=0.40-0.74; ptrend=0.0001; Figure 2 and Supplemental Table 5).

Discussion

We found some oral microbial population characteristics to be independently associated with all-cause mortality in this large, representative cohort of the 20- to 69-year-old US population. The alpha diversity measures incorporating richness and evenness (i.e., the Shannon and Simpson indices) were significantly inversely associated with mortality. Some significant associations were also observed with the beta diversity matrices which incorporate the relative abundance of the ASVs (i.e., Bray-Curtis and weighted UniFrac) which suggests that the oral microbial community itself may be related to health and disease. At the genus-level, a higher relative abundance of *Granulicatella* and *Lactobacillus* were associated with increased all-cause mortality, while compared with individuals without detectable *Bacteroides*, a high relative

abundance of *Bacteroides* was associated with decreased all-cause mortality. These three genera are common oral species, but *Granulicatella* has been associated with endocarditis and bacteremia,¹³ *Lactobacillus* has been associated with increased risk of lung cancer,⁴ and *Bacteroides* has been implicated in periodontal disease¹ and health.¹⁴

This study has important strengths as it is representative of the adult US population and prospective in nature with relatively long follow-up from NDI linkage. We included thorough adjustment for many confounders, particularly measured periodontal disease status and edentulism, which provides strong evidence for independent microbiome associations. However, there are some limitations to note. Due to the use of 16S rRNA gene sequencing, we were only able to accurately assign the bacterial taxa to the genus-level. The use of shotgun metagenomics in future studies would allow for more granular and imputed functional associations. There were also relatively few deaths over the follow-up, which precluded the study of cause-specific mortality outcomes. Finally, there is the possibility for residual confounding from unmeasured or uncontrolled factors.

In conclusion, some measures of the oral microbiome were independently associated with all-cause mortality in a nationally representative US cohort. Our findings highlight the importance of the oral microbiome and oral health to overall human health and more research could lead to a greater understanding of these associations.

	Died over follow-up N = 426		Alive at the end of follow-up N = 7,295			
	Ν	(Weighted %)	Ν	(Weighted %)	HR	(95% CI)
Age groups (years)						· · · ·
20-29	22	5.44%	1,637	22.26%	Ref	
30-39	25	6.30%	1,545	20.85%	1.13	(0.56, 2.28)
40-49	58	15.73%	1,532	22.24%	2.38	(1.10, 5.15)
50-59	124	33.42%	1,318	20.80%	5.33	(2.39, 11.86)
60-69	197	39.11%	1,263	13.85%	9.47	(4.41, 20.36)
Sex						
Male	271	65.51%	3,552	48.10%	Ref	
Female	155	34.49%	3,743	51.90%	0.48	(0.37, 0.61)
Self-reported race and ethnicity						
Mexican American	55	6.50%	1,217	9.20%	0.62	(0.42, 0.91)
Other Hispanic	35	2.83%	768	6.00%	0.47	(0.27, 0.82)
Non-Hispanic Black	125	15.16%	1,681	11.72%	0.90	(0.66, 1.23)
Non-Hispanic White	186	71.36%	2,738	64.81%	Ref	
Non-Hispanic other race including multiracial	25	4.15%	891	8.28%	0.45	(0.26, 0.78)
Education						
Less than high school	161	28.46%	1,703	15.76%	Ref	
Completed high school/GED	99	24.61%	1,581	20.57%	0.89	(0.56, 1.39)
More than high school ^A	166	46.93%	4,011	63.66%	0.80	(0.58, 1.11)
Marital status						
Never married	68	16.95%	1,693	21.06%	Ref	
Married/living with partner ^A	224	59.74%	4,328	63.57%	0.65	(0.39, 1.09)

Table 1. Selected characteristics of participants and associations with overall mortality in the linked oral microbiome and mortality NHANES files 2009-2012

Divorced/widowed/separated	134	23.31%	1,274	15.37%	0.63	(0.35, 1.17)
Income-to-poverty ratio ^A						
High poverty (<1)	128	22.23%	1,660	15.56%	1.78	(1.32, 2.40)
Medium poverty (1-1.999)	135	23.10%	1,723	17.99%	1.47	(1.05, 2.05)
Low poverty (2-2.999)	45	13.74%	853	12.86%	1.39	(0.84, 2.30)
Very low poverty (≥3)	88	34.14%	2,459	47.03%	Ref	
Missing	30	6.80%	600	6.56%	1.19	(0.68, 2.09)
Body mass index (kg/m ²)						
<18.5 (Underweight)	12	3.86%	112	1.59%	3.49	(1.31, 9.33)
18.5-24.999 (Normal weight) ^B	109	21.16%	2,091	29.99%	Ref	
25-29.999 (Overweight)	126	31.15%	2,368	32.98%	1.18	(0.77, 1.80)
30-34.999 (Class 1 obesity)	83	19.41%	1,510	20.46%	1.09	(0.75, 1.60)
≥35 (Class 2 & 3 obesity)	96	24.41%	1,214	14.97%	2.01	(1.35, 2.99)
Cigarette smoking history						
Never smoker	152	34.74%	4,221	57.36%	Ref	
Current ^C smokers ≤10 cigs/day	81	17.21%	1,065	12.67%	1.85	(1.20, 2.86)
Current ^C smokers 11-20 cigs/day	69	16.79%	497	7.08%	1.89	(1.17, 3.05)
Current ^C smokers >20 cigs/day	22	5.42%	146	2.37%	1.53	(0.74, 3.15)
Current ^C smokers, intensity unknown	10	3.31%	156	2.34%	1.67	(0.83, 3.33)
Former smokers, quit 1-10 years ago, smoked ≤10						
cigs/day ^D	19	3.59%	293	4.10%	1.28	(0.54, 3.05)
Former smokers, quit 1-10 years ago, smoked 11-		5 1 2 2	105	1 500/		
20 cigs/day	14	5.13%	105	1.59%	3.44	(1.71, 6.91)
Former smokers, quit 1-10 years ago, smoked >20	12	3 5/1%	68	0.96%	1 20	(0.46, 3.12)
Former smokers quit >10 years ago smoked <10	12	5.5470	00	0.7070	1.20	(0.40, 5.12)
cigs/day ^E	34	7.49%	380	5.28%	1.35	(0.80, 2.29)
Former smokers, quit ≥ 10 years ago, smoked >10						· · · ·
cigs/day	13	2.79%	364	6.25%	0.28	(0.13, 0.62)
Alcohol consumption						

Never drinker	34	7.18%	807	8.50%	Ref	
Drinks 0 drinks/week in past 12 months	143	29.30%	969	11.95%	1.49	(0.92, 2.41)
Drinks >0-<1 drink/week in past 12 months	83	22.15%	2,145	28.75%	0.84	(0.47, 1.50)
Drinks 1-<8 drinks/week in past 12 months	63	15.47%	1,752	28.08%	0.64	(0.32, 1.26)
Drinks 8-<14 drinks/week in past 12 months	24	6.85%	484	8.02%	0.82	(0.39, 1.71)
Drinks ≥14 drinks/week in past 12 months	54	13.46%	485	7.19%	1.26	(0.75, 2.11)
Missing	25	5.58%	653	7.51%	1.00	(0.47, 2.12)
Periodontal disease/edentulism						
No periodontal disease	72	19.61%	2,473	40.53%	Ref	
Mild periodontal disease	20	5.77%	343	4.43%	1.92	(0.86, 4.32)
Moderate periodontal disease	121	29.20%	1,597	18.81%	1.45	(0.94, 2.26)
Severe periodontal disease	68	14.30%	608	5.66%	1.98	(1.08, 3.61)
Not eligible for periodontal assessment (<30 years						
old)	22	5.44%	1,637	22.26%	NA	
Edentulous ^F	69	14.17%	202	2.15%	2.81	(1.61, 4.90)
Missing	54	11.51%	435	6.15%	1.62	(0.95, 2.76)

^A The income-to-poverty ratio is based on self-reported family income and the Department of Health and Human Services poverty guidelines ^B Includes missing (missing <5%)

^C Includes individuals who quit <1 year ago

^D Includes intensity unknown

^E Includes intensity unknown or missing quit years

^F Periodontal disease severity is calculated only for dentate adults

The weighted percentage estimates may be imprecise for small cell sizes. Hazard ratio (HR) and 95% confidence interval (CI) calculated from a Cox proportional hazards model using linked mortality data including all listed variables. The HR for the individuals who were not eligible for the periodontal disease examination is not presented because it is the same as the 20-29 year age group variable.

Figure legends

Figure 1. Forest plot of the estimated hazard ratios (HR) with 95% confidence intervals and p-values for the associations of alpha diversity (i.e., observed amplicon sequence variants [ASV], Faith's Phylogenetic Diversity [PD], Shannon Weiner, and Simpson indices), beta diversity (i.e., Bray-Curtis, unweighted UniFrac, and weighted UniFrac) principal coordinate analysis (PCoA) vectors, and specific genera with all-cause mortality in the linked oral microbiome and mortality NHANES files 2009-2012. The Cox proportional hazards models included adjustment for age, age splines, sex, self-reported race and ethnicity as categorized by NHANES, education, marital status, income-to-poverty ratio, body mass index category, detailed smoking history, alcohol consumption, periodontal disease/edentulism, prevalent diabetes or hypertension, and use of prescription medications for antibiotics, antilipidemic drugs, inhaled respiratory drugs, and drugs for treatment of gastroesophageal reflux disease. Alpha diversity, beta diversity, relative abundance, and centered log-ratio (CLR) transformed abundance associations were modeled as the per standard deviation change in the estimate. For the presence analysis, no genera were significantly associated with all-cause mortality at a Bonferroni adjusted p-value (p-value < 0.05/133 = 0.0004), but genera with a pvalue less than 0.01 are presented. All of the calculated genus-level presence associations are included in Supplemental Table 2. For the relative abundance analysis, two genera were statistically significant at the Bonferroni adjusted p-value (p-value < 0.05/103 = 0.0005) and genera with a p-value less than 0.01 are presented. All of the calculated genus-level relative abundance associations are included in Supplemental Table 3. For the CLR-transformed abundance analysis, no genera were significantly associated with all-cause mortality at a Bonferroni adjusted p-value (p-value < 0.05/103 = 0.0005), but genera with a p-value less than

0.01 are presented. All of the calculated genus-level CLR-transformed abundance associations are included in Supplemental Table 4.

Figure 2. Forest plot of the estimated hazard ratios (HR) with 95% confidence intervals and p-values for the associations of the combined genus-level presence and relative abundance with all-cause mortality in the oral microbiome and linked mortality NHANES files 2009-2012. The Cox proportional hazards models included adjustment for age, age splines, sex, self-reported race and ethnicity as categorized by NHANES, education, marital status, income-to-poverty ratio, body mass index category, detailed smoking history, alcohol consumption, periodontal disease/edentulism, prevalent diabetes or hypertension, and use of antibiotics, anti-lipidemic drugs, inhaled respiratory drugs, and prescription drugs for treatment of gastroesophageal reflux disease. The reference value is the absence of the specific genus compared to the calculated tertiles of relative abundance for that genus. For the p-value of the trend, only *Bacteroides* with a p-trend of 0.0001 was statistically significant at the Bonferroni adjusted p-value (p-value < 0.05/83 = 0.0006), but genera with a p-value less than 0.01 for the association with any tertile are presented. All of the calculated combined genus-level associations are included in Supplemental Table 5.

References

1. Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL, Jr. Microbial complexes in subgingival plaque. *J Clin Periodontol*. Feb 1998;25(2):134-44. doi:10.1111/j.1600-051x.1998.tb02419.x

2. Costalonga M, Herzberg MC. The oral microbiome and the immunobiology of periodontal disease and caries. *Immunology letters*. 2014;162(2):22-38.

3. Maki KA, Ganesan SM, Meeks B, et al. The role of the oral microbiome in smokingrelated cardiovascular risk: a review of the literature exploring mechanisms and pathways. *Journal of translational medicine*. 2022;20(1):1-26.

4. Vogtmann E, Hua X, Yu G, et al. The Oral Microbiome and Lung Cancer Risk: An Analysis of 3 Prospective Cohort Studies. *J Natl Cancer Inst*. Nov 14 2022;114(11):1501-1510. doi:10.1093/jnci/djac149

5. Nieminen MT, Salaspuro M. Local Acetaldehyde-An Essential Role in Alcohol-Related Upper Gastrointestinal Tract Carcinogenesis. *Cancers (Basel)*. Jan 5 2018;10(1)doi:10.3390/cancers10010011

6. Garcia RI, Krall EA, Vokonas PS. Periodontal disease and mortality from all causes in the VA Dental Longitudinal Study. *Annals of Periodontology*. 1998;3(1):339-349.

7. Liu J, Zong X, Vogtmann E, et al. Tooth count, untreated caries and mortality in US adults: a population-based cohort study. *Int J Epidemiol*. Aug 10 2022;51(4):1291-1303. doi:10.1093/ije/dyac072

8. Vogtmann E, Chaturvedi AK, Blaser MJ, et al. Representative oral microbiome data for the US population: the National Health and Nutrition Examination Survey. *The Lancet Microbe*. 2023;4(2):e60-e61.

9. National Center for Health Statistics. About the National Health and Nutrition Examination Survey. Updated 9/15/2017. Accessed 10/5, 2022. https://www.cdc.gov/nchs/nhanes/about_nhanes.htm

10. National Center for Health Statistics. NCHS Data Linked to NDI Mortality Files. Accessed 7/15, 2024. <u>https://www.cdc.gov/nchs/data-linkage/mortality.htm</u>

11. Gillison ML, Broutian T, Pickard RK, et al. Prevalence of oral HPV infection in the United States, 2009-2010. *Jama*. Feb 15 2012;307(7):693-703. doi:10.1001/jama.2012.101

12. National Center for Health Statistics. National Health and Nutrition Examination Survey 2009-2010 and 2011-2012 Oral Microbiome Data Documentation. Updated 10/2022. Accessed 11/2, 2023. <u>https://wwwn.cdc.gov/nchs/data/nhanes/omp/OralMicrobiomeDataDocumentation-508.pdf</u>

13. Cargill JS, Scott KS, Gascoyne-Binzi D, Sandoe JA. Granulicatella infection: diagnosis and management. *Journal of medical microbiology*. 2012;61(6):755-761.

14. Leys EJ, Lyons SR, Moeschberger ML, Rumpf RW, Griffen AL. Association of Bacteroides forsythus and a novel Bacteroides phylotype with periodontitis. *Journal of clinical microbiology*. 2002;40(3):821-825.

				_
Alpha diversity	Observed ASVs			0.1056
	Faith's PD			0.2928
	Shannon Weiner	_		0.0288
	Simpson			0.0824
Beta diversity				
Bray–Curtis	PCoA 1			0.9246
	PCoA 2	—		0.0022
	PCoA 3			0.8785
	PCoA 4			0.7022
Unweighted UniFrac	PCoA 1			0.4952
	PCoA 2			0.1246
	PCoA 3			0.8841
	PCoA 4			0.8528
Weighted UniFrac	PCoA 1	•		0.0263
	PCoA 2			0.5616
	PCoA 3			0.1781
	PCoA 4			0.6480
Presence	Bacteroides			0.0056
	Anaerococcus		•	— 0.0025
	Alysiella			0.0061
Relative abundance	Unknown genus of Sporichthyaceae	●		0.0045
	Granulicatella	_		< 0.0001
	Lactobacillus	_		< 0.0001
	Faecalibacterium			0.0044
	Anaeroglobus	_ _		0.0097
CLR-transformed abundance	Bacteroides	· ·		0.0019
	Haemophilus	_ _		0.0037
		0.5 1.0 HR	2.0	

Bacteroides	Absent (Ref)		0.0004	
	T1		$p_{\rm trend} = 0.0001$	0.8530
	Τ2			0.2099
	Т3	· · · · · · · · · · · · · · · · · · ·		0.0003
Unknown genus of WCHB1-69	Absent (Ref)			
	T1		$p_{\text{trend}} = 0.3243$	0.0015
	Т2			0.4009
	ТЗ	•		0.2068
Unknown genus of Lachnospiraceae	Absent (Ref)			
	T1		$p_{\text{trend}} = 0.1076$	0.5164
	Τ2			0.0045
	ТЗ			0.2506
Megasphaera	Absent (Ref)			
	T1		p _{trend} = 0.4271	0.0249
	Τ2			0.0081
	ТЗ			0.1599
Unknown genus of Fusobacteriales	Absent (Ref)			
	T1		$p_{\rm trend} = 0.2791$	0.1359
	Т2			- 0.0055
	Т3	•		0.8550
Kingella	Absent (Ref)			-)
	T1		p _{trend} = 0.1451	0.7338
	T2			0.0032
	ТЗ			0.4318
				-
		0.3 1.0	3.0	

HR