1 Integration of serum androgens and Sex Hormone-Binding Globulin for optimized early detection of aggressive prostate cancer 2 Olga Lazareva^{1,2,3,4}, Anja Riediger^{1,5,6,7}, Oliver Stegle^{2,3,4}, Holger Sültmann^{6,8}, 3 Markus Hohenfellner⁵, Magdalena Görtz^{1,5,6} 4 5 ¹Junior clinical cooperation unit 'Multiparametric Methods for Early Detection of 6 Prostate Cancer', German Cancer Research Center (DKFZ), Heidelberg, Germany 7 ²Division of Computational Genomics and Systems Genetics, German Cancer 8 Research Center (DKFZ), Heidelberg, Germany 9 10 ³European Molecular Biology Laboratory, Genome Biology Unit, Heidelberg, 11 Germany 12 ⁴Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, UK ⁵Department of Urology, University Hospital Heidelberg, Heidelberg, Germany 13 ⁶National Center for Tumor Diseases (NCT), Heidelberg, Germany 14 ⁷Faculty of Biosciences, Heidelberg University, Heidelberg, Germany 15 ⁸Division of Cancer Genome Research, German Cancer Consortium (DKTK), 16 German Cancer Research Center (DKFZ), Heidelberg, Germany 17 18 19 20 Send correspondence to: Magdalena Görtz 21 magdalena.goertz@med.uni-heidelberg.de 22 23 24 25 26

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

28 Blood markers associated with prostate cancer severity

29

- 30 Keywords:
- 31 Androstenedione, Dehydroepiandrosterone-Sulfate, Early Detection of Cancer,
- 32 Prostatic Neoplasms, Prostate-Specific Antigen, Risk Assessment, Sex Hormone-
- 33 Binding Globulin
- 34
- 35
- 36 Word count abstract: 298 / 300
- 37 Word count text: 3000 / 3000

38

40 **ABSTRACT**

41

42 Background

Aggressive prostate cancer (PC) represents a significant health concern worldwide. Conventional initial screening methods, primarily based on prostate-specific antigen (PSA) levels, lack specificity, leading to a high rate of unnecessary biopsies and an urgent need for more accurate diagnostic tools. This study addresses the gap by exploring the potential of integrating clinical and routine blood laboratory parameters including a comprehensive hormone assessment to enhance the non-invasive prediction of aggressive PC.

50 Methods

In a pilot study of 578 patients who were scheduled for a prostate biopsy due to suspicion of PC, we analyzed an extensive panel of 28 laboratory values alongside data on family history, diet, and lifestyle. A logistic regression classifier was developed, and model performance was evaluated using repeated k-fold crossvalidation on the complete dataset (n=282). Participants were histologically categorized into three risk groups: healthy, moderate PC (ISUP 1-2 PC), and aggressive PC (ISUP 3-5 PC).

58 **Results**

59 Significant associations were found between PC aggressiveness and lower levels of 60 androstenedione, Dehydroepiandrosterone-Sulfate (DHEA-S) and free PSA 61 percentage, as well as higher levels of Sex Hormone Binding Globulin (SHBG). The 62 integration of these serum markers with clinical parameters into a new multi-stage 63 risk classifier for PC prediction significantly improved the predictive accuracy. The 64 risk model outperformed PSA-only methods, demonstrating higher sensitivity and 65 specificity in predicting aggressive PC.

66 Conclusions

67	Incorporating serum markers DHEA-S, androstenedione, and SHBG into a novel risk
68	classifier can improve early detection of aggressive PC. These widely available and
69	cost-effective blood biomarkers could reduce reliance on invasive prostate biopsies
70	and expensive magnetic resonance imaging by providing a more targeted approach
71	to non-invasive prediction of aggressive PC following PSA testing. Our pilot study
72	lays the groundwork for larger-scale research to further explore the integration of
73	androgens and SHBG in future risk stratification models for improved clinical decision
74	making.
75	
76	
77	
78	
79	
80	
81	
82	
83	
84	

85 **INTRODUCTION**

86 Prostate cancer (PC) is the second most prevalent cancer in men, representing 27% 87 of all new cancer diagnoses in males (1). It is a major cause of mortality worldwide, 88 with projections of the Lancet Commission on PC indicating a doubling of new cases 89 between 2020 and 2040, particularly in low-income countries (2). Early diagnosis 90 through screening can offer numerous benefits such as a higher likelihood of cure, 91 less aggressive treatment options, reduced progression to metastatic stages, and 92 improved quality of life (3–5). Prostate-specific antigen (PSA) screening is a widely 93 used method for early detection of PC. Recent results from the Rotterdam Section of 94 the European Randomized Study of Screening for PC indicate that PSA screening 95 can substantially reduce long-term mortality rates (6). However, PSA testing has 96 limited specificity, frequently resulting in unnecessary biopsies and the detection of 97 indolent PC (7).

To address the limitations of PSA testing, the American Urological Association (AUA) 98 99 guidelines recommend integrating risk calculators into the traditional decision-making 100 process for PC screening (8). The primary tool is the PSA blood test, and for cases 101 with elevated PSA levels, the AUA guidelines advise using risk calculators that 102 include PSA, free PSA %, age, ethnicity, and family history for personalized risk 103 assessment. Clinicians are then encouraged to engage in shared decision-making 104 with patients, discussing the value of additional diagnostic data from biomarkers or 105 multiparametric magnetic resonance imaging (mpMRI) and considering the 106 advisability of proceeding with a biopsy. Similarly, the European Association of 107 Urology (EAU) recommends using risk stratification nomograms for patients with 108 elevated PSA, including age, family history, digital rectal examination (DRE), prostate 109 volume, and PSA density (PSAD) (9). This method helps to identify men at low risk

who may only require clinical follow-up, thereby reducing the need for expensive orinvasive tests such as mpMRI and biopsy.

mpMRI and fusion-targeted biopsy of suspicious mpMRI lesions are advanced diagnostic methods for PC, significantly enhancing detection accuracy (10,11). High cost and restricted availability of mpMRI, especially in low-resource settings, limit its routine use as a screening tool (12). The economic and logistical challenges of MRI highlight the need for cost-effective and accessible biomarkers to facilitate broader implementation across diverse healthcare settings, improving early detection capabilities.

119 Recent research increasingly focuses on modifiable risk factors, such as lifestyle and 120 diet, associated with PC risk (13,14). Metabolic changes linked to many diseases, 121 including cancer, are reflected in human blood metabolomic patterns, which provide 122 valuable insights into underlying physiological states (15). Systemic information from 123 blood parameters, such as C-reactive protein, amino acids, and glycated hemoglobin, 124 has been predictive of multiple cancers (16). Additionally, adrenal androgens 125 influence the aetiopathogenesis of PC and are associated with aggressive PC 126 (17),(18).

127 The study aimed to improve non-invasive prediction of aggressive PC by integrating 128 44 laboratory and clinical parameters into a predictive model. Our findings 129 demonstrate that incorporating these parameters after PSA testing significantly 130 improved diagnostic accuracy. This approach could reduce unnecessary biopsies, 131 decrease the number of costly MRIs, and more accurately predict PC 132 aggressiveness, aligning with precision medicine principles. By validating a 133 multifaceted diagnostic strategy over traditional PSA-only methods, our research 134 supports a more nuanced approach to PC screening and decision-making.

135 MATERIALS AND METHODS

136 Study population

137 565 men who received a prostate biopsy at the University Hospital Heidelberg 138 between June 2021 and December 2023 were recruited for supplementary clinical 139 and, if a preoperative blood sample had to be taken and patient consent was given, 140 laboratory values assessment. They were eligible for study inclusion in case of 141 suspicion of PC with a PSA level \geq 3 ng/ml and/or suspicious DRE and no previous 142 treatment for PC. Additionally, 13 healthy men with a PSA <2.5 ng/ml who presented 143 to the Urology Clinic at Heidelberg University Hospital for benign conditions, were 144 recruited to enlarge the healthy control group. Data was collected prospectively, and 145 institutional review board approval was obtained (S-130/2021). Clinical information 146 regarding family history (defined as having a first-degree relative with PC), lifestyle, 147 etc. was assessed via questionnaire. Laboratory parameters included, among others, 148 lipid metabolism, inflammatory markers, hormones, and vitamins (Supplementary Table 1). After obtaining informed consent, blood samples were collected via vein 149 150 puncture between 8 and 11 AM in fasting patients. All blood samples were analyzed 151 in the central laboratory of Heidelberg University Hospital, in addition to the standard preoperative lab values. The central laboratory of Heidelberg University Hospital has 152 153 been accredited according to DIN EN ISO 15189 since 2005.

154 **Prostate biopsy**

Prebiopsy mpMRI was performed on 543 patients with suspicion of PC and eligibility for mpMRI. All mpMRI scans followed the PI-RADS recommendations, according to European Society of Urogenital Radiology guidelines (19). Men underwent transperineal fusion-targeted biopsy using a UroNav system (Philips Invivo, Gainsville, FL, USA) of MRI-suspicious lesions and saturation biopsy adjusted to prostate volume, as previously described (20,21). Histopathological analyses were performed according to International Society of Urological Pathology (ISUP) standards. The participants were categorized into three risk groups: healthy patients (PSA <2.5 ng/ml or no PC in biopsy), moderate PC (ISUP 1-2 PC), and aggressive PC (ISUP 3-5 PC)(10).

165 Statistical analysis

In this study, 28 laboratory and 16 clinical parameters were analyzed to determine their impact on PC risk. Each parameter was initially tested for its ability to discriminate between the three predefined risk groups: healthy patients (n=253), moderate PC (n=229), and aggressive PC (n=96) through statistical analysis. For continuous variables, an ordinary least squares (OLS) methodology was employed, fitting models for each parameter to predict the risk groups and subsequently testing the significance of the slope coefficients $\beta_1^{\ k}$:

173
$$y = \beta_0^k + \beta_1^k x^k$$
,

where $y \in \{0,1,2\}$ is a risk category corresponding to healthy patients, ISUP 1-2 and ISUP 3-5 PC, $\beta_0^{\ k}$ and $\beta_1^{\ k}$ are intercept and slope coefficients for the *k*-th tested parameter x^k .

For categorical variables the Chi-square test was used (22). All p-values were reported and subsequently adjusted for multiple comparisons using the Benjamini & Hochberg correction (23) with a family-wise error rate at 0.1 level. Complete results are provided in **Supplementary Table 1**, with significant findings highlighted in **Figure 1**.

182 Risk Modeling

183 Risk modeling was performed on 282 patients who had available data for clinical and 184 laboratory parameters, showing a significant association with ISUP grade. The 185 logistic regression model was developed in a staged approach:

- 1. Baseline model: Initiated with PSA, the standard marker for PC screening,
 to establish a benchmark for model performance.
- 2. Serum marker integration: Added laboratory measurements that showed a statistically significant association with PC risk, including Androstenedione, free PSA %, Dehydroepiandrosterone-sulfate (DHEA-Sulfate), and Sex Hormone Binding Globulin (SHBG) to evaluate improvements in sensitivity and specificity.
- Olinical parameter integration: Included clinical factors from international
 guidelines (age, family history, DRE results, prostate volume, and PSAD) to
 assess their combined predictive value.
- 4. MRI results: Added PIRADS scores to capture additional risk information
 provided by imaging.
- 5. Advanced baseline model: Combined MRI and PSA.

Model performance was assessed at each stage by calculating the F1 score, Precision, Recall, and Accuracy. The model was evaluated using a repeated k-fold cross-validation (k = 10, repeated 10 times) approach, and all performance metrics were reported based on the test set (n=282). We used 'L2' regularization to decrease overfitting probability with the default regularization strength ($\lambda = 1$) using scikit-learn models (24).

205 Interpretative Analysis Using SHAP Values

To understand the influence of individual variables on predictive outcomes, SHAP (SHapley Additive exPlanations) methodology was applied, following the approach proposed by Lundberg and Lee (25). This method quantitatively measured the impact of each parameter included in the model, providing clear insights into how specific factors contribute to overall risk prediction.

211 **RESULTS**

212 Patient cohort description

The study included a total of 578 patients. During the initial phase of the analysis, when statistically significant markers were selected, all available measurements were used without restricting to a fully overlapping cohort, maximizing the use of the data. Each laboratory parameter was represented by a minimum of 314 and a maximum of 353 measurements, while other parameters ranged from 314 to 578 measurements. A comprehensive list of these parameters and their respective counts is provided in **Supplementary Table 1**.

Following feature selection, the cohort was refined to include only patients with complete data across all parameters identified as significant (post-multiple hypothesis correction). This subset comprised 282 patients. Detailed descriptions of these patients and their corresponding data are presented in **Table 1**.

224 Role of blood biomarkers in prostate cancer

225 Significant associations were found between hormonal markers and PC severity as 226 classified by ISUP grades (**Figure 1**). Lower levels of androstenedione and DHEA-227 Sulfate were significantly associated with more aggressive forms of PC (ISUP grades

3-5) (corrected p-values 0.013 and 0.03, respectively). This finding supports thehypothesis that reductions in certain androgens may signal advanced disease.

The bioavailability of circulating androgens is largely determined by the concentration
of SHBG. Accordingly, in our study higher levels of SHBG were associated with
increased PC severity (corrected p-value 0.037).
Higher PSA levels were positively correlated with PC severity (corrected p-value
0.03). Additionally, a lower percentage of free PSA % indicated more aggressive
cancer (corrected p-value 0.025). These results reinforce the role of PSA and free
PSA percentage as critical biomarkers in assessing PC aggressiveness.

Incorporating clinical and lab values into new risk models for non-invasive prostate cancer prediction

In developing a PC risk classifier, we included the five significant laboratory values, along with clinical features and imaging parameters, that were strongly associated with ISUP grades. Age, DRE results, PSAD, PIRADS scores, and prostate volume showed statistically significant associations with ISUP grade severity, all with corrected p-values <0.05. Although a family history of PC did not show a significant statistical association with ISUP grades (corrected p-value 0.552), it was included in the model to adhere to AUA and EAU guidelines (8,9).

The performance of the risk classifier was evaluated through a multi-stage approach: 1. Baseline model initiated with PSA, 2. Enhancement with laboratory biomarkers significantly associated with PC risk (Androstenedione, free PSA %, DHEA-Sulfate, and SHBG), 3. Integration of clinical factors (Age, family history, DRE results, prostate volume, and PSAD), 4. Integration of MRI results (PIRADS scores) and 5. Advanced baseline model with solely PSA and MRI as comparison. This stepwise model enhancement resulted in gradual improvements across all metrics from stage

1-4, including F1 score, precision, recall, and accuracy. The progression of median
scores through each stage is depicted in Figure 2A, with exact values provided in
Supplementary Table 2.

256 The regression coefficient distribution was analyzed to assess the prognostic 257 significance of each parameter (Figure 2B). PSAD emerged as a particularly 258 influential factor in identifying healthy patients. Established diagnostic parameters like 259 PSA, DRE results and PIRADS also showed high importance for non-invasive prediction of PC presence. However, with these parameters, coefficients for ISUP 260 261 grades 1-2 and 3-5 PC often shared the same sign, indicating challenges in 262 differentiating these two categories. Serum androgens displayed variable signs for 263 ISUP 1-2 versus ISUP 3-5 PC, suggesting their potential impact on the model's 264 predictive capability.

265 Influence of individual variables on predictive outcomes using SHAP Values

266 To understand the influence of individual variables on predictive outcomes for each patient, we employed SHAP analysis to obtain per-patient SHAP values. These 267 268 values help to identify the specific factors contributing to each patient's prediction (PC 269 versus no PC) and provide insight into the influence of each variable on the 270 predictive outcomes. The model was developed to assist urologists in making 271 informed decisions regarding prostate biopsies, especially when the complexity of 272 individual patient profiles challenges clinical judgement. Values from two study 273 patients are shown where biopsies didn't reveal PC, and the model correctly 274 predicted the absence of PC, demonstrating the practical utility of SHAP analysis as 275 a transparent, data-driven foundation for clinicians (Figure 3A and 3B).

276 **DISCUSSION**

This study aimed to improve the non-invasive prediction of aggressive PC by integrating clinical and laboratory parameters into a comprehensive risk model. By identifying novel, cost-effective serum markers, we sought to improve decisionmaking following PSA testing and reduce reliance on invasive procedures and expensive MRI use.

several results. 282 Our study revealed First, levels of DHEA sulfate and 283 androstenedione were significantly reduced in newly diagnosed aggressive PC 284 compared to healthy controls. In line with these results, higher levels of SHBG were 285 significantly associated with increased PC aggressiveness. These markers had a stronger association with aggressive PC detection than established variables advised 286 287 for use in risk stratification nomograms, such as family history (8,9). Thus, androgens 288 have potential as diagnostic markers for early detection of aggressive PC, offering a 289 complementary diagnostic tool alongside PSA testing. Androgens were particularly 290 effective in distinguishing between ISUP 1-2 and ISUP 3-5 PC, highlighting their 291 potential in differentiating moderate from aggressive PC, e.g. for treatment 292 approaches such as active surveillance (26). ISUP 1-2 PC patients may be offered 293 active surveillance instead of radical therapy due to the indolent nature of their 294 disease (27). Existing biomarkers expressed by all PC, including indolent tumors, 295 offer limited potential to selectively detect high-grade disease (28). Novel biomarkers 296 distinguishing high-grade from low-grade PC are urgently needed, aligning with 297 guidelines emphasizing a focus on high-grade PC due to the indolent nature of low-298 grade PC (8).

Second, integrating clinical findings and laboratory values into the decision-making process for recommending MRI scans has the potential to improve the selection of patients who benefit most from imaging. MRI remains time- and resource-intensive

302 and may pose considerable costs for patients, apart from limitations in availability and 303 inter-rater reliability. Barriers in healthcare delivery limit the wide adoption of MRI in 304 real-world practice as a diagnostic test following PSA. The decision to use MRI and 305 fusion biopsy can be supported by relatively inexpensive routine blood values and 306 clinical parameters, minimizing unnecessary MRIs for PSA-positive patients with a 307 low likelihood of PC. Commercially available blood values are straightforward to 308 implement and less resource-intensive, allowing urologists to more selectively apply 309 MRI and/or biopsy (29.30). Our results support implementing integrated decision-310 making frameworks in clinical routine to optimize MRI and biopsy use in PC 311 diagnostics.

Third, the application of an explainable decision support concept in our study demonstrates an approach to guide urologists and patients in making informed decisions regarding prostate biopsies. A transparent and comprehensive assessment of each patient's clinical scenario, including evaluation of clinical, lab values, and, if applicable, MRI findings, allows urologists and patients to make trustworthy and clinically justified decisions.

Our findings align with previous research, such as Severi et al., who found high levels of adrenal androgens like androstenedione and DHEA Sulfate associated with a reduced risk of aggressive PC (31). Their study of 17,049 men reported that double concentrations of androstenedione were associated with almost half the risk of aggressive PC, and double concentrations of DHEA-Sulfate were associated with a 37% lower risk. Additionally, Travis et al. described an inverse association of the androstenedione concentration with advanced PC (32).

325 Our finding of higher SHBG levels in ISUP 3-5 PC complements previous results that 326 proposed SHBG as a marker of aggressive PC in early-stage disease. Preoperative 327 serum SHBG levels were an independent predictor of extraprostatic extension in localized PC, suggesting a role of SHBG in PC progression (33). A study by Price et
al. demonstrated that germline genetic variations of androgen-related pathway genes
are associated with serum androgen and SHBG concentrations and PC risk (34).

331 Routine blood test results harbor valuable information that may not be immediately 332 apparent to clinicians. Podnar et al. demonstrated the feasibility of brain tumor 333 diagnosis from routine blood tests using a machine learning predictive model with 334 diagnostic accuracy comparable to imaging studies (35). Regular screening with 335 blood tests for disease prevention is already recommended in many countries (36). 336 This study's results indicate the potential of simple clinical parameters combined with 337 comprehensive laboratory predictors to estimate aggressive PC risk. Highlighting 338 androgens as complementary diagnostic tools at minimal cost and emphasizing an 339 integrated approach to MRI utilization could lead to more efficient screening 340 strategies for aggressive PC.

341 Several promising additional biomarkers could complement the current diagnostic 342 pathway for PC, including PSA derivatives and genetic markers (37). Combining the 343 prostate health index and the 4K score (38), if available, with widely accessible and 344 cost-effective clinical and lab values may offer a comprehensive and accurate 345 approach for early non-invasive prediction of aggressive PC. Recognizing the low 346 specificity of PSA as a challenge for PC screening programs, Srivastava et al. 347 recently proposed a "more is more" approach in PC screening programs with the 348 addition of biomarkers in a multistep approach (30). This strategy could help to 349 optimize biopsy decisions, reduce unnecessary biopsies, and improve patient 350 outcomes.

351 Several limitations must be acknowledged. The pilot study's sample size may limit 352 the generalizability and statistical power of the findings. Validation in independent, 353 larger cohorts is necessary to ensure the classifier's robustness and generalizability.

354 The study population was from a single center, lacking diversity in ethnicity and 355 geographic factors, which may affect the applicability of results to all populations. 356 Identified serum markers, DHEA, SHBG, and androstenedione, need further 357 investigation to validate their efficacy and reliability across diverse populations and 358 their potential for prevention, detection and prognosis of early-stage, aggressive PC. 359 Additionally, the study's nature did not allow determining causal relationships among 360 variables. Longitudinal studies are necessary to establish relationships between hormone levels and PC development or progression. Lastly, the definition of PC that 361 362 is aggressive and requires active treatment rather than active surveillance is 363 controversial, with varying definitions of aggressive PC. As ISUP 1-2 PC patients can 364 be offered active surveillance due to the indolent nature of the disease, histological 365 assessment was used to stratify patients into moderate (ISUP 1-2) and aggressive 366 (ISUP 3-5) PC (10,21).

367 CONCLUSIONS

Incorporating serum markers DHEA-S, androstenedione, and SHBG into a novel risk classifier can improve early detection of aggressive PC. These widely available and cost-effective blood biomarkers could reduce reliance on invasive prostate biopsies and expensive MRI by providing a more targeted approach to non-invasive prediction of aggressive PC following PSA testing. Our pilot study lays the groundwork for larger-scale research to further explore the integration of androgens and SHBG in future risk stratification models for improved clinical decision making.

375

376

378

380 ABBREVIATIONS

381	AUA	American Urological Association		
382	DHEA	Dehydroepiandrosterone		
383	DRE	Digital rectal examination		
384	EAU	European Association of Urology		
385	ISUP	International Society of Urological Pathology		
386	mpMRI	Multiparametric magnetic resonance		
387		imaging		
388	PC	Prostate cancer		
389	PSA	Prostate-specific antigen		
390	PSAD	Prostate-specific antigen density		
391	SHBG	Sex Hormone Binding Globulin		
392				

393 **DECLARATIONS**

394 **Conflicting interests:** None.

Funding: This work was realized through support by the German Federal Ministry for
Economic Affairs and Climate Action (funding # 01MT21004A) and the Dieter
Morszeck Foundation.

398 Ethical approval: The study was approved by the ethical committee of the University
399 of Heidelberg (Approval No. S-130/2021) and was performed in accordance with the
400 Declaration of Helsinki.

401 Contributorship: Conceptualization: M.G.; Data curation: A.R., O.L.; Formal
402 analysis: O.L.; Funding acquisition: M.G., M.H.; Investigation: H.S.; M.G., O.L;
403 Methodology: H.S.; M.G., O.L.; Project Administration: M.G.; Resources: M.G., M.H.,
404 O.S.; Software: O.L., O.S.; Supervision: M.G.; Validation: A.R., H.S., O.L.;

- 405 Visualization: O.L.; Writing original draft: M.G., O.L.; Writing review & editing: A.R.,
- 406 H.S., M.H., O.S.
- 407 All authors critically reviewed, edited, and added to the manuscript as well as
- 408 approved the final version of the manuscript.
- 409 Acknowledgements:
- 410 We sincerely thank Daniela Janscho for her invaluable assistance in handling patient
- 411 samples.

412 Data availability statement:

- 413 The data sets generated during and analyzed during the current study are available
- 414 from the corresponding author on reasonable request.
- 415 **Declaration of generative AI and AI-assisted technologies in the writing** 416 **process:**
- 417 During the preparation of this work the authors used ChatGPT-40 in order to improve
- 418 language and readability. After using this tool, the authors reviewed and edited the
- 419 content as needed and take full responsibility for the content of the publication.
- 420
- 421
- 422

423 **REFERENCES**

Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA Cancer
 J Clin. 2022 Jan;72(1):7–33.

426 2. James ND, Tannock I, N'Dow J, Feng F, Gillessen S, Ali SA, et al. The Lancet

427 Commission on prostate cancer: planning for the surge in cases. The Lancet

428 [Internet]. 2024 Apr 27 [cited 2024 Jul 23];403(10437):1683–722. Available from:

429 https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(24)00651-2/fulltext

430 3. Schröder FH, Hugosson J, Roobol MJ, Tammela TLJ, Zappa M, Nelen V, et

al. Screening and prostate cancer mortality: results of the European Randomised

432 Study of Screening for Prostate Cancer (ERSPC) at 13 years of follow-up. Lancet

433 Lond Engl. 2014 Dec 6;384(9959):2027–35.

434 4. Schröder FH, Hugosson J, Carlsson S, Tammela T, Määttänen L, Auvinen A,

435 et al. Screening for prostate cancer decreases the risk of developing metastatic

436 disease: findings from the European Randomized Study of Screening for Prostate

437 Cancer (ERSPC). Eur Urol. 2012 Nov;62(5):745–52.

438 5. Van Poppel H, Albreht T, Basu P, Hogenhout R, Collen S, Roobol M. Serum

439 PSA-based early detection of prostate cancer in Europe and globally: past, present

440 and future. Nat Rev Urol. 2022 Sep;19(9):562–72.

441 6. de Vos II, Meertens A, Hogenhout R, Remmers S, Roobol MJ, ERSPC

442 Rotterdam Study Group. A Detailed Evaluation of the Effect of Prostate-specific

443 Antigen-based Screening on Morbidity and Mortality of Prostate Cancer: 21-year

444 Follow-up Results of the Rotterdam Section of the European Randomised Study of

445 Screening for Prostate Cancer. Eur Urol. 2023 Oct;84(4):426–34.

446 7. Roobol MJ, Steyerberg EW, Kranse R, Wolters T, van den Bergh RCN,

447 Bangma CH, et al. A risk-based strategy improves prostate-specific antigen-driven

detection of prostate cancer. Eur Urol. 2010 Jan;57(1):79–85.

449	8. Wei JT, Barocas D, Carlsson S, Coakley F, Eggener S, Etzioni R, et al. Early				
450	Detection of Prostate Cancer: AUA/SUO Guideline Part II: Considerations for a				
451	Prostate Biopsy. J Urol [Internet]. 2023 Jul [cited 2024 May 29];210(1):54–63.				
452	ailable from: https://www.auajournals.org/doi/10.1097/JU.0000000000003492				
453	9. Van Poppel H, Roobol MJ, Chapple CR, Catto JWF, N'Dow J, Sønksen J, et				
454	al. Prostate-specific Antigen Testing as Part of a Risk-Adapted Early Detection				
455	Strategy for Prostate Cancer: European Association of Urology Position and				
456	Recommendations for 2021. Eur Urol [Internet]. 2021 Dec 1 [cited 2024 Aug				
457	7 15];80(6):703–11. Available from:				
458	https://www.sciencedirect.com/science/article/pii/S0302283821019278				
459	10. Ahmed HU, El-Shater Bosaily A, Brown LC, Gabe R, Kaplan R, Parmar MK, et				
460	al. Diagnostic accuracy of multi-parametric MRI and TRUS biopsy in prostate cancer				
461	(PROMIS): a paired validating confirmatory study. Lancet Lond Engl. 2017 Feb				
462	25;389(10071):815–22.				
463	11. Rouvière O, Puech P, Renard-Penna R, Claudon M, Roy C, Mège-				
464	Lechevallier F, et al. Use of prostate systematic and targeted biopsy on the basis of				
465	multiparametric MRI in biopsy-naive patients (MRI-FIRST): a prospective,				
466	multicentre, paired diagnostic study. Lancet Oncol. 2019 Jan;20(1):100–9.				
467	12. van der Leest M, Israël B, Cornel EB, Zámecnik P, Schoots IG, van der Lelij H,				
468	et al. High Diagnostic Performance of Short Magnetic Resonance Imaging Protocols				
469	for Prostate Cancer Detection in Biopsy-naïve Men: The Next Step in Magnetic				
470	Resonance Imaging Accessibility. Eur Urol [Internet]. 2019 Nov 1 [cited 2024 May				
471	29];76(5):574–81. Available from:				
472	https://www.sciencedirect.com/science/article/pii/S0302283819304361				
473	13. Peisch SF, Van Blarigan EL, Chan JM, Stampfer MJ, Kenfield SA. Prostate				

474 cancer progression and mortality: a review of diet and lifestyle factors. World J Urol.

- 475 2017 Jun;35(6):867–74.
- 476 14. Plym A, Zhang Y, Stopsack KH, Delcoigne B, Wiklund F, Haiman C, et al. A
- 477 Healthy Lifestyle in Men at Increased Genetic Risk for Prostate Cancer. Eur Urol.
- 478 2023 Apr;83(4):343–51.
- 479 15. Deelen J, Kettunen J, Fischer K, van der Spek A, Trompet S, Kastenmüller G,
- 480 et al. A metabolic profile of all-cause mortality risk identified in an observational study
- 481 of 44,168 individuals. Nat Commun [Internet]. 2019 Aug 20 [cited 2024 May
- 482 29];10(1):3346. Available from: https://www.nature.com/articles/s41467-019-11311-9
- 483 16. Kok DEG, van Roermund JGH, Aben KKH, den Heijer M, Swinkels DW,
- 484 Kampman E, et al. Blood lipid levels and prostate cancer risk; a cohort study.
- 485 Prostate Cancer Prostatic Dis. 2011 Dec;14(4):340–5.
- 486 17. Watts EL, Perez Cornago A, Fensom GK, Smith Byrne K, Noor U, Andrews
- 487 CD, et al. Circulating free testosterone and risk of aggressive prostate cancer:
- 488 Prospective and Mendelian randomisation analyses in international consortia. Int J
- 489 Cancer [Internet]. 2022 Oct 1 [cited 2024 May 29];151(7):1033–46. Available from:
- 490 https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7613289/
- 491 18. Dobbs RW, Malhotra NR, Greenwald DT, Wang AY, Prins GS, Abern MR.
- 492 Estrogens and prostate cancer. Prostate Cancer Prostatic Dis. 2019 May;22(2):185–
- 493 94.
- 494 19. PI-RADS Prostate Imaging Reporting and Data System: 2015, Version 2 -
- 495 PubMed [Internet]. [cited 2024 May 29]. Available from:
- 496 https://pubmed.ncbi.nlm.nih.gov/26427566/
- 497 20. Görtz M, Radtke JP, Hatiboglu G, Schütz V, Tosev G, Güttlein M, et al. The
- Value of Prostate-specific Antigen Density for Prostate Imaging-Reporting and Data
- 499 System 3 Lesions on Multiparametric Magnetic Resonance Imaging: A Strategy to

- 500 Avoid Unnecessary Prostate Biopsies. Eur Urol Focus [Internet]. 2021 Mar 1 [cited
- 501 2024 May 29];7(2):325-31. Available from: https://www.eu-
- 502 focus.europeanurology.com/article/S2405-4569(19)30353-0/fulltext
- 503 21. Radtke JP, Schwab C, Wolf MB, Freitag MT, Alt CD, Kesch C, et al.
- 504 Multiparametric Magnetic Resonance Imaging (MRI) and MRI-Transrectal Ultrasound
- 505 Fusion Biopsy for Index Tumor Detection: Correlation with Radical Prostatectomy
- 506 Specimen. Eur Urol. 2016 Nov;70(5):846–53.
- 507 22. Pearson K. X. On the criterion that a given system of deviations from the
- probable in the case of a correlated system of variables is such that it can be
- reasonably supposed to have arisen from random sampling. Lond Edinb Dublin
- 510 Philos Mag J Sci [Internet]. 1900 Jul 1 [cited 2024 Jun 5];50(302):157–75. Available
- 511 from: https://doi.org/10.1080/14786440009463897
- 512 23. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical
- and Powerful Approach to Multiple Testing. J R Stat Soc Ser B Methodol [Internet].
- 514 1995 [cited 2024 Jun 7];57(1):289–300. Available from:
- 515 https://onlinelibrary.wiley.com/doi/abs/10.1111/j.2517-6161.1995.tb02031.x
- 516 24. Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al.
- 517 Scikit-learn: Machine Learning in Python. J Mach Learn Res [Internet]. 2011 [cited
- 518 2024 Sep 2];12(85):2825–30. Available from:
- 519 http://jmlr.org/papers/v12/pedregosa11a.html
- 520 25. Lundberg SM, Lee SI. A Unified Approach to Interpreting Model Predictions.
- 521 In: Advances in Neural Information Processing Systems [Internet]. Curran
- 522 Associates, Inc.; 2017 [cited 2024 May 31]. Available from:
- 523 https://papers.nips.cc/paper_files/paper/2017/hash/8a20a8621978632d76c43dfd28b
- 524 67767-Abstract.html
- 525 26. Ploussard G, Isbarn H, Briganti A, Sooriakumaran P, Surcel CI, Salomon L, et

- 526 al. Can we expand active surveillance criteria to include biopsy Gleason 3+4 prostate
- 527 cancer? A multi-institutional study of 2,323 patients. Urol Oncol. 2015
- 528 Feb;33(2):71.e1-9.
- 529 27. Albuquerque-Castro Â, Macedo-Silva C, Oliveira-Sousa R, Constâncio V,
- 530 Lobo J, Carneiro I, et al. Redefining prostate cancer risk stratification: a pioneering
- 531 strategy to estimate outcome based on Ki67 immunoscoring. Biomark Res [Internet].
- 532 2024 [cited 2024 Aug 15];12(1). Available from:
- 533 https://link.springer.com/epdf/10.1186/s40364-024-00627-4
- 534 28. Tosoian JJ, Zhang Y, Xiao L, Xie C, Samora NL, Niknafs YS, et al.
- 535 Development and Validation of an 18-Gene Urine Test for High-Grade Prostate
- 536 Cancer. JAMA Oncol. 2024 Jun 1;10(6):726–36.
- 537 29. Jiao B, Gulati R, Hendrix N, Gore JL, Rais-Bahrami S, Morgan TM, et al.
- 538 Economic Evaluation of Urine-Based or Magnetic Resonance Imaging Reflex Tests
- in Men With Intermediate Prostate-Specific Antigen Levels in the United States.
- 540 Value Health [Internet]. 2021 Aug 1 [cited 2024 Jul 23];24(8):1111–7. Available from:
- 541 https://www.sciencedirect.com/science/article/pii/S1098301521001649
- 542 30. Srivastava A, Lewicki P, Morgan TM. Biomarkers in Prostate Cancer
- 543 Screening: Sometimes "More is More." Eur Urol [Internet]. 2024 Jun 18 [cited 2024
- 544 Jul 23]; Available from:
- 545 https://www.sciencedirect.com/science/article/pii/S030228382402428X
- 546 31. Severi G, Morris HA, MacInnis RJ, English DR, Tilley W, Hopper JL, et al.
- 547 Circulating steroid hormones and the risk of prostate cancer. Cancer Epidemiol
- 548Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol. 2006
- 549 Jan;15(1):86–91.
- 550 32. Travis RC, Key TJ, Allen NE, Appleby PN, Roddam AW, Rinaldi S, et al.
- 551 Serum androgens and prostate cancer among 643 cases and 643 controls in the

- 552 European Prospective Investigation into Cancer and Nutrition. Int J Cancer [Internet].
- 553 2007 [cited 2024 Jul 23];121(6):1331–8. Available from:
- 554 https://onlinelibrary.wiley.com/doi/abs/10.1002/ijc.22814
- 555 33. Lee SE, Chung JS, Han BK, Park CS, Moon KH, Byun SS, et al. Preoperative
- 556 Serum Sex Hormone-Binding Globulin as a Predictive Marker for Extraprostatic
- 557 Extension of Tumor in Patients with Clinically Localized Prostate Cancer. Eur Urol
- 558 [Internet]. 2008 Dec 1 [cited 2024 Aug 15];54(6):1324–32. Available from:
- 559 https://www.sciencedirect.com/science/article/pii/S0302283808002649
- 560 34. Price DK, Chau CH, Till C, Goodman PJ, Leach RJ, Johnson-Pais TL, et al.
- 561 Association of Androgen Metabolism Gene Polymorphisms with Prostate Cancer
- 562 Risk and Androgen Concentrations: Results from the Prostate Cancer Prevention
- 563 Trial. Cancer [Internet]. 2016 Aug 1 [cited 2024 Jul 23];122(15):2332–40. Available
- from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4956504/
- 565 35. Podnar S, Kukar M, Gunčar G, Notar M, Gošnjak N, Notar M. Diagnosing
- 566 brain tumours by routine blood tests using machine learning. Sci Rep [Internet]. 2019
- 567 Oct 9 [cited 2024 Jun 7];9(1):14481. Available from:
- 568 https://www.nature.com/articles/s41598-019-51147-3
- 569 36. nhs.uk [Internet]. 2023 [cited 2024 Jun 7]. NHS Health Check. Available from:
- 570 https://www.nhs.uk/conditions/nhs-health-check/
- 571 37. Eickelschulte S, Riediger AL, Angeles AK, Janke F, Duensing S, Sültmann H,
- 572 et al. Biomarkers for the Detection and Risk Stratification of Aggressive Prostate
- 573 Cancer. Cancers [Internet]. 2022 Dec 11 [cited 2024 Jun 5];14(24):6094. Available
- from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC9777028/
- 575 38. Duffy MJ. Biomarkers for prostate cancer: prostate-specific antigen and
- 576 beyond. Clin Chem Lab Med. 2020 Feb 25;58(3):326–39.1

578 Table 1 – Patient characteristics (modeling cohort n = 282) in the healthy, ISUP

579 1-2 PC vs. ISUP 3-5 PC groups for significant parameters. For continuous

580 parameters, median and IQR are provided, otherwise, count per group and

- 581 percentage are shown.
- 582

Parameter	Unit	Healthy patients (n=122)	ISUP 1-2 PC (n=117)	ISUP 3-5 PC (n=43)
Age	Years	63 (57-69)	66 (61-73)	72 (64.50-75)
PSA	ng/dl	5.75 (4.47-8.06)	5.81 (4.2-8.41)	10.11 (6.67-15.49)
Androstenedione	µg/l	70 (59-88)	66 (53-78)	58 (48-67)
Free PSA %	%	16 (0.30-25)	11 (0.24-20)	8 (0.18-17.50)
DHEA-Sulfate	µg/ml	1.23 (0.76-1.75)	1.06 (0.69-1.47)	0.76 (0.54-1.14)
Sex Hormone Binding				
Globulin	nmol/l	43 (32-53)	46 (35-58)	48 (35.5-62)
Prostate volume	ml	61.32 (45-81.97)	46.37 (34-66.45)	45 (33.94-60)
PSAD	ng/ml/ml	0.1 (0.07-0.13)	0.13 (0.08-0.19)	0.21(0.16-0.35)
Positive family history	Count, %	12 (9.83)	14 (11.97)	7 (16.28)
Positive DRE	Count, %	10 (8.20)	72 (61.54)	21 (48.84)
PIRADS ≤2	Count, %	19 (15.57)	6 (5.13)	0 (0)
PIRADS 3	Count, %	44 (36.07)	23 (19.66)	1 (2.33)
PIRADS 4	Count, %	53 (43.44)	58 (49.57)	14 (32.56)
PIRADS 5	Count, %	6 (4.92)	30 (25.64)	28 (65.12)

583

585 Figure 1: Statistically significant associations with the presence of moderate or

586 aggressive PC

- A Statistically significant association between laboratory parameters and the presence of ISUP 1-2 PC, ISUP 3-5 PC, or healthy patients. **B** Statistically significant associations between clinical and imaging parameters and the presence of ISUP 1-2
- 590 PC, ISUP 3-5 PC or healthy patients.
- 591

592 Figure 2: Risk modeling of PC

- 593 A Performance metric of a logistic classifier based on five sets of features: 1. PSA, 2.
- 594 Integration of additional laboratory biomarkers, 3. Integration of clinical factors, 4.
- 595 Integration of MRI results and 5. Solely PSA and MRI as comparison. **B** Feature
- importance for the three groups (healthy patients, ISUP 1-2 PC and ISUP 3-5 PC).
- 597

598 Figure 3: SHAP analysis to obtain per-patient SHAP values

- 599 SHAP values for two patients of the cohort who had no PC in the prostate biopsy:
- 600 blue color indicates a reduction of the probability of having PC, and orange color
- 601 indicates an increased probability of having PC.
- 602







в

Α