

# Effect of background therapy with Non-steroidal anti-inflammatory drugs (NSAIDs) and other anti-inflammatory agents on COVID-19 outcomes

Lucio Miele<sup>1\*</sup>, San Chu<sup>2,3</sup>, William Hillegass<sup>4</sup>, Claudine Jurkovitz<sup>5</sup>, William Beasley<sup>6</sup>, David Chen<sup>5</sup>, A Jerrod Anzalone<sup>7</sup>, Daniel Fort<sup>8</sup>, John Kirwan<sup>2</sup>, Brian Melancon<sup>2</sup>, Sally Hodder<sup>9</sup>, Ronald Horswell<sup>2</sup>, on behalf of the National COVID Cohort Collaborative Consortium<sup>^</sup>

<sup>1</sup> Department of Genetics, Louisiana State University Health Sciences Center, New Orleans, LA; Louisiana Center and Translational Science (LA CaTS) Center, Baton Rouge, LA.

<sup>2</sup> Pennington Biomedical Research Centre, Baton Rouge, Louisiana; LA CaTS Center, Baton Rouge, LA.

<sup>3</sup> Louisiana State University, Baton Rouge, Louisiana; LA CaTS Center, Baton Rouge, LA.

<sup>4</sup> Departments of Data Science and Medicine, Mississippi Center for Clinical and Translational Research, University of Mississippi Medical Center, Jackson, MS.

<sup>5</sup> Institute for Research in Equity and Community Health (iREACH), ChristianaCare Health Services, Inc., Wilmington, DE.

<sup>6</sup> Biomedical and Behavioral Methodology Core, University of Oklahoma, Norman, OK; Oklahoma Clinical and Translational Science Institute, Oklahoma City, OK.

<sup>7</sup> Department of Neurological Sciences, College of Medicine, University of Nebraska Medical Center, Omaha, NE; Great Plains IDeA-CTR, Omaha, NE.

<sup>8</sup> Ochsner Center for Outcomes Research, Division of Academics, Ochsner Health, New Orleans, LA; Louisiana Center and Translational Science (LA CaTS) Center, Baton Rouge, LA.

<sup>9</sup> West Virginia University School of Medicine, Morgantown, WV; West Virginia Clinical and Translational Sciences Institute, Morgantown, WV.

## Corresponding Author:

E-mail [lmiele@lsuhsc.edu](mailto:lmiele@lsuhsc.edu) (LM)

<sup>^</sup>Membership of the National COVID Cohort Collaborative Consortium is provided in the Acknowledgments

## 35 Abstract

36 **Background:** Inflammation plays a complex, incompletely understood role in the  
37 pathogenesis of acute COVID-19 and Post-Acute Sequelae of SARS-CoV-2 infection (PASC or  
38 “Long COVID”). Systemic acute inflammation resulting in cytokine storm, hypercoagulability and  
39 endothelial damage is thought to be a central mechanism for severe morbidity and mortality in  
40 acute COVID-19. Anti-inflammatory medications taken routinely for chronic conditions prior to  
41 contracting COVID-19 (“background medications”) may modulate acute COVID-19 outcomes.

42 **Methods:** Using data from the National COVID Cohort Collaborative (N3C) enclave, we  
43 estimated effects of six classes of background medications on acute COVID outcomes.  
44 Medication classes included aspirin, celecoxib, other NSAIDs, steroids, immune suppressants,  
45 and antidepressants. Acute COVID outcomes included probability of hospital admission,  
46 inpatient mortality, and mortality among diagnosed COVID patients. Each medication class was  
47 compared to benzodiazepines (excluding midazolam) which served as a comparator/control.  
48 Only adult COVID patients with pre-existing osteoarthritis and without any diagnosed  
49 autoimmune disease were included in the analyses. Random effects logistic regression models  
50 were used to adjust for covariates and data contributing organization. Medication effects also  
51 were estimated for COVID-negative cases.

52 **Results:** Non-aspirin NSAIDs were associated with lower mortality among diagnosed  
53 COVID-19 patients: adjusted Odds Ratio (aOR)=0.32 (p=.032) for celecoxib; aOR=0.51 (p<.001)  
54 for NSAIDs other than aspirin and celecoxib. For inpatient mortality: aOR=0.34 (p=.060) for  
55 celecoxib and aOR=0.74 (p=.200) for other non-aspirin NSAIDs. Similar effects were observed  
56 for COVID-negative cases, including for inpatient mortality: aOR=0.21 (p<.001) for celecoxib and

57 aOR=0.34 (p<.001) for other non-aspirin NSAIDS. Secondary analyses examined alternative  
58 explanations for results.

59 **Discussion:** Protective effects were observed for non-aspirin NSAIDS, especially  
60 celecoxib. However, those estimated effects implicitly assume the medication classes did not  
61 differ on the probability a true COVID-19 case was diagnosed. The similarity of COVID-positive  
62 and COVID-negative results suggest possible missing covariates. However, such similarity  
63 plausibly could stem from a medication having both “direct” and “indirect” effects on COVID  
64 outcomes. Adjudicating among the alternative interpretations would require data beyond those  
65 available. However, the effects observed for non-aspirin NSAIDS, while possibly biased,  
66 rationalize further investigation using study designs constructed to overcome the limitations of  
67 existing datasets.

68 **Keywords:** Anti-inflammatory, NSAIDs, celecoxib, aspirin, immune suppressant, steroid,  
69 antidepressant, benzodiazepine, COVID19, NC3.

70

71

72

## 73 1. Introduction

74 Inflammation plays a complex and incompletely understood role in the pathogenesis of acute  
75 COVID-19 [1-4] and Post-Acute Sequelae of SARS-CoV-2 infection (PASC or “Long COVID”) [5].  
76 Systemic acute inflammation resulting in cytokine storm, hypercoagulability and endothelial  
77 damage is thought to be a central mechanism for severe morbidity and mortality in acute  
78 COVID-19 [1-4].

79 Several classes of anti-inflammatory agents have been proposed as COVID-19 treatments. A  
80 meta-analysis of 7 randomized controlled trials (RCT) showed that corticosteroids  
81 (dexamethasone, hydrocortisone or methylprednisolone) decrease mortality when used to  
82 treat critically ill COVID-19 patients [6].

83 Non-steroidal anti-inflammatory drugs (NSAIDs) have been considered potentially unsafe in  
84 COVID-19 [7] due to their cardiovascular and renal side effects and the potential to decrease  
85 anti-viral responses [7], though very little information is available on their safety in COVID-19  
86 patients [8]. At least two selective serotonin re-uptake inhibitors (SSRIs), fluvoxamine and  
87 sertraline, were suggested to be effective as adjunctive therapy for COVID-19 based on their  
88 anti-inflammatory effects [9].

89 The anti-inflammatory agents of interest here were used not as therapy for acute events but  
90 rather as “background medications” (i.e., medications taken routinely for chronic conditions).  
91 Here we examine the possible effects on COVID outcomes of several background medication  
92 groups, all of which have anti-inflammatory activity.

93 Using the National COVID Collaborative Cohort (N3C) database [10], Reese et. al. [11] focused  
94 on analyzing the effect of NSAIDs (including celecoxib, diclofenac, droxicam, etodolac,

95    ketorolac, ibuprofen, indomethacin, lornoxicam, meloxicam, naproxen, piroxicam, tenoxicam)  
96    on patients with hospital admissions for COVID-19, finding a statistically significant relationship  
97    between NSAIDs use prior to admission and inpatient mortality (odds ratio = 0.51), but noted  
98    the potential vulnerability of that estimate to missing covariates.

99    The research question that our analyses seek to address is: Is there an association between  
100   background chronic treatment with anti-inflammatory medications and COVID-19 outcomes?  
101   To address that question, we conducted analyses using N3C data, as described below, for  
102   osteoarthritis patients who did not have concomitant autoimmune disorders, as these patients  
103   are most likely to be taking prescription anti-inflammatory agents and are not subject to the  
104   possible confounding effects of autoimmune disorders. Within the osteoarthritis group, we also  
105   considered patients taking immune-suppressant medications (e.g., transplant recipients) and  
106   patients taking SSRIs or systemic, oral steroids, based on the literature on these agents in  
107   COVID-19.

108   The estimation of background medication treatment effects is challenging for several reasons,  
109   leading to estimated treatment effects whose interpretation is ambiguous. Much of this paper  
110   is devoted to development of strategies to help clarify interpretation of estimated medication  
111   effects.

112   The analyses and results described in this paper were generated by using the National Institutes  
113   of Health’s National COVID Cohort Collaborative (N3C) database. N3C is a centralized, secure  
114   harmonized electronic health record database containing U.S. nationwide COVID-19-related  
115   data. As of April 2023, the N3C database included data from 74 data partner (DP) healthcare  
116   organizations. Each DP has contributed data for two types of individuals: (1) Individuals with at

117 least one positive COVID-19 test result or provider diagnosis (here called “COVID-positive”  
118 patients), and (2) individuals who were tested for COVID-19 one or more times but with no  
119 positive test results on record (called “COVID-negative” patients).

## 120 2. Methods

### 121 2.1. Eligibility

122 Each DP contributed clinical data for all its identified COVID-19 positive cases. Each DP also  
123 contributed a sample of its COVID-19 negative cases, with that sample generated by matching  
124 to COVID-19 positive cases on age, gender, race, and ethnicity using a positive-to-negative  
125 matching ratio of 1:2. The N3C data were first made available for research in August 2020 and  
126 have been subsequently updated weekly. The data sent to N3C by each DP include  
127 demographics, symptoms, lab test results, diagnoses, procedures, medications, medical  
128 conditions, physical measurements. Of the 74 DPs contributing data to N3C, data contributions  
129 from 70 DPs were of potential use. As of February 12, 2022 (the date of data extraction for this  
130 project) those 70 DPs had contributed data for 12,093,403 patients. Data from 70 of the 74 DPs  
131 were used. The four excluded DPs’ data could not be used, as those DPs’ data lacked explicit  
132 dates of events (e.g., outcomes events) that the analysis required.

133 The diagram in Fig 1 summarizes how an ultimate group of 485,779 osteoarthritis patients was  
134 selected from those 12,093,403 patients. To avoid the possible confounding effects of  
135 vaccination on COVID outcomes, only data covering events during the time span 01/01/2020  
136 through 03/31/2021 (i.e., before widespread vaccination implementation) were used. To assure  
137 the cohort included only patients with well-documented clinical histories, only adult patients

138 with outpatient visits spanning 30+ days and prescription information spanning 30+ days were  
139 considered potentially eligible.

140 **Figure 1: Selection of patients meeting eligibility criteria.**

141 Some of the N3C DPs do not have inpatient facilities, and their data contributions include little  
142 (inpatient COVID-19 rate  $\leq 0.001$ ) or no inpatient data. Therefore, data from four (of 74) DP  
143 organizations were excluded, because the outcomes measures used in our analyses cannot be  
144 defined without use of inpatient information. Exclusions for age  $<35$  and for missing  
145 demographic information left 2,396,594 individuals eligible before clinical exclusions.

146 To achieve a more nearly clinically homogenous cohort of patients, but a cohort with  
147 substantial use of anti-inflammatory agents, eligibility was further restricted to patients with  
148 diagnosed osteoarthritis but without autoimmune disease. Specifically excluded for  
149 autoimmune disease were those diagnosed with rheumatoid arthritis, lupus, juvenile  
150 rheumatoid arthritis, inflammatory bowel disease, as well as certain other less prevalent  
151 autoimmune disorders. Further, regardless of diagnosis codes present, any patients using  
152 medications strongly associated with autoimmune disease treatment were excluded. The  
153 medications whose use led to exclusion included DMARDs, Methotrexate, cytokine-targeted  
154 biologics, non-anti-inflammatory biologics, and anti-allergy biologics.

155 The above inclusion/exclusion criteria led to a final group of 485,779 osteoarthritis patients, of  
156 which 140,260 were “COVID-positive” and 345,519 were “COVID-negative.”

157 ***2.2. Background medication (treatment) groups***

158 The anti-inflammatory medications considered represent six distinct medication groupings  
159 groups, as shown in Table 1. In addition, a seventh medication group (benzodiazepine

160 sedatives) was defined to serve as a control group. Benzodiazepines were selected because to  
161 our knowledge they lack clinically meaningful anti-inflammatory or anti-viral effects. As  
162 depicted by the solid arrows in Fig 2, the analyses described here compare each of the six anti-  
163 inflammatory medication groups to the control group. Inclusion in a medication group required  
164 evidence (in the form of a prescription date) that the medication was used both within the 30  
165 days prior to COVID-19 diagnosis and also more than 30 days prior to the COVID-19 diagnosis.

166 **Table 1: Background Medication Groups**

<b>Medication group</b>	<b>Descriptions</b>
<b>Non-specific NSAIDs</b>	Non-steroidal anti-inflammatory drugs (excludes celecoxib and aspirin)
<b>Celecoxib</b>	Celecoxib was considered individually due to its selectivity for COX-2 as opposed to other prescription NSAIDs
<b>Aspirin</b>	Aspirin was considered individually as it is often prescribed for its anti-platelet activity to patients at significant risk for cardiovascular or cerebrovascular disease.
<b>Immune suppressants</b>	Small-molecule, non-steroid immune suppressants often prescribed to prevent rejection in transplant recipients were included due to their potential suppressive effects on anti-viral immunity.
<b>Steroids</b>	Steroids were included due to their potential suppressive effects on anti-viral immunity.



<b>Antidepressants</b>	Antidepressants (SSRIs and tricyclic medications) have anti-inflammatory properties and can be prescribed for osteoarthritic pain.
<b>Benzodiazepine sedatives (excluding midazolam)</b>	<b>(This group serves as a comparator /control group.)</b> As a small-molecule control class, we selected benzodiazepine sedatives. These are prescribed as anxiolytics or hypnotics. To our knowledge, this class of medications has no known clinically significant anti-inflammatory or anti-viral effects.

167

168 **Figure 2: Pairwise Comparisons used in the Primary Analyses.**

### 169 *2.3. Primary analyses*

170 Each of the six pairings with the control medication group shown in Fig 2 led to six distinct  
171 statistical models, which estimated six comparison measures which for convenience we refer to  
172 as M1, M2, M3, M4, M5, and M6. After adjustment for demographics and known comorbidities,  
173 three of the measures compare outcomes among COVID-positive patients, and three compare  
174 outcomes among COVID-negative patients. Specifically, the COVID-positive measures are:

175 **M1** = the adjusted odds ratio (aOR) comparing **90-day death rates**, defined as death or  
176 discharge to hospice within the 90 days following the first date on which the subject was  
177 known to be COVID-positive.

178 **M2** = the adjusted odds ratio comparing **90-day admission rates**, defined as a hospital  
179 admission within the 90 days following the first date on which the subject was known to  
180 be COVID-positive.

181 **M3** = the adjusted odds ratio comparing **death rate among hospitalized patients**, defined as  
182 death or discharge to hospice within the 90 days following a hospital admission, with  
183 that admission required to have occurred within +/- seven days of the subject's first  
184 known COVID-positive date.

185 The three corresponding measures for COVID-negative patients are:

186 **M4** = the adjusted odds ratio comparing **90-day death rates**, defined as death or discharge to  
187 hospice within the 90 days following an encounter date randomly selected from the  
188 subject's encounters.

189 **M5** = the adjusted odds ratio comparing **90-day admission rates**, defined as a hospital  
190 admission within the 90 days following an encounter date randomly selected from the  
191 subject's encounters.

192 **M6** = the adjusted odds ratio comparing **death among hospitalized patients**, defined as death  
193 or discharge to hospice within 90 days of a hospital admission, with the hospital  
194 admission randomly selected from among the subject's hospitalizations.

195 Because many patients use multiple medications, each of the above six comparison measures  
196 was estimated using the exclusion logic depicted in Fig 3. For example, when comparing aspirin  
197 users to benzodiazepine users (the control medication group), those using both of those  
198 medications were excluded. Also excluded were those using any of the other Table 1  
199 medications.

200 **Figure 3: Medication-related Inclusion/Exclusion Logic for Pairwise Comparisons.**

201 We also conducted direct outcomes comparisons for all the other pair-wise comparisons of  
202 medication groups, such as the comparison between celecoxib and aspirin. Those results

203 appear in the S1 Supplemental Material. Those modeling results, however, are strongly  
204 transitive. For example, the result of comparing celecoxib and aspirin can be derived with  
205 considerable accuracy from the comparisons of celecoxib to the control and aspirin to the  
206 control.

207 Logistic regression mixed models were used to estimate M1, M2, M3, M4, M5, and M6 for each  
208 of the medication group comparisons defined in Fig 2 (a total of 36 models). In each of those  
209 models, the medication effect was defined as a binary treatment variable (e.g., celecoxib versus  
210 the control group, aspirin versus the control group, etc.). Covariates used in the models  
211 included rurality (rural, urban), gender (female, male), race (black, white, and other), ethnicity  
212 (Hispanic, non- Hispanic), age group (35-44, 45-54, 55-64, 65-74, and 75+), and Charlson  
213 Comorbidity Index [12] score (range 0 to 31), and subjects are nested within DP, and DP was  
214 incorporated into the models as a random effect. Because age was a covariate in the model, it  
215 was adjusted from the Charlson Comorbidity Index formula for this project as follow:  $CCI\_Index$   
216  $= mi + chf + pvd + stroke + dementia + pulmonary + rheumatic + pud + livermild +$   
217  $dm + 2*dmcx + 2*paralysis + 2*renal + 2*cancer + 3*liverSevere + 6*mets + 6*hiv$ . Each single  
218 disease's contribution to this formula was also tested with stepwise logistic regression models  
219 and have no single significant effect.

220 Fig 4 summarizes the eligibility logic underlying the statistical modeling. Eligibility required at  
221 least one outpatient clinic visit and at least one medication prescription over a time span that  
222 ended 30 days prior to the subject's reference date. As shown in Fig 4, we distinguished  
223 between (a) background medication use within 30 days prior to a subject's reference date and  
224 (b) background medication use earlier than 30 days prior to the reference date. For analysis

225 purposes, background medication group (e.g., aspirin users) included only those with evidence  
226 of background medication usage in both of those time spans (e.g., aspirin prescriptions in both  
227 time spans.)

228 **Figure 4: Medication Comparisons Analysis Logic.**

## 229 *2.4. Secondary Analyses*

230 Our secondary analyses are included to help clarify interpretation of the primary analysis  
231 results. In particular, the secondary analyses attempt to address, at least to some extent, two  
232 interpretation questions:

233 **Question 1:** Do background medication effects estimated using data from only diagnosed  
234 COVID cases logically reflect the true effects of the medications?

235 **Question 2:** Do the primary analyses medication effects, estimated as comparisons to the  
236 control medication group, reflect effects actually generated by the background  
237 medications; or do those estimated effects simply reflect unknown risk differences that  
238 where among the factors originally determining background medication choices?

239 As described below, Question #1 and #2 above, both reflecting interpretation ambiguities, arise  
240 for two main reasons: (1) the fact that the background medication user groups were formed, by  
241 definition, prior to exposure to COVID, and (2) the possibility that the medication user groups  
242 differ systematically on unknown risk factors.

### 243 2.4.1. Assessing for Possible Confounding due to Pre-existing Treatment Groups

244 The medication groups compared were implicitly formed prior to the subjects' development of  
245 COVID. Therefore, it is plausible that the medications may have differentially affected the  
246 severity of COVID illness prior to diagnosis, leading to possible differences in the probability

247 that an active COVID case was diagnosed. As described below, the result can be interpretation  
248 ambiguity.

249 Here we refer to a target medication of interest (e.g., aspirin) as the “Medication A” cohort, and  
250 the control medication users as the “Medication B” cohort. Fig 5 conceptually depicts the  
251 nested Medication A patient groups that are involved (either implicitly or explicitly) in our  
252 primary analyses. The full A1 ellipse represents the full cohort of background medication A  
253 users who meet the Fig 4 eligibility criteria, including those whose data are not on the N3C  
254 enclave, because they were never diagnosed. The A2 ellipse represents those who contracted  
255 COVID. A3 represents those who were diagnosed with COVID-19. A4 represents those who had  
256 hospital admissions due to COVID-19. And, finally, A5 represents those Medication A users who  
257 died within 90 days of COVID diagnosis. Note that, while not shown in the figure, there are  
258 analogous groups (B1, B2, B3, B4, and B5) for Medication B users (the control group.) The  
259 COVID-positive cases represented on the N3C enclave for Medication A include only those in A3  
260 (which also implicitly includes A4 and A5). Among the Medication B cohort, the N3C enclave  
261 includes only those in B3 (which includes B4 and B5.)

262 **Figure 5: Conceptual Diagram of Nested Groups within the COVID Positive Treatment**  
263 **Medication Cohort.**

264 Our primary analysis measure M1 compares the Medication A and B groups on mortality among  
265 diagnosed COVID cases. The relative risk (RR) version of M1 in terms of the Fig 5 the nested  
266 patient groups is:

267 (Equation 1)  $M1_{RR} = RR(90\text{-day mortality} | \text{COVID diagnosis})_{A/B} = \frac{P(A5|A3)}{P(B5|B3)} = \frac{N_{A5}/N_{A3}}{N_{B5}/N_{B3}}$

268 (Note: Relative Risk measures are used here, because the mathematical rationale for our  
269 secondary analyses is more efficiently described in relative risk terms.)  
270  $M1_{RR}$  (and M1) ignore any possible medication group difference in the probability of being  
271 diagnosed with COVID. In terms of Fig 5,  $M1_{RR}$  (and M1) use only the cases falling within the A3  
272 and B3 sets. By contrast, a measure  $M7_{RR}$  comparing medication cohorts on full-cohort  
273 mortality would compare values of  $P(A5|A1)$  and  $P(B5|B1)$ ; that is:

274 (Equation 2) 
$$M7_{RR} = RR(90\text{-day mortality})_{A/B} = \frac{P(A5|A1)}{P(B5|B1)} = \frac{N_{A5}/N_{A1}}{N_{B5}/N_{B1}} = \left(\frac{N_{A5}}{N_{B5}}\right)\left(\frac{N_{B1}}{N_{A1}}\right)$$

275  $M7_{RR}$  also has interpretation problems; for example, it could be distorted by medication group  
276 differences in the probability of contracting COVID. However, **if it could be determined that**  
277  **$M1_{RR} \approx M7_{RR}$  holds, that would strongly suggest (albeit not conclusively prove) that pre-**  
278 **diagnosis medication effects are not distorting the medication comparison** and, therefore,  
279  $M1_{RR}$  (and its adjusted odds ratio form M1) is a valid comparison of medication effects on  
280 COVID mortality, and is unlikely to be meaningfully distorted by medication differences in the  
281 probability of COVID diagnosis.

282 But measure  $M7_{RR}$  cannot be estimated directly, because  $N_{B1}$  and  $N_{A1}$  cannot be directly  
283 calculated from COVID-positive N3C data. However, estimating  $M7_{RR}$  does not necessarily  
284 require an ability to estimate  $N_{A1}$  and  $N_{B1}$  individually. All that is required is the ability to  
285 calculate the ratio  $N_{A1}/N_{B1}$ . The Supplemental Material describes an approach to estimating the  
286 ratio  $N_{B1}/N_{A1}$  using the COVID-negative cases appearing on the N3C enclave. The results for  
287 measure  $M7_{RR}$  appearing in Section 3.0 below were obtained using that method.

288 Note that the above describes two different mortality constructs.  $M1_{RR}$  is a comparison of  
289 mortality among those diagnosed with COVID, while  $M7_{RR}$  is a comparison of mortality among  
290 all members of a medication usage cohort. If  $M1_{RR} \approx M7_{RR}$  holds, the two mortality constructs  
291 are numerically very similar, in which case  $M1_{RR}$  can be viewed as a good surrogate for  $M7_{RR}$ .  
292 If  $M1_{RR}$  and  $M7_{RR}$  differ substantially, it might be thought that they still are both valid measures,  
293 each being a measure for its own mortality constructs. That is true. However, as described in  
294 the Supplementary Material, it is not clear that  $M1_{RR}$  has any substantive meaning other than  
295 possibly qualifying as a surrogate for  $M7_{RR}$ . The reason is that the value of  $M1_{RR}$  alone (or  $M1$   
296 alone) does not of itself carry any implications regarding background medication choice.  $M1_{RR}$   
297 (and  $M1$ ) may show a post-diagnosis mortality advantage for background medication A. But  
298 pre-diagnosis medication effects may negate or reverse that advantage.

299 Note that once  $M7_{RR}$  is estimated as described above, the measure  $M8_{RR}$ , the ratio of  $M7_{RR}$  to  
300  $M1_{RR}$ , compares the two medication groups on the probability of being diagnosed with COVID:

301 (Equation 3) 
$$M8_{RR} = \frac{N_{A3}/N_{A1}}{N_{B3}/N_{B1}} \approx \frac{N_{A5}/N_{A1}}{N_{B5}/N_{B1}} \div \frac{N_{A5}/N_{A3}}{N_{B5}/N_{B3}} = \frac{M7_{RR}}{M1_{RR}}$$

302 This measure also is useful in helping to clarify interpretation.

### 303 2.4.2. Possible Distortion due to Missing Covariates

304 Each patient in a particular background medication group (e.g., the Medication A users) is  
305 presumably in the medication group for primarily clinical reasons, although patient preference,  
306 insurance coverage, and chance are contributing factors as well. It is, therefore, quite plausible  
307 that the available data are not sufficient to fully adjust for all the clinical factors underlying  
308 medication decisions.

309 However, in discussing the effects of missing covariates, it is helpful to first describe another  
310 potential COVID-related data characteristic; namely, that a medication’s effect on COVID  
311 outcomes may be either “indirect” or “direct.” An indirect (non-COVID-specific) effect is present  
312 if the target medication alters (reduces or increases) the risk of non-COVID outcomes by  
313 mitigating (or exacerbating) general health risk. By contrast, a direct effect on COVID outcomes  
314 is present if the target medication alters COVID outcomes risk without altering general health  
315 risk.

316 As described in the Supplemental Material, the medication treatment effect in a primary  
317 outcomes model (for example, the M1 effect) is a composite effect that conceptually includes:

- 318 a) Any effects of unknown risk factors that are orthogonal to the known risk factors  
319 included in the model,
- 320 b) Any indirect effects of medication A, stemming from medication A affecting the levels of  
321 risk factors, including both known and unknown risk factors, and
- 322 c) Direct effects of medication A on COVID outcomes.

323 The Supplemental Material describes two scenarios under which the “direct effects” can be  
324 estimated:

325 Scenario 1: Estimate direct effects assuming there may be unknown risk factors, but there are  
326 no indirect medication effects.

327 Scenario 2: Estimate direct effects assuming there may be indirect medication effects but no  
328 unknown risk factors.

329 As shown in Section 3.2 below (and described in the Supplemental Material), the estimated  
330 direct effect varies greatly depending on which of the above scenarios is assumed to hold, and



331 estimates based on the above scenarios serve to conceptually bound the influence of potential  
332 unknown covariates and indirect medication effects. However, it must be kept in mind that the  
333 estimation of a medication's direct effect is not fully satisfactory, because a medication's  
334 indirect effect also is a true medication effect.

## 335 2.5 Ethics Statement

336 N3C operates under the authority of the National Institutes of Health IRB, with Johns Hopkins  
337 University serving as the central IRB (IRB00249128). The N3C Enclave contains de-identified,  
338 retrospective data collected by data partners (DP) under a single IRB-approved protocol  
339 through IRB reliance agreements. No data can be downloaded from the N3C enclave. Analytical  
340 code is uploaded onto the enclave, and only results can be downloaded. The study described  
341 herein was also approved by the Pennington Biomedical Research Center's IRB (20211-016-  
342 PBRC). The N3C Data Access Committee (RP-504BA5) approved the study "COVID-19  
343 Treatments Associated with Lower Mortality". No informed consent was obtained from  
344 individual patients because the study used a limited data set already stripped of direct  
345 identifiers in compliance with the HIPAA Privacy Rule.

## 346 **3. Results**

### 347 3.1 Primary Analysis Results

348 Fig 6 and Table 2 summarize the results from the primary analysis models. Non-specific NSAIDs  
349 as a group were associated with significantly lower risks of admission in both the COVID-  
350 positive and the COVID-negative cohorts and with significantly lower risk of inpatient death for

351 the COVID-negative but not the COVID-positive cohort. (For COVID-positive hospitalization rate  
 352 aOR = 0.79, and for COVID-negative hospitalization rate aOR = 0.78)).

353 **Figure 6: Estimate treatment effects (adjusted odds ratios) from the primary analysis.**

354 **Table 2: Adjusted odds ratios from the primary analysis**

Measure	Med group A	aOR	95% CI	p_value
<b>M1</b>	NSAID	0.51	0.35, 0.74	<0.001
	Celecoxib	0.32	0.09, 0.81	0.032
	Aspirin	1.32	1.04, 1.68	0.024
	Immune suppressant	1.13	0.72, 1.75	0.6
	Steroid	1.15	0.93, 1.45	0.2
	Anti-depressant	0.93	0.72, 1.19	0.5
<b>M2</b>	NSAID	0.79	0.67, 0.92	0.002
	Celecoxib	0.83	0.59, 1.14	0.2
	Aspirin	1.62	1.40, 1.88	<0.001
	Immune suppressant	1.86	1.45, 2.40	<0.001
	Steroid	1.33	1.17, 1.50	<0.001
	Anti-depressant	1.1	0.96, 1.26	0.2
<b>M3</b>	NSAID	0.74	0.45, 1.20	0.2
	Celecoxib	0.14	0.01, 0.71	0.06
	Aspirin	1.08	0.79, 1.49	0.6
	Immune suppressant	1.15	0.68, 1.94	0.6
	Steroid	1.23	0.93, 1.66	0.2
	Anti-depressant	1.01	0.73, 1.40	>0.9
<b>M4</b>	NSAID	0.25	0.18, 0.34	<0.001
	Celecoxib	0.11	0.04, 0.27	<0.001

	Aspirin	0.86	0.73, 1.03	0.1
	Immune suppressant	0.37	0.22, 0.60	<0.001
	Steroid	0.77	0.66, 0.91	0.001
	Anti-depressant	0.65	0.53, 0.79	<0.001
<b>M5</b>	NSAID	0.78	0.71, 0.86	<0.001
	Celecoxib	0.95	0.80, 1.11	0.5
	Aspirin	1.17	1.06, 1.28	0.001
	Immune suppressant	0.92	0.75, 1.12	0.4
	Steroid	1.26	1.16, 1.36	<0.001
	Anti-depressant	0.93	0.85, 1.02	0.12
<b>M6</b>	NSAID	0.34	0.26, 0.43	<0.001
	Celecoxib	0.21	0.12, 0.32	<0.001
	Aspirin	0.66	0.58, 0.76	<0.001
	Immune suppressant	0.49	0.35, 0.68	<0.001
	Steroid	0.79	0.70, 0.89	<0.001
	Anti-depressant	0.67	0.58, 0.78	<0.001

355

356 Celecoxib showed no significant association for hospitalization in either for COVID-positive

357 hospitalization rate aOR = 0.83, or for COVID-negative hospitalization rate aOR = 0.95.

358 Conversely, aspirin was associated with significantly higher aOR of hospitalization in COVID-

359 positive aOR = 1.62 and to a lesser extent in COVID-negative patients aOR = 1.17.

360 Immune suppressants were associated with significantly higher aOR of hospitalization in COVID-

361 positive aOR = 1.86 but not in COVID-negative cases aOR = 0.92.

362 Background steroid treatment was associated with significantly higher aOR of hospitalization in  
363 both groups, with very similar point estimates. For COVID-positive hospitalization rate, aOR =  
364 1.33, and for COVID-negative hospitalization rate aOR = 1.26.

365 No significant associations were observed for antidepressants as a group. (For COVID-positive  
366 hospitalization rate aOR = 1.1, and for COVID-negative hospitalization rate aOR = 0.94).

367 The medication group that showed the largest difference between COVID-positive and COVID-  
368 negative cases was immune suppressants (1.86 versus 0.92, respectively), consistent with the  
369 notion that patients on these medications are at higher risk for infectious complications,  
370 including COVID-19 [12].

371 When we examined the risk of death in all comers (inpatients and outpatients), non-specific  
372 NSAIDs and celecoxib were associated with significantly lower aOR of mortality in both COVID-  
373 positive (for NSAIDs' mortality rate aOR = 0.51, and for celecoxib's mortality rate aOR = 0.32)  
374 and COVID-negative cases (for NSAIDs' mortality rate aOR = 0.25, and for celecoxib's mortality  
375 rate aOR = 0.11), while aspirin was associated with significantly higher aOR of mortality in  
376 COVID-positive aOR = 1.32 but not in COVID-negative aOR = 0.86. Similarly, background steroid  
377 treatment was associated with significantly higher aOR of mortality in COVID-positive aOR =  
378 1.15 but not in COVID-negative cases aOR = 0.77 while antidepressants showed no significant  
379 association with aOR of mortality in COVID-positive aOR = 0.93 and a significant negative  
380 association in COVID-negative aOR = 0.65.

### 381 3.2 Secondary Analysis Results

382 The secondary analyses focus on the interpretation questions posed in Section 2.4. Fig 7  
383 compares values of measure  $M1_{RR}$  (relative risk of death given a COVID diagnosis) to values of

384  $M7_{RR}$  (relative risk of COVID death among all medication cohort members.) Results for the two  
385 measures are quite similar for all the medications except celecoxib and the immune  
386 suppressants, for which  $M1_{RR}$  and  $M7_{RR}$  differ approximately by a factor of two, but in different  
387 directions. The implication is that the diagnosis-denominated measure,  $M1_{RR}$ , may not  
388 adequately reflect the COVID mortality experience of those two medication groups. However,  
389 the sample sizes for celecoxib and for immune suppressants are too small to draw any  
390 confident conclusions.

391 **Figure 7: Relative Risk of Death | COVID Diagnosis and Relative Risk of Death | in Medication**  
392 **Cohort.**

393 Fig 8 shows estimates of medication direct effects under two different assumptions: an  
394 assumption that some relevant covariates are unknown but there are no indirect medication  
395 effects, versus the assumption that indirect medication effects may be present but there are  
396 not missing risk factors. The adjusted odds ratios for our primary analysis measure  $M1$  are  
397 graphed by the solid black line. The dashed red line graphs  $M1$  after adjustment for the  
398 potential effects of unknown risk factors. The dashed blue line graphs  $M1$  after adjustment for  
399 the potential indirect medication effects. Of course, bias may be present from both unknown  
400 covariates and indirect medication effects. The potential bias from indirect medication effects is  
401 much less than the potential bias from unknown covariates. Further, the actual bias present  
402 from either source may be much less than the full potential bias quantified in Fig 8.

403 **Figure 8: Medication Effects after adjustment for unknown covariates of adjustment for**  
404 **indirect effects.**

405 In Fig 8, results under an assumption of indirect medication effects seem more plausible than  
406 results assuming possible missing covariates.

## 407 4. Discussion

408 With the interpretation caveats described above, given osteoarthritis patients who had not  
409 been diagnosed with autoimmune disorders as a target population, background NSAIDs except  
410 aspirin were associated with lower hospitalization rates as compared to the non-anti-  
411 inflammatory control group (sedatives). In contrast, background aspirin and systemic steroids  
412 were associated with higher admission rates compared to control.

413 The results described in this study in the N3C enclave provide potentially useful information for  
414 clinicians treating patients at risk for COVID-19. It is important to point out that these results do  
415 not necessarily imply that pharmacological activities of the agents we studied directly affect  
416 COVID-19 outcomes. This is because of the interpretation challenges described under Methods,  
417 and because groups of patients identified by treatment with each of these medication classes  
418 do not have identical baseline risks of hospitalization and death, nor do they have identical  
419 baseline risk profiles for SARS-CoV2 infection or symptomatic COVID-19 disease. Rather, each  
420 medication class can be seen as identifying a group of patients potentially at higher or lower  
421 risk of hospitalization or death, and COVID-19 adds to these different baseline risk profiles.

422 Among patients with osteoarthritis, background prescription NSAIDs, either non-specific or  
423 COX-2 selective, do not appear to contribute to the risk of COVID-19 hospitalization or death,  
424 with the prominent exception of aspirin. Patients on background prescription aspirin appear to  
425 be at higher risk of hospitalization when COVID-positive, and to a lesser extent when COVID-  
426 negative, and to be at significantly higher risk of non-inpatient death (presumably from acute  
427 complications prior to hospitalization) when COVID-positive but not when COVID-negative. A  
428 possible explanation for this is protopathic effect, meaning that aspirin is used to treat

429 conditions that increase the risk of hospitalization and death in COVID-positive patients. Unlike  
430 other NSAIDs, in addition to its uses as an anti-inflammatory agent, aspirin is used alone or in  
431 combination with other agents as part of anti-platelet therapy to prevent thrombosis in  
432 patients at risk after acute cardiovascular events or procedures, e.g. patients with percutaneous  
433 coronary intervention (PCI), [13], total hip arthroplasty [14], or limb revascularization [15].  
434 Furthermore, prescription aspirin is used for the primary prevention of coronary artery disease  
435 [16]. These conditions are likely to increase the baseline risk of hospitalization and sudden  
436 death in COVID-positive patients. Cardiovascular disease (CVD) is a recognized risk factor for  
437 COVID-19 morbidity and mortality [17]. Our data suggest that patients on chronic prescription  
438 aspirin who are still unvaccinated against currently circulating strains of SARS-CoV-2 should  
439 consider COVID-19 vaccination and mitigate the risk of infection.

440 Our results for NSAIDs are consistent with the results obtained by Reese et al [11],  
441 whose estimated effect of NSAIDs, including Celecoxib, on inpatient mortality (OR = 0.51) lies  
442 between our estimated effects for Celecoxib (OR = 0.14) and non-specific NSAIDs (OR = 0.74).

443 Celecoxib deserves separate consideration, as background treatment with it was  
444 associated with reduced relative risk of death in both COVID-positive and COVID-negative  
445 patients. We do not know whether these effects are due to a hidden variable common to  
446 celecoxib-treated patients or can be attributed to the medication itself. If celecoxib simply  
447 masked symptoms of mild COVID-19, preventing diagnosis of such cases, one would expect the  
448 opposite effect, i.e., a higher risk of severe outcomes in celecoxib-treated patients. COX-2  
449 inhibition had been proposed as a possible treatment for COVID-19 [18] and a small clinical trial  
450 in mild or moderate COVID-19 supports this notion [19]. In that study, lymphocyte counts were

451 increased in patients taking celecoxib. Furthermore, a large real-world evidence observational  
452 study by Liao [20], based on claims data from 2,935,415 unvaccinated and 189,692 vaccinated  
453 patients, identified NSAIDs including celecoxib-aspirin combinations, celecoxib-ibuprofen  
454 combinations and celecoxib single agent as having repurposing potential for COVID-19  
455 treatment [20]. Unlike our results, aspirin was also associated with a lower risk of COVID-19  
456 mortality in that study. Significant differences between the two studies are the inclusion  
457 criteria, with in our study are restricted to osteoarthritis without autoimmune disorders, and  
458 the fact that risk adjustment in Liao's study used a proprietary Optum Episode Risk Group Score  
459 [18]. A different theoretical approach, using multi-evidence deep graph neural networks [21],  
460 predicted anti-COVID-19 activities for aspirin and celecoxib. A molecular modeling study  
461 identified celecoxib as a potential inhibitor of the SARS-CoV-2 main protease, thus  
462 hypothesizing a direct antiviral activity [22]. Besides this putative antiviral activity, COX-2, the  
463 primary target of celecoxib, is a key player in inflammation, and its major product, PGE<sub>2</sub>,  
464 suppresses T-cell-mediated cellular immunity while promoting Th2 responses [23]. It is possible  
465 that COX-2 inhibition may improve immune responses to SARS-CoV-2. The apparent association  
466 of background celecoxib with lower risk of mortality in COVID-negative patients is more difficult  
467 to explain. Given that celecoxib is contraindicated in patients with history of cardiovascular  
468 events, including myocardial infarction and PCI or other coronary revascularization, it is possible  
469 that the baseline risk of cardiovascular events in patients on celecoxib may be lower than that  
470 of patients on other medication groups we studied. However, the PRECISION trial results  
471 showed that at currently recommended doses, the cardiovascular and renal safety profile of  
472 celecoxib is favorable compared to non-specific NSAIDs ibuprofen and naproxen [24]. It's worth



473 noting that PGE<sub>2</sub> also suppresses tumor immunity [25, 26], some studies support the use of  
474 celecoxib as an adjuvant to cancer treatment [27] and a vast literature supports the use of  
475 NSAIDs [28-31], and celecoxib in particular [32] as cancer chemo-preventive agents. Initial  
476 enthusiasm for selective COX-2 inhibitors was dampened by their cardiovascular side effects  
477 [33]. Our data, within the limitations of the N3C dataset and the analytical strategy we used,  
478 are consistent with the notion that background use of celecoxib is associated with a decreased  
479 risk of all-cause mortality in COVID-negative and -positive patients. This data supports the  
480 predictions made by Liao [20] and Hsieh et al. [21].

481 Background steroid medications were association with increased risk of inpatient death and  
482 all-comers death in COVID-19 positive but not in COVID-19 negative patients. Despite the fact  
483 that dexamethasone is effective in treating severe COVID-19, it is possible that chronic use of  
484 steroids may dampen protective immunity against SARS-CoV-2.

485 Immune-suppressants (e.g. rapalogs) were associated with increased risk of hospitalization  
486 but not death in COVID-19 positive but not negative cases. These medications are usually  
487 prescribed to transplant recipients, who have been identified as being at higher risk for severe  
488 COVID-19 [32]. A possible explanation for our findings is that patients on immune-suppressants  
489 on average adopted precautions to avoid SARS-CoV-2 exposure, but once infected were more  
490 likely to be hospitalized. The lack of significantly increased risk of death is an encouraging  
491 finding.

## 492 5. Conclusions

493 In conclusion, our analysis of N3C enclave data suggests that patients on prescription non-  
494 specific NSAIDs or celecoxib do not face increased risks of severe outcomes in COVID-19, with  
495 the possible exception of patients on aspirin. Celecoxib was associated with decreased risk of  
496 mortality in both COVID-negative and COVID-positive cases. Unvaccinated patients on chronic  
497 systemic steroids and particularly immune-suppressive medications may face increased risk of  
498 severe COVID-19 outcomes. Our data do not support the notion that background treatment  
499 with antidepressants or fluvoxamine affect COVID-19 outcomes in the patient categories we  
500 studied.

501

## 502 Acknowledgements

503 National COVID Cohort Collaborative (N3C) Consortium membership includes:

504 1. Christopher G. Chute

505 Johns Hopkins University

506 ORCID: 0000-0001-5437-2545

507 chute@jhu.edu

508 Contributions: clinical data model expertise, data curation, data integration, data quality

509 assurance, data security, funding acquisition, governance, N3C Phenotype definition, project

510 evaluation, project management, and regulatory oversight/admin.

### 511 Individual Acknowledgements for Core Contributors

512 We gratefully acknowledge the following core contributors to N3C: Adam B. Wilcox, Adam M.

513 Lee, Alexis Graves, Alfred (Jerrod) Anzalone, Amin Manna, Amit Saha, Amy Olex, Andrea Zhou,

514 Andrew E. Williams, Andrew M. Southerland, Andrew T. Girvin, Anita Walden, Anjali

515 Sharathkumar, Benjamin Amor, Benjamin Bates, Brian Hendricks, Brijesh Patel, G. Caleb

516 Alexander, Carolyn T. Bramante, Cavin Ward-Caviness, Charisse Madlock-Brown, Christine

517 Suver, Christopher G. Chute, Christopher Dillon, Chunlei Wu, Clare Schmitt, Cliff Takemoto, Dan

518 Housman, Davera Gabriel, David A. Eichmann, Diego Mazzotti, Donald E. Brown, Eilis Boudreau,

519 Elaine L. Hill, Emily Carlson Marti, Emily R. Pfaff, Evan French, Farrukh M Korashy, Federico

520 Mariona, Fred Prior, George Sokos, Greg Martin, Harold P. Lehmann, Heidi Spratt, Hemalkumar

521 B. Mehta, J.W. Awori Hayanga, Jami Pincavitch, Jaylyn Clark, Jeremy Richard Harper, Jessica

522 Yasmine Islam, Jin Ge, Joel Gagnier, Johanna J. Loomba, John B. Buse, Jomol Mathew, Joni L.

523 Rutter, Julie A. McMurry, Justin Guinney, Justin Starren, Karen Crowley, Katie Rebecca Bradwell,

524 Kellie M. Walters, Ken Wilkins, Kenneth R. Gersing, Kenrick Cato, Kimberly Murray, Kristin  
525 Kostka, Lavance Northington, Lee Pyles, Lesley Cottrell, Lili M. Portilla, Mariam Deacy, Mark M.  
526 Bissell, Marshall Clark, Mary Emmett, Matvey B. Palchuk, Melissa A. Haendel, Meredith Adams,  
527 Meredith Temple-O'Connor, Michael G. Kurilla, Michele Morris, Nasia Safdar, Nicole Garbarini,  
528 Noha Sharafeldin, Ofer Sadan, Patricia A. Francis, Penny Wung Burgoon, Philip R.O. Payne,  
529 Randeep Jawa, Rebecca Erwin-Cohen, Rena C. Patel, Richard A. Moffitt, Richard L. Zhu,  
530 Rishikesan Kamaleswaran, Robert Hurley, Robert T. Miller, Saiju Pyarajan, Sam G. Michael,  
531 Samuel Bozzette, Sandeep K. Mallipattu, Satyanarayana Vedula, Scott Chapman, Shawn T.  
532 O'Neil, Soko Setoguchi, Stephanie S. Hong, Steven G. Johnson, Tellen D. Bennett, Tiffany J.  
533 Callahan, Umit Topaloglu, Valery Gordon, Vignesh Subbian, Warren A. Kibbe, Wenndy  
534 Hernandez, Will Beasley, Will Cooper, William Hillegass, Xiaohan Tanner Zhang. Details of  
535 contributions available at [covid.cd2h.org/core-contributors](https://covid.cd2h.org/core-contributors).  
536 The authors received direct benefit from discussions and presenting the project to the N3C –  
537 Rural Health Domain teams. The authors would like to thank Sharon Patrick and the West  
538 Virginia Clinical and Translational Science Institute (WVCTSI) for project planning and  
539 management.

## 540 References

541 1. Vollbracht C, Kraft K. Oxidative Stress and Hyper-Inflammation as Major Drivers of  
542 Severe COVID-19 and Long COVID: Implications for the Benefit of High-Dose Intravenous  
543 Vitamin C. *Front Pharmacol.* 2022;13:899198. Epub 20220429. doi:

- 544 10.3389/fphar.2022.899198. PubMed PMID: 35571085; PubMed Central PMCID:  
545 PMCPMC9100929.
- 546 2. Montazersaheb S, Hosseiniyan Khatibi SM, Hejazi MS, Tarhriz V, Farjami A, Ghasemian  
547 Sorbeni F, et al. COVID-19 infection: an overview on cytokine storm and related interventions.  
548 Virol J. 2022;19(1):92. Epub 20220526. doi: 10.1186/s12985-022-01814-1. PubMed PMID:  
549 35619180; PubMed Central PMCID: PMCPMC9134144.
- 550 3. Eskandarian Boroujeni M, Sekrecka A, Antonczyk A, Hassani S, Sekrecki M, Nowicka H, et  
551 al. Dysregulated Interferon Response and Immune Hyperactivation in Severe COVID-19:  
552 Targeting STATs as a Novel Therapeutic Strategy. Front Immunol. 2022;13:888897. Epub  
553 20220517. doi: 10.3389/fimmu.2022.888897. PubMed PMID: 35663932; PubMed Central  
554 PMCID: PMCPMC9156796.
- 555 4. Bahmani M, Chegini R, Ghanbari E, Sheykhsaran E, Shiri Aghbash P, Leylabadlo HE, et al.  
556 Severe acute respiratory syndrome coronavirus 2 infection: Role of interleukin-6 and the  
557 inflammatory cascade. World J Virol. 2022;11(3):113-28. doi: 10.5501/wjv.v11.i3.113. PubMed  
558 PMID: 35665236; PubMed Central PMCID: PMCPMC9150027.
- 559 5. Newell KL, Waickman AT. Inflammation, immunity, and antigen persistence in post-  
560 acute sequelae of SARS-CoV-2 infection. Curr Opin Immunol. 2022;77:102228. Epub 20220524.  
561 doi: 10.1016/j.coi.2022.102228. PubMed PMID: 35724449; PubMed Central PMCID:  
562 PMCPMC9127180.
- 563 6. Group WHOREAfC-TW, Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, et al.  
564 Association Between Administration of Systemic Corticosteroids and Mortality Among Critically  
565 Ill Patients With COVID-19: A Meta-analysis. JAMA. 2020;324(13):1330-41. doi:

- 566 10.1001/jama.2020.17023. PubMed PMID: 32876694; PubMed Central PMCID:  
567 PMCPMC7489434.
- 568 7. Zhang W, Qin C, Fei Y, Shen M, Zhou Y, Zhang Y, et al. Anti-inflammatory and immune  
569 therapy in severe coronavirus disease 2019 (COVID-19) patients: An update. Clin Immunol.  
570 2022;239:109022. Epub 20220425. doi: 10.1016/j.clim.2022.109022. PubMed PMID: 35477027;  
571 PubMed Central PMCID: PMCPMC9040414.
- 572 8. Micallef J, Soeiro T, Jonville-Bera AP, French Society of Pharmacology T. Non-steroidal  
573 anti-inflammatory drugs, pharmacology, and COVID-19 infection. Therapie. 2020;75(4):355-62.  
574 Epub 20200507. doi: 10.1016/j.therap.2020.05.003. PubMed PMID: 32418728; PubMed Central  
575 PMCID: PMCPMC7204680.
- 576 9. Eteraf-Oskouei T, Najafi M. The relationship between the serotonergic system and  
577 COVID-19 disease: A review. Heliyon. 2022;8(5):e09544. Epub 20220526. doi:  
578 10.1016/j.heliyon.2022.e09544. PubMed PMID: 35652122; PubMed Central PMCID:  
579 PMCPMC9132783.
- 580 10. Haendel MA, Chute CG, Bennett TD, et al. The National COVID Cohort Collaborative  
581 (N3C): Rationale, design, infrastructure, and deployment. J Am Med Inform Assoc.  
582 2021;28(3):427-443. doi:10.1093/jamia/ocaa196
- 583 11. Reese JT, Coleman B, Chan L, Blau H, Callahan TJ, Cappelletti L, et al. NSAID use and  
584 clinical outcomes in COVID-19 patients: a 38-center retrospective cohort study. Virol J. 2022  
585 May 15;19(1):84. doi: 10.1186/s12985-022-01813-2. PMID: 35570298; PMCID: PMC9107579.

- 586 12. Quan-Charlson paper here: Quan H, Li B, Couris CM, et al. Updating and validating the  
587 Charlson comorbidity index and score for risk adjustment in hospital discharge abstracts using  
588 data from 6 countries. *Am J Epidemiol*. 2011;173(6):676-682. doi:10.1093/aje/kwq433
- 589 13. Danziger-Isakov L, Blumberg EA, Manuel O, Sester M. Impact of COVID-19 in solid organ  
590 transplant recipients. *Am J Transplant*. 2021;21(3):925-37. Epub 20210226. doi:  
591 10.1111/ajt.16449. PubMed PMID: 33319449; PubMed Central PMCID: PMCPMC9800718.
- 592 14. Nso N, Nassar M, Zirkiyeva M, Mbome Y, Lyonga Ngonge A, Badejoko SO, et al. Factors  
593 Impacting Stent Thrombosis in Patients With Percutaneous Coronary Intervention and Coronary  
594 Stenting: A Systematic Review and Meta-Analysis. *Cureus*. 2022;14(4):e23973. Epub 20220409.  
595 doi: 10.7759/cureus.23973. PubMed PMID: 35547463; PubMed Central PMCID:  
596 PMCPMC9089933.
- 597 15. Borton ZM, Bhangoo NS, Quah CS, Stephen AB, Howard PW. Aspirin monotherapy is a  
598 suitable standard thromboprophylactic agent following total hip arthroplasty. *Hip Int*.  
599 2022;32(3):286-90. Epub 20210218. doi: 10.1177/1120700021990544. PubMed PMID:  
600 33601921.
- 601 16. Ipema J, Brand AR, GJ DEB, JP DEV, Unl UC. Antiplatelet and anticoagulation therapy  
602 after revascularization for lower extremity artery disease: a national survey and literature  
603 overview. *J Cardiovasc Surg (Torino)*. 2021;62(1):59-70. Epub 20201028. doi: 10.23736/S0021-  
604 9509.20.11402-2. PubMed PMID: 33112125.
- 605 17. Ujjawal A, Gupta M, Ghosh RK, Jain V, Bandyopadhyay D, Qamar A, et al. Aspirin for  
606 Primary Prevention of Coronary Artery Disease. *Curr Probl Cardiol*. 2021;46(3):100553. Epub  
607 20200214. doi: 10.1016/j.cpcardiol.2020.100553. PubMed PMID: 32222319.

- 608 18. Vasudeva R, Challa A, Al Rifai M, Polana T, Duran B, Vindhya M, et al. Prevalence of  
609 cardiovascular diseases in COVID-19 related mortality in the United States. *Prog Cardiovasc Dis.*  
610 2022;74:122-6. Epub 20221021. doi: 10.1016/j.pcad.2022.09.002. PubMed PMID: 36279944;  
611 PubMed Central PMCID: PMC9585886.
- 612 19. Baghaki S, Yalcin CE, Baghaki HS, Aydin SY, Daghan B, Yavuz E. COX2 inhibition in the  
613 treatment of COVID-19: Review of literature to propose repositioning of celecoxib for  
614 randomized controlled studies. *Int J Infect Dis.* 2020;101:29-32. Epub 20200930. doi:  
615 10.1016/j.ijid.2020.09.1466. PubMed PMID: 33007455; PubMed Central PMCID:  
616 PMC97525269.
- 617 20. Ghaznavi H, Mohammadghasemipour Z, Shirvaliloo M, Momeni MK, Metanat M,  
618 Gorgani F, et al. Short-term celecoxib (celebrex) adjuvant therapy: a clinical trial study on  
619 COVID-19 patients. *Inflammopharmacology.* 2022;30(5):1645-57. Epub 20220714. doi:  
620 10.1007/s10787-022-01029-4. PubMed PMID: 35834150; PubMed Central PMCID:  
621 PMC9281238.
- 622 21. Liao Y. Identification of potential new COVID-19 treatments via RWD-driven drug  
623 repurposing. *Sci Rep.* 2023;13(1):14586. Epub 20230904. doi: 10.1038/s41598-023-40033-8.  
624 PubMed PMID: 37666866; PubMed Central PMCID: PMC977169.
- 625 22. Hsieh K, Wang Y, Chen L, Zhao Z, Savitz S, Jiang X, et al. Drug repurposing for COVID-19  
626 using graph neural network and harmonizing multiple evidence. *Sci Rep.* 2021;11(1):23179.  
627 Epub 20211130. doi: 10.1038/s41598-021-02353-5. PubMed PMID: 34848761; PubMed Central  
628 PMCID: PMC8632883.



- 629 23. Gimeno A, Mestres-Truyol J, Ojeda-Montes MJ, Macip G, Saldivar-Espinoza B, Cereto-  
630 Massague A, et al. Prediction of Novel Inhibitors of the Main Protease (M-pro) of SARS-CoV-2  
631 through Consensus Docking and Drug Reposition. *Int J Mol Sci.* 2020;21(11). Epub 20200527.  
632 doi: 10.3390/ijms21113793. PubMed PMID: 32471205; PubMed Central PMCID:  
633 PMCPMC7312484.
- 634 24. Kalinski P. Regulation of immune responses by prostaglandin E2. *J Immunol.*  
635 2012;188(1):21-8. doi: 10.4049/jimmunol.1101029. PubMed PMID: 22187483; PubMed Central  
636 PMCID: PMCPMC3249979.
- 637 25. Obeid S, Libby P, Husni E, Wang Q, Wisniewski LM, Davey DA, et al. Cardiorenal risk of  
638 celecoxib compared with naproxen or ibuprofen in arthritis patients: insights from the  
639 PRECISION trial. *Eur Heart J Cardiovasc Pharmacother.* 2022;8(6):611-21. doi:  
640 10.1093/ehjcvp/pvac015. PubMed PMID: 35234840.
- 641 26. Liu B, Qu L, Yan S. Cyclooxygenase-2 promotes tumor growth and suppresses tumor  
642 immunity. *Cancer Cell Int.* 2015;15:106. Epub 20151105. doi: 10.1186/s12935-015-0260-7.  
643 PubMed PMID: 26549987; PubMed Central PMCID: PMCPMC4635545.
- 644 27. Jin K, Qian C, Lin J, Liu B. Cyclooxygenase-2-Prostaglandin E2 pathway: A key player in  
645 tumor-associated immune cells. *Front Oncol.* 2023;13:1099811. Epub 20230127. doi:  
646 10.3389/fonc.2023.1099811. PubMed PMID: 36776289; PubMed Central PMCID:  
647 PMCPMC9911818.
- 648 28. Ye SY, Li JY, Li TH, Song YX, Sun JX, Chen XW, et al. The Efficacy and Safety of Celecoxib in  
649 Addition to Standard Cancer Therapy: A Systematic Review and Meta-Analysis of Randomized  
650 Controlled Trials. *Curr Oncol.* 2022;29(9):6137-53. Epub 20220825. doi:

- 651 10.3390/currenocol29090482. PubMed PMID: 36135051; PubMed Central PMCID:  
652 PMCPMC9497539.
- 653 29. Patrono C. Cyclooxygenase Inhibitors and Cancer: The Missing Pieces. *J Pharmacol Exp*  
654 *Ther.* 2023;386(2):181-9. Epub 20230606. doi: 10.1124/jpet.122.001631. PubMed PMID:  
655 37280092.
- 656 30. Rashid G, Khan NA, ElSORI D, Rehman A, Tanzeelah, Ahmad H, et al. Non-steroidal anti-  
657 inflammatory drugs and biomarkers: A new paradigm in colorectal cancer. *Front Med*  
658 (Lausanne). 2023;10:1130710. Epub 20230306. doi: 10.3389/fmed.2023.1130710. PubMed  
659 PMID: 36950511; PubMed Central PMCID: PMCPMC10025514.
- 660 31. Newman P, Muscat J. Potential Role of Non-Steroidal Anti-Inflammatory Drugs in  
661 Colorectal Cancer Chemoprevention for Inflammatory Bowel Disease: An Umbrella Review.  
662 *Cancers (Basel).* 2023;15(4). Epub 20230209. doi: 10.3390/cancers15041102. PubMed PMID:  
663 36831446; PubMed Central PMCID: PMCPMC9954537.
- 664 32. Gadi V, Shetty SR. Potential of Anti-inflammatory Molecules in the Chemoprevention of  
665 Breast Cancer. *Recent Adv Inflamm Allergy Drug Discov.* 2022;16(2):60-76. doi:  
666 10.2174/2772270816666220829090716. PubMed PMID: 36043708.
- 667 33. Abdelhaleem EF, Kassab AE, El-Nassan HB, Khalil OM. Recent advances in the  
668 development of celecoxib analogs as anticancer agents: A review. *Arch Pharm (Weinheim).*  
669 2022;355(12):e2200326. Epub 20220822. doi: 10.1002/ardp.202200326. PubMed PMID:  
670 35996360.
- 671 34. El-Malah AA, Gineinah MM, Deb PK, Khayyat AN, Bansal M, Venugopala KN, et al.  
672 Selective COX-2 Inhibitors: Road from Success to Controversy and the Quest for Repurposing.

673    Pharmaceuticals (Basel). 2022;15(7). Epub 20220703. doi: 10.3390/ph15070827. PubMed  
674    PMID: 35890126; PubMed Central PMCID: PMC9318302.  
675    35.    Eyitemi J, Thomas B, Ramos Y, Feng X, Ezekwesili C. SARS-CoV-2: Review of Conditions  
676    Associated With Severe Disease and Mortality. Int J Prev Med. 2022;13:109. Epub 20220808.  
677    doi: 10.4103/ijpvm.IJPVM\_640\_20. PubMed PMID: 36247195; PubMed Central PMCID:  
678    PMC9564226.

## 679    **Supporting information**

### 680           **S1 Supplemental Material**




Fig 1

Non-specific NSAIDs

Celecoxib

Aspirin

Immune Suppressants

Anti-depression meds

Steroids

**Sedatives**  
**(control)**

A diagram consisting of six blue lines that originate from the left side of the image and converge towards a single point on the right. The lines are arranged in a fan shape, with the top line being the longest and the bottom line being the shortest. The lines are positioned between the text labels on the left and the central text on the right.

Fig 2




Fig 3




Fig 4







Fig 5






M1, M2, and M3 are the adjusted aOR values comparing **90-day death rates, 90-day admission rates, and death rate among hospitalized patients** for COVID positive subjects. M4, M5, and M6 are the adjusted aOR values comparing **90-day death rates, 90-day admission rates, and death rate among hospitalized patients** for COVID negative subjects.


Each adjusted odds ratio (aOR) in the figure is an estimate medication effect from one model – adjusted for demographics, Charlson Comorbidity Index, with data partner as a random effect.

Fig 6



Dashed line: relative risk of death given a COVID diagnosis  
 Solid Line: relative risk of death among all people in the medication cohort  
 Bars are 95% confidence intervals  
 Numbers above bars are pvalues for difference in the two relative risk values.

Fig 7



Lower curve: Measure M1 values from primary analysis.  
 Upper curve: Measure M1 values after adjustment for unknown covariates.  
 Middle curve: Measure M1 values with indirect medication effects removed.

Fig 8