It is made available under a CC-BY 4.0 International license .

Algorithm Refinement of the National Early Warning Score 2 (NEWS2): Study Protocol

Chris Plummer, BSc, PhD, BM, BCh, FRCP ^{1,2,*} Madison Milne-Ives, BAS, MSc^{,2,5}, Cen Cong, BEng, MA,³ Lynsey Threlfall, MBBS, MRCP,^{1,2} Edward Meinert, MA, MSc, MBA, MPA, MPH, PhD, CEng FBCS EUR ING^{3,4}

¹The Newcastle Upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, NE7 7DN, UK

² NIHR Newcastle Biomedical Research Centre, Newcastle University, Newcastle upon Tyne, NE1 7RU, UK

³ Translational and Clinical Research Institute, Newcastle University, Newcastle upon Tyne, UK

⁴ Department of Primary Care and Public Health, Imperial College London, London, UK

⁵ Centre for Health Technology, School of Nursing and Midwifery, University of Plymouth, Plymouth, UK

Correspondence: [*] Correspondence to: Chris Plummer, National Institute for Health and Care Research Newcastle Biomedical Research Centre, Newcastle University, Newcastle, UK. Tel: <u>+44(0) 191 2081148</u>; e-mail; <u>Chris.Plummer@newcastle.ac.uk</u>

Abstract

Introduction: The second iteration of the National Early Warning Score (NEWS2) has been widely adopted for predicting patient deterioration in healthcare settings using routinely collected physiological observations. The use of NEWS2 has been shown to reduce in-hospital mortality, but it has limited accuracy in the prediction of clinically important outcomes, especially over longer time periods. The increasing implementation of digital patient observations and health records presents an opportunity to investigate whether the addition of individual patient characteristics and information about their care-setting, would improve the predictive accuracy of the score.

Methods and analysis: This protocol describes the work to determine whether the performance of the current NEWS2 system could be improved by the use of additional variables. The project has been designed after an extensive scoping review of existing literature on NEWS2 and an exploration of retrospective cohort data in The Newcastle upon Tyne Hospitals NHS Foundation Trust, with input from key clinical stakeholders.

Ethics and dissemination: The project has received competitive funding following peer-review, from the NIHR Newcastle Biomedical Research Centre as an Interdisciplinary Research Award. Ethical approval has been requested. Findings are expected to be produced by June 2025, and will be disseminated at symposia, conferences and in journal publications.

Keywords: National Early Warning Score, clinical deterioration, performance, proof-of-concept

Strengths and limitations of this study

Strengths:

- This work highlights the importance of investigating the use of additional clinical variables to those used in NEWS2, in the development of a new early warning score
- The study design was informed by an evidence synthesis of the literature

Limitations:

- Some retrospective data sets may be of low quality and/or incomplete
- External validation will be needed to test algorithm generalisability

It is made available under a CC-BY 4.0 International license .

1. Introduction

The second National Early Warning Score (NEWS2) system is a widely used tool for the systematic documentation and identification of clinical deterioration. It has been demonstrated to be effective at reducing in-hospital mortality [1–4], it facilitates effective communication between clinicians and enables timely interventions to improve patient outcomes, but NEWS2 has limited positive and negative predictive accuracy [5–7], particularly in predicting adverse events beyond 24 hours [8–11].

The use of digital technologies in healthcare presents an opportunity to evaluate whether the inclusion of additional routinely collected variables would improve the predictive accuracy of an early warning score and thus reduce patient risk. The same technologies could then be used to provide clinical decision support using artificial intelligence-derived algorithms that would not be practical using paper observation charts and healthcare records [1,11].

This further personalisation of risk prediction is increasingly important in our ageing population. Older adults are particularly vulnerable to sudden changes in physiological status [13], but this group was not included in the development of NEWS2 [14]. Improved predictive accuracy in this growing frail and vulnerable group could significantly improve patient outcomes [14–16].

In an evaluation of the performance of the original NEWS, it demonstrated consistent predictive accuracy across multiple patient groups [5]. However, the performance of NEWS2 varies between care settings, and between patients with different characteristics [11,17,18]. NEWS2 has been found to have significantly lower specificity, sensitivity, and positive predictive value for those who were at risk of type II respiratory failure (T2RF), but higher specificity and positive predictive value for those with documented T2RF status, then the original NEWS [17]. The accuracy of NEWS has also been found to vary between care settings [18]. These observations suggest that the inclusion of individualised patient and care setting characteristics may improve the predictive accuracy, sensitivity and specificity of the system and thus improve outcomes in a range healthcare settings and in diverse patient groups.

To determine which variables may improve the risk prediction of NEWS2 we have reviewed six databases (CINAHL, PubMed, Embase, ScienceDirect, Cochrane Library and Web of Science) [19]. Our preliminary findings show the demographic variables including age and ethnicity [6,20], trend data [21–23], and adjustments to component weighting [12,24] may improve on the performance of NEWS2 in predicting clinical deterioration.

The principal aim of this work is to develop an algorithm that can improve on the predictive accuracy and thus clinical value of the NEWS2 system. The objectives of this study are:

- 1. To determine which additional variables have been previously used to enhance the accuracy of clinical early warning scores, including NEWS2;
- 2. To collect a large set of patient data that links observations, demographic, and admissions data with key deterioration-related outcomes;

- 3. To use these data to train an artificial intelligence (AI) model to determine which variables, and what weightings, provide optimal predictive accuracy;
- 4. To test the accuracy, sensitivity, and specificity of the algorithm;
- 5. To consider how the new model could be applied across a wide range of healthcare settings, including different physiological observation equipment, different electronic patient record systems, or in paper-based clinical record systems.

2. Methods and Analysis

The study is a retrospective cohort study using approximately 7 years (2017-2023) of historical, anonymised patient data from the Newcastle upon Tyne Hospitals NHS Foundation Trust to explore ways in which a new alerting system could improve on NEWS2. We will recruit PPIE members to join the project steering committee through well-established PPIE processes in the Newcastle BRC Informatics and Precision Care for an Ageing Population theme and Newcastle University's Clinical Trials Unit. PPIE members will be included in all project management group meetings and will provide their lived experience with hospital monitoring to highlight any concerns or barriers they foresee in the adoption and implementation of a modified NEWS system. We will recruit at least one older adult to ensure that we are including a perspective that has been missing in the development of previous NEWS systems.

Data collection: We will use a large number of routinely collected patient variables linked with individual patient outcomes and use these linked data to determine which variables, and with which algorithm, best predicts outcomes. The Newcastle Hospitals NHS Foundation Trust has an established information governance system for the use of anonymised patient data for research, which has been successfully used in the ADMISSION [32] and AI-Multiply [33] studies. A data scientist at the Trust will exclude all data from patients who have opted out of their data being used for research purposes through local or national processes. The data will then be de-identified before the research team can access it within the Trust's IT network for training and testing the algorithm.




Figure 1. Study logic diagram

It is made available under a CC-BY 4.0 International license .

Data analysis: Deidentified data from the Trust's clinical data marts, including patient observations, admissions information, demographic data, and outcomes data, will be divided into training and testing subsets. Using these data sets, we will train and test an algorithm that optimises the variables and their weightings to predict the risk of key clinical outcomes, including mortality, ITU admission, sepsis and cardiac arrest, to demonstrate a proof of concept for a modified scoring system. By the end of this study in April 2025, we will have trained and tested an algorithm based on 7 years of historical patient data. We will then test the new system on data in other NHS Trusts across the UK to assess its generalisability. In 2026, we will conduct implementation studies in Newcastle and then in other NHS Trusts to assess the adoption of the system and gather initial data on patient outcomes to construct a learning healthcare system to evaluate real-world performance of the algorithm and facilitate continual improvement. By 2029, we aim to implement the system in NHS Trusts across the UK and conduct large-scale trials to evaluate its effectiveness at improving patient outcomes.

Intended outputs: The aim of this project is to improve on the predictive accuracy of the NEWS2 scoring system, particularly its accuracy over more than 24-hours, and its predictive value in older patients and children. It will investigate whether using the currently collected data differently and the inclusion of additional data would result in an improved algorithm. There are two main ways in which this could improve patient outcomes: 1) Improving the system's ability to accurately predict patients who will deteriorate within and beyond 24-hours would give healthcare professionals the opportunity to optimise their care and prevent deterioration, and 2) reducing false positive alerts would allow care to be concentrated where it will achieve the greatest benefit.

To be of national and international importance, the tool must meet the needs of patients throughout their life-course. It is clearly important to examine the predictive accuracy of any new early warning system in older people, but there is also limited outcome data for use of such systems in children. On 3 November 2023, NHS England announced the roll-out of a national Paediatric Early Warning System (PEWS) [25], and we will collaborate with the PEWS team in Newcastle Hospitals to develop a new evidence-based clinical alerting system for deteriorating children.

The tool must also be usable in the widest possible range of healthcare settings. As many hospitals worldwide do not have a high degree of digital maturity [26,27] and rely on paper-based documentation [28,29], we aim to develop a scalable paper- or smart-phone application-based system that can achieve the highest possible predictive accuracy and clinical value from an improved algorithm [30].

The overall aim is to incorporate an improved early warning system in an integrated Learning Health System. This requires the technical capacity to routinely collect outcomes data and provide structured opportunities to test and improve the algorithm for a changing population. It also needs robust mechanisms for capturing and using clinical feedback on its implementation, successes, and barriers in a wide range of care contexts [31].

It is made available under a CC-BY 4.0 International license .

3. Ethics and Dissemination

An application for the ethical approval of this study has been submitted to the UK The Health Research Authority Integrated Research Approval System and is under consideration at the time of manuscript submission. The study findings are expected to be produced in April 2025 and will be disseminated symposia, conferences, clinical meetings and academic peer-reviewed publications. PPIE members will help guide the dissemination of the results of the study to the population through publicly accessible forums. Further funding applications will be prepared to continue this programme of work by applying the new algorithm to other Trusts across the UK and other hospital systems around the world, to test its generalisability and refine its performance. Patient data will be stored only on secure NHS Trust systems. Other data available to the research team will be stored securely on Newcastle University systems until the study is complete. Analyses generated by the study will be available upon publication for 10 years from the PI on reasonable request.

4. Conclusion

This work has identified potential areas of improvement to the NEWS2 system and outlined a project for improving on its ability to predict adverse outcomes through the use of additional patient data and/or changes to the scoring process. The project's outputs will be shared widely to facilitate collaboration between healthcare professionals, researchers, and policymakers to support the implementation of improved clinical early warning systems in diverse populations.

It is made available under a CC-BY 4.0 International license .

Conflicts of Interest

The authors have no conflict of interest to report.

Abbreviations

NEWS = National Early Warning Score

Funding

This work was supported by the National Institute for Health and Care Research (NIHR) Newcastle Biomedical Research Centre (BRC) based at the Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle University, and the Cumbria, Northumberland, and Tyne and Wear (CNTW) NHS Foundation Trust. The views expressed in this publication are those of the authors and not necessarily those of the NIHR or BRC, or any of the authors' affiliated universities. The open access publication fee was paid by the Newcastle BRC. The funding body was not involved in the study design, data collection or analysis, or the writing and decision to submit the article for publication.

CRediT author statement

Chris Pummer: Conceptualisation, Funding acquisition, Writing - review & editing
Madison Milne-Ives: Methodology, Writing - Original Draft
Cen Cong: Writing - Original Draft, Writing - Review & Editing
Lynsey Threlfall: Writing - review & editing
Edward Meinert: Conceptualisation, Supervision, Writing - review & editing

It is made available under a CC-BY 4.0 International license .

References

- 1 Holland M, Kellett J. The United Kingdom's National Early Warning Score: should everyone use it? A narrative review. *Intern Emerg Med*. 2023;18:573–83. doi: 10.1007/s11739-022-03189-1
- 2 Finnikin S, Wilke V. What's behind the NEWS? National Early Warning Scores in primary care: National Early Warning Scores in primary care. *Br J Gen Pract*. 2020;70:272–3. doi: 10.3399/bjgp20x709361
- 3 Myrstad M, Ihle-Hansen H, Tveita AA, *et al.* National Early Warning Score 2 (NEWS2) on admission predicts severe disease and in-hospital mortality from Covid-19 - a prospective cohort study. *Scand J Trauma Resusc Emerg Med.* 2020;28:66. doi: 10.1186/s13049-020-00764-3
- 4 Scott LJ, Redmond NM, Garrett J, *et al.* Distributions of the National Early Warning Score (NEWS) across a healthcare system following a large-scale roll-out. *Emerg Med J.* 2019;36:287–92. doi: 10.1136/emermed-2018-208140
- 5 Price C, Prytherch D, Kostakis I, *et al.* Evaluating the performance of the National Early Warning Score in different diagnostic groups. *Resuscitation*. 2023;193:110032. doi: 10.1016/j.resuscitation.2023.110032
- Carr E, Bendayan R, Bean D, et al. Evaluation and improvement of the National Early Warning Score (NEWS2) for COVID-19: a multi-hospital study. BMC Med. 2021;19:23. doi: 10.1186/s12916-020-01893-3
- 7 Williams B. The National Early Warning Score: from concept to NHS implementation. *Clin Med*. 2022;22:499–505. doi: 10.7861/clinmed.2022-news-concept
- 8 Holland M, Kellett J. A systematic review of the discrimination and absolute mortality predicted by the National Early Warning Scores according to different cut-off values and prediction windows. *Eur J Intern Med*. 2022;98:15–26. doi: 10.1016/j.ejim.2021.12.024
- 9 Wei S, Xiong D, Wang J, *et al.* The accuracy of the National Early Warning Score 2 in predicting early death in prehospital and emergency department settings: a systematic review and meta-analysis. *Annals of Translational Medicine*. 2023;11. doi: 10.21037/atm-22-6587
- 10 Mcneill H, Khairat S. Impact of Intensive Care Unit Readmissions on Patient Outcomes and the Evaluation of the National Early Warning Score to Prevent Readmissions: Literature Review. *JMIR Perioperative Medicine*. 2020;3:e13782. doi: 10.2196/13782
- 11 Zhang K, Zhang X, Ding W, *et al.* The Prognostic Accuracy of National Early Warning Score 2 on Predicting Clinical Deterioration for Patients With COVID-19: A Systematic Review and Meta-Analysis. *Front Med.* 2021;8:699880. doi: 10.3389/fmed.2021.699880
- 12 Clarke J, Gallifant J, Grant D, *et al.* Predictive value of the National Early Warning Score 2 for hospitalised patients with viral respiratory illness is improved by the addition of inspired oxygen fraction as a weighted variable. *BMJ Open Respir Res.* 2023;10:e001657. doi:

It is made available under a CC-BY 4.0 International license .

10.1136/bmjresp-2023-001657

- 13 Clegg A, Young J, Iliffe S, *et al.* Frailty in Older People. *Lancet*. 2013;381:752. doi: 10.1016/S0140-6736(12)62167-9
- 14 Vardy ER, Lasserson D, Barker RO, *et al.* NEWS2 and the older person. *Clin Med* . 2022;22:522–4. doi: 10.7861/clinmed.2022-0426
- 15 Kemp K, Alakare J, Harjola V-P, *et al.* National Early Warning Score 2 (NEWS2) and 3-level triage scale as risk predictors in frail older adults in the emergency department. *BMC Emerg Med.* 2020;20:1–9. doi: 10.1186/s12873-020-00379-y
- 16 Sze S, Pan D, Williams CML, *et al.* Letter to the Editor: Variability but not admission or trends in NEWS2 score predicts clinical outcome in elderly hospitalised patients with COVID-19. *J Infect.* 2021;82:159. doi: 10.1016/j.jinf.2020.05.063
- 17 Pimentel MAF, Redfern OC, Gerry S, et al. A comparison of the ability of the National Early Warning Score and the National Early Warning Score 2 to identify patients at risk of in-hospital mortality: A multi-centre database study. *Resuscitation*. 2019;134:147–56. doi: 10.1016/j.resuscitation.2018.09.026
- 18 Zaidi H, Bader-El-Den M, McNicholas J. Using the National Early Warning Score (NEWS/NEWS 2) in different Intensive Care Units (ICUs) to predict the discharge location of patients. *BMC Public Health*. 2019;19:1231. doi: 10.1186/s12889-019-7541-3
- 19 Milne-Ives M, Riccalton V, Plummer C, *et al.* Modifications to the National Early Warning Score: a scoping review protocol. *BMJ Open*. 2024;14:e089061. doi: 10.1136/bmjopen-2024-089061
- 20 Nissen SK, Candel BGJ, Nickel CH, *et al.* The impact of age on predictive performance of National Early Warning Score at arrival to emergency departments: Development and external validation. *Ann Emerg Med.* 2022;79:354–63. doi: 10.1016/j.annemergmed.2021.09.434
- 21 Forster S, McKeever TM, Churpek M, et al. Predicting outcome in acute respiratory admissions using patterns of National Early Warning Scores. *Clin Med.* 2022;22:409–15. doi: 10.7861/clinmed.2022-0074
- 22 Gonem S, Taylor A, Figueredo G, *et al.* Dynamic early warning scores for predicting clinical deterioration in patients with respiratory disease. *Respir Res.* 2022;23:203. doi: 10.1186/s12931-022-02130-6
- 23 Zhu Y, Chiu Y-D, Villar SS, *et al.* Dynamic individual vital sign trajectory early warning score (DyniEWS) versus snapshot national early warning score (NEWS) for predicting postoperative deterioration. *Resuscitation*. 2020;157:176–84. doi: 10.1016/j.resuscitation.2020.10.037
- 24 Malycha J, Farajidavar N, Pimentel MAF, *et al.* The effect of fractional inspired oxygen concentration on early warning score performance: A database analysis. *Resuscitation*. 2019;139:192–9. doi: 10.1016/j.resuscitation.2019.04.002
- 25 England NHS. NHS England » New NHS warning system to spot deterioration in children's health faster.

https://www.england.nhs.uk/2023/11/new-nhs-warning-system-to-spot-deterioration-in-childrens-health-faster/ (accessed 3 November 2023)

- 26 Koebe P, Bohnet-Joschko S. What's next in hospital digitization? A Delphi-based scenario study. *Eur J Futures Res.* 2023;11:1–18. doi: 10.1186/s40309-023-00222-0
- Geissler A, Hollenbach J, Haring M, et al. A nationwide digital maturity assessment of hospitals Results from the German DigitalRadar. *Health Policy Technol*. 2024;13:100904. doi: 10.1016/j.hlpt.2024.100904
- 28 Hufnagl C, Doctor E, Behrens L, *et al.* DIGITISATION ALONG THE PATIENT PATHWAY IN HOSPITALS. Published Online First: 2019.
- 29 Fröhlich D, Bittersohl C, Schroeder K, *et al.* Reliability of paper-based routine documentation in psychiatric inpatient care and recommendations for further improvement. *Front Psychiatry*. 2019;10:954. doi: 10.3389/fpsyt.2019.00954
- 30 Ma M-W, Gao X-S, Zhang Z-Y, *et al.* Extracting laboratory test information from paper-based reports. *BMC Med Inform Decis Mak*. 2023;23:251. doi: 10.1186/s12911-023-02346-6
- 31 Hardie T, Horton T, Thornton-Lee N, *et al.* Developing learning health systems in the UK: Priorities for action. The Health Foundation 2022.
- 32 ADMISSION Collaborative, Newcastle University https://research.ncl.ac.uk/admissioncollab/
- 33 AI-MULTIPLY Newcastle University https://ai-multiply.co.uk/