

1

2

3 **From Traits to Puffs: The Interplay of Personality, Pandemic Stress, and Smoking Behaviors**

4

5 Ying Tian ^{1¶}, Weiyi Xiang^{1¶}, Silvia Dzhugaryan,^{1&} Dayoung Bae,¹³ Jessica Barrington-Trimis,¹³ Terry
6 Church ^{2*}

7

8

9 ¹University of Southern California Keck School of Medicine, Institution of Addiction Science, Los
10 Angeles, CA, USA

11 ²University of Southern California, Alfred E. Mann, School of Pharmacy and Pharmaceutical Sciences,
12 Los Angeles, CA, USA

13 ³University of Southern California, Keck School of Medicine, Department of Population and Public
14 Health Sciences, Los Angeles, CA, USA

15

16 *Corresponding author
17 Email: tdchurch@usc.edu

18 ¶ These authors contributed equally to this work.

19 & These authors also contributed equally to this work.

20

21 **Abstract**

22 Smoking, a leading cause of chronic diseases, is often used to cope with stress, which has been
23 heightened by the pandemic due to health and economic concerns. Studies have shown that the Big Five
24 personality traits are linked to smoking behavior, suggesting that different personality traits influence
25 nicotine use in varying ways. However, there remains a significant gap in understanding how individuals
26 with different personalities respond to nicotine use under stress. This study aims to investigate how
27 nicotine dependence changes for different Big Five personalities under the pandemic stress and whether
28 other stress-related factors influence nicotine dependence during COVID-19. This cross-sectional study
29 collect data from randomly selected adults aged 18-30 in the US. The Big Five Personality Model
30 assessed personality traits, and nicotine dependence was measured with the Hooked-on Nicotine
31 Checklist. Stress was evaluated using the Perceived Stress Scale, while demographics and other
32 pandemic-related stressors were gathered through structured questions. Correlation and multiple logistic
33 regression models were used for data analysis. The main findings showed that both before ($r = -.25$, p
34 $< .001$) and during ($r = -.19$, $p < .001$) the pandemic, agreeableness was significantly negatively
35 associated with nicotine dependence, indicating that higher agreeableness was linked to lower nicotine
36 dependence. Similarly, conscientiousness was negatively correlated with nicotine dependence both before
37 ($r = -.123$, $p < .001$) and during COVID-19 ($r = -.19$, $p < .001$). Although no direct association was found
38 between perceived stress, personality traits, and smoking behavior, the analysis identified that external
39 stressors played a moderating role. These findings emphasize the importance of understanding how
40 different personality traits influence young people's dependence on nicotine under stress. The outcome
41 can guide the design of targeted nicotine withdrawal interventions and inform effective public health
42 strategies.

43

44

45 **Introduction**

46 Tobacco use has been one of the most serious public health problems. Studies have shown that
47 tobacco use can cause cardiovascular damage, respiratory conditions like emphysema and chronic
48 obstructive pulmonary disease (COPD), and mouth cancer (1). By 2020, 22.3% of the global population
49 uses tobacco, and nearly 8 million people die from tobacco use each year (2). However, among the fatal
50 diseases, tobacco-related diseases can be effectively reduced or even artificially avoided, and reducing
51 tobacco use can effectively reduce the global burden of disease (3). Therefore, it is important to
52 understand the motivations and mechanisms underlying smoking behavior and nicotine dependence.

53 Personality traits have been shown to play an important role in smoking behavior, and there are
54 significant differences in smoking behavior among people with different personalities. Studies have found

55 that people with high levels of extraversion and neuroticism are more likely to start smoking. This may be
56 due to their impulsivity, social tendencies, or emotional instability (4, 5). On the other hand, individuals
57 who score high in conscientiousness and agreeableness tend to smoke less. These traits are often linked to
58 better self-control and respect for social norms(4). Openness, however, is associated with higher nicotine
59 dependence. This suggests that people who are more open to experiences may be more willing to
60 experiment with substances like tobacco(5). Different personality traits not only influence the individual's
61 propensity to smoke but may also determine their level of nicotine dependence and ease of quitting. For
62 men, those with lower neuroticism and extraversion and higher openness were more likely to be
63 successful in quitting, while women with lower conscientiousness were more likely to quit (6). These
64 findings suggest that individual personality traits may play a key role in the development and
65 maintenance of smoking behavior.

66 Moreover, stress is also an important factor that influences smoking behavior. People often
67 smoke as a means of coping with negative emotions and to relieve major stressful events in their lives(7).
68 Studies have shown that when smokers' stress levels or negative emotions increase, their daily tobacco
69 consumption, time spent smoking, and frequency of smoking also increase significantly(8-10). This
70 phenomenon may be due to the transient relaxing effect of the nicotine component of cigarettes. By
71 influencing the central nervous system, nicotine controls the production of dopamine, resulting in a
72 transient pleasure experience and a calming effect (11). High levels of stress are also one of the main
73 reasons why smokers relapse. Studies have shown that stress significantly increases the likelihood of
74 relapse by making smokers more inclined to remember the environment associated with smoking and
75 enhancing their memory bias for smoking-related items(12).

76 In the specific context of the COVID-19 pandemic, the complexity of smoking behavior has been
77 further exacerbated by the unprecedented amplification of psychological stress. Surveys have shown that
78 the COVID-19 epidemic has disrupted people's daily, increased anxiety of isolation, unemployment, and
79 health anxiety(13, 14). One study noted that during the COVID-19 pandemic, 28% of participants
80 (n=3396) reported an increase in their smoking, with increased stress, prolonged time spent at home, and

81 boredom cited as the main reasons(15). In addition, 16.8% of young nicotine e-cigarette users in 2020
82 reported that they had increased their smoking frequency due to epidemic-related stressors(16).

83 Although a large number of studies have validated personality traits and stress as a cause of
84 smoking, their interaction has rarely been noted. Thus, the objective of this study is to investigate how
85 stress from the COVID-19 pandemic influenced nicotine dependence based on the Big Five personality
86 traits (e.g., neuroticism, extraversion, openness, agreeableness, and conscientiousness) within nicotine
87 users vs non-nicotine users in young adults. We have three hypotheses: (1) Personality is related to
88 nicotine. (2) people with neuroticism and extraversion personalities show an increase in nicotine
89 dependence, whereas people with conscientious personalities do not show significant changes or
90 decreases in nicotine dependence. (3) Other stress-related factors influence nicotine dependence during
91 COVID-19, such as health concerns, access to nicotine, economic factors, mental health, routine
92 disruption, social networking, and traumatic events.

93 **Methods**

94 This study has been approved and exempted by the University of Southern California Institutional
95 Review Board and participant authorization was obtained in the form of posters and questionnaires (IRB
96 Study ID: UP-23-01234). Our observational cross-sectional survey was conducted from February 2024 to
97 May 13, 2024. Participants aged 18–30, able to provide informed consent, were recruited through flyers at
98 the University of Southern California, social media platforms (Instagram and Snapchat), and Amazon
99 Mechanical Turk (MTurk). Incentives included an Apple AirPods, eight Apple AirTags, and a >5% chance
100 to win a \$10 Amazon gift card. Of the 433 respondents, <30% of data were excluded due to ineligibility
101 or incomplete responses, yielding a final sample of 323 participants. Inclusion criteria required
102 participants to be aged 18–30 and able to complete the survey. Participants who have used any nicotine
103 product in their lifetime answered the following questions measuring sociodemographics, personality
104 (Big Five) if they used any nicotine products, what type of nicotine product, nicotine consumption hooked
105 on nicotine checklist (HONC), other influences of nicotine behavior, and retrospectively reported
106 perceived stress (PSS) levels both before COVID-19 (2018-2019) and during COVID-19 (2020-2021).

107 Non-nicotine users only answered the following questions: measuring, sociodemographics, personality
108 (Big Five), and whether they have ever used any nicotine products. All measurements had the same
109 variables and scoring in conjunction with the following timeline.

110 **Measures**

111 Personality traits were measured using the 44-item Big Five Personality Inventory (Goldberg,
112 1993), which assesses five dimensions: Extraversion, Neuroticism, Agreeableness, Conscientiousness,
113 and Openness. Items were rated on a 5-point Likert scale ranging from 1 = “Strongly Disagree” to 5 =
114 “Strongly Agree.” Some items required reverse coding, and final scores were calculated by summing all
115 items within each dimension, with higher scores indicating stronger traits. This scale has demonstrated
116 high reliability (Cronbach’s $\alpha = 0.70\text{--}0.90$).

117 Nicotine dependence was assessed using the Hooked on Nicotine Checklist (HONC), a 10-item
118 tool designed to evaluate the loss of autonomy over nicotine use. Responses were scored as 1 = “Yes” and
119 0 = “No,” with total scores ranging from 0 to 10. Dependence levels were categorized as follows: 0 = No
120 dependence, 1–2 = Low dependence, 3–5 = Moderate dependence, 6–8 = High dependence, and 9–10 =
121 Severe dependence. Participants who reported lifetime nicotine use (yes/no) were further queried about
122 specific products used, including e-cigarettes, cigars, hookah, smokeless tobacco, non-combustibles, and
123 dissolvables. Only nicotine users completed the HONC and related measures of nicotine behaviors.

124 Stress levels were measured using the Perceived Stress Scale (PSS), a 10-item questionnaire
125 assessing perceived stress in the past month. Items were rated on a 5-point Likert scale from 0 = “Never”
126 to 4 = “Very Often.” Scores were calculated by summing all item responses, with higher scores indicating
127 greater stress. Several items required reverse coding before summing. The PSS has demonstrated high
128 reliability (Cronbach’s $\alpha = 0.70\text{--}0.90$).

129 To further explore the relationship between stress regulation, personality, and nicotine use
130 behavior, an additional stress-related factors questionnaire was administered. This measure assessed the

131 influence of factors such as stress relief, relaxation, boredom, social situations, and academic pressure on
132 nicotine use. Participants rated these factors on a 5-point Likert scale ranging from 1 = “Not at all” to 5 =
133 “Extremely,” with higher scores indicating a stronger influence. This questionnaire, completed only by
134 nicotine users, aimed to identify specific stressors contributing to nicotine consumption and demonstrated
135 high internal consistency (Cronbach’s $\alpha = 0.92$). Non-nicotine users were excluded from this assessment.

136 Demographic information, including age, gender, race/ethnicity, financial status, and health
137 conditions, was collected to contextualize findings and assess sociodemographic influences on nicotine
138 use. Participants provided responses regarding their age, assigned sex at birth, gender identity,
139 race/ethnicity, and personal financial situation, alongside self-reported health conditions diagnosed by a
140 healthcare provider. Together, these measures provided a detailed understanding of how stress influences
141 the relationship between personality and nicotine use.

142 **Data Analyze**

143 This study used SPSS for descriptive statistics, correlations, and multiple regression models to
144 ensure the accuracy of experimental data analysis and examine relationships between key variables.

145 First, descriptive statistics were used to summarize and describe participants' demographic
146 characteristics, health status, and economic status to provide a clear base profile for subsequent analyses.

147 The 'Crosstabs' function was used in SPSS to compare nicotine users and non-users, generating
148 contingency tables that provide an overview of participant diversity.

149 Subsequently, correlations were conducted to test the hypothesis of the moderate effect of
150 pandemic stress on the relationship between personality traits and nicotine dependence. In SPSS, the
151 ‘Pearson correlation’ function assessed the strength and direction of associations between variables,
152 providing a reliable measure of their linear relationships.

153 Moreover, multiple logistic regression analyses were performed using the built-in "Logistic
154 Regression" tool in SPSS to effectively assess the pathways of nicotine dependence across personality
155 traits and pandemic-related stress. This approach explores the moderate effects of COVID-related
156 concerns between personality traits and nicotine dependences.

157 **Results**

158 **Descriptive results, by Nicotine Use Status**

159
160 Demographic data were collected on age, sex at birth, race/ethnicity, sexual orientation, gender
161 identity, diagnosed health diseases, and financial income as "Table 1". The study includes 324
162 participants, with 269 nicotine users and 54 non-nicotine users. The average age of all participants is
163 26.10 years; nicotine users averaged 26.78 years, while non-nicotine users averaged 22.69 years. There
164 are more males and fewer females among the nicotine users (57.4% male and 33.1% female) than the
165 non-nicotine users (33.3% male and 66.9% female). Nicotine users have a lower percentage of
166 Heterosexual/Straight individuals (63.6%), and a higher percentage of Bisexual individuals (30.5%)
167 compared to non-nicotine users (81.5% and 11.1%, respectively). In terms of health diagnoses, there are,
168 more nicotine users were diagnosed with a disease (66.5%) compared to non-nicotine users (16.7%),
169 including 28.6% of nicotine users with a physical disability. Financially, more nicotine users lived
170 comfortably (54.6%) compared to non-nicotine users (42.6%). A higher percentage of non-nicotine users
171 met their needs with little left over (33.3% vs. 26.0%) or just met basic expenses (20.4% vs. 19.0%).

172 As shown in "Table 2", nicotine users had higher extraversion scores (25.73 vs. 23.96, $p < .05$)
173 and lower agreeableness (29.23 vs. 32.85, $p < .001$) and conscientiousness scores (29.03 vs. 31.25, p
174 $< .001$) compared to non-nicotine users. For nicotine users, we also assessed HONC and PSS scores
175 before and during COVID-19. The average HONC score slightly decreased during COVID-19 (4.71, SD
176 = 4.00) compared to before (4.85, SD = 3.96). PSS scores were stable, with an average of 26.77 (SD =
177 3.50) during COVID-19 and 26.62 (SD = 3.63) before COVID-19.

178 **Association of Personality with Nicotine Dependence in Nicotine product users**

179 As “Table 3” indicates, for all nicotine users, agreeableness was negatively associated with
180 nicotine dependence both before ($r = -.25, p < 0.001$) and during COVID-19 ($r = -.19, p < 0.001$).
181 Conscientiousness was also negatively associated before COVID-19 ($r = -.123, p < 0.001$). To examine
182 the relationship between personality traits and nicotine dependence under COVID-19 stress, participants
183 were categorized into low and high-stress groups based on their PSS scores relative to the mean. Those
184 scoring above the mean were placed in the high-stress group, while those below were in the low-stress
185 group. Before COVID-19, agreeableness was negatively associated with nicotine dependence in the low-
186 stress group ($r = -.19, p < 0.05$). In the high-stress group, extraversion showed a positive association with
187 nicotine dependence ($r = .23, p < 0.05$), and agreeableness a negative one ($r = -.25, p < 0.05$). During
188 COVID-19, no significant correlations between personality traits and nicotine dependence were observed.

189 “Table 4” used three regression models to examine the relationships between personality traits
190 and nicotine dependence, personality traits and COVID-19 concerns, and COVID-19 concerns and
191 nicotine dependence. Model one is to test the regression between personalities and nicotine dependence.
192 Among the five personality traits, agreeableness was significantly negatively associated with nicotine
193 dependence ($B = -.170, p < .05$), indicating that higher agreeableness is linked to lower nicotine
194 dependence.

195 In model two, which tests the regression between personalities and COVID-19 concerns,
196 agreeableness was consistently negatively associated with various COVID-19-related concerns, including
197 health ($B = -.020, p < .001$), economic ($B = -.034, p < .001$), mental health stress ($B = -.018, p < .05$),
198 routine disruption ($B = -.023, p < .001$), social network ($B = -.017, p < .05$), and traumatic events ($B =$
199 $-.021, p < .05$). Extraversion was positively linked to concerns about access to nicotine ($B = .019, p$
200 $< .05$), while neuroticism was negatively associated with health ($B = -.019, p < .001$) and economic

201 concerns ($B = -.016, p < .05$). These findings underscore the significant roles of agreeableness,
202 extraversion, and neuroticism in shaping responses to COVID-19-related concerns.

203 The regression between COVID-19 concerns and nicotine dependence was tested in model three.
204 The regression analysis identified significant predictors of nicotine dependence during COVID-19.
205 Access to nicotine was positively associated with increased dependence ($B = 2.666, p < .05$), as were
206 routine disruption ($B = 2.571, p < .05$) and traumatic events ($B = 2.139, p < .05$). These findings suggest
207 that these stress factors contributed to higher nicotine dependence during the pandemic. Other factors,
208 such as health concerns ($B = -.469, p = .557$), economic factors ($B = -1.663, p = .076$), mental health
209 stress ($B = 1.204, p = .148$), and social network concerns ($B = -.771, p = .414$), were not significant
210 predictors of nicotine dependence.

211 **Discussion**

212 This study revealed significant correlations between personality traits, stress, and nicotine
213 dependence levels among young adults. Notably, extraversion, conscientiousness, and agreeableness
214 showed significant differences between nicotine users and non-users. Moreover, specific stress factors
215 related to the COVID-19 pandemic, such as access to nicotine, routine disruption, and traumatic events,
216 were potential predictors of increased nicotine dependence. The demographic characteristics of our
217 sample provide us with critical insights into the relationship between personal traits and smoking
218 behavior. The first significance we observed is gender difference, with a higher percentage of males
219 among the nicotine users, suggesting that males may have different stress-coping strategies compared to
220 females, possibly leading to higher smoking prevalence. Another notable finding related to the health of
221 the participants was that most nicotine users were diagnosed with a medical condition, with 69.5%
222 diagnosed with a disease, compared to 16.7 % of non-nicotine users; suggesting that individuals with
223 physical health risks are more likely to be associated with smoking behavior. Additionally, the younger
224 average age of non-nicotine users might reflect a generational shift in smoking behaviors due to targeted

225 public health campaigns. Our findings are consistent with previous studies that have reported higher rates
226 of smoking among men and older age groups “Table 1-2”. Also, we found that a higher percentage of
227 nicotine users were bisexual compared to non-nicotine users, a novel finding that warrants further study.
228 An explanation for this is that this population may face unique stressors or social stigma that led to higher
229 dependence on nicotine, which is consistent with the findings on the effects of minority stress on
230 substance use “Table 3”.

231 Compared to non-nicotine users, active users showed higher extraversion scores and lower
232 agreeableness and conscientiousness scores. The positive correlation between extraversion and smoking
233 behavior is consistent with previous research suggesting that extroverts are more likely to engage in social
234 smoking behaviors. At the same time, lower levels of agreeableness and conscientiousness among
235 nicotine users suggest these group populations are more likely to use nicotine products, potentially due to
236 a lack of self-regulation and empathy for others.

237 Regression analyses provided greater insight into the relationships among personality traits,
238 COVID-19 worry, and nicotine dependence. Among the five personality traits, agreeableness was
239 significantly negatively correlated with nicotine dependence ($B = -.170, p < .05$), indicating that higher
240 agreeableness was associated with lower nicotine dependence. This is consistent with the negative
241 correlation between agreeableness and nicotine dependence observed before and during COVID-19 ($r =$
242 $-.25, p < .001$; $r = -.19, p < .001$) in “Table 4”. The mechanisms underlying the negative association
243 between agreeableness and nicotine dependence may be manifold. Easy-going people tend to be more
244 cooperative, empathetic, and willing to compromise for the benefit of others. These characteristics may
245 make them less likely to engage in behaviors that are considered harmful or socially undesirable, such as
246 smoking. At the same time, agreeableness tends to be associated with higher levels of social support and
247 better stress-coping strategies. During times of stress, such as the COVID-19 pandemic, easygoing
248 individuals may be better at seeking and utilizing social support, thus reducing the likelihood that they
249 will turn to nicotine as a coping mechanism. This hypothesis is supported by consistent negative
250 correlations between agreeableness and nicotine dependence before and during the pandemic.

251 Contrary to our preconceived hypothesis, we did not find an effect on smoking behavior across
252 personalities under the influence of COVID-19 stress from the Perceived Stress Scale. In this case, our
253 study examined the impact of a more comprehensive set of stressors on smoking behavior and found that
254 agreeableness was negatively associated with all seven of our redesigned stressors, suggesting that
255 individuals with higher levels of these traits reported fewer health worries during the epidemic. In
256 addition, neuroticism was negatively associated with health and financial concerns as well, suggesting
257 that these personality traits may alleviate some of the stress associated with the epidemic. However, we
258 also found that extraversion was positively associated with concerns related to access to nicotine, which
259 indicates extroverts felt more pressure during the epidemic from social interactions and restricted access
260 to nicotine products. These results suggest that specific personality traits lead to altered nicotine
261 dependence and that stress plays a moderating role.

262 There are several limitations to this study. First, the cross-sectional design and the sample
263 selection limited the ability to infer causal relationships. Second, the reliance on self-reported data may be
264 subject to recall bias and introduce social desirability bias. In addition, the sample was limited to young
265 adults aged 18-30 years, and some specific racial groups are heavily represented, limiting the findings'
266 generalizability to other age groups. Nonetheless, Although the Big Five Inventory is valid over time,
267 personality is a dynamic trait. The scale can only reflect an individual's behavioral trends to a certain
268 extent, and it has limited explanatory power for the complexity of specific behaviors. Finally, a slight
269 adjustment of the time frame of the scale to investigate the topic of stress during the COVID-19 pandemic
270 may have caused our study to lose certain validity.

271 **Conclusion**

272 In conclusion, this study highlights the significant role that personality traits and stress play in
273 influencing young people's dependence on nicotine. Contrary to our assumptions, however, we did not
274 find an effect of stress on the correlation between personality and nicotine dependence within the
275 pandemic context. Regardless of our initial hypothesis, these findings provide valuable insights for the

276 development of targeted smoking cessation interventions and contribute to a broader understanding of
277 smoking behavior within the context of personality psychology. Addressing specific stressors and
278 demographic factors could improve the effectiveness of public health strategies aimed at reducing
279 nicotine dependence.

280 Future studies should consider longitudinal designs to better understand the causal relationship
281 between personality traits, stress, and nicotine dependence. Also adopting a randomized sampling
282 approach involving samples from different age groups and cultural backgrounds would enhance the
283 generalizability of the findings. Meanwhile, the use of some bioindicator measures can further enhance
284 the accuracy of the results, such as utilizing the cotinine test. In addition, exploring the relationship
285 between personality and other substance dependencies can further the understanding of the critical nature
286 of personal traits in addiction.

287

288 **References**

- 289 1. Bartal M. Health effects of tobacco use and exposure. *Monaldi Arch Chest Dis.* 2001;56(6):545-
290 54.
- 291 2. Organization WH. World Health Organization; 2023 [Available from: [https://www.who.int/news-](https://www.who.int/news-room/fact-sheets/detail/tobacco)
292 [room/fact-sheets/detail/tobacco](https://www.who.int/news-room/fact-sheets/detail/tobacco).
- 293 3. Kelly EV, Grummitt L, Teesson M, Newton NC. Associations between personality and uptake of
294 tobacco smoking: Do they differ across adolescence? *Drug Alcohol Rev.* 2019;38(7):818-22.
- 295 4. Hakulinen C, Hintsanen M, Munafò MR, Virtanen M, Kivimäki M, Batty GD, et al. Personality
296 and smoking: individual-participant meta-analysis of nine cohort studies. *Addiction.* 2015;110(11):1844-
297 52.
- 298 5. Munafò MR, Black S. Personality and smoking status: a longitudinal analysis. *Nicotine Tob Res.*
299 2007;9(3):397-404.
- 300 6. Jin W, Xian B, Zhao L, Li C. Association between personality traits and smoking cessation
301 among Chinese adults. *BMC Psychol.* 2023;11(1):398.
- 302 7. Todd M. Daily processes in stress and smoking: effects of negative events, nicotine dependence,
303 and gender. *Psychol Addict Behav.* 2004;18(1):31-9.
- 304 8. Buhelt LP, Pisinger C, Andreasen AH. Smoking and stress in the general population in Denmark.
305 *Tob Prev Cessat.* 2021;7:27.
- 306 9. Wang Y, Chen X, Gong J, Yan Y. Relationships Between Stress, Negative Emotions, Resilience,
307 and Smoking: Testing a Moderated Mediation Model. *Subst Use Misuse.* 2016;51(4):427-38.
- 308 10. Conklin CA, Perkins KA. Subjective and reinforcing effects of smoking during negative mood
309 induction. *J Abnorm Psychol.* 2005;114(1):153-64.
- 310 11. Rowell PP. *Current Concepts on the Effects of Nicotine on Neurotransmitter Release in the*
311 *Central Nervous System.* New York: Plenum Press; 1987.
- 312 12. Lee JH, Kang S, Maier SU, Lee SA, Goldfarb EV, Ahn WY. Acute Stress Enhances Memory and
313 Preference for Smoking-Related Associations in Smokers. *Nicotine Tob Res.* 2024;26(3):333-41.
- 314 13. Pfeifer LS, Heyers K, Ocklenburg S, Wolf OT. Stress research during the COVID-19 pandemic
315 and beyond. *Neurosci Biobehav Rev.* 2021;131:581-96.
- 316 14. Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, et al. Impact of COVID-19 pandemic on
317 mental health in the general population: A systematic review. *J Affect Disord.* 2020;277:55-64.
- 318 15. Yingst JM, Krebs NM, Bordner CR, Hobkirk AL, Allen SI, Foulds J. Tobacco Use Changes and
319 Perceived Health Risks among Current Tobacco Users during the COVID-19 Pandemic. *Int J Environ*
320 *Res Public Health.* 2021;18(4).
- 321 16. Parks MJ, Fleischer NL, Patrick ME. Increased nicotine vaping due to the COVID-19 pandemic
322 among US young adults: Associations with nicotine dependence, vaping frequency, and reasons for use.
323 *Prev Med.* 2022;159:107059.
- 324

Table 1. Descriptive statistics for participants, demographic information, and the P (chi-square or t-test) value between nicotine users (n =269) and non-nicotine users (n = 54).

Variable	Nicotine users	Non-nicotine users	P (chi-square or t-test)
Mean Age	26.78	22.69	< .001**
Sex at birth			< .001**
Male	170 (57.4%)	18 (33.3%)	
Female	98 (33.1%)	36 (66.9%)	
Intersex	1 (0.3%)	0 (0.0%)	
Race/ethnicity			< .001**
Asian or Pacific Islander	47 (17.5%)	40 (74.1%)	
Black or African American	6 (2.2%)	1 (1.9%)	
Hispanic or Latino	8 (3.0%)	6 (11.1%)	
Native American/Alaskan Native	9 (3.3%)	0 (0.0%)	
White or Caucasian	192 (71.4%)	6 (11.1%)	
Middle Eastern	5 (1.9%)	0 (0.0%)	
Another (not listed)	2 (0.7%)	1 (1.9%)	
Sexual identity			< .05*
Heterosexual/Straight	171 (63.6%)	44 (81.5%)	
Homosexual	13 (4.8%)	3 (5.6%)	
Bisexual	82 (30.5%)	6 (11.1%)	
Pansexual	1 (0.4%)	0 (0.0%)	
Asexual	1 (0.4%)	0 (0.0%)	
Another (not listed)	0 (0.0%)	0 (0.0%)	
Prefer not to say	1 (0.4%)	1 (1.9%)	
Diagnosed health disease			< .001**
Chronic Mental Illness	56 (20.8%)	5 (9.3%)	
Development Disability	49 (18.2%)	0 (0.0%)	
Physical Disability	77 (28.6%)	1 (1.9%)	
Neurological Disability	35 (13.0%)	0 (0.0%)	
Learning Disability	35 (13.0%)	1 (1.9%)	
Communication Disability	29 (10.8%)	0 (0.0%)	
Autism Disorder	24 (7.4%)	0 (0.0%)	
N/A	90 (33.5%)	45 (83.3%)	
Another (not listed)	6 (2.2%)	2 (3.7%)	
Prefer not to say	9 (3.3%)	1 (1.9%)	
Financial Situation			< .05*
Live comfortably	147 (54.6%)	23 (42.6%)	
Meet needs with a little left	70 (26.0%)	18 (33.3%)	
Just meet basic expenses	51 (19.0%)	11 (20.4%)	
Don't meet basic expenses	1 (0.4%)	2 (3.7%)	

p < .05*, p < .001**

medRxiv preprint doi: <https://doi.org/10.1101/2024.12.11.24318860>; this version posted December 12, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Table 2. Relationship between personality traits, and stress levels amongst nicotine users vs. nonusers during COVID-19.

Variable	Nicotine users	Non-nicotine users	P (chi-square or t-test)
Personality			
Extraversion	25.73	23.96	< .05*
Agreeableness	29.23	32.85	< .001**
Conscientiousness	29.03	31.25	< .001**
Neuroticism	24.82	25.33	.417
Openness	35.10	34.83	.700
HONC (before COVID-19)	4.85 (sd = 3.96)	N/A	N/A
HONC (during COVID-19)	4.71 (sd = 4.00)	N/A	N/A
PSS (before COVID-19)	26.62 (sd = 3.63)	N/A	N/A
PSS (during COVID-19)	26.77 (sd = 3.50)	N/A	N/A

medRxiv preprint doi: <https://doi.org/10.1101/2024.12.14.24318860>; this version posted December 14, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Nicotine users (n = 269) and non-nicotine users (n = 54). The *p*-values indicate the significance level based on chi-square tests or t-tests comparing the groups of nicotine users and non-nicotine users.

Table 3. The correlation coefficients (r) and significance levels (p) between Big Five Personalities and HONC scores under three conditions: for all nicotine users (n = 269), high-stress, and low-stress groups. The stress groups are determined based on PSS scores compared to the average PSS scores before and during COVID-19.

Personalities	All Nicotine Users (r, p)	High-Stress Group (r, p)	Low-Stress Group (r, p)
Before COVID-19			
Extraversion	.096, .117	.228, < .05*	.050, .585
Agreeableness	-.245, < .001**	-.248, < .05*	-.191, < .05*
Conscientiousness	-.123, < .001**	-.031, .722	-.136, .131
Neuroticism	-.002, .974	-.123, .152	.017, .852
Openness	.005, .939	.124, .149	-.141, .119
During COVID-19			
Extraversion	.032, .605	.150, .073	.030, .757
Agreeableness	-.190, < .001**	-.039, .645	-.161, .097
Conscientiousness	-.114, .061	-.102, .223	.014, .882
Neuroticism	.034, .573	-.031, .709	.027, .784
Openness	-.010, .873	.052, .532	-.005, .958

Independent variables: Big Five Personality traits. Dependent variable: HONC scores.

Sample sizes: High-stress (n = 137) and low-stress (n = 124) groups before COVID-19. High-stress (n = 107) and low-stress (n = 144) groups during COVID-19.

p < .05*, *p* < .001**

Table 4. Results of linear regression models.

Model	Predictor/Dependent Variable	B	S.E.	Beta	t	Sig.
1 ^a	HONC During Covid					
	Constant	10.689	3.401	N/A	3.196	< .05*
	Extraversion	.083	.071	.082	1.166	.244
	Agreeableness	-.170	.064	-.186	-2.670	< .05*
	Conscientiousness	-.062	.066	-.068	-.945	.345
	Openness	.010	.053	.013	.189	.851
	Neuroticism	-.021	.062	-.023	-.343	.732
2 ^b	During COVID-19 Health Concerns					
	Constant	1.686	.319	N/A	5.287	< .001**
	Extraversion	.008	.007	.084	1.224	.222
	Agreeableness	-.020	.006	-.218	-3.213	< .05*
	Conscientiousness	-.008	.006	-.094	-1.357	.176
	Neuroticism	-.019	.006	-.218	-3.269	< .05*
	Openness	.008	.005	.104	1.548	.123
	During COVID-19 access to Nicotine Concerns					
	Constant	.959	.300	N/A	3.192	< .05*
	Extraversion	.019	.006	.204	3.053	< .05*
	Agreeableness	-.030	.006	-.347	-5.263	< .001**
	Conscientiousness	-.005	.006	-.054	-.798	.425
	Neuroticism	-.003	.005	-.034	-.532	.595
	Openness	.007	.005	.107	1.636	.103
	During COVID-19 economic Factors Concerns					
	Constant	1.563	.348	N/A	4.489	< .001**
	Extraversion	.012	.007	.110	1.618	.107
	Agreeableness	-.034	.007	-.335	-5.019	< .001**
	Conscientiousness	-.009	.007	-.091	1.330	.185
	Neuroticism	-.016	.006	-.170	-2.596	<.05*
	Openness	.003	.005	.039	.583	.560
	During Covid Mental Health Stress Concerns					
	Constant	1.632	.315	N/A	5.182	< .001**
	Extraversion	.007	.007	.075	1.076	.283
	Agreeableness	-.018	.006	-.209	-3.024	< .05*
	Conscientiousness	-.007	.006	-.087	-1.224	.222
	Neuroticism	-.008	.006	-.093	-1.371	.172
	Openness	.000	.005	.006	.091	.928
	During COVID-19 Routine Disruption Concerns					
	Constant	1.366	.342	N/A	3.997	< .001**
	Extraversion	.013	.007	.129	1.855	.065
	Agreeableness	-.023	.007	-.245	-3.570	< .001**

medRxiv preprint doi: <https://doi.org/10.1101/2024.12.11.24318860>; this version posted December 12, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

	Conscientiousness	-.003	.007	-.029	-.410	.682
	Neuroticism	-.003	.006	-.030	-.440	.660
	Openness	.000	.005	.005	-.069	.945
	During Covid Social Network Concerns					
	Constant	1.431	.289	N/A	4.948	< .001**
	Extraversion	.005	.006	.060	.857	.392
	Agreeableness	-.017	.006	-.215	-3.087	< .05*
	Conscientiousness	-.002	.006	-.024	-.331	.741
	Neuroticism	-.006	.005	-.075	-1.092	.276
	Openness	.001	.004	.018	-.260	.795
	During COVID-19 Traumatic Event Concerns					
	Constant	1.520	.349	N/A	4.349	< .001**
	Extraversion	.014	.007	.128	1.833	.068
	Agreeableness	-.021	.007	-.212	-3.098	< .05*
	Conscientiousness	-.010	.007	-.104	-1.482	.140
	Neuroticism	-.010	.006	-.103	-1.536	.126
	Openness	.004	.005	.046	.675	.500
3^c	HONC During COVID-19					
	Constant	1.383	.713	N/A	1.941	.053
	Health concerns	-.469	.797	-.046	-.588	.557
	Economic Factors Concerns	2.666	.839	.252	3.180	< .05*
	Mental Health Stress Concerns	1.633	.934	-.180	-1.781	.076
	Routine Disruption Concerns	2.571	1.035	.265	2.485	< .05*
	Social Network Concerns	-.771	.942	-.066	-.818	.414
	Traumatic Event Concerns	2.139	.942	.226	2.272	.024

medRxiv preprint doi: <https://doi.org/10.1101/2024.12.11.24318860>; this version posted December 12, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

^aModel 1 is to test the regression between personalities and nicotine dependence.

^bModel 2 is to test the regression between personalities and COVID concerns.

^cModel 3 is to test the regression between Covid concerns and nicotine dependence.

B: Unstandardized regression coefficient. *S.E.*: Standard error of the coefficient. *Beta*: Standardized regression coefficient. *t*: t-value for the hypothesis test of whether the coefficient is different from zero.

Sig.: Significance level of the test. *N/A*: Not applicable for the constant in the regression model.

n = 269 for Model 1, *n* = 254 for Model 2 and Model 3 $p < .05^*$, $p < .001^{**}$.