

1 *Analyzing the determinants for using health research evidence* 2 *in health planning in Tanzania: a cross-sectional study.*

3 Pius Kagoma*^{1&2}, Richard Mongi¹, Albino Kalolo^{3,4}

4 ¹*Department of Public Health and Community Nursing, the University of Dodoma, Dodoma,*
5 *Tanzania;*

6 ²*Division of Health, Social Welfare and Nutrition Services, President's Office-Regional*
7 *Administration Local Government, Dodoma, Tanzania;*

8 ³*Department of Public Health, St. Francis University College of Health and Allied Sciences,*
9 *Morogoro, Tanzania;*

10 ⁴*Centre for Reforms, Innovation, Health Policies, and Implementation Research,*
11 *Dodoma, Tanzania.*

12 *Corresponding author: Pius Kagoma; +255 (0) 76 5909942; P.O. Box 1923 Dodoma, Tanzania;
13 piuskagoma@gmail.com

14 Richard Mongi; +255 (0) 67 6953510; P.O. Box 259, Dodoma, Tanzania;
15 richiemongi@gmail.com

16 Albino Kalolo; +255 (0) 764340819; P.O. Box 175, Ifakara, Morogoro, Tanzania;
17 kaloloa@gmail.com

18 **Abstract**

19 **Introduction:** Achieving Universal Health Coverage (UHC) requires utilizing research
20 evidence to inform the decision-making process. However, little information is available on the
21 determinants for using research evidence in planning in Lower Middle-Income Countries
22 (LMICs), including Tanzania. This paper aims to investigate the determinants of using health
23 research evidence in health planning in Tanzania.

24
25 **Materials and methods:** This study employed a cross-sectional study design. Data on health
26 research evidence and its determinants were collected using a structured questionnaire from 422
27 respondents from 9 regions of Tanzania from October to December 2023. The data were

28 analyzed using STATA version 18 for descriptive and inferential statistics. The association
29 between variables was determined using a chi-square test at a 95% confidence level.

30 **Results:** The study revealed that 66.2% of participants strongly agreed to use health research
31 evidence during planning. However, significant barriers were identified, including lack of
32 dissemination (74.5%), inadequate human and non-human resources (70.0%), and insufficient
33 knowledge and training in research (63.7%). A chi-square test confirmed significant associations
34 between these barriers and the reduced use of research evidence ($p<0.05$). Conversely, more than
35 70% of respondents identified opportunities such as the availability of research coordinators,
36 university partnerships, available research budgets, and internet access, all significantly
37 associated with increased health research evidence use. More than 50% of participants reported
38 motivational factors that like continuous quality improvement agenda in the healthcare sector,
39 availability of short and long-term courses, on-the-job training, and provision of incentives like
40 extra duty allowances were positively linked to research utilization.

41 **Conclusion:** The study found that 66.2% of participants used health research evidence in
42 planning, but barriers like lack of dissemination, resource shortages, and inadequate training
43 persisted. Interventions should focus on improving dissemination, resources, and training. Future
44 research should explore strategies for enhancing these interventions.

45 **Keywords:** Determinants, Research evidence use, Health planning, Tanzania.
46
47

48 INTRODUCTION

49 Health research evidence plays an essential role in shaping effective health policies and
50 interventions, which are crucial for improving public health outcomes (1). In resource-limited
51 settings like Tanzania, where health challenges are complex and resources are constrained, the
52 strategic use of research evidence is particularly important. It can guide decision-makers in
53 identifying priority areas, optimizing resource allocation, and implementing interventions that
54 are both cost-effective and impactful. Moreover, the use of evidence can help health planning
55 teams avoid repeating past failures and introduce innovative solutions (2). Despite its recognized

56 importance, the extent to which health research evidence is utilized by health planning teams in
57 Tanzania remains unclear. Understanding the factors that influence this utilization is essential for
58 promoting evidence-based decision-making, which is critical for achieving the ambitious health
59 targets set forth in the Sustainable Development Goals (SDGs), particularly Universal Health
60 Coverage (UHC)(3).

61 Efforts to promote the integration of research evidence into health planning and decision-making
62 have been made in various settings. These include the establishment of knowledge translation
63 platforms, research dissemination strategies, health conferences, and capacity-building initiatives
64 targeting health planning teams. However, the success of these efforts varies significantly across
65 contexts, largely depending on factors such as institutional support, the capacity of the health
66 workforce, and access to reliable data. In Tanzania, while some initiatives aim to bridge the gap
67 between research and practice, the actual impact of these efforts on health planning remains
68 under-explored.

69 Research across different settings has demonstrated that the use of health research evidence in
70 decision-making is influenced by a variety of factors. These include organizational culture, the
71 availability of resources, the expertise of the planning teams, and access to research outputs [4].
72 In a study conducted by Kagoma et al (4), it was found that health planning teams in Tanzania
73 consist of professionals from various disciplines, including Administrators (9%), Medical
74 doctors (25.36%), Nurses (23.46%), Laboratory scientists (8.53%), Pharmacists (6.64%),
75 Radiographers (1.66%), Environmental health officers (3.08%), Nutrition officers (3.08%),
76 Social welfare officers (4.74%), and others (14.46%). Their primary responsibility is to plan,
77 implement, and evaluate health interventions [4]. However, several factors may limit their ability
78 to fully integrate research evidence into their work (5). These include inadequate dissemination
79 of research findings, lack of access to research evidence, limited skills in interpreting research
80 data, and insufficient organizational support (3,6–8).

81 While previous studies have explored various factors influencing the use of health research
82 evidence in policy and planning, there remains a gap in understanding the specific determinants
83 that affect the utilization of health research evidence among health planning teams in
84 Tanzania(4,9). This study seeks to fill this gap by analyzing the key factors that influence

85 evidence use among these teams at both the regional and council levels, guided by the COM-B
86 Model (Capability, Opportunity, and Motivation) as the theoretical framework (10), which has
87 been widely used in understanding evidence-based decision-making processes. The COM-B
88 model is particularly suited to the Tanzanian context because it focuses on the essential
89 components of behavior change Capability, Opportunity, and Motivation which are critical for
90 understanding and addressing the factors influencing the use of health research evidence by
91 planning teams. By examining these elements, the model helps identify specific barriers and
92 enablers that can be targeted to improve evidence-based decision-making in Tanzania's resource-
93 limited. By applying this theoretical lens, the study aims to provide a deeper understanding of the
94 barriers and facilitators to using research evidence in health planning, with the following specific
95 objectives: 1) Analyze the current usage of health research evidence among planning team
96 members at the regional and council levels, 2) Analyze the capacity of health planning team
97 members to utilize health research evidence, 3) Identify opportunities for enhancing the use of
98 health research evidence in health planning, 4) Identify opportunities for enhancing the use of
99 health research evidence in health planning, 5) Explore the motivations for using health research
100 evidence among health planning members at both regional and council levels. The findings of
101 this study are expected to contribute to the development of strategies that enhance the integration
102 of research evidence into health planning processes, ultimately leading to more effective health
103 interventions and improved health outcomes in Tanzania.

104 **Conceptual framework**

105 The conceptual framework for this study was developed after several consultative meetings with
106 different stakeholders and researchers. The theory was adopted and modified from the
107 (Capability, Opportunity, and Motivation) COM-B model. The COM- B Model at the center of a
108 proposed framework is a behavior system involving three essential conditions: Capability,
109 Opportunity, and Motivation, what we term the 'COM-B system(10,11). This study investigated
110 how the capability, opportunity, and motivation determinants can influence the behavior of the
111 use of health research evidence during health planning as shown in Figure 1

112

113

114

115

Insert figure 1

116

117

Figure 1: The domains of the COM-B Model modified from Michie et al.,2011(10)

118

Materials and Methods

119

Study setting

120

This study was conducted between October and December 2023 in the United Republic of Tanzania, a Lower and Middle low-income country located in East Africa, with a population of about 62 million people. The allocated budget for the country’s Ministry of Health in 2023/2024 was estimated to be 443.6 million US dollars, with 1.2 million US dollars (0.27%) allocated for evidence production. The health research evidence users in Tanzania involve three important Ministries, namely; the Ministry of Health (MoH), The President’s Office Regional Administration and Local Government (PO-RALG), and Health management. The health evidence producers in Tanzania are the National Institute for Medical Research (NIMR), the Tanzania Commission for Science and Technology (COSTECH), Public and Private Universities, Health related institutions or authorities, local and international non-governmental organizations (NGOs), and civil society organizations as shown in Table 1.

131

132

Table 1: A list of health research evidence producers and users in Tanzania

133

134

SN	Evidence users
1	Ministry of Health
2	President’s Office Regional Administration and Local Government
3	Ministry of Education Science and Technology (MEST)
4	Prime Minister’s Office
5	Regional Health Management Teams

6	Council Health Management Teams
7	Healthcare workers
8	Non-Governmental Organizations (NGOs)
9	Civil Society Organizations (CSOs)
10	Private Sector
Evidence producers	
1	National Institute for Medical Research (NIMR)
2	Tanzania Commission for Science and Technology (COSTECH)
3	Public Universities; The Muhimbili University of Health and Allied Sciences (MUHAS), University of Dar es Salaam (UDSM), University of Dodoma (UDOM)
4	Private Universities; Catholic University of Health and Allied Sciences (CUHAS), Kilimanjaro Christian Medical Centre (KCMC), Hubert Kairuki Memorial University (HKMU), St. John's University of Tanzania (SJUT), St. Augustine University of Tanzania (SAUT), Kampala International University (KIU), St. Francis University College of Health and Allied Sciences (SFUCHAS)
5	Kilimanjaro Clinical Research Institute (KCRI)
6	Ifakara Health Institute (IHI)
7	Non-Governmental Organizations (NGOs)
8	Civil society organizations (CSOs)
9	Private sector

135

136

137 **The study sites**

138 The study site was eighteen (18) Councils of the Local Government Authorities within the nine
139 (9) regions out of 26 Regions of Tanzania Mainland from nine (9) geographical zones as
140 summarized in Table 2 and Figure 2. The nine zones are selected to seek the country's
141 geographical representation. Together, these regions have a total population of 21,119,700,
142 which represents 35.7% of the Tanzanian population. These regions are heterogeneous in
143 population size, distribution of health facilities, human resources for health, and institutions
144 carrying out health research activities. A similar approach has been used in major Tanzanian
145 health studies(4,5,12); This approach provided a comprehensive understanding of the status quo
146 for the use of health research evidence in health planning, and heterogeneous study population

147 for random sampling hence acting as a representative snapshot of all regions in Tanzania. The
 148 study was conducted in 63 randomly selected health facilities, nine (9) Regional Referral
 149 Hospitals (RRHs), eighteen (18) District Hospitals (DHs), Eighteen (18) Health Centres (HCs),
 150 and eighteen (18) Dispensaries. To ensure the inclusion of urban and rural health facilities,
 151 stratification was conducted followed by random sampling.

152
 153
 154
 155
 156

157 **Table 2:** Summary of study sites

ZONE	REGION	COUNCIL
Northern		Siha District
	Kilimanjaro	Moshi Municipal
Central zone		Chamwino District
	Dodoma	Dodoma City
Dar es Salaam		Kigamboni Municipal
	Dar es Salaam	Dar es Salaam City
Eastern		Morogoro Municipal
	Morogoro	Gairo District
Western		Kigoma District
	Kigoma	Kigoma Municipal
Southwest highland		Mbeya City
	Mbeya	Kyela District
Lake		Kwimba District
	Mwanza	Mwanza City
Southern		Mtwara Municipal
	Mtwara	Nanyamba
Southern Highland	Iringa	Iringa Municipal
		Iringa District

158
 159
 160

161 Insert figure 2

162

163 **Figure 2:** Distribution of sampled facilities by zones, regions, and districts

164

165 **The health planning landscape in Tanzania**

166 The health planning process in Tanzania is conducted at two levels; At the Council level, the
167 health facility prepares its plans as well and the Council Health Management Team (CHMT)
168 prepares its plan, whereby plans from the facility and CHMT are later consolidated to form
169 Comprehensive Council plan (CCHP). The preparation of health facility plans is guided by the
170 Health facility planning guidelines and CCHP guidelines.(13). At the regional level, the Regional
171 Health Management Teams (RHMTs) prepare their plans using the RHMT planning guide and the
172 Regional Referral Hospitals (RRHs) plans using the Comprehensive Operational Plan Guide
173 (CHOP)(14). When the plans are completed are sent to the Ministry of Health and President’s
174 Office-Regional Administration and Local Government for final assessment before being sent to
175 the Ministry of Finance for funding. At all levels, we are planning to use the guidelines and the
176 routine data collected from each level.(14).

177 **Study Design**

178 The study adopts a quantitative cross-sectional design focusing on nine regions of Tanzania.
179 (15,16).

180 **Sample size and Sampling procedures**

181 The sample size involved the health planning team members from the RRHs, DHs, HCs, and
182 Dispensaries randomly selected from the nine regions. The sample size was 422 calculated from
183 the Cochran formula (1977). To date, in Tanzania there is no cited reference for the percentage
184 of the use of health research evidence in health planning therefore the marginal error of 50% was
185 used (17).

$$n = \frac{Z^2 * p (1 - p)}{d^2} \text{ (Cochran (1977))}$$

186
187

188 Where by

189 N-sample size

190 Z-confidence interval 95% 1.96, P-
191 proportional from previous study d = margin
192 of error, which is approximately 5% Thus,

$$193 \quad n = \frac{(1.96)^2 \times 50(100 - 50)}{(5)^2}$$

195 =384 including 10% of non-response sample 384 plus 10%=38+384=422

196

197 This study employed a multistage sampling technique for the selection of the study units as
198 summarized in Figure 3. The sampling stages are zones, regions, councils, and public primary
199 health facilities. The first stage was a random selection of one region from each of the nine
200 Zones of the country. In the second stage, in each selected region, councils were clustered into
201 rural and urban, and then one rural and one urban council will be selected from each region
202 followed by a random selection of the health facilities, see Figure 3. The technique is convenient
203 for studying large and diverse populations (5). The sampling stages were zones, regions,
204 councils, and public primary health facilities.

205

206 Insert figure 3

207

208

209 **Variables and their measurements.**

210 **Dependent variable**

211 The dependent variable of this study i.e. the use of health research evidence was obtained from a
212 set of four (4) questions that were measured. The questions were a mixture of binary and
213 multiple-choice questions. Those questions were divided into four areas. The mean score was
214 calculated from those sets of questions; the score had a value of 0 and 1. Those respondents who
215 scored 1 were regarded to have used the health research evidence, while those who scored 0 were

216 regarded to have not used the health research evidence. For multiple choice questions, the
217 respondents were provided with a set of predefined options, allowing them to select the one that
218 best represents their response, which will capture varying degrees or categories on the use of
219 health research evidence.

220 **Independent Variables**

221 The independent variables in this study encompass numerous factors including demographic
222 information such as sex, age, and education level; professional attributes such as position within
223 the health facility and years of schooling; and contextual elements including the type of health
224 facility, stakeholders group represented district, and region. The study aimed to gain insights into
225 the determinants that influence the use of health research evidence during health planning. The
226 determinants had a combination of three constructs which are, capability determinants,
227 opportunity determinants, and motivation determinants both derived from the COM-B Model.

228 **Capability determinants** were obtained from a set of twelve (12) Likert scale questions with a
229 scale of 1 to 5 (1 = strongly agree, 2 = somewhat agree, 3 = neither agree nor disagree, 4 =
230 somewhat disagree, 5 = strongly disagree) and one multiple choice question. The mean and
231 standard deviation of the responses to assess central tendency and variability were calculated.
232 For a multiple-choice question, the frequency distribution of each response category was
233 analyzed. Additionally, a chi-square test was calculated to determine if the distribution of
234 responses was significantly different from what would be expected by chance.

235 **Opportunity determinants**, were obtained from a set of seventeen (17) Likert scale questions
236 with a scale of 1 to 5 (1 = strongly agree, 2 = somewhat agree, 3 = neither agree nor disagree, 4 =
237 somewhat disagree, 5 = strongly disagree). We assigned numerical values (1 to 5) to these
238 responses, where lower values indicate stronger agreement. Next, the mean or median was
239 calculated of these numerical values to quantify the overall tendency. The multiple regression
240 analysis was used to assess the relationship between the independent variable and dependent
241 variables.

242 **Motivation determinants** were obtained from a set of eight (8) Likert scale questions with a
243 scale of 1 to 5 (1 = strongly agree, 2 = somewhat agree, 3 = neither agree nor disagree, 4 =
244 somewhat disagree, 5 = strongly disagree). We assigned numerical values (1 to 5) to these

245 responses, where lower values indicate stronger agreement. Next, the mean or median was
246 calculated of these numerical values to quantify the overall tendency. The multiple regression
247 analysis was used to assess the relationship between the independent variable and dependent
248 variables.

249 **Data collection procedures and tools**

250 Quantitative data was collected face to face person by using the Swahili version of the survey
251 after translation of the English version of the survey and document review checklist (see
252 supplemental file) to planning team members at national, regional, council, and health facility
253 levels. The questions from the tool were adopted and modified from previous studies (18,19).
254 The survey collected information on the use of health research evidence and its determinants
255 from the sampled participants using Open Data Kit software (ODK).

256 The document review checklist was used to guide the document review to see whether the
257 available plans have any evidence of being prepared using health research evidence. The
258 checklist contains a list of questions that will help the review of the plans made at the facility,
259 council, regional, and national levels. The checklist is used to collect information on the
260 available health plans if they are prepared by using health research evidence. Data was collected
261 by a research assistant, who was trained for three days in data collection methods, tools, and
262 ethics. They were selected according to their experience in the health system. Pilot testing of the
263 tools was conducted at Dodoma RHMT and Chamwino District Council at Chamwino CHMT,
264 Chamwino Council Hospital, Chamwino Health Centre, and Buigiri Dispensary to enhance the
265 validity, reliability, and effectiveness of the tool in capturing the intended information.

266 **Data analysis**

267 Responses were reported on all survey items in all response categories and summarized using
268 frequencies and percentages for categorical variables and means and standard deviation for non-
269 categorical variables were computed. Given that the outcome variable had two categories
270 (0=Negative, 1=Positive), A binary logistic regression model was used to assess factors
271 associated with the perception of implementers on the prime vendor system. The model results
272 are regression parameter estimates and odds ratios (OR). The data analysis was conducted using

273 STATA version 18, and the significance of all statistical tests was established at a 5%
274 significance level.

275

276 **Data validity and reliability**

277 To ensure validity and reliability, the validity of quantitative data collection tools was reviewed
278 by independent subject matter experts from the University of Dodoma (UDOM). The reliability
279 assessment of the questionnaire was done using the internal consistency test, with the alpha
280 reliability coefficient being the statistic. (20,21). The range of the alpha coefficient, also known
281 as Cronbach's alpha was 0.71 acceptable to this study. Moreover, exploratory factor analysis to
282 establish the construct validity of the questionnaire and pilot study conducted at Chamwino DC
283 before field data collection to ensure clarity of the data collection tool.

284

285 **Ethics and Dissemination**

286 The study was granted ethical approval by the University of Dodoma Ethical Committee with
287 **Ref. No. MA.84/261/102/A/64/91**. Permission to conduct the study and consent to participate
288 in the study was sought from relevant authorities and participants, respectively. Participants
289 received information about the purpose of the study and data protection. The findings of the
290 study were disseminated to relevant stakeholders through collaborative communication with the
291 Ministry of Health, RHMT/CHMT, Tanzania Health Summit as well as publications to target
292 researchers, practitioners, implementers, and policymakers.

293

294 **Results**

295 The finding from this study is presented in three parts using tables, percentages, and charts, Part
296 one shows the respondents' demographic characteristics and stakeholder group. Part two shows
297 the proportion of the respondents who reported using Health research evidence in health
298 planning and, part three shows the association between health research use in planning and the
299 COM-B constructs (capability, Motivation, and opportunity).

300 **Demographic characteristics**

301 Table 1 describes the respondents' characteristics and the stakeholder group they represent.
 302 There was an almost equal number of female 50.2 % and male 49.8% respondents with a mean
 303 age of 39.51 ± 7.86 years old. Most of them (42.7%) had undergraduate education followed by a
 304 Diploma (36.3%) and less than one-third (14.0%) had a master's degree and above. Few (7.1%)
 305 respondents 'had certificate-level education. Professional background results revealed that the
 306 majority 107 (25.4%) of the respondents were medical doctors followed by nurses 99 (23.5%).
 307 Physiotherapists and panning offices were very few each accounting for 0.24% of the
 308 respondents.

309

310 Table 3: Demographic characteristics of respondents

Variable	Frequency	Percent	Mean \pmSD
Age			39.51 \pm 7.86
<25	3	0.7	
26-35	159	37.7	
36-45	156	37.0	
46-55	93	22.4	
>46	11	2.6	
Region			
Mbeya	48	11.4	
Dodoma	50	11.9	
Iringa	41	9.7	
Mtwara	49	11.6	
Mwanza	50	11.85	
Kigoma	51	12.1	
Kilimanjaro	45	10.7	
Morogoro	48	11.4	
Dar es salaam	40	9.5	
Type of health facility			
Dispensary	37	8.8	
Health center	82	19.4	
Council Hospital	97	23.0	
Regional referral hospital	81	19.2	
RHMT	87	20.6	
CHMT	38	9.0	
Gender			
Male	210	49.76	
Female	212	50.24	
Professional background			
Doctor	107	25.36	

Nurse	99	23.46	
Laboratory scientist	36	8.53	
Pharmacist	28	6.64	
Radiographer	7	1.66	
Environmental Health officer	13	3.08	
Nutrition officer	13	3.08	
Social Welfare Officer	20	4.74	
Physiotherapist	1	0.24	
Biomedical engineer	3	0.71	
Planning officer	1	0.24	
Health Secretary	38	9.00	
Other	56	13.3	
Stakeholders' group			
RHMT	49	11.6	
HMT	280	66.4	
CHMT	90	21.3	
FBO	1	0.2	
HGFC	2	0.5	
Highest level of education			
Certificate	30	7.1	
Diploma Certificate	153	36.3	
Undergraduate degree	180	42.7	
Master's Degree	56	13.3	
PhD	3	0.7	
The position holds currently in health planning.			
Chairperson	30	7.1	
Secretary	45	10.7	
Technical Advisor	11	2.6	
Member	335	79.4	
Other	1	0.2	
Years participate			5.08 ±3.96
<5	283	67.1	
6-10	103	24.4	
>10	36	8.5	

311

312 **Proportion of the use of health research evidence**

313 **Table 4** provides data on the use of health research evidence in a particular context. It shows that
 314 the majority of participants (96.7%) have used routine data, with 49.5% reporting a high extent
 315 of use. Additionally, evidence from routine data was the most commonly used type of evidence
 316 during health planning, with 98.8% of participants reporting its use. Most participants (66.2%)
 317 have used health research evidence as shown in Figure 4, and a significant majority (98.8%)

318 have used health planning guidelines. Moreover, the majority of participants (98.8%) also
 319 indicated the importance of the use of health research evidence, with 82.6% considering it very
 320 important. These findings suggest a strong reliance on routine data and a high recognition of the
 321 importance of health research evidence in the context studied.

322 Table 4: The proportion of the use of health research evidence during health planning

Variable	Frequency	Percent
Ever used routine data		
No	14	3.3
Yes	408	96.7
Extent of use of routine data		
Low	41	10.1
Medium	165	40.4
High	202	49.5
Type of evidence used during health planning (Multiple response)		
Evidence from routine data	403	98.8
Policy documents	280	68.6
Research publications (general)	133	32.6
Systematic reviews/ meta-analysis	41	10.1
Randomized Control trials	17	4.2
Experimental studies	30	7.4
Non-experimental studies	20	4.9
Expert opinions	271	66.4
Policy beliefs	162	39.7
Other	17	4.2
Use of health research evidence		
No	138	33.8
Yes	270	66.2
Use of the Health planning guidelines		
No	5	1.2
Yes	403	98.8
Use of the Ruling party manifesto		
No	44	10.8
Yes	364	89.2
Use of the Policy documents		
No	137	33.6
Yes	271	66.4
Importance of the use of health research evidence		
No	5	1.2
Yes	403	98.8
Level of importance of the use of health research evidence		
Very unimportant	5	1.2
Unimportant	2	0.5

Neutral	14	3.4
Important	50	12.3
Very important	337	82.6

323

324

325

326

.....

327

Insert figure 4

328

.....

329

330 **Figure 4:** Proportion of the use of health research evidence.

331 **The factors associated with the use of health research evidence during health**
332 **planning.**

333 The table presents the results of a binary logistic regression analysis examining factors
334 associated with the use of research evidence during health planning. The analysis includes
335 variables such as age, region, type of health facility, gender, professional background,
336 stakeholder group, highest level of education, position held in health planning, and years of
337 participation. The adjusted odds ratios (OR), confidence intervals (CI), and p-values are provided
338 to indicate the strength and statistical significance of associations.

339 *Significant Variables and Interpretation*

- 340 • **Region:** The analysis compares various regions to the reference category, Kilimanjaro.
341 The odds ratios (OR) show how likely individuals from other regions are to use health
342 research evidence in health planning compared to those from Kilimanjaro. For instance, if
343 a region has an OR greater than 1, individuals from that region are more likely to use
344 research evidence than those from Kilimanjaro. If the OR is less than 1, they are less
345 likely. The p-values indicate whether these differences are statistically significant.

346 • **Level of Education:** Compared to individuals with a diploma (reference category), those
347 with higher education levels were more likely to use research evidence:

348 ➤ Undergraduate degree (OR = 2.396, CI: [1.094, 5.249], p-value = 0.0290):
349 Those with an undergraduate degree are over twice as likely to use
350 research evidence compared to diploma holders.

351 ➤ Master's degree (OR = 2.278, CI: [0.921, 5.631], p-value = 0.0747):
352 Similar to undergraduates, master's degree holders are also more likely to
353 use research evidence in health planning although were not significant.

354 These OR values reflect the increased likelihood of individuals with higher education
355 levels using research evidence compared to the reference group, with p-values indicating
356 the strength of this relationship.

357 • **Years of Participation:**

358 ➤ 6–10 years (OR = 1.092, CI: [0.677,1.761], p-value = 0.7184): Individuals
359 with 6–10 years of participation in health planning show a higher
360 likelihood of using research evidence compared to those with fewer years,
361 though this was not statistically significant.

362 ➤ More than 10 years (OR = 1.399, CI: [0.648,3.019], p-value = 0.3924):
363 Those with over 10 years of participation have a 40% higher likelihood of
364 using research evidence compared to those with fewer years of experience,
365 and this difference was also not statistically significant.

366 The adjusted odds ratios reflect the likelihood of each category of the variable influencing the
367 use of research evidence, controlling for other variables in the model. Significance is determined
368 by both the size of the OR and the p-value, with confidence intervals providing a range within
369 which the true effect likely falls.

370 Table 5: Binary logistic regression for factors associated with the use of research evidence during
371 health planning.

372

Variable	Not use	Use	Unadjusted logistic analysis		Adjusted logistic analysis	
	N (%)	N (%)	OR [95%CI]	p-value	OR [95%CI]	p-value
Age						
<35	52(32.1)	110(67.9)	ref			
36-45	56(35.9)	100(64.1)	0.844[0.530,1.343]	0.4747		
>45	35(33.7)	69(66.4)	0.932[0.552,1.573]	0.7919		
Region						
Kilimanjaro	26(57.8)	19(42.2)	ref		ref	
Mbeya	17(35.4)	31(64.6)	2.495[1.081,5.760]	0.0322	2.604[1.074,6.315]	0.0342
Dodoma	17(34.0)	33(66.0)	2.656[1.156,6.104]	0.0214	2.514[1.053,6.001]	0.0378
Iringa	10(24.4)	31(75.6)	4.242[1.680,10.71]	0.0022	4.707[1.772,12.499]	0.0019
Mtwara	21(42.9)	28(57.1)	1.825[0.805,4.137]	0.1499	1.801[0.727,4.466]	0.2039
Mwanza	12(24.0)	38(76.0)	4.333[1.801,10.43]	0.0011	4.794[1.893,12.140]	0.0009
Kigoma	10(19.6)	41(80.4)	5.611[2.259,13.94]	0.0002	5.922[2.224,15.769]	0.0004
Morogoro	17(35.4)	31(64.6)	2.495[1.081,5.760]	0.0322	2.180[0.901,5.274]	0.0839
Dar es salaam	13(32.5)	27(67.5)	2.842[1.170,6.904]	0.0211	3.229[1.211,8.611]	0.0192
Type of health facility						
Dispensary	17(46.0)	20(54.1)	ref		ref	
Health center	25(30.5)	57(69.5)	1.938[0.871,4.311]	0.1048	2.029[0.829,4.964]	0.1211
Council Hospital	45(46.4)	52(53.6)	0.982[0.459,2.100]	0.9631	0.809[0.323,2.026]	0.6511
RRH	19(23.5)	62(76.5)	2.774[1.214,6.335]	0.0155	2.663[1.895,7.920]	0.0383
RHMT	27(31.0)	60(69.0)	1.889[0.857,4.163]	0.1147	0.350[0.031,3.902]	0.3932
CHMT	10(26.3)	28(73.7)	2.380[0.903,6.273]	0.0795	5.647[0.602,52.94]	0.3167
Gender						
Male	71(33.8)	139(66.2)	1.007[0.673,1.507]	0.9736		
Female	72(34.0)	140(66.0)	ref			
Professional background						
Doctor	37(34.6)	70(65.4)	ref			
Nurse	35(35.4)	64(64.7)	0.967[0.545,1.714]	0.9073		
Lab scientist & Pharmacist	25(35.2)	46(64.8)	0.973[0.518,1.825]	0.9310		
Non-medical	28(31.5)	61(68.5)	1.152[0.633,2.096]	0.6444		
Other	18(32.1)	38(67.7)	1.116[0.561,2.220]	0.7548		
Stakeholders' group						
RHMT	13(26.5)	36(73.5)	1.609[0.816,3.171]	0.1697	0.843[0.177,4.012]	0.8297
CHMT	26(28.9)	64(71.1)	1.430[0.854,2.396]	0.1741	5.647[0.602,52.93]	0.1295
HMT	104(36.8)	179(63.3)	ref		ref	
Highest level of education						
Certificate	15(50.0)	15(50.0)	ref		ref	
Diploma	57(37.3)	96(62.8)	1.684[0.767,3.700]	0.1943	1.609[0.670,3.867]	0.2874
Undergraduate	53(29.4)	127(70.6)	2.396[1.094,5.249]	0.0290	1.824[0.710,4.685]	0.2120

Master's Degree	18(30.5)	41(69.5)	2.278[0.921,5.631]	0.0747	2.082[0.684,6.337]	0.1968
Position holds currently in health planning						
Secretary	19(42.2)	26(57.8)	ref		ref	
Chairperson	8(26.7)	22(73.3)	2.009[0.738,5.475]	0.1724	1.933[0.647,5.780]	0.2381
Technical	4(36.4)	7(63.6)	1.279[0.327,5.000]	0.7236	0.789[0.173,3.599]	0.7600
Member	112(33.3)	224(66.7)	1.462[0.776,2.754]	0.2404	1.328[0.651,2.708]	0.4354
Years participate						
<5	99(35.0)	184(65.0)	ref			
6-10	34(33.0)	69(67.0)	1.092[0.677,1.761]	0.7184		
>10	10(27.8)	26(72.2)	1.399[0.648,3.019]	0.3924		

373

374

375 **Capability: The knowledge and skills of health research evidence use**

376 Table 6 presents the findings from a survey on the capacity to use health research evidence
 377 (knowledge and skills). Each row in the table represents a different variable related to the use of
 378 health research evidence, and the columns display the responses categorized into five levels of
 379 importance: Very Unimportant (VUIM), Unimportant (UIM), Neutral, Important (IM), and Very
 380 Important (VIM).

381 From the data, it's clear that the majority of respondents perceive the use of health research
 382 evidence as very important or important across the various variables. In the variable "Gives the
 383 latest information," the majority of respondents rated it as Very Important (65.7%) or Important
 384 (24.5%). Similarly, in the variable "Help the policymakers understand a specific problem," a
 385 substantial proportion of respondents indicated that it is Very Important (66.7%) or Important
 386 (24.3%). Overall, the data suggests that there is a strong recognition of the significance of health
 387 research evidence in various aspects of policymaking and intervention implementation.

388

389 Table 6: Knowledge and skills of health research evidence use

Variable	VUIM n (%)	UIM n (%)	Neutral n (%)	IM n (%)	VIM n (%)
Gives the latest information	5(1.2)	3(1.2)	32(7.8)	100(24.51)	268(65.7)
Help the policymakers understand a specific problem	3(0.7)	4(1.0)	30(7.4)	99(24.3)	272(66.7)
Help the policymakers to	3(0.7)	5(1.2)	24(5.9)	113(27.7)	263(64.5)

implement various health interventions					
Avoid repeating the failures of others	3(0.7)	3(0.7)	19(4.7)	101(24.8)	282(69.1)
Introduces health planners to new ideas	3(0.7)	2(0.5)	19(4.7)	107(26.2)	277(67.9)

390

391 Table 7 provides information on the obstacles hindering the utilization of health research
 392 evidence. The variables include Lack of necessary knowledge and training in research,
 393 Inadequate human and non-human resources, Difficulty in accessing health research evidence,
 394 Perceptions that health research is for academic purposes, Lack of involvement in research
 395 activities, and Traditional ways of planning, as well as the minimal usage of health research
 396 evidence among planning team members at the regional and council levels, and lack of
 397 dissemination.

398 The data is presented in terms of the number and percentage of participants who strongly
 399 disagree (SDSA), disagree (DSA), are neutral, agree (AG), and strongly agree (SAGR) with each
 400 obstacle. This data provides insight into the specific challenges that impact the utilization of
 401 health research evidence within the context of the study. The highest percentage is for
 402 "Perceptions that health research is for academic purposes" at 22.6%, and the lowest percentage
 403 is for "Lack of necessary knowledge and training in research" at 2.9%.

404

405 **Table 7: Obstacles that hinder the utilization of health research evidence**

Variable	SDSA n (%)	DSA n (%)	Neutral n (%)	AG n (%)	SAGR n (%)
Lack of necessary knowledge and training in research	12(2.94)	18(4.4)	7(1.7)	67(16.4)	304(74.5)
Inadequate human and non-human resources	16(3.92)	18(4.4)	7(1.7)	81(19.9)	286(70.1)
Difficult in accessing health research evidence	36(8.82)	41(10.1)	24(5.9)	110(27.0)	197(48.3)
Perceptions that health research	92(22.55)	40(9.8)	16(3.9)	81(19.9)	179(43.9)

is for academic purposes					
Lack of involvement in research activities	15(3.68)	25(6.1)	11(2.7)	93(22.8)	264(64.7)
Traditional ways of planning, and Minimal usage of health research evidence among planning team members at the regional and council levels	36(8.82)	21(5.2)	13(3.2)	92(22.6)	246(60.3)
lack of dissemination	11(2.70)	15(3.7)	17(4.2)	105(25.7)	260(63.7)

406

407

408 **Opportunities for the use of health research evidence**

409 Table 8 indicates the opportunities for the use of health research evidence, helping stakeholders
 410 to identify areas for improvement and investment to optimize the use of health research
 411 evidence. The table presents the opportunities for utilizing health research evidence, categorized
 412 into physical and social opportunities.

413 For physical opportunities, the table highlights the availability of resources such as research
 414 coordinators, equipment, internet access, a planned budget for research, and the presence of
 415 symposiums for research discussions. The percentages indicate the distribution of responses,
 416 ranging from strongly disagree to strongly agree, reflecting the varying levels of agreement
 417 among the respondents regarding these physical opportunities. In the physical opportunities
 418 category, the highest percentage is 76.3%, representing the availability of the internet, while the
 419 lowest percentage is 0.7%, indicating the availability of Internet.

420 In the social opportunities section, the table emphasizes the presence of disparities among
 421 different populations, the involvement of community members in health planning teams, and the
 422 enhancement of health literacy in society. The percentages provide an insight into the attitudes
 423 and perceptions of the respondents toward these social opportunities. In the social opportunities
 424 category, the highest percentage is 55.9%, reflecting improved health literacy in society, and the

425 lowest percentage is 2.4%, which represents the presence of disparities among different
 426 populations allowing for more equitable representation.

427 Table 8: Opportunities for the use of health research evidence

Variable	SDSA n (%)	DSA n (%)	Neutral n (%)	AG n (%)	SAGR n (%)
Physical opportunities					
Enabling the utilization of information generated from the healthcare system	6(1.4)	9(2.1)	18(4.3)	130(30.8)	259(61.4)
Availability of research coordinators	9(2.1)	18(4.3)	22(5.2)	93(22.0)	280(66.4)
Availability of equipment	4(1.0)	16(3.8)	21(5.0)	88(20.9)	293(69.4)
Availability of internet	3(0.7)	9(2.1)	19(4.5)	69(16.4)	322(76.3)
Availability of planned budget for research	11(2.6)	17(4.0)	27(6.4)	64(15.2)	303(71.8)
Presence of symposiums for research discussions	9(2.1)	17(4.0)	23(5.5)	106(25.1)	267(63.3)
Social opportunities					
The presence of disparities among different populations allows for more equitable	10(2.4)	24(5.7)	38(9.0)	147(34.8)	203(48.1)
The presence of community members in the health planning teams	11(2.6)	28(6.6)	55(13.0)	119(28.2)	209(49.5)
Improved health literacy in the society, empowering individuals to make informed	4(1.0)	19(4.5)	18(4.3)	145(34.4)	236(55.9)

428

429 **Motivations for the use of health research evidence**

430 Table 9, based on the data presented, we can observe that the automatic motivations for the use
 431 of health research evidence are primarily centered around the availability of transparency and
 432 accountability mechanisms 66.4% strongly agreeing, and 23.2% agreeing with the statement

433 availability of job training 73.2% strongly agreeing and 21.3% agreeing respectively. Provision
 434 of incentives to health planners, with 72.4% strongly agreeing and 15.4% agreeing respectively.
 435 In contrast, reflective motivations such as continuous quality improvement in the healthcare
 436 system and the presence of active stakeholder engagement in health planning also garnered
 437 significant support, with 63.51% and 59.48% strongly agreeing or agreeing, respectively. This
 438 indicates a strong emphasis on the proactive involvement of various stakeholders and the
 439 continuous enhancement of the healthcare system based on research evidence.

440

441 Table 9: Motivations of the use of health research evidence

Variable	SDSA n (%)	DSA n (%)	Neutral n (%)	AG n (%)	SAGR n (%)
Automatic motivations					
Provision of incentives to health planners	19(4.5)	21(5.0)	13(3.1)	65(15.4)	304(72.0)
Availability of job training	3(0.7)	4(1.0)	16(3.8)	90(21.3)	309(73.2)
Availability of Short-term courses	5(1.2)	13(3.1)	21(5.0)	75(17.8)	308(73.0)
Availability of Long-term course	18(4.3)	24(5.7)	24(5.7)	105(24.9)	251(59.5)
Reflective motivations					
Presence of interaction between policymakers, implementers, researchers, and academicians	3(0.7)	17(4.0)	21(5.0)	111(26.3)	270(64.0)
Presence of active stakeholder engagement in health planning	6(1.4)	16(3.8)	33(7.8)	116(27.5)	251(59.5)
Continuous quality improvement in the healthcare system	1(0.2)	10(2.4)	12(2.8)	131(31.0)	268(63.5)
Availability of transparency and accountability mechanisms	5(1.2)	12(2.8)	27(6.4)	98(23.2)	280(66.4)

442

443 Findings from document review checklist

444 The analyzed data from Table 10 indicates that the majority of planning teams incorporate
 445 research evidence into their health plans. A substantial 97.8% of respondents confirmed having a
 446 current fiscal year health plan, and 98.9% reported having a planning guideline. Most plans

447 include key health frameworks and guides, with 70.8% having the CCHP, 83.2% using the HF
 448 Planning Guide, and 89.9% incorporating situational analysis. Scientific engagement is
 449 evidenced by 75.3% of teams receiving invitation letters for conferences. However, challenges
 450 remain, as only 22.5% have a budget for research, and only 23.6% allocate budget specifically
 451 for research within the health plan. Additionally, just 28.1% include a research coordinator in the
 452 planning team, and 43.8% of the plans are well-cited. This highlights a strong foundation in
 453 research usage but also gaps in financial and personnel support for research-driven planning.

454 Table 10: looking for evidence that planning teams use health research evidence in their plans

Variable	No	Yes	Not applicable
	n (%)	n (%)	n (%)
Do you have a current fiscal year health plan?	1(1.1)	87(97.8)	1(1.1)
Do you have a planning guideline?	1(1.12)	88(98.9)	0(0.0)
Presence of CCHP	12(13.5)	63(70.8)	14(15.7)
HF Planning Guide	13(14.6)	74(83.2)	2(2.3)
RHMT guide	26(29.2)	20(22.5)	43(48.3)
CHOP	30(33.7)	24(27.0)	35(39.3)
Presence of situational analysis in the plan	8(9.0)	80(89.9)	1(1.1)
Presence of invitation letters to attend various scientific conferences	22(24.7)	67(75.3)	0(0.0)
Are the well-cited plans	50(56.2)	39(43.8)	0(0.0)
Is there a budget for research available	69(77.5)	20(22.5)	0(0.0)
Is there a research-allocated budget in their health plan?	68(76.4)	21(23.6)	0(0.0)
Is the research coordinator among the planning team members who prepared the plan	64(71.9)	25(28.1)	0(0.0)

455

456

457 **Discussion**

458 This study represents the first comprehensive analysis of the use of health research evidence in
459 Tanzania. It aimed to analyze the determinants of the use of health research evidence by deep-
460 diving into the three determinants influencing behavior i.e. Capability, Opportunity, and
461 Motivation. The findings from this study indicated that all three determinants significantly
462 impact the ability of health planners to use health research evidence during health planning.
463 However, we found that the use of research evidence is associated with having a degree, more
464 than 10 years of involvement in health planning activities, and prior knowledge in research. We
465 found several facilitator determinants such as organizational support infrastructures like, the
466 availability of computers and the internet, the availability of research experts and research
467 coordinators, availability of academic institutions that give a chance for dissemination of
468 research. On the other hand, barrier determinants included lack of human resources, lack of
469 funding for research activities, lack of necessary knowledge and training, perceptions that
470 research is for academic purposes, and lack of dissemination.

471 **Proportion of the use of health research evidence**

472 The findings of this study on the proportion of the use of health research evidence demonstrate a
473 high level of reliance on health research evidence, particularly routine data, during health
474 planning. The majority of participants (96.7%) reported using routine data, with nearly half
475 (49.5%) using it extensively, which is consistent with findings from similar studies. For instance,
476 a study by Lavis et al (22) found that routine data and health statistics are frequently utilized in
477 policy-making and health-planning processes, underscoring the central role of such data in
478 evidence-based decision-making. Moreover, the widespread use of health planning guidelines
479 (98.8%) and the high proportion of participants (98.8%) recognizing the importance of health
480 research evidence align with findings from studies on the use of research evidence in health
481 decision-making process which emphasize the role of guidelines and evidence in shaping health
482 policies (23,24). The high rate of health research evidence use (66.2%) in this study also mirrors
483 the findings which reported similar levels of research evidence use in healthcare settings(25).
484 These results highlight the growing recognition and integration of research evidence in health
485 planning, reflecting broader trends in evidence-informed policymaking across health systems in
486 Tanzania and globally.

487 **Capability: The knowledge and skills of health research evidence use**

488 Explicitly, capability determinants such as knowledge and skills were found to be critical, with
489 health planning team members who had higher levels of education (Undergraduate, a master
490 level, and above) and access to training more likely to use health research evidence to almost
491 70%, with the capability constraints faces more those with certificate and diploma level, lack of
492 necessary knowledge and training in research accounts up to 70%, being the most barrier to
493 utilization of health research evidence, inadequate human and non-human resources, difficult in
494 accessing research evidence, perception that research is for academic purposes and lack of
495 dissemination. These findings are consistent with systematic reviews conducted in other low-
496 and middle-income countries, where similar capability constraints such as poor access to good
497 quality relevant research, and lack of timely research output have been identified as major
498 barriers to evidence-based health planning (26,27).

499

500 **Opportunities for the use of health research evidence**

501 In comparing our results with existing literature on Opportunity determinants (both physical and
502 social), it is evident that the availability of resources and institutional support plays a crucial role
503 in enabling the use of research evidence. Our study revealed that health planning teams with
504 better access to physical opportunities such as health information system, availability of research
505 coordinators, equipment, internet, research budgets, presence of symposiums for research
506 discussion, the existence of academic institutions such as universities, and social opportunities
507 like the presence of community members representatives within the health planning teams and
508 improved health literacy in the community were more inclined to utilize research findings in
509 their work. This aligns with findings from two studies conducted in LMICs (28,29), which
510 highlighted the importance of institutional support in facilitating the use of research evidence in
511 health policy. However, our study also noted a distinct challenge in the Tanzanian context with
512 limited access to relevant health research evidence due to infrastructural constraints, a barrier
513 less emphasized in studies from more resource-rich settings (30,31).

514 **Motivations for the use of health research evidence**

515 Particularly the automatic motivations to utilize health research evidence were also found to be
516 significant, especially on the provision of incentives to health planners to greater than 72%.

517 Other motivations include job training, and short and long-term courses that influenced the
518 motivations to use the health research evidence. The automatic motivation factors included the
519 presence of interaction between policymakers, implementers, researchers, and academicians, and
520 continuous quality improvement agenda in the health sector. Health planners who perceived the
521 use of research evidence as beneficial and aligned with their professional goals were more likely
522 to engage with it actively. This is in line with the COM-B model, which underscores the role of
523 motivation in behavior change (10). Our findings add to the body of literature by highlighting the
524 specific motivational barriers in Tanzania, such as a lack of a clear roadmap or framework for
525 the utilization of health research evidence, which can reduce the enthusiasm for utilizing research
526 in health planning (32)(33).

527 **Document review checklist**

528 The findings from the documents review checklist revealed a high level of integration of research
529 evidence in health planning aligning with similar studies that underscore the importance of
530 research-informed frameworks in public health planning (34)(35). The near-universal adoption
531 of fiscal-year health plans (97.8%) and planning guidelines (98.9%) is consistent with global
532 trends in evidence-based health planning, which emphasizes standardized planning processes
533 (36). The presence of documents like CCHP and HF Planning guide mirrors findings from an
534 urban strategic planning using blue-green infrastructure (37), who highlighted their role in
535 improving health outcomes, the limited financial resources dedicated to health research in this
536 study 22.5% suggests a common barrier. Similarly, the low inclusion of research coordinators
537 (28.1%) echoes observations by a study designed for research approach during learning (38) on
538 the limited interdisciplinary engagement in health planning. These gaps in budget allocation and
539 personnel highlight persistent structural challenges to fully operationalizing research evidence-
540 backed plans, despite a broad commitment to evidence-based frameworks.

541 The study provided valuable insights into the factors influencing the use of health research
542 evidence during health planning in Tanzania. The results from the binary logistic regression
543 analysis highlight significant associations between region, level of education, and years of
544 participation in health planning. Notably, region was a significant predictor, with participants
545 from regions other than Kilimanjaro demonstrating varying likelihoods of using research

546 evidence, as indicated by the odds ratios. Educational attainment also emerged as a key
547 determinant; individuals with undergraduate degrees were over twice as likely to use health
548 research evidence compared to those with a diploma, with statistically significant results (p-value
549 = 0.029). However, while master's degree holders showed a similar trend, the relationship was
550 not statistically significant (p-value = 0.0747). The analysis of years of participation revealed
551 that those with more than 10 years of experience were 40% more likely to use health research
552 evidence, though this association was not statistically significant. These findings align with other
553 studies that emphasize the importance of education and experience in evidence-based health
554 planning (24,26). The adjusted odds ratios and confidence intervals provide a nuanced
555 understanding of the strength and variability of these associations, contributing to a growing
556 body of literature on the determinants of research utilization in public health policy and planning.

557 The implications of these findings are essential for improving health planning in Tanzania. To
558 enhance the use of health research evidence, interventions must address all three determinants:
559 enhancing the capability of health planners through capacity building, improving opportunities
560 by ensuring better access to human and non-human resources and organizational support, and
561 fostering motivation by recognizing and rewarding health planners that use evidence-based
562 decision-making. These strategies align with recommendations from similar studies in other
563 LMICs, suggesting that a multifaceted approach including the development of middle range
564 theory (MRT) outlining the main facilitators for utilization of evidence for malaria treatment
565 policy in Uganda and the importance of measuring total brain activity in neuroimaging are
566 essential for promoting the use of research in health planning (39)(40).

567 **Conclusion**

568 This study provides a comprehensive assessment of the determinants influencing the use of
569 health research evidence in Tanzania, focusing on capability, opportunity, and motivation. The
570 findings demonstrate that all three determinants significantly impact the behavior of health
571 planners, highlighting those factors such as higher education levels, years of experience in health
572 planning, and prior research knowledge contribute to greater use of health research evidence.
573 Organizational infrastructure and institutional support further facilitate this process, while

574 barriers include insufficient human resources, lack of funding, inadequate training, and
575 perceptions that research is solely for academic purposes.

576 The study also reveals a high reliance on routine data and health planning guidelines, indicating
577 the growing recognition of research evidence in health decision-making. However, addressing
578 the barriers to research evidence use will require targeted interventions that strengthen the
579 capability of health planners, improve access to resources and institutional support, and enhance
580 motivation through incentives and professional development.

581 Future research should explore the long-term impact of these interventions, as well as contextual
582 factors like political will and leadership, to fully understand how these determinants collectively
583 shape health planning behavior. A multifaceted approach i.e. a generic framework for
584 strengthening Knowledge Management in Tanzania (KMT) that will have a triad pathway:
585 Capacity building on healthcare workers and researchers, infrastructure development, and
586 funding that will improve access to health research database and resources within the healthcare
587 system, and organizational support through the establishment of guidelines for the use of health
588 research evidence in the decision-making process. This is essential for promoting the use of
589 health research evidence, ultimately improving health outcomes in Tanzania and other low- and
590 middle-income countries with similar challenges (41,42)

591

592 **Abbreviations**

CCHP	Comprehensive Council Health Planning
CHOP	Comprehensive Hospital Operational Plan
LGAs	Local Government Authorities
LMICs	Lower-and Middle-Income Countries
MoH	Ministry of Health
PO-RALG	President's Office-Regional Administration and Local Government
RHMT	Regional Health Management Team

SDGs Sustainable Development Goals

UHC Universal Health Coverage

WHO World Health Organization

593

594 **Acknowledgments**

595 We would like to acknowledge the contribution of the University of Dodoma faculty members
596 from the School of Nursing and Public Health for their input in the process of development of
597 this manuscript. We would like to express our sincere thanks to the study facilities, the
598 RHMTs, and the CHMTs for their support. We are grateful to the research assistants (Edmund
599 Bunyaga, Theresia Ngungulu, Jestina Nyondo, Devina Mafuta Micky Masanyaji, and Ally
600 Kinyaga) for their assistance during data collection.

601 **Author Contributions**

602 PK; Conceptualization, design of the work, data collection and analysis, preparing the
603 manuscript draft, developing the methodology, results, discussion, and conclusion. RM,
604 Conceptualization, critically reviewing the methodology and critically reviewing the manuscript,
605 AK, Conceptualization, critically reviewing the methodology. All authors revised the work and
606 approved the paper to be submitted for publication.

607 **REFERENCES**


- 608 1. Khalid FJI. The Impact of Poor Planning and Management on the Duration of Construction
609 Projects. Multi-Knowl Electron Compr J Educ Sci Publ [Internet]. 2019;(2):1–20. Available
610 from: www.mecsjs.com
- 611 2. Onwujekwe O, Etiaba E, Mbachu C, Arize I, Nwankwor C, Ezenwaka U, et al. Does
612 improving the skills of researchers and decision-makers in health policy and systems
613 research lead to enhanced evidence-based decision making in Nigeria?—A short term
614 evaluation. Oelke N, editor. PLOS ONE [Internet]. 2020 Sep 3 [cited 2024 Feb
615 16];15(9):e0238365. Available from: <https://dx.plos.org/10.1371/journal.pone.0238365>
- 616 3. Henriksson DK, Peterson SS, Waiswa P, Fredriksson M. Decision-making in district health
617 planning in Uganda: Does use of district-specific evidence matter? Health Res Policy Syst.
618 2019 Jun 6;17(1).

- 619 4. Kagoma P, Mongi R, Kapologwe NA, Kengia J, Kalolo A. Health research evidence: its
620 current usage in health planning, determinants and readiness to use knowledge translation
621 tools among health planning teams in Tanzania—an exploratory mixed-methods study
622 protocol. *BMJ Open* [Internet]. 2024 Jun [cited 2024 Jun 27];14(6):e081517. Available from:
623 <https://bmjopen.bmj.com/lookup/doi/10.1136/bmjopen-2023-081517>
- 624 5. Kengia JT, Kalolo A, Barash D, Chwa C, Hayirli TC, Kapologwe NA, et al. Research
625 capacity, motivators and barriers to conducting research among healthcare providers in
626 Tanzania’s public health system: a mixed methods study. *Hum Resour Health* [Internet].
627 2023 Sep 5 [cited 2024 Feb 9];21(1):73. Available from: [https://human-resources-](https://human-resources-health.biomedcentral.com/articles/10.1186/s12960-023-00858-w)
628 [health.biomedcentral.com/articles/10.1186/s12960-023-00858-w](https://human-resources-health.biomedcentral.com/articles/10.1186/s12960-023-00858-w)
- 629 6. Van De Goor I, Hämäläinen RM, Syed A, Juel Lau C, Sandu P, Spitters H, et al.
630 Determinants of evidence use in public health policy making: Results from a study across six
631 EU countries. *Health Policy* [Internet]. 2017 Mar [cited 2024 Feb 26];121(3):273–81.
632 Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0168851017300192>
- 633 7. Van Kammen J, Savigny DD, Sewankambo N. Using knowledge brokering to promote
634 evidence-based policy-making: the need for support structures. *Bull World Health Organ*.
635 2006;84(8).
- 636 8. Guilbert JJ. The World Health Report 2006: Working together for health [1]. *Educ Health*
637 *Change Learn Pract*. 2006;19(3):385–7.
- 638 9. Mboera LEG, Rumisha SF, Mbata D, Mremi IR, Lyimo EP, Joachim C. Data utilisation and
639 factors influencing the performance of the health management information system in
640 Tanzania. *BMC Health Serv Res*. 2021;21(1):4–11.
- 641 10. Michie S, van Stralen MM, West R. The behaviour change wheel: A new method for
642 characterising and designing behaviour change interventions. *Implement Sci*. 2011 Apr
643 23;6(1).
- 644 11. De Leo A, Bayes S, Bloxsome D, Butt J. Exploring the usability of the COM-B model and
645 Theoretical Domains Framework (TDF) to define the helpers of and hindrances to evidence-
646 based practice in midwifery. *Implement Sci Commun* [Internet]. 2021 Dec [cited 2024 Aug
647 12];2(1):7. Available from:
648 [https://implementationsciencecomms.biomedcentral.com/articles/10.1186/s43058-020-](https://implementationsciencecomms.biomedcentral.com/articles/10.1186/s43058-020-00100-x)
649 [00100-x](https://implementationsciencecomms.biomedcentral.com/articles/10.1186/s43058-020-00100-x)
- 650 12. Kapologwe NA, Kalolo A, Kibusi SM, Chaula Z, Nswilla A, Teuscher T, et al.
651 Understanding the implementation of Direct Health Facility Financing and its effect on
652 health system performance in Tanzania: A non-controlled before and after mixed method
653 study protocol. *Health Res Policy Syst*. 2019 Jan 30;17(1).
- 654 13. United THE, Of R. Comprehensive Council Health Planning Guideline Fifth Edition. 2019;
- 655 14. Ministry of Health. Guideline for Developing Comprehensive Hospital Operational Plan (
656 CHOP) for Regional Referral Hospitals. 2016;(August).

- 657 15. Mwilike B, Nalwadda G, Kagawa M, Malima K, Mselle L, Horiuchi S. Knowledge of
658 danger signs during pregnancy and subsequent healthcare seeking actions among women in
659 Urban Tanzania: a cross-sectional study. *BMC Pregnancy Childbirth* [Internet]. 2018 Dec
660 [cited 2024 Sep 21];18(1):4. Available from:
661 <https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/s12884-017-1628-6>
- 662 16. Lyimo AA, Guo J, Mushy SE, Mwilike BE. Use of contraceptives and associated factors
663 among male adolescents in rural secondary schools, Coast Region, Tanzania: a school-based
664 cross-sectional study. *Contracept Reprod Med* [Internet]. 2024 Mar 1 [cited 2024 Sep
665 21];9(1):8. Available from:
666 <https://contraceptionmedicine.biomedcentral.com/articles/10.1186/s40834-024-00268-w>
- 667 17. Wahed T, Kaukab SST, Saha NC, Khan IA, Khanam F, Chowdhury F, et al. Knowledge of,
668 attitudes toward, and preventive practices relating to cholera and oral cholera vaccine among
669 urban high-risk groups: findings of a cross-sectional study in Dhaka, Bangladesh. *BMC*
670 *Public Health* [Internet]. 2013 Dec [cited 2024 Feb 9];13(1):242. Available from:
671 <https://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-13-242>
- 672 18. Subedi D. Explanatory sequential mixed method design as the third research community of
673 knowledge claim. *Am J Educ Res*. 2016;4(7):570–7.
- 674 19. Shiyanbola OO, Rao D. Using an exploratory sequential mixed methods design to adapt an
675 Illness Perception Questionnaire for African Americans with diabetes : the mixed data
676 integration process. 2021;9(1):796–817.
- 677 20. Surucu L, Maslakci A. Validity and Reliability in Quantitative Research. *Bus Manag Stud*
678 *Int J*. 2020;8(3):2694–726.
- 679 21. Jiyenze MK, Tundui C, Sirili N, Mollel H. Effects of implementing a health sector strategic
680 plan : Eidence from Tanzania. *East Afr Med J*. 2022;99(9):5147–55.
- 681 22. Lavis J. Assessing country-level efforts to link research to action. *Bull World Health Organ*
682 [Internet]. 2006 Aug 1 [cited 2024 Oct 4];84(8):620–8. Available from:
683 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2627430/pdf/16917649.pdf/>
- 684 23. Hanney SR, Kanya L, Pokhrel S, Jones TH, Boaz A. How to strengthen a health research
685 system: WHO’s review, whose literature and who is providing leadership? *Health Res Policy*
686 *Syst* [Internet]. 2020 Dec [cited 2024 Oct 4];18(1):72. Available from: [https://health-policy-](https://health-policy-systems.biomedcentral.com/articles/10.1186/s12961-020-00581-1)
687 [systems.biomedcentral.com/articles/10.1186/s12961-020-00581-1](https://health-policy-systems.biomedcentral.com/articles/10.1186/s12961-020-00581-1)
- 688 24. Orton L, Lloyd-Williams F, Taylor-Robinson D, O’Flaherty M, Capewell S. The Use of
689 Research Evidence in Public Health Decision Making Processes: Systematic Review. Ross
690 JS, editor. *PLoS ONE* [Internet]. 2011 Jul 26 [cited 2024 Oct 4];6(7):e21704. Available
691 from: <https://dx.plos.org/10.1371/journal.pone.0021704>
- 692 25. Innvær S, Vist G, Trommald M, Oxman A. Health policy-makers’ perceptions of their use of
693 evidence: a systematic review. *J Health Serv Res Policy* [Internet]. 2002 Oct 1 [cited 2024
694 Oct 3];7(4):239–44. Available from: <https://doi.org/10.1258/135581902320432778>

- 695 26. Oliver K, Innvar S, Lorenc T, Woodman J, Thomas J. A systematic review of barriers to and
696 facilitators of the use of evidence by policymakers. *BMC Health Serv Res* [Internet]. 2014
697 Dec [cited 2024 Apr 8];14(1):2. Available from:
698 <https://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-14-2>
- 699 27. Koon AD. A scoping review of the uses and institutionalisation of knowledge for health
700 policy in low- and middle-income countries. 2020;
- 701 28. Lavis JN, Paulsen EJ, Oxman AD, Moynihan R. Evidence-informed health policy 2 – Survey
702 of organizations that support the use of research evidence. *Implement Sci*. 2008;3(1):1–17.
- 703 29. Lavis JN, Wilson MG, Grimshaw JM, Haynes RB, Hanna S, Raina P, et al. Effects of an
704 evidence service on health-system policy makers’ use of research evidence: A protocol for a
705 randomised controlled trial. 2011;
- 706 30. Moat KA, Lavis JN. 10 best resources for . . . evidence-informed health policy making.
- 707 31. Gavine A. Maximising the availability and use of high-quality evidence for policymaking:
708 collaborative, targeted and efficient evidence reviews. *PALGRAVE Commun*. 2018;
- 709 32. Armstrong R, Waters E, Dobbins M, Anderson L, Moore L, Petticrew M, et al. Knowledge
710 translation strategies to improve the use of evidence in public health decision making in local
711 government: intervention design and implementation plan. *Implement Sci* [Internet]. 2013
712 Dec [cited 2024 Feb 16];8(1):121. Available from:
713 <http://implementationscience.biomedcentral.com/articles/10.1186/1748-5908-8-121>
- 714 33. Armstrong JS. *Guidelines for Science: Evidence and Checklists*.
- 715 34. Smith JD, Hasan M. Quantitative approaches for the evaluation of implementation research
716 studies. *Psychiatry Res* [Internet]. 2020 Jan [cited 2024 Oct 28];283:112521. Available from:
717 <https://linkinghub.elsevier.com/retrieve/pii/S0165178119307024>
- 718 35. Roberts-Lewis SF, Baxter HA, Mein G, Quirke-McFarlane S, Leggat FJ, Garner HM, et al.
719 The Use of Social Media for Dissemination of Research Evidence to Health and Social Care
720 Practitioners: Protocol for a Systematic Review. *JMIR Res Protoc* [Internet]. 2023 May 12
721 [cited 2024 Oct 28];12:e45684. Available from:
722 <https://www.researchprotocols.org/2023/1/e45684>
- 723 36. Brown G, Kyttä M, Reed P. Using community surveys with participatory mapping to
724 monitor comprehensive plan implementation. *Landsc Urban Plan* [Internet]. 2022 Feb [cited
725 2024 Oct 28];218:104306. Available from:
726 <https://linkinghub.elsevier.com/retrieve/pii/S0169204621002693>
- 727 37. Sörensen J, Persson AS, Olsson JA. A data management framework for strategic urban
728 planning using blue-green infrastructure. *J Environ Manage* [Internet]. 2021 Dec [cited 2024
729 Oct 28];299:113658. Available from:
730 <https://linkinghub.elsevier.com/retrieve/pii/S0301479721017205>

- 731 38. Lim FV, Nguyen TTH. Design-based research approach for teacher learning: a case study
732 from Singapore. *ELT J* [Internet]. 2022 Oct 14 [cited 2024 Oct 28];76(4):452–64. Available
733 from: <https://academic.oup.com/eltj/article/76/4/452/6316630>
- 734 39. Nabyonga-Orem J, Ssengooba F, Macq J, Criel B. Malaria treatment policy change in
735 Uganda: what role did evidence play? 2014;
- 736 40. Hyder F, Rothman DL. Evidence for the importance of measuring total brain activity in
737 neuroimaging.
- 738 41. Bennett NJ. Using perceptions as evidence to improve conservation and environmental
739 management. *Conserv Biol*. 2016;30(3).
- 740 42. Hollingworth SA. What do we need to know? Data sources to support evidence-based
741 decisions using health technology assessment in Ghana. . *Epidemiology*. 2020;
- 742



Figure

DISTRIBUTION OF SAMPLED FACILITIES BY ZONES/ REGIONS/DISTRICTS.



**Legend
Facilities**

- Regional Hospital
- Council Hospital
- Dispensary
- Health Centre

- Regions
- Selected Regions
- Water Bodies



Figure





Fig.3 Sampling design



Figure