

Title: Undiagnosed hypertension and its associated factors in India: A rural-urban contrast from the National Family Health Survey (2019-21)

Short title – Rural-urban differences in undiagnosed hypertension in India

Authors and Affiliations:

Aritrik Das¹, Yukti Bhandari¹, Angad Singh², Jugal Kishore³, Sonu Goel^{4*}

¹Department of Community Medicine and School of Public Health, Postgraduate Institute of Medical Education and Research, Chandigarh, India

²Department of Biostatistics and Epidemiology, International Institute for Population Sciences Mumbai, India

³Department of Community Medicine, Vardhman Mahavir Medical College and Safdarjung Hospital, New Delhi, India

⁴School of Medicine, University of Limerick, Ireland.

Corresponding Author

*Adjunct Clinical Associate Professor, Public Health Masters Program, School of Medicine, University of Limerick, Ireland.

Email ID - sonu.goel@ul.ie

Author's Contributions

AD participated in conceptualization, methodology and variable selection, data analysis and writing of manuscript draft. YB participated in conceptualization, methodology and variable selection, writing of manuscript draft, and reviewing and editing manuscript. AS participated in conceptualization, methodology and variable selection, data analysis and reviewing and editing manuscript. JK participated in writing of manuscript draft, reviewing and editing manuscript and final approval of manuscript. SG participated in conceptualization, methodology and variable selection, reviewing and editing manuscript draft and final approval of manuscript.

Abstract

Undiagnosed hypertension extracts significant social cost, with money spent on complications already accounting for one-fifths of total health expenditure. Widespread socioeconomic disparities and inequity in health care access between rural and urban areas is expected. It is important to identify the different factors associated with undiagnosed hypertension in the working age population (15-49-years) residing in urban and rural areas, both of whom are vital to the economic development of our country. Data from the National Family Health Survey-5 (2019-21) for men and women aged 15-49 years was extracted and analysed. Operational definitions were prepared to identify known and undiagnosed hypertension. Distribution of undiagnosed hypertension according to sociodemographic, anthropometric and health-related behaviour was studied using frequencies and weighted proportions. Choropleth maps were used to depict state-wise distribution of undiagnosed hypertension. Multivariable logistic regression was used to find risk factors and protective factors for undiagnosed hypertension for men and women in rural and urban areas. The prevalence of undiagnosed hypertension was 11.7% among men and 7.2% among women. The proportion of men with undiagnosed hypertension (66.3%) was significantly higher than the proportion of women (41.4%). Urban-rural differences were noted in various states. Education and empowerment of rural women through provision of means of socioeconomic enhancement and strengthening of community-based screening and referral under the national programme were some of the major policy implications of our findings. Future research is warranted in areas such as health insurance coverage, working away from home, owning a mobile telephone and other interventions to improve health-seeking behaviour in the rural areas.

Keywords

Hypertension, Noncommunicable diseases, Delayed diagnosis, Health care seeking behaviour, Diagnostic screening programs

Introduction

Hypertension is the commonest cardiovascular disorder and a major public health problem across the world. Hypertension is usually essential or primary, silent, and asymptomatic [1]. It has a significant social cost, with money spent on their complications accounting for one-fifths of total health expenditure [2]. According to the World Health Organization, an estimated 1.28 billion adults aged 30-79 years worldwide have hypertension, among whom two-thirds live in low and middle-income countries [3]. Hypertension is a major cause of premature death worldwide. Only one in five adults with hypertension have it under control [3]. The rule of halves was coined in 1970s in the USA [4] and has been reported previously from surveys in India [5,6]. Pooled estimates from population representative studies from across the world, show that 51% among men and 41% among women with hypertension, were undiagnosed. The proportion of undiagnosed hypertension increased to 55% among men and 46% among women in East and South-east Asia [7].

The Global Burden of Hypertension study has highlighted that of the global burden of 212 million Disability-adjusted life years (DALYs) related to hypertension, 18% occurred in India in 2015 [8]. It is a leading risk factor for cardiovascular disease, which accounted for 23% of total deaths and 32% of adult deaths in 2010–2013 [9]. India has committed to meet the Sustainable Development Goals (SDG) target of reducing premature mortality from non-communicable diseases (NCDs) by one-third by 2030 when compared to 2015 [10]. Another nationwide study found that 42.3% of the individuals with hypertension were previously undiagnosed and the proportion was 12.4% higher in rural areas as compared to urban areas [11]. Among Indians suffering from hypertension, only one-fourth in rural areas and one-third in urban areas, are being treated while only one-tenth in rural and one-fifth in urban areas, have their blood pressure under control [6,12,13].

Since 2018, India's population has entered the phase where the working age population is larger than the dependent population, and this opportunity of demographic dividend will last until 2055 [13]. At the same time, rapid changes in dietary habits and lifestyle has resulted in an upsurge of non-communicable diseases over the past three decades [14]. The COVID-19 pandemic may have compounded the situation with more resources being diverted toward COVID management and away from non-COVID services which includes non-communicable disease management [15]. It is important to identify the factors associated with undiagnosed hypertension in this population who are vital to the economic development of our country and will also contribute to the growing burden on the health care system in the years to come. Given the heterogeneity in the sociodemographic conditions as well as the variability in primary health care delivery across states in India, it is hypothesized that there would be considerable inter-state variations in prevalence of undiagnosed hypertension. Widespread socioeconomic disparities and inequity in health care access between rural and urban areas in a state is also expected. Hence, estimates from both rural and urban areas in a state are required to identify priority areas for intervention, evaluate national programme delivery and guide policy formulation. The overall aim of this study was to determine the burden of undiagnosed hypertension, its associated factors and its distribution across rural and urban areas among men and women aged 15-49 years in India (2019-21).

Methods

Study Design

A cross-sectional analytical study involving secondary data from the fifth round of National Family Health Survey (NFHS) [16].

Setting

The NFHS is a large-scale, multi-round survey conducted in a representative sample of households throughout India. It is conducted under the stewardship of the Ministry of Health and Family Welfare, Government of India. International Institute for Population Sciences, Mumbai, is the nodal agency for all the rounds of NFHS. Technical assistance for NFHS-5 was provided by the ICF International (Maryland USA) through the Demographic and Health Surveys Program, which is supported by USAID. First phase of NFHS-5 was conducted from June 17, 2019 to January 30, 2020 covering 17 states and 5 union territories and second phase was from January 2, 2020 to April 30, 2021 covering 11 states and 3 union territories. The survey covered 636 699 households, 724 115 women (15-49 years), and 101 839 men (15-54 years).

The NFHS-5 sample is a stratified two-stage sample. Villages in rural areas and census enumeration blocks in urban areas were the primary sampling units. Four questionnaires (titled household, woman, man, and biomarker) were developed in 18 local languages and data collection was done using computer assisted personal interviewing. This enabled International Institute for Population Sciences and ICF to run extensive data quality checks on the data from the field and to provide real-time feedback to field agencies and teams to help improve data quality.

Blood pressure was measured for all women and men aged between 15 and 49 years using an Omron® blood pressure monitor to determine the prevalence of hypertension. Blood pressure measurements for each respondent were taken three times with an interval of five minutes between readings.

Study Population

For state-wise prevalence of undiagnosed hypertension, we included all men and women in the 15-49 years age group from NFHS-5 (2019-21). For proportion of undiagnosed hypertension

among men and women with hypertension and its associated factors, we included the subpopulation with hypertension (diagnosed and undiagnosed).

Operational Definitions of Undiagnosed and Known Hypertension

We derived known hypertension as presence of any one of the following three criteria: i) currently reported as having hypertension, ii) told by any healthcare provider that they had high blood pressure (BP) on two or more occasions, or iii) currently taking medication to lower BP.

We derived undiagnosed hypertension if the respondents fulfilled all the following four: i) average of second and third systolic BP reading ≥ 140 and/or diastolic BP reading ≥ 90 mm Hg, ii) not reported as currently having hypertension, iii) not told by any healthcare provider previously to have had high BP, and iv) reported as not taking any medication to lower BP.

Data Variables

We extracted the variables required to classify the status of hypertension (known/undiagnosed). In addition, socio-demographic characteristics, anthropometric data and data on health-related behaviours was extracted.

Analysis

Software

Database was constructed, cleaned with Microsoft Excel (Microsoft, Redmond, WA, USA) and imported to STATA (version 16, copyright 1985–2011 Stata Corp LP USA) and Datawrapper developed by Datawrapper GmbH for analysis [17].

Statistics

We depicted the prevalence of undiagnosed hypertension using percentages and 95% CI. Descriptive analysis of sociodemographic, anthropometric and health-related behaviour variables was expressed for men and women residing in rural and urban areas with undiagnosed hypertension separately. Log binomial regression model has been applied to determine the

factors associated with undiagnosed hypertension among all people with hypertension. The associations were summarized using adjusted odds ratios and 95% CI. Owing to the large sample size, a cut-off for public health significance was set for the adjusted odds ratios [aOR ≤ 0.8 & ≥ 1.3]. Choropleth figures have been used to depict state-wise proportion of undiagnosed hypertension in rural and urban areas among all men and women with hypertension. All the analysis has been weighted for the multistage sampling design and hence, weighed estimates have been provided.

Ethical Statement

The study was ethically approved by the Institute's Ethical Committee, Postgraduate Institute of Medical Education and Research (PGIMER), Chandigarh (IEC-08/2022-2535 dated 17.08.2022). Data was accessed and extracted on 3rd September, 2022. The authors did not have access to any individual identifiers.

Results

Burden of undiagnosed hypertension

Out of a total of 1,01,839 men, the prevalence of undiagnosed hypertension was 11.7%. The proportion of men with undiagnosed hypertension among all men with hypertension was 66.3% (Figure 1). The proportion was similar between rural (66.2%) and urban (66.5%) areas.

Figure 1 - : Burden of Undiagnosed Hypertension among men and women aged 15-49 years from NFHS 5 (2019-2021)

Table 1 describes the distribution of men with undiagnosed hypertension according to various socio-demographic, anthropometric and health related behaviour variables.

Out of 7,24,115 women, the prevalence of undiagnosed hypertension was 7.2%. The proportion of undiagnosed hypertension among all women with hypertension was 41.4% (Figure 1). The proportion was also similar between rural (41.3%) and urban areas (41.8%). Table 2 describes the distribution of women with undiagnosed hypertension according to various socio-demographic, anthropometric and health related behaviour variables.

Rural-urban differences in distribution of undiagnosed hypertension among men

The proportion of undiagnosed hypertension was higher among the 15-19 year (75.9% vs 62.2%) group in the urban areas as compared to the rural areas. Proportion of undiagnosed hypertension among Muslim men was lower (58.4%), especially in the rural areas (54%). The proportion of men belonging to scheduled tribes with undiagnosed hypertension was higher in rural areas than in urban areas (75% vs 66.9%). The proportion of undiagnosed hypertension was higher among men from rural areas who had been widowed/separated/divorced as compared to urban areas (75.5% vs 52%). A higher proportion of undiagnosed hypertension was found among men who were uneducated (72.6% vs 66.4%) or unemployed (72.6% vs 64.2%) in the urban areas as compared to the rural areas. In the East region, proportion of undiagnosed hypertension was higher among men from rural areas (54.3% vs 43.3%). It was higher among men who haven't used any type of health facility in recent times (68.5%). The proportion was higher among men who had used a public facility in urban areas as compared to the rural areas (66.1% vs 61.5%).

<p>Table 1: Proportional distribution (with 95% confidence interval) of men with undiagnosed hypertension according to various socio-demographic, anthropometric and health related behaviour variables among all men with hypertension in the 15-49 years age group in India NFHS-5, 2019-21.</p>

Variables	Proportion of men with undiagnosed hypertension (95% CI)			Total (N)
	Total	Rural	Urban	
Total	66.3 (65-67.6)	66.2 (64.8-67.6)	66.5 (63.7-69.1)	17936
Age group				
15-19	66.9 (60.6-72.7)	62.2 (55.8-68.3)	75.9 (63-85.3)	895
20-29	69.9 (67-72.5)	68.9 (65.9-71.7)	71.7 (65.7-77)	3659
30-39	69.2 (66.8-71.5)	69.3 (66.9-71.5)	69.1 (63.9-73.8)	6090
40-49	62.3 (60.2-64.3)	63 (60.7-65.2)	61.1 (56.9-65)	7292
Cast/Tribe				
Scheduled caste	68.4 (65.4-71.3)	67.1 (63.9-70.1)	70.8 (64.3-76.6)	3418
Scheduled tribe	73.5 (70.3-76.5)	75 (71.8-78)	66.9 (56.2-76.1)	3662
OBC	66.4 (64.3-68.4)	66.8 (64.7-68.8)	65.7 (61.5-69.6)	6614
Others	62.3 (59.5-64.9)	60.5 (57.4-63.5)	64.7 (59.7-69.3)	4242
Religion				
Hindu	67.5 (66-68.9)	67.6 (66.1-69.1)	67.2 (64-70.3)	13419
Muslim	58.4 (54.5-62.2)	54 (49.2-58.8)	63.3 (56.9-69.3)	1748
Christian	69.2 (63.2-74.6)	70.4 (63.2-76.8)	67.1 (56.4-76.4)	1516
Others	66 (59.7-71.8)	67.5 (60.5-73.8)	63 (50.3-74.2)	1253
Marital Status				
Never in union	70.3 (67.3-73.1)	69.2 (66.2-72.1)	71.8 (66-76.9)	3704
Currently married	65.2 (63.7-66.7)	65.3 (63.7-66.9)	65.1 (62-68.2)	13898
Widowed/Divorced /Separated	69.5 (60-77.6)	75.5 (64.1-84.1)	52 (35.4-68.2)	334
Education				
No education/Pre- primary education	67.8 (64.3-71.2)	66.4 (62.6-70.1)	72.6 (64.2-79.7)	2125
Primary	67.1 (63.4-70.6)	67 (63.1-70.8)	67.2 (58.5-74.9)	2214
Secondary Education	66.7 (64.9-68.4)	66.8 (64.9-68.6)	66.5 (62.7-70.1)	10273
Higher Education	63.8 (60.5-66.9)	62.9 (59.2-66.5)	64.5 (59.1-69.5)	3324
Occupation				
No Occupation	67.8 (63.3-71.9)	64.2 (59.5-68.7)	72.6 (64.1-79.8)	1729
Professional/ Clerical	62.7 (57.7-67.4)	66.7 (61.1-71.8)	60.1 (52.9-66.9)	1514
Sales	63.9 (59.4-68.2)	60.2 (54.6-65.6)	66.9 (60.2-72.9)	1801
Services/household and domestic	67.5 (62-72.6)	70.5 (64.4-76)	65 (56.2-73)	1454
Agricultural	67 (64.9-69)	67 (64.8-69.1)	67.2 (57.5-75.7)	5880
Skilled, unskilled manual and Others	66.7 (64.3-69)	66.2 (63.6-68.6)	67.4 (63-71.5)	5558
Wealth Index				
Poorest	63.4 (60.3-66.5)	63.7 (60.5-66.7)	59.1 (39.7-76.1)	2981
Poorer	67.7 (65-70.3)	67.5 (64.8-70.1)	68.7 (58.3-77.6)	3723
Middle	66.7 (64-69.3)	66 (63.2-68.7)	68.4 (62.1-74.2)	3906
Richer	67.1 (64.2-69.9)	68.2 (65-71.3)	65.8 (60.7-70.6)	3773
Richest	65.7 (62.3-68.9)	64.9 (60.1-69.3)	66 (61.7-70.1)	3553
Work at home or away				

At home	67.6 (66.2-69)	67.6 (66.1-69.1)	67.6 (64.7-70.4)	15328
Away	58.1 (54.5-61.6)	58.6 (55.1-62)	56.6 (47.6-65.2)	2608
Region				
North	66.6 (64.8-68.4)	66.6 (64.3-68.8)	66.6 (63.5-69.5)	3743
Central	71.3 (69.5-73.1)	71.2 (69.2-73)	71.8 (67.7-75.5)	3881
East	51.3 (48.1-54.5)	54.3 (50.8-57.7)	43.3 (36.7-50.2)	2338
Northeast	55 (52.2-57.8)	54.8 (51.6-58)	55.7 (50-61.3)	3167
West	74.8 (71.1-78.3)	74.9 (71.2-78.3)	74.8 (67.6-80.8)	1614
South	71.2 (68.9-73.4)	71.6 (69-74.1)	70.6 (66.4-74.5)	3193
Use of Internet				
Never	67.3 (65.5-69.1)	67.4 (65.5-69.3)	67 (62.4-71.4)	8032
Yes	66 (64-68)	65.4 (63.2-67.5)	66.6 (63.1-70)	9427
Owns a mobile telephone				
Yes	66 (61.4-70.4)	66.1 (61.1-70.8)	65.6 (52.8-76.4)	1364
No	66.7 (65.2-68.1)	66.6 (65.1-68)	66.8 (63.9-69.6)	16095
Usage of any type of tobacco				
No	68.2 (66.4-69.9)	67.5 (65.6-69.4)	69.2 (65.9-72.4)	9286
Yes	64.1 (62.1-66.1)	64.9 (62.9-66.9)	62.4 (57.8-66.8)	8650
Current alcohol usage				
No	66.1 (64.5-67.7)	65.8 (64.1-67.5)	66.7 (63.4-69.8)	11531
Less than once a week	64.4 (60.3-68.4)	63.9 (59.7-67.8)	65.3 (56.4-73.3)	2274
Once a week	69.7 (66.4-72.9)	70.1 (66.6-73.4)	69.1 (62.2-75.2)	2924
Everyday	63.9 (57.6-69.7)	66.2 (60.7-71.3)	59.9 (45.8-72.5)	1207
Covered by Health Insurance				
No	66 (64.2-67.7)	65.2 (63.3-67)	67.2 (63.7-70.5)	10977
Yes	66.9 (64.8-68.9)	67.8 (65.7-69.9)	65.1 (60.6-69.3)	6959
Type of health facility recently used				
None	68.5 (66.8-70.1)	68.6 (66.9-70.2)	68.4 (64.9-71.6)	12261
Public facility	63.1 (60-66.1)	61.5 (58.3-64.6)	66.1 (59.5-72)	3513
Private facility	61 (57.2-64.6)	62.5 (58.5-66.4)	58.7 (51.4-65.6)	1999
Other	59.8 (46.4-71.8)	57.2 (42.1-71)	69.3 (42.4-87.4)	163
Anaemia				
Severe	50 (31.1-69)	48.8 (29.5-68.5)	80.7 (27.3-97.9)	50
Moderate	59.4 (52.7-65.7)	60 (52.3-67.2)	57.6 (44.1-70.1)	600
Mild	62.9 (59.5-66.2)	63.7 (60.1-67.1)	61 (53.1-68.4)	2709
Not anaemic	68.3 (66.8-69.8)	68.2 (66.6-69.7)	68.6 (65.4-71.5)	13659
BMI				
Underweight	68.4 (63.8-72.7)	67 (62.4-71.3)	72.7 (60.4-82.2)	1296
Normal weight	66.6 (64.9-68.4)	66.5 (64.6-68.3)	67 (63.1-70.6)	9871
Overweight	68.8 (66.5-71)	67.9 (65.3-70.4)	70 (65.8-73.8)	5290
Obese	57.8 (51.7-63.7)	60.3 (53.9-66.3)	55.6 (45.6-65.1)	1318

Waist to Hip Ratio				
Less or equal to 0.9 for men	67.6 (65.7-69.5)	66.6 (64.5-68.6)	69.5 (65.4-73.4)	7886
(Cut offs > 0.9 for men)	65.9 (64.1-67.7)	66.4 (64.5-68.3)	65.2 (61.5-68.7)	9917

Rural-urban differences in distribution of undiagnosed hypertension among women

Table 2 also depicts the rural-urban differences in proportion of undiagnosed hypertension among women. The proportion of women with undiagnosed hypertension was higher across all age groups (67.9% vs 37.5% in the 20-29 year age group) in rural areas as compared to urban areas. The proportion of undiagnosed hypertension was lower among women belonging to the richest wealth index in the rural areas as compared to the urban areas (35.6% vs 40.5%). The proportion was higher among women using any type of tobacco (46.1% vs 40.3%) or alcohol (65.2% vs 54.2%) in the rural areas than in the urban areas. There was a higher proportion of undiagnosed hypertension among unemployed women in the urban areas as compared to the rural areas (42.4% vs 38.4%).

Table 2: Proportional distribution (with 95% confidence interval) of women with undiagnosed hypertension according to various socio-demographic, anthropometric and health related behaviour variables among all women with hypertension in the 15-49 years age group in India NFHS-5, 2019-21.				
Variables	Proportion of women with undiagnosed hypertension (95% CI)			Total
	Total	Rural	Urban	
Total	41.4 (41.1-41.8)	41.3 (40.9-41.7)	41.8 (41-42.5)	126564
Age group				
15-19	42 (40.4-43.6)	59.6 (57.9-61.3)	46.3 (42.6-50.1)	6584
20-29	33.7 (32.9-34.5)	67.9 (67-68.7)	37.5 (35.8-39.3)	26562
30-39	43 (42.3-43.7)	57.7 (57-58.5)	44.5 (43.1-45.9)	40204
40-49	43.9 (43.3-44.5)	54.5 (53.8-55.1)	41.1 (40-42.3)	53214
Cast/Tribe				
Scheduled caste	39.9 (39-40.7)	39.6 (38.7-40.5)	40.4 (38.7-42.2)	24569
Scheduled tribe	55.9 (54.7-57.1)	57 (55.8-58.3)	50.2 (46.5-53.9)	22796
OBC	41.2 (40.6-41.7)	40.8 (40.2-41.4)	42 (40.9-43.1)	46997
Others	38.7 (37.9-39.5)	37 (36.1-37.9)	41.2 (39.8-42.6)	32202
Religion				

Hindu	42 (41.5-42.4)	42.2 (41.7-42.7)	41.5 (40.6-42.4)	92359
Muslim	39.9 (38.8-41)	36.8 (35.5-38.1)	44.2 (42.3-46.1)	15844
Christian	43.8 (41.4-46.2)	45.4 (42.7-48.1)	41.1 (36.7-45.7)	9451
Others	34.3 (32.7-36)	33.8 (32.1-35.4)	35.6 (31.8-39.7)	8910
Marital Status				
Never in union	46.7 (45.4-47.9)	46 (44.6-47.4)	47.9 (45.4-50.3)	12236
Currently married	40.6 (40.2-41)	40.5 (40-40.9)	40.9 (40.1-41.8)	105922
Widowed/Divorced/Separated	45 (43.5-46.6)	46.1 (44.4-47.8)	43.2 (40.4-46.1)	8406
Education				
No education/Pre-primary education	45.2 (44.6-45.9)	45.7 (45-46.4)	43.3 (41.6-45.1)	38628
Primary	42.8 (41.8-43.9)	42.5 (41.4-43.6)	43.7 (41.4-46)	18254
Secondary Education	39.5 (38.9-40)	38.2 (37.5-38.8)	41.6 (40.4-42.7)	56146
Higher Education	37.3 (36.1-38.5)	34.3 (32.8-35.9)	39.5 (37.8-41.3)	13536
Occupation				
No Occupation	39.8 (38.6-41.1)	38.4 (37-39.7)	42.4 (40-44.9)	12344
Professional/ Clerical	45.5 (39.1-52)	46.4 (38.9-54)	44.8 (35.2-54.8)	625
Sales	46.6 (38.4-55.1)	46 (37.5-54.6)	47.2 (33.7-61.2)	456
Services/household and domestic	47.4 (41.9-53.1)	48.1 (41-55.2)	46.9 (38.7-55.4)	769
Agricultural	51 (48.6-53.3)	51 (48.6-53.4)	50.8 (40.6-61)	3254
Skilled, unskilled manual and Others	42.8 (39.3-46.3)	43.5 (39.4-47.7)	41.6 (35.5-48)	1543
Wealth Index				
Poorest	43.9 (43-44.7)	43.9 (43.1-44.8)	42.7 (37.9-47.7)	23203
Poorer	41.8 (41-42.6)	41.4 (40.6-42.3)	44.3 (41.4-47.3)	26910
Middle	41.8 (41-42.6)	41.4 (40.5-42.3)	42.9 (41-44.9)	26884
Richer	41.4 (40.5-42.2)	40.6 (39.5-41.6)	42.3 (40.9-43.7)	25718
Richest	39 (38.1-39.9)	35.6 (34.4-36.8)	40.5 (39.4-41.7)	23849
Work at home or away				
At home	43 (42-44.1)	42.6 (41.4-43.8)	43.9 (41.7-46)	17506
Away	34.9 (31.4-38.5)	33.8 (30.3-37.5)	36.8 (29.7-44.5)	1485
Region				
North	34.4 (33.7-35.1)	33.5 (32.7-34.3)	35.9 (34.7-37.1)	27838
Central	42.5 (41.8-43.1)	42.7 (42-43.4)	41.8 (40.3-43.3)	29848
East	34.1 (33.2-34.9)	34.8 (33.9-35.7)	31.7 (29.8-33.6)	20039
Northeast	37.5 (36.5-38.6)	38 (36.8-39.1)	35.9 (33.7-38.3)	19857
West	52.5 (51-53.9)	53.6 (52-55.2)	51.1 (48.6-53.6)	10328
South	48.1 (47.1-49)	49.1 (48-50.3)	46.7 (45.2-48.3)	18654
Use of Internet				
Never	43.8 (42.6-44.9)	43.1 (41.9-44.4)	45.6 (43-48.3)	13847
Yes	38.4 (36.3-40.5)	36.7 (34.3-39.2)	39.9 (36.6-43.3)	5144
Owns a mobile telephone				
Yes	46.5 (45-48)	45.6 (44-47.1)	49.6 (46-53.2)	8112
No	39.2 (37.8-40.6)	38.2 (36.6-39.8)	40.6 (38.1-43.1)	10879
Usage of any type of tobacco				

No	41.2 (40.8-41.6)	40.9 (40.5-41.4)	41.8 (41-42.6)	116150
Yes	44.9 (43.4-46.3)	46.1 (44.5-47.6)	40.3 (36.6-44.2)	10414
Current alcohol usage				
No	41.3 (40.9-41.7)	41.1 (40.6-41.5)	41.7 (41-42.5)	123097
Less than once a week	53.6 (49.4-57.7)	55.3 (50.9-59.7)	45.4 (33.5-57.9)	1525
Once a week	57.4 (53.1-61.6)	59.2 (54.8-63.5)	45.5 (32.2-59.5)	1413
Everyday	63.7 (57.5-69.4)	65.2 (59.1-70.9)	54.2 (34.2-73)	529
Covered by Health Insurance				
No	40.7 (40.3-41.2)	40.1 (39.6-40.6)	41.9 (41-42.8)	85298
Yes	43.1 (42.4-43.7)	43.8 (43.1-44.6)	41.4 (40.1-42.8)	41266
Type of health facility recently used				
None	44.8 (44.3-45.3)	45.1 (44.5-45.6)	44.4 (43.4-45.4)	78135
Public facility	37 (36.2-37.7)	36.4 (35.5-37.2)	38.5 (36.8-40.1)	30984
Private facility	35.3 (34.3-36.3)	34 (32.9-35.1)	37.1 (35.3-39)	17005
Other	33.5 (28-39.6)	30.8 (24.7-37.7)	39.8 (28.6-52.2)	440
Anaemia				
Severe	31.5 (29-34.1)	30.9 (28.3-33.6)	32.7 (27.4-38.5)	2723
Moderate	38.4 (37.7-39.2)	38.5 (37.7-39.3)	38.2 (36.7-39.8)	31898
Mild	40.3 (39.5-41.1)	40.2 (39.4-41.1)	40.5 (38.9-42.1)	28685
Not anaemic	44.6 (44-45.1)	44.1 (43.4-44.7)	45.5 (44.3-46.6)	57550
BMI				
Underweight	39.6 (38.5-40.7)	39.5 (38.3-40.6)	40.1 (37.1-43.2)	12735
Normal weight	42.4 (41.8-42.9)	42.2 (41.6-42.8)	42.7 (41.5-44)	62736
Overweight	44.5 (43.7-45.3)	44.5 (43.6-45.5)	44.5 (43.1-45.9)	31618
Obese	40.4 (39.3-41.6)	40.3 (38.9-41.8)	40.5 (38.7-42.3)	13881
Waist to Hip Ratio				
Less or equal to 0.9 for men and 0.8 for women	41.2 (40.6-41.8)	41.2 (40.6-41.9)	41.1 (39.8-42.4)	45257
(Cut offs > 0.9 for men and >0.8 for women)	42.1 (41.6-42.6)	41.7 (41.2-42.3)	42.8 (41.9-43.8)	80156

State-wise distribution of undiagnosed hypertension among men

State-wise distribution of proportion of men with undiagnosed hypertension among all men with hypertension is visualized in Figure 1. Chhattisgarh reported the highest proportion of 83.2% (Rural-86.9%; Urban-70.3%), closely followed by Madhya Pradesh with 81.7% (Rural-82.5%; Urban-79.8%). As many as 13 states & Union Territories reported greater than 70% proportion of undiagnosed hypertension among males with hypertension in the 15-49 year age group. Several states show wide rural-urban divides. The widest urban-rural divides are seen in NCT of Delhi (Rural-93.7%; Urban-70.3%), Puducherry (Rural-88.4%; Urban-55.5%),

Nagaland (Rural-70.9%; Urban-48%), Bihar (Rural-45.9%; Urban-29.7%), Andaman and Nicobar islands (Rural-68.6%; Urban-42%) and Goa (Rural-54.5%; Urban-68.4%). States with the lowest proportion of men with undiagnosed hypertension are Ladakh, Bihar, Tripura, and Jammu & Kashmir.

Figure 2: State-wise proportional distribution of men with undiagnosed hypertension among all men with hypertension in the 15-49 year age group in India;

A-Overall; B-Urban; C-Rural

State-wise distribution of undiagnosed hypertension among women

State-wise distribution of proportion of women with undiagnosed hypertension among all women with hypertension is visualized in Figure 2. Chhattisgarh reported the highest proportion of 64.3% (Rural-67.7%; Urban-54.5%) followed by Karnataka with 56.2% (Rural-56.3%; Urban-56.1%) and Lakshadweep with 56.2% (Rural-42.8%; Urban-59.1%).

Maharashtra, Telengana, Gujarat and Arunachal Pradesh reported greater than 50% proportion of undiagnosed hypertension among women with hypertension in the 15-49 year age group. The rural-urban divide is also prominent in this distribution as seen in states like Chhattisgarh (Rural-67.7%; Urban-54.5%), NCT of Delhi (Rural-30.4%; Urban-46.2%), Sikkim (Rural-50.4%; Urban-33.5%), Nagaland (Rural-51.9%; Urban-33%), Mizoram (Rural-53.2%; Urban-39%) and Ladakh (Rural-23.5%; Urban-7.2%). States with the lowest proportion of women with undiagnosed hypertension are Ladakh, Bihar, Haryana, Himachal Pradesh and Punjab.

Figure 3: State-wise proportional distribution of women with undiagnosed hypertension among all women with hypertension in the 15-49 year age group in India;

A-Overall; B-Urban; C-Rural

Factors associated with undiagnosed hypertension

The proportion of men with undiagnosed hypertension (66.3%) was significantly higher than the proportion of women (41.4%) (p-value <0.001). Multivariable logistic regression analysis was done to calculate adjusted odds for factors associated with undiagnosed hypertension in men and women in the 15-49 year age group. Table 3 summarizes the protective factors and risk factors associated with undiagnosed hypertension among men and women across rural and urban areas.

Among men, age, usage of tobacco products, recent usage of public or private health facility and health insurance coverage were found to be significant predictors for undiagnosed hypertension in urban areas. In rural areas, the important predictors were caste/tribe, religion, anaemia, working away from home, owning a mobile telephone and recent usage of public or private health facility. Among women, caste/tribe, marital status, wealth index, anaemia and recent usage of public or private health facility were found to be significant predictors for undiagnosed hypertension in urban areas. In rural areas, the important predictors were caste/tribe, daily alcohol usage, age, religion, higher education, wealth index, anaemia and recent usage of any health facility.

Table 3: Summary of risk factors and protective factors associated with undiagnosed hypertension among men and women aged 15-49 years from NFHS 5 (2019-21)				
			Rural aOR (95% C.I.)**	Urban aOR (95% C.I.)**
Men	Risk factors	Age		
		40-49 years	Ref	
		15-19 years	0.78 (0.61-1.01)	1.37 (0.88-2.13)
		20-29 years	1.26 (1.11-1.45)	1.43 (1.15-1.79)*
		30-39 years	1.27 (1.15-1.4)	1.39 (1.2-1.61)*
		Caste/Tribe		
	Others	Ref		
	Scheduled Tribe	1.45 (1.25-1.68)*	0.92 (0.7-1.21)	
	Scheduled Caste	0.95 (0.83-1.09)	1.02 (0.83-1.24)	
	OBC	1.0 (0.89-1.12)	1.12 (0.95-1.32)	
	Protective factors	Usage of any tobacco product	0.89 (0.79-1.01)	0.77 (0.64-0.92)*
		Anemia		
Not anemic		Ref		

		Mild	0.86 (0.77-0.96)	0.76 (0.63-0.92)*			
		Moderate	0.74 (0.6-0.91)*	0.65 (0.45-0.93)*			
		Severe	0.39 (0.2-0.75)*	0.56 (0.07-4.33)			
		Religion					
		Hindu	Ref				
		Muslim	0.61 (0.52-0.71)*	0.97 (0.79-1.19)			
		Christian	0.95 (0.79-1.14)	1.02 (0.75-1.38)			
		Others	0.83 (0.7-0.97)	0.81 (0.6-1.09)			
		Works away from home					
		Works at home	Ref				
		Works away from home	0.75 (0.68-0.84)*	0.87 (0.72-1.05)			
		Owns a mobile telephone					
		No	Ref				
		Yes	0.8 (0.69-0.94)*	1.32 (0.92-1.88)			
		Recent usage of health facility					
		None	Ref				
		Public	0.68 (0.62-0.76)*	0.74 (0.63-0.87)*			
		Private	0.67 (0.58-0.76)*	0.61 (0.5-0.73)*			
		Others	0.87 (0.59-1.28)	0.98 (0.41-2.35)			
		Health insurance coverage					
		No	Ref				
		Yes	1.1 (1.01-1.2)	0.8 (0.71-0.93)*			
		Women	Risk factors	Caste/Tribe			
				Others	Ref		
				Scheduled Tribe	2 (1.91-2.1)*	1.3 (1.17-1.42)*	
Scheduled Caste	1 (0.95-1.04)			0.98 (0.92-1.06)			
OBC	1.15 (1.11-1.19)			1.11 (1.05-1.17)			
Current alcohol usage							
None	Ref						
Less than once a week	1.19 (1.05-1.35)			1.02 (0.78-1.32)			
Once a week	1.21 (1.07-1.37)			1.08 (0.76-1.53)			
Everyday	1.39 (1.13-1.7)*			1.46 (0.8-2.65)			
Protective factors	Age						
	40-49 years		Ref				
	15-19 years		0.64 (0.59-0.7)*	0.93 (0.8-1.08)			
	20-29 years		0.64 (0.61-0.67)*	0.83 (0.77-0.9)			
	30-39 years		0.91 (0.88-0.94)	1.06 (1-1.12)			
	Marital Status						
	Never in union		Ref				
	Currently married		0.6 (0.56-0.64)*	0.68 (0.62-0.76)*			
	Widowed/Separated/Divorced		0.58 (0.53-0.63)*	0.67 (0.59-0.77)*			
	Religion						
Hindu	Ref						
Muslim	0.76 (0.72-0.8)*	0.93 (0.86-0.99)					
Christian	0.8 (0.79-0.9)*	0.81(0.72-0.91)					
Others	0.82 (0.77-0.87)	0.81 (0.73-0.91)					
Education							

	No education/Pre-primary education	Ref	
	Primary education	0.89 (0.85-0.93)	0.98 (0.9-1.07)
	Secondary education	0.82 (0.79-0.85)	0.88 (0.82-0.94)
	Higher education	0.76 (0.71-0.81)*	0.85 (0.78-0.93)
	Wealth Index Quintile		
	Poorest	Ref	
	Poorer	0.87 (0.84-0.91)	0.87 (0.74-1.03)
	Middle	0.83 (0.79-0.87)	0.81 (0.69-0.95)
	Richer	0.79 (0.75-0.83)*	0.77 (0.66-0.9)*
	Richest	0.75 (0.71-0.8)*	0.73 (0.63-0.86)*
	Anemia		
	Not anemic	Ref	
	Mild	0.84 (0.81-0.87)	0.82 (0.77-0.87)
	Moderate	0.78 (0.75-0.81)*	0.76 (0.72-0.8)*
	Severe	0.54 (0.49-0.6)*	0.5 (0.42-0.59)*
	Recent usage of health facility		
	None	Ref	
	Public	0.72 (0.7-0.75)*	0.78 (0.73-0.83)*
	Private	0.61 (0.59-0.64)*	0.69 (0.65-0.74)*
	Others	0.56 (0.43-0.71)*	0.7 (0.47-1.05)

*Factors of relevant public health significance

**Adjusted for all socioeconomic, anthropometric and health-related behaviour variables

Discussion

Burden of undiagnosed hypertension

The prevalence of undiagnosed hypertension was found to be 11.7% among men and 7.2% among women. The proportion of undiagnosed hypertension among all those with hypertension was 66.3% among men and 41.4% among women. This means that a large section of the population with hypertension have remained undiagnosed in comparison to a proportion of less than 10% in countries with high treatment coverage [7]. Around 25-30% of women and men with hypertension in many sub-Saharan African and in some countries in south, and southeast Asia had uncontrolled hypertension who were not diagnosed or treated. Pooled analysis of several studies from India showed that around 20% men and women with uncontrolled hypertension had not been diagnosed or treated [7]. The differences between the estimates can

be attributed to the difference in study definitions as well as the difference in study population, where the pooled analyses reported data for ages between 30 and 79 years. The findings indicate that it is possible to bring the proportion of undiagnosed hypertension far below current levels with expansion of screening, diagnostic and treatment coverage. It also reinforces the need to understand the factors behind undiagnosed hypertension so as to be able to bring more people under the umbrella of treatment and control.

Previous analysis of nationwide data on undiagnosed hypertension is scarce in India. An analysis of undiagnosed hypertension among women from NFHS-4 has since been retracted due to errors in classification [18]. A study on hypertension screening, awareness, treatment and control from the NFHS 4 dataset by Prenissl et al. found that 55.3% of those with hypertension were unaware of their status [6]. Analysis of the Longitudinal Ageing Study of India, Wave-1 (2017-18) which included adults above the age of 45 years, found the proportion of undiagnosed hypertension among all those with hypertension to be 42.3% (48.5% for men and 37.5% for women) [11]. The difference in findings can be explained by the different age groups of the respective study populations, but once again, it indicates that a sizeable proportion among even the elderly have remained undiagnosed.

Rural-urban context in undiagnosed hypertension

Our study found an even distribution of undiagnosed hypertension between urban and rural areas in both men and women. This contrasts with the findings from the analysis of the LASI data [11], where prevalence of undiagnosed hypertension was 12.4% higher in rural areas as compared to urban areas. However, our state-level analysis found rural-urban differences in distribution of undiagnosed hypertension for both men and women with Delhi, Puducherry, Nagaland, Chhattisgarh, Bihar and Sikkim reporting around 20% higher prevalence of undiagnosed hypertension in rural areas.

The predictors for undiagnosed hypertension in urban and rural areas varied. Among men, ages 20-29 and 30-39 years were more likely to have undiagnosed hypertension in the urban areas. This may be an indication of the effect of the increasing adoption of unhealthy lifestyles and diet among the urban youth and middle-aged male population. This may also be due to the existing perception that hypertension is primarily a disease of the elderly, and the younger adults do not feel the need to be screened or tested. Conversely, rural women in the 15-19- and 20-29-year age groups were less likely to have undiagnosed hypertension. The fact that all pregnant women receiving antenatal care are screened for hypertension may have contributed to the higher proportions of known hypertension in these age groups. Though, community based and opportunistic screening for hypertension has been implemented across the country through the National Programme for Prevention & Control Of Cancer, Diabetes, Cardiovascular Diseases & Stroke (NPCDCS) for those above the age of 30 years [19], the high proportions, especially in the 30-39 year age groups indicate that the coverage is nowhere near the necessary levels even within the urban healthcare system. Scheduled tribes have consistently higher odds of having undiagnosed hypertension in the rural areas among both men and women. This reinforces previous policy priorities and highlights the need to continue to treat tribal health as a priority if we are to achieve universal health coverage [20]. Having ever been married was associated with lower odds of undiagnosed hypertension among both urban and rural women. Similar findings have been reported from previous studies in India [6,11,21]. This may indicate better health seeking behaviour among ever married women as compared to those who had never been in a union. Higher educational attainment was associated with lower odds of undiagnosed hypertension among women. The association was of public health significance in rural women. There has been previous evidence for the same [6,11,21]. The findings reiterate the importance of education among women and it's likely impact on their health-seeking behaviour, especially in the rural areas. Higher wealth index

was associated with lower odds of having undiagnosed hypertension among women in both urban and rural areas. Similar association was not observed among men. This may indicate a link between socioeconomic status and women empowerment, which also ties in with health-seeking behaviour [22].

Being diagnosed with moderate or severe anaemia was associated with lower odds of undiagnosed hypertension among men and women in both rural and urban areas, which may be due to their increased contact with the health system for associated treatment. The effectiveness of opportunistic screening for hypertension has been well documented. [23,24]

Usage of tobacco is associated with lower odds of undiagnosed hypertension among urban men. This may be due to higher contact of the tobacco users with the health system due to other health issues, which may have led to opportunistic screening and diagnosis of hypertension. This also indicates better healthcare seeking behaviour among this population – as compared to the rural men where the consumption of tobacco is less often associated with health risks and social taboos [25]. Current daily alcohol usage was associated with higher odds of undiagnosed hypertension among rural women. Poor health-seeking behaviour has been previously reported among alcohol users and this may explain our finding [23,24].

Working away from home was associated with lesser odds of having undiagnosed hypertension among rural men. While occupation did not show an association of public health significance, employment was significantly associated with lower odds of undiagnosed hypertension (Table S1). This may be because of the existing occupational health initiatives by public and private entities to ensure provision of preventive, promotive and curative care for the employed [26].

Owning a mobile telephone was also associated with lesser odds of having undiagnosed hypertension among rural men. Having access to mobile phones has been previously known to have a positive effect on disease awareness and health-seeking behaviour [27]. Having health insurance coverage was associated with lower odds of having undiagnosed hypertension among

urban men. Literature indicates that having health insurance coverage encourages increased medical check-ups [28]. However, the pattern is not seen in rural areas and among the poor [29], which needs to be further explored. Visit to public or private health facility was significantly associated with lower odds of undiagnosed hypertension among both men and women in both urban and rural areas. While this indicates a good performance of the facility-based screening programme, it calls into question the effectiveness of community-based risk assessment and screening for non-communicable diseases.

Strengths and Limitations

To the best of our knowledge, this is the first study to analyse the NFHS-5 dataset for factors associated with undiagnosed hypertension and their distribution pattern across rural and urban areas. The large sample size and the uniform methodology of the nationwide survey makes the results reliable and generalizable. State-level analysis of the data along with an urban-rural breakdown allowed for the highlighting of high priority areas where interventions may need to be stepped up or alternative approaches may need to be considered.

This study has various limitations. It is based upon secondary data available from a cross-sectional survey that assesses relationships at one point in time. This also means that causation cannot be inferred and only possible associations can be discussed. There is a possibility of recall bias and low truth quotient associated with all survey instruments. Application of American Heart Association thresholds of 130/80 for stage 1 hypertension, would have substantially increased the number of people with undiagnosed hypertension. The survey population was restricted to men and women aged 15–49 years and this means that we have not been able to account for a major proportion of those affected i.e., men and women over 50 years of age. NFHS-5 was conducted in two phases, one before the COVID-19 pandemic, and one during the pandemic. It is not possible to estimate how the disruption of regular services may have affected the burden of undiagnosed hypertension.

Implications for Policy and Practice

Improvement of health seeking behaviour among women may be brought about through education and empowerment by means of livelihood and increased earning potential. Special focus may be needed to improve health seeking behaviour of those who had never been married. Health-seeking behaviour among current alcohol users needs to be improved, through de-stigmatization of the topic, especially among women and through easy availability of cessation services. Screening of all those who report current alcohol or tobacco usage for hypertension may be incorporated as part of the national programme guidelines. High quality screening, diagnostic and treatment services for hypertension needs to be made accessible at all healthcare facilities at primary, secondary and tertiary levels. Community-based screening and referral under NPCDCS needs to be strengthened to reduce the burden of undiagnosed hypertension, especially in the rural areas of States identified with a high rural-urban gap in prevalence. Health promotion through the platform of mobile messaging and social media may be one the ways forward to draw the attention of the rural and migrant youth towards adoption of healthy lifestyles and the importance of early screening and diagnosis of hypertension. Improvement of access to health insurance coverage among the rural population is also likely to reduce the burden of undiagnosed hypertension.

Future Research

Health technology assessment studies to explore the possibilities of changes to the NPCDCS guidelines for screening, diagnosis and treatment of hypertension, may be considered to expand the coverage of hypertension awareness and treatment, with focus on the urban youth. The findings of this analysis open avenues for further research on interventions to improve health-seeking behaviour in the rural areas, as well as among current alcohol users. Predictors such as health insurance coverage, working away from home and owning a mobile telephone needs to be explored as possible areas of intervention to further reduce the burden of undiagnosed

hypertension in the future. Potential improvement in quality of life, increase in life expectancy and reduction in DALYs due to undiagnosed hypertension may be estimated to better understand the benefits that would be accrued as a result of reducing the burden of undiagnosed hypertension.

Conclusion

Our analysis of nationally representative data from the NFHS 5 (2019-21) revealed that two-thirds of men and two-fifths of women with hypertension were previously undiagnosed. State-level analysis highlighted multiple states with higher proportions of undiagnosed hypertension, both in urban and rural areas. Guidelines for population-based screening of common NCDs including hypertension has been implemented through the NPCDCS under the National Health Mission.²⁶ It is also operationalised through the Ayushman Bharat Health and Wellness Centres as part of the comprehensive primary health care package [30]. The findings of this study show the ground that needs to be covered in terms of diagnosing all who have hypertension. Only when one is aware of their status, will it be possible to bring them under treatment and ultimately ensure effective control of disease. Several predictors identified in this study such as age, caste, education, wealth index, alcohol and tobacco use, health insurance coverage, working away from home and owning a mobile telephone, all provide interesting avenues for further research, in order to inform necessary change in policy and practices in the future. The problem of undiagnosed hypertension is not restricted to only the rural areas but has important predictors in urban areas as well, which need to be explored further and acted upon. There exists an urgent need to plan interventions with an understanding of the health seeking behaviour of different sub-populations. Strengthening and expansion of community-based screening for hypertension appears to be the most important intervention to be able to identify apparently healthy individuals who are unlikely to visit the health facilities.

Acknowledgement

The authors would like to acknowledge the contribution and mentoring received from Dr. Hemant D Shewade, Scientist E (Medical), Division of Health Systems Research, ICMR-National Institute of Epidemiology, Chennai, Tamil Nadu, India for this research proposal and manuscript writing. The authors would also like to thank the team of Resource Center for Tobacco Control, Department of Community Medicine and School of Public Health, PGIMER, Chandigarh.

Competing interests

Authors have no competing interests to declare.

Financial disclosure statement

The author(s) received no specific funding for this work

References


1. Moore J. Hypertension: catching the silent killer. *The Nurse Practitioner*. 2005 Oct 1;30(10):16-35.
2. Faizi N, Ahmad A, Khalique N, Shah MS, Khan MS, Maroof M. Existence of rule of halves in hypertension: An exploratory analysis in an Indian village. *Materia socio-medica*. 2016 Apr;28(2):95.
3. World Health Organization. Hypertension Fact Sheet. Last updated 25th August 2021. Available on <https://www.who.int/news-room/fact-sheets/detail/hypertension>.
4. Wilber JA, Barrow JG. Hypertension-A community Problem. *Am J Med*. 1972;52:653–63.

5. Deepa R, Shanthirani CS, Pradeepa R, Mohan V. Is the ‘Rule of Halves’ in Hypertension Still Valid?-Evidence from the Chennai Urban Population Study. *J Assoc Phy India*. 2003;51:153–7.
6. Preissl J, Manne-Goehler J, Jaacks LM, Prabhakaran D, Awasthi A, Bishops AC, Atun R, Bärnighausen T, Davies JI, Vollmer S, Geldsetzer P. Hypertension screening, awareness, treatment, and control in India: a nationally representative cross-sectional study among individuals aged 15 to 49 years. *PLoS medicine*. 2019 May 3;16(5):e1002801.
7. Zhou B, Carrillo-Larco RM, Danaei G, Riley LM, Paciorek CJ, Stevens GA et al. Worldwide trends in hypertension prevalence and progress in treatment and control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104 million participants. *The Lancet*. 2021 Sep 11;398(10304):957-80.
8. Forouzanfar MH, Liu P, Roth GA, et al. Global burden of hypertension and systolic blood pressure of at least 110 to 115 MM Hg, 1990-2015. *JAMA*2017;317:165-82.
9. Office of the Registrar General. Registrar General of India report on medical certification of cause of death 2013. New Delhi, India: Office of the Registrar General, 2015.
10. Lim SS, Allen K, Bhutta ZA, Dandona L, Forouzanfar MH, Fullman N, Gething PW, Goldberg EM, Hay SI, Holmberg M, Kinfu Y. Measuring the health-related Sustainable Development Goals in 188 countries: a baseline analysis from the Global Burden of Disease Study 2015. *The Lancet*. 2016 Oct 8;388(10053):1813-50.
11. Boro B, Banerjee S. Decomposing the rural–urban gap in the prevalence of undiagnosed, untreated and under-treated hypertension among older adults in India. *BMC Public Health*. 2022 Dec;22(1):1-6.


12. Swain PK, Behera B, Das D. Association between area-level socio-economic status and hypertension in EAG states of India: an insight from NFHS-4 (2015-16). *Int J Agricult Stat Sci.* 2019;15:39–52.
13. United Nations Population Fund India. Harnessing India’s Demographic Dividend. Policy Brief. July 2018. Available at https://india.unfpa.org/sites/default/files/pub-pdf/harnessing_demographic_dividend_in_india_-_final.pdf.
14. Misra A, Singhal N, Sivakumar B, Bhagat N, Jaiswal A, Khurana L. Nutrition transition in India: Secular trends in dietary intake and their relationship to diet-related non-communicable diseases. *Journal of diabetes.* 2011 Dec;3(4):278-92.
15. Nadar SK, Tayebjee MH, Stowasser M, Byrd JB. Managing hypertension during the COVID-19 pandemic. *Journal of Human Hypertension.* 2020 Jun;34(6):415-7.
16. International Institute for Population Sciences (IIPS) and ICF. 2021. National Family Health Survey (NFHS-5), 2019-21: Volume I. Mumbai: IIPS.
17. Datawrapper. Datawrapper GmbH. Available on <https://app.datawrapper.de/>.
18. Talukdar D, Tripathi M, Tripathi V, Teelucksingh S. Retraction Note to: Prevalence and associated factors of undiagnosed hypertension among women aged 15-49 years in India: an analysis of National Family Health Survey-4 data. *Journal of human hypertension.* 2022 Oct;36(10):944.
19. National Programme For Prevention & Control Of Cancer, Diabetes, Cardiovascular Diseases & Stroke. National Health Mission. MOHFW. GoI. Available at <https://www.nhm.gov.in/index1.php?lang=1&level=2&sublinkid=1048&lid=604>
20. Tribal Health in India. Bridging the gap and a Roadmap for the future. Report of the Expert Committee on Tribal Health. MOHFW and Ministry of Tribal Affairs. Government of India. Available at https://nhm.gov.in/nhm_components/tribal_report/Executive_Summary.pdf

21. Mukhopadhyay K, Mukherjee S, Barkandaj B, Chatterjee C. Association of different socio-economic factors with hypertension prevalence, awareness, treatment and control in India: A demographic analysis Of NFHS-4. *Int J Res Med Sci*. 2019 Mar;7(3):815-21.
22. Anik AI, Islam MR, Rahman MS. Do women's empowerment and socioeconomic status predict the adequacy of antenatal care? A cross-sectional study in five South Asian countries. *BMJ open*. 2021 Jun 1;11(6):e043940.
23. Ogunyemi AO, Egemba TO, Onigbogi OO. Health-seeking Behaviour and Self-rated Health of Adult Men in an Urban Local Government Area in Lagos, Nigeria. *Annals of Health Research*. 2021 May 28;7(2):153-64.
24. Ayalew TL, Wale BG, Zewudie BT. Burden of undiagnosed hypertension and associated factors among adult populations in Wolaita Sodo Town, Wolaita Zone, Southern Ethiopia. *BMC cardiovascular disorders*. 2022 Dec;22(1):1-9.
25. Ruhil R. Sociodemographic determinants of tobacco use in India: Risks of risk factor—An analysis of global adult tobacco survey India 2016-2017. *SAGE Open*. 2019 Apr;9(2):2158244019842447.
26. Saha RK. Occupational Health In India. *Annals Of Global Health*. 2018;84(3):330.
27. Oshima SM, Tait SD, Thomas SM, Fayanju OM, Ingraham K, Barrett NJ, Hwang ES. Association of smartphone ownership and internet use with markers of health literacy and access: cross-sectional survey study of perspectives from project PLACE (Population Level Approaches to Cancer Elimination). *Journal of medical Internet research*. 2021 Jun 9;23(6):e24947.
28. Al-Hanawi MK, Mwale ML, Kamninga TM. The effects of health insurance on health-seeking behaviour: evidence from the kingdom of Saudi Arabia. *Risk Management and Healthcare Policy*. 2020;13:595.


29. Lee J, Wilkens J, Meijer E, Sekher TV, Bloom DE, Hu P. Hypertension awareness, treatment, and control and their association with healthcare access in the middle-aged and older Indian population: A nationwide cohort study. *PLoS medicine*. 2022 Jan 4;19(1):e1003855.
30. Operational guidelines. Comprehensive primary health care through Health and Wellness centres. Ayushman Bharat. NHSRC. 2018.




Figure



medRxiv preprint doi: <https://doi.org/10.1101/2024.12.18.24319257>; this version posted December 20, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).



Source: National Family Health Survey 2019-21 • Map data: © OSM • Created with Datawrapper



medRxiv preprint doi: <https://doi.org/10.1101/2024.12.18.24319257>; this version posted December 20, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

