

The Way Home: A Scoping Review of Public Health Strategies to Increase the Utilization of Home Dialysis in Chronic Kidney Disease Patients

Authors: Natcha Yongphiphatwong^a, Yot Teerawattananon^b, Pitsinee Supapol^{b**}, Denla Pandejpong^{b**}, Tanainan Chuanchaiyakul^{b**}, Jiratorn Sutawong^{b**}, Naline Gandhi^{c**}, Nutthawongse Kiatkrissada^d, Saudamini Vishwanath Dabak^b, Thunyarat Anothaisintawee^{e*}

Affiliations:

^a Thailand Development Research Institute (TDRI), Bangkok, Thailand

^b Health Intervention and Technology Assessment Program (HITAP), Ministry of Public Health, Nonthaburi, Thailand

^c DUKE-NUS Medical School, Singapore, Singapore

^d Faculty of Pharmaceutical Science, Chulalongkorn University, Thailand

^e Department of Clinical Epidemiology and Biostatistics, Department of Family Medicine, Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

*Corresponding author

Email: thunyarat.ano@mahidol.ac.th

Address: Faculty of Medicine, Ramathibodi Hospital, Mahidol University, Bangkok, Thailand

**These authors made equal contributions

ABSTRACT

Introduction: Home dialysis (HoD) remains underutilized, despite evidence showing it provides comparable mortality rates to in-center hemodialysis (ICHD) while offering advantages such as improved quality of life and lower overall costs. This scoping review comprehensively evaluates the impact of public health interventions on increasing the use of HoD, including both Peritoneal Dialysis (PD) and Home Hemodialysis (HHD).

Methods: Relevant studies were searched in the Web of Science, Medline, Embase, Scopus, EBSCOhost, and EconLit databases from their inception through May 2024. Studies were eligible for review if they assessed the effectiveness of public health interventions in terms of utilization and retention rates for general HoD, PD, and HHD.

Results: Forty-three studies were included, with interventions categorized into three main types: educational programs, service provision improvements, and modifications to payment structures. Our findings indicate that educational interventions—aimed at enhancing knowledge about dialysis options and promoting shared decision-making among patients, families, and healthcare providers—and service provision improvements, such as assisted PD and nephrologist-performed catheter insertions, could significantly increase the initiation, utilization, and retention rates of HoD. However, the impact of payment interventions on HoD outcomes differed across different contexts.

Conclusion: Education and service provision enhancements may represent the most effective public health interventions for increasing initiation, utilization, and retention rates of HoD in dialysis requiring patients. However, these findings are predominantly based on evidence from observational studies; further experimental studies with rigorous methodology are warranted to validate the effectiveness of these interventions in promoting HoD utilization.

PLAIN TEXT SUMMARY

Kidney dialysis is a life-sustaining therapy that can be offered both at home and in medical centres, however, home dialysis is underutilised globally. This scoping review gathers evidence from around the world to identify and assess the effectiveness of public health interventions to improve home dialysis utilization. The interventions we found were mainly related to improving patient knowledge, redesigning service provision, or adjusting payment/reimbursement conditions. Our results suggest that educating patients about their dialysis options to support shared decision-making before they require dialysis and offering assisted peritoneal dialysis at home can help increase the number of patients starting and staying on home dialysis. However, adjusting payment and reimbursement policies showed mixed results.

INTRODUCTION

Chronic Kidney Disease (CKD) is a significant public health burden with a global prevalence of 13.4% (11.7-15.1%)¹. CKD is associated with an increased risk of all-cause and cardiovascular mortality alongside substantial impacts on patients' daily lives. As kidney function declines, health-related quality of life (HRQoL) deteriorates, affecting the physical, emotional, and social well-being of patients^{2,3}. According to the Kidney Outcomes Quality Initiative (KDOQI) guideline⁴, CKD can be classified into five stages using the estimated glomerular function (eGFR) parameter and/or evidence of structural renal changes e.g. proteinuria. CKD stage 5 is defined as eGFR of less than 15 mL/min/1.73 m² and is also referred to as end-stage kidney disease (ESKD). At this stage patients typically require Kidney Replacement Therapy (KRT) including kidney transplantation (KT) or dialysis –to replace lost kidney function. While KT yields the highest HRQoL among available treatment options, access to KT is limited due to a shortage of both living and cadaveric donors⁵⁻⁸. Consequently, most ESKD patients depend on dialysis for ongoing care.

Dialysis options include in-centre hemodialysis (ICHD), peritoneal dialysis (PD), and home hemodialysis (HHD), with the latter two offering the flexibility of home-based care, meaning that they can be carried out by patients or their caregivers in the comfort of their homes. Findings from systematic reviews and meta-analyses suggest that PD had comparable mortality risk to ICHD⁹. Additionally, PD patients experience fewer cardiovascular events¹⁰ and report a better health-related quality of life (HRQoL) compared to those on ICHD^{11,12}. In terms of value for money, evidence from high-income countries (HICs) indicates that PD is more cost-effective than ICHD^{7,13-15}. Moreover, in low- and middle-income countries (LMICs), a cost-effectiveness analysis conducted in Thailand also found that when compared to palliative care, the average incremental cost-effectiveness ratio (ICER) for initial treatment with PD was lower than that for ICHD¹⁶.

Although PD is associated with lower costs and improved patient HRQoL compared to ICHD, it remains significantly underutilized, particularly in LMICs¹⁷. A global survey highlighted the disparity, revealing that the utilization of PD in low-income countries (LICs) is 60 times lower than in HICs, with PD use at just 0.9 per million population (pmp) (95% CI: 0.7–1.5) in LICs, compared to 53.0 pmp (95% CI: 40.6–89.8) in HICs¹⁷. Several barriers limit the utilization of PD in both HICs and LMICs. These include insufficient education on the available KRT options, leading to a lack of shared decision-making between patients and healthcare providers¹⁸. Additionally, inadequate support for PD services—such as limited PD expertise and insufficient clinical training for physicians and nurses¹⁹—low provider reimbursement²⁰, and unsuitable home environments for PD²¹ further hinder its use. These barriers pose challenges to both international and local recommendations aimed at enhancing home-based treatments for dialysis-requiring patients.

In Thailand, PD utilization declined dramatically, following the 2022 shift from the “PD-First” policy to one where patients may select their preferred dialysis modality. While the new policy offers greater autonomy to patients, its aftermath included lowered ICHD quality due to service capacity overload, a sharp increase in the dialysis budget, and a severely threatened PD ecosystem due to reduced patient volumes^{22,23}. In an effort to mitigate these effects, a government-commissioned working group (WG) in Thailand has recommended increasing PD utilization from 15% to 50%. Therefore, to inform the WG, we conducted a scoping review of the effectiveness of public health interventions in increasing the utilization of HoD, including both PD and HHD. This scoping review aims to assess the effects of the interventions on the uptake and retention of HoD utilization across the world. HHD was included in this review as the lessons learned from HHD provision in HICs may also apply to PD provision in lower- and middle-income contexts. Beyond informing the WG,

the findings of this review can also provide valuable insights for the global kidney community.

RESULTS

A comprehensive search yielded 25,067 studies, as shown in Figure 1. After removing 7,774 duplicates, the title and abstract of the remaining 17,283 studies were screened, resulting in 726 studies whose full texts were assessed for eligibility. Of the 726 full texts assessed, 42 studies met the inclusion criteria, and a thorough review of the reference lists of the selected studies further identified one additional study. Thus, 43 studies were included in this scoping review²⁴⁻⁶⁶.

Characteristics of the included studies are presented in Table 1. All the included studies were published within the last two decades with 49% of included studies published from 2020 onwards^{34-40,48-55,61-66}, and 42% published in the 2010s^{24-33,44-47,57-60}. The majority of studies were conducted in the Region of the Americas (AMR) and Western Pacific Region (WPR) according to the World Health Organization regions, with 15 of 43 (35%) studies conducted in the United States of America (USA)^{25,30,31,33,36,37,39,41,53,58-61,64,66}. Additionally, 39 of the 43 included studies (91%) were conducted in HICs^{24-33,35-43,45,47-56,58-66}, as defined by the World Bank's income group, while the remaining four studies were conducted in upper-middle-income countries (i.e., China^{44,46} and Thailand^{34,57}).

The interventions are classified into three main groups: education, service provision, and payment. The most common intervention types among the included studies were education (17 of 43; 40%)²⁴⁻⁴⁰ followed by service provision (12 of 43; 30%)^{24,41,43-52}, and payment (11 of 43; 26%)⁵⁶⁻⁶⁶. Additionally, three studies assessed the effect of combined education with service-provision interventions (3 of 43; 7%)⁵³⁻⁵⁵.

All educational and service-related interventions were provided by nephrologists and nurses, or a multidisciplinary care team consisting of a combination of nephrologists and nurses, together with other relevant professionals, such as renal dietitians, trained kidney educators, social workers, pharmacists, and psychologists. In one study, the educational interventions were also led by existing patients who had the experience of undertaking HoD²⁴, and in another study, the educational intervention was led by a government healthcare payer (i.e., Medicare)³⁶. For payment, the majority of these interventions were led by the government, except for one study⁶², which examined the impact of private insurance, where patients themselves paid for the insurance, on HoD utilization.

The reported outcomes focused on the initiation and utilization of PD/HoD, and HHD. PD/HoD initiation refers to the number of CKD patients who started PD/HoD as their first dialysis option, while PD/HoD utilization refers to the number of CKD patients currently using PD/HoD at the time of outcome measurement. HHD initiation and HHD utilization were reported in the same manner. Outcomes related to HoD dialysis retention were only reported for PD but not for HHD. PD retention is defined as the number of PD patients who did not switch to ICHD or KT. In studies where the PD drop-off or technique failure rates were reported, the inverse was calculated to express the outcomes homogenously as PD retention to facilitate comparison between studies.

Education

Out of 17 studies evaluating the effectiveness of education, seven studies reported outcomes related to the initiation of PD/HoD, and three studies focused on PD/HoD utilization outcomes. Three studies measured both the initiation and utilization of PD/HoD. Two studies reported on both PD and HHD utilization. Additionally, one study reported on PD retention, and one study covered both PD initiation and PD retention outcomes.

Educational interventions for CKD patients primarily aim to equip them with the knowledge necessary to navigate KRT options. These programs provided comprehensive information on KRT, covering dialysis techniques and the advantages and disadvantages of each option. Education was delivered by a multidisciplinary care team—including nurses and experienced PD patients—to offer varied perspectives. A range of teaching methods, such as face-to-face sessions, simulation-based teaching, videos, and web-based platforms, were used to improve patient engagement and understanding. Ultimately, these programs supported patients in making informed, collaborative decisions with their dialysis team regarding the best KRT method for their individual needs. The effectiveness of educational interventions from each study is presented in Table 2.

Regarding the outcome of PD/HoD initiation^{25,28–30,33–37,40}, six of the ten studies reporting this outcome found that providing education about PD/HoD significantly increased the initiation of PD/HoD^{25,28,35–37,40}. The remaining four studies also observed an increase in PD/HoD initiation in patients receiving this intervention, but these effects were not statistically significant^{29,30,33,34}. Additionally, two studies^{27,35} evaluated the outcome of PD retention, with all of them showing that educational interventions increased the retention rate of PD, although only one study reached statistical significance in this regard³⁵.

Nine studies reported outcomes on the utilization of PD/HoD^{24,26,31–33,36–39}. All of these studies found that educational interventions significantly increased the utilization of PD/HoD compared to no intervention. However, Castledine et al.²⁴ found impact varied according to the modality of education delivery (e.g., via home visits, group sessions, video materials, and patients having the experience of performing PD). Specifically, they found that among the education delivery methods investigated, only providing education intervention using home visits significantly increased the rate of PD/HoD utilization in dialysis-requiring patients.

For HHD utilization, results were conflicting between the two studies reporting this outcome. Findings from Blankenship et al. demonstrated a significant benefit of educational interventions in increasing HHD utilization, while results from Dubin et al. found a non-significant benefit of educational intervention in increasing HHD utilization^{31,39}.

Service Provision

Among the 12 studies evaluating the effectiveness of service provision interventions, three studies reported on the initiation of PD/HoD, while four studies focused on the utilization of PD/HoD outcomes. One study measured both PD/HoD initiation and utilization, and four studies assessed the PD retention rate.

In contrast to educational interventions which focus on pre-dialysis and support the decision-making process, service provision interventions are aimed at enhancing the delivery of dialysis care. These interventions included assisted PD where nurse or other health care providers help patients perform PD at home, catheter insertion performed by nephrologists rather than surgeons, and quality improvement programs, which often involved a multidisciplinary care team.

Regarding four studies reporting the outcome of PD/HoD initiation, all of which involved interventions such as assisted PD and improving PD care quality through a multidisciplinary care team^{45,48,50,52}. These studies found that service provision interventions significantly increased the rate of PD initiation compared to no intervention.

For four studies reporting on PD retention outcomes, each assessing the impact of improving the quality of care using different techniques^{44,46,47,51}. Two studies provided closely integrated services between hospital and home, and both found that this approach significantly helped patients continue using PD^{44,46}. Another study employed telehealth to support patients in performing dialysis at home, which resulted in an increased rate of PD

retention⁴⁷. The fourth study evaluated the impact of increasing centre volume on PD retention; however, the study found no significant difference in retention rates between large and small centre volumes⁵¹.

Among the five studies focused on the outcome of PD/HoD utilization, two studies^{43,50} investigated the effect of home care-assisted PD, and three studies^{24,41,49} assessed the impact of catheter insertion by nephrologists. Of the two studies assessing home care-assisted PD, one found a significant benefit in increasing PD utilization⁵⁰, while one found no significant effect⁴³. The results concerning catheter insertion by nephrologists were also inconsistent: two studies reported a significant increase in PD utilization^{41,49}, while another found no significant benefit from this intervention²⁴.

Combined Education and Service Provision

Among the three studies evaluating the impact of combined education and service provision interventions⁵³⁻⁵⁵, one study assessed PD/HoD utilization outcomes⁵⁴, while another examined both PD/HoD initiation and utilization⁵⁵. The third study reported on PD/HoD initiation and utilization as well as HHD initiation and utilization⁵³.

Two studies^{53,55} that reported on PD/HoD initiation outcomes observed an increase in initiation rates among patients receiving the combined interventions, though this benefit was not statistically significant⁵⁵. All three studies that evaluated PD/HoD utilization outcomes⁵³⁻⁵⁵ consistently showed an increase in PD/HoD utilization rates with combined interventions; however, this effect did not reach statistical significance in any of the studies. For the study reporting HHD initiation and utilization outcomes, this study found no significant benefit from the combined interventions in increasing HHD initiation or utilization rates⁵³.

Payment

Of the eleven studies assessing the effectiveness of payment interventions^{56–66}, five studies reported outcomes related to PD/HoD utilization^{56,59,63,64,66}, while two studies examined both PD/HoD initiation and utilization^{57,61}. One study⁶⁰ assessed outcomes for PD/HoD initiation, utilization, and PD retention, and another study⁶⁵ focused on PD/HoD utilization and PD retention. Additionally, one study measured outcomes for PD/HoD initiation, utilization, and HHD utilization⁶².

Payment-related interventions include bundled payments (e.g., Medicare Prospective Payment System, henceforth Medicare PPS^{58–60}), capitation (e.g., Thailand’s PD-First policy⁵⁷), fee-for-service (e.g., physician fee in Canada^{56,63}), pay-for-performance (e.g., End-Stage Renal Disease Treatment Choices Model, henceforth ETC model^{64,66}), and private payments (e.g., private health insurance⁶²).

Among the four studies reporting outcomes in terms of PD initiation, two studies investigating the impact of the Medicare scheme in the US—specifically, the Medicare PPS and coverage for PD catheters, found a significant increase in PD initiation following the implementation of these payment interventions^{60,61}. Thailand’s PD-First policy also led to a statistically significant rise in PD initiation⁵⁷. In Australia, however, access to private health insurance was associated with a lower likelihood of PD initiation, and this effect was statistically significant⁶². Two studies reported on PD retention with inconsistent results^{60,65}. Sloan et al., investigating, found that the payment system with the US Medicare PPS was associated with higher rates of PD retention⁶⁰. On the other hand, Chang found that Taiwan’s PD-encouraging reimbursement policy was associated with lower PD retention rates⁶⁵.

PD/HoD utilization was reported in eleven studies^{56–66}. Interventions that were associated with a significant increase in PD/HoD utilization were the US Medicare PPS, Taiwan’s PD-encouraging reimbursement policy, and Thailand’s PD-First policy^{57,59–61,65}.

However, Medicare's home dialysis training add-on was not associated with a significant increase in PD/HoD utilization⁵⁹. Mixed results were found for the ETC model: two studies^{64,66} found that the model was associated with an increase in HoD utilization, but the impact was only statistically significant in one study⁶⁶. The patient having supplementary private health insurance in Australia⁶² and the increase of the PD fee-for-service for nephrologists to be equivalent to HD⁶³ in Canada were not associated with significant increases in PD utilization.

Regarding the impact of payment interventions on HHD utilization^{58,62}, one study⁶² found that providing supplementary private health insurance significantly increased HHD utilization. However, the implementation of the Medicare Prospective Payment System (PPS) did not result in an increased HHD utilization rate.

DISCUSSION

This scoping review provides a comprehensive analysis of public health interventions aimed at enhancing the initiation, utilization, and retention of HoD, including both PD and HHD. Our findings indicate that education and service provision interventions can effectively increase initiation, utilization, and retention rates of HoD in patients requiring dialysis, with benefits observed across various types of these interventions. However, the impact of payment interventions on HoD initiation, utilization, and retention varied, showing inconsistent effects depending on the specific type of payment intervention^{60,63}.

The decision-making process for selecting a dialysis modality is complex and involves balancing multiple factors, including physician expertise and practices, patient and family values, and the patient's autonomy and self-management capability⁶⁷. This complexity contributes to the low utilization of PD, despite previous evidence showing that patients on PD and ICHD experience similar mortality outcomes^{68,69}. Barriers to HoD utilization can be categorized as those impacting patients—such as limited knowledge, lack of social support,

and living in remote areas—as well as barriers within healthcare providers (e.g., reimbursement issues) and the healthcare system (e.g., limited PD catheter access and late referrals to nephrologists⁷⁰). Addressing these barriers through pre-dialysis education, adjustments in service provision, and modifications to payment structures may increase HoD utilization among dialysis-requiring patients.

Our review found that most of the studies assessing the effectiveness of educational interventions show a significant benefit in increasing the utilization and retention of HoD in dialysis-requiring patients. Successful educational programs often stemmed from the pre-dialysis education initiatives that provided comprehensive information on KRT options. To illustrate, healthcare providers may help patients through an exercise where they draw out how different dialysis modalities may be incorporated into their weekly timetable²⁸. Additionally, patients may be asked to state the pros and cons of each dialysis modality and assign weights to each factor based on their personal preference²⁸. Beginning this process well in advance of when patients require dialysis ensures ample time for shared decision-making among patients, families, and healthcare providers^{24–26,28,31–33,36,37,40}. In addition, nearly half of the educational interventions that achieved statistically significant increases in the utilization and retention of HoD were led by multidisciplinary care teams^{28,31,32,35,37}. These findings emphasize the importance of incorporating multidisciplinary personnel in improving the effectiveness of the interventions.

The mode of education delivery also plays a critical role; for example, the results from Castledine et al. suggest that providing education via home visits has proven more effective than providing video-based education²⁴. Therefore, further investigation into the specific benefits of different educational delivery methods is necessary to draw more meaningful conclusions.

Service provision interventions included assisted PD, which enables patients to perform PD at home with support from nurses or a multidisciplinary care team. Other service provision interventions involved having nephrologists, rather than surgeons, handle PD catheter insertions and implementing mobile or telehealth systems to monitor and assist patients in managing HoD. Our study found that most studies evaluating these approaches reported significant benefits in increasing HoD initiation and utilization rates, especially through assisted PD and catheter insertions performed by nephrologists. A possible explanation for the increased PD uptake rates when nephrologists handle catheter insertions is the reduced delay in starting PD. When surgeons manage catheter insertions, scheduling challenges, and the prioritization of emergency cases often result in delayed PD initiation⁷¹⁻⁷³, especially when patients need to be referred to a different healthcare facility to undergo this procedure.

Our study indicates that assisted PD can enhance the utilization and retention of HoD, especially among elderly and physically dependent patients requiring dialysis⁷⁴. These patients often face distinctive obstacles to self-managed dialysis, including a higher prevalence of comorbidities compared to younger patients and a loss of independence due to increasing frailty, which leads to a greater need for caregiver assistance. Providing an assisted PD program for these individuals could be an effective approach to increasing PD use within this group.

Unlike educational and service provision interventions, which show consistent results across various interventions in the same group, the effectiveness of payment interventions found in our review varied depending on the specific type of payment intervention used as well as the context of the health system in which the policy was applied. Illustratively, the 2008 PD-First policy in Thailand was the payment intervention demonstrating the highest impact, with an OR of 5.89 for PD initiation and 3.47 for PD utilization⁵⁷. This significant

impact arose from making dialysis services accessible to previously underserved populations and designating PD as the first line of treatment. Conversely, an initiative to promote home dialysis among patients already accessing other forms of dialysis did not achieve similar success: raising nephrologist fee-for-service to match HD fees in Canada, where national health insurance covers both PD and HD services, did not lead to significant change in PD usage⁶³.

In contexts where the cost of PD provision is lower than that of ICHD, such as the US and Taiwan, bundled payments were successful at increasing HoD usage^{15,75,76}. Studies showed that the Medicare PPS correlated with a significant increase in HoD use, although this effect was not statistically significant for the training add-on⁵⁸⁻⁶⁰. Taiwan's bundled payment, subject to a global budget, has been effective in increasing PD utilization, yet it has also led to a lower PD retention rate⁶⁵. The odds of PD drop-off were 1.33 times higher in clinics compared to medical-centre hospitals, suggesting that inadequate medical knowledge may contribute to reduced retention⁶⁵.

Interestingly, the relationship between private health insurance and home dialysis modality utilization revealed that supplementary private health insurance was associated with higher odds of HHD utilisation but lower odds of PD initiation⁶². However, this study did not control for income as a confounder; those who can afford private health insurance are often better off financially and may be more likely to utilize HHD due to better living conditions⁶².

Overall, education, service and payment-related interventions can contribute to higher home dialysis initiation, retention, and utilization. However, only three studies⁵³⁻⁵⁵ investigated interventions in more than one of these three groupings. Therefore, the synergetic effects of these interventions could not be clearly understood. Additionally, public health interventions to increase home dialysis usage may be achieved more than via

education, service provision, or payment, for example through amending regulations or legislation^{77,78}, but their effectiveness are not assessed in the literature. For example, while the Advancing American Kidney Health Executive Order⁷⁹ explicitly supports the use of HoD, we did not find any studies which examines its impact—likely due to the technical difficulties associated with quantitatively assessing high-level interventions such as an executive order. Nevertheless, studies assessing the impact of the End-Stage Renal Disease Treatment Choices (ETC) payment model, which arose as a result of the executive order, were included in our review^{64,66}.

Our scoping review has several strengths. Firstly, we provide a comprehensive review of the effectiveness of various public health interventions on the initiation, utilization, and retention of HoD. Additionally, we considered both PD and HHD as outcomes of interest. The evidence on HHD utilization offers valuable insights, as lessons learned from HHD provision in HICs may also apply to PD provision in LMICs.

However, our study has some limitations. A key limitation is the inconsistency in measures of intervention effects, which complicates comparisons of intervention effectiveness across studies. Additionally, most included studies were observational studies and used pre-intervention data as historical controls, which may introduce confounding bias; therefore, further studies with rigorous methodologies are needed to confirm our findings. In addition, our review did not include studies from grey literature, which may lead to publication bias in our findings. Lastly, the studies included in this review were primarily conducted in HICs. This focus underscores a significant gap in evidence from resource-limited settings.

In conclusion, this scoping review suggests that enhancing education and service provision may be the most effective public health strategies for improving initiation,

utilization, and retention rates of HoD among dialysis-requiring patients. These findings provide valuable insights for prioritizing policy interventions to support the initiation, uptake, and sustained use of home dialysis both in Thailand and globally.

The findings from this scoping review were presented to a dialysis policy working group and the results were submitted as policy recommendations to the National Health Security Office (NHSO)—the government body managing Thailand’s Universal Health Coverage program. Looking ahead, future research should focus on evaluating these recommended interventions to systematically assess their impacts on dialysis policy.

METHODS

This scoping was conducted and reported according to the PRISMA-ScR (PRISMA extension for Scoping Reviews)⁸⁰.

Study identification

Relevant studies were identified through a comprehensive search of six databases including Web of Science, PubMed, Embase, Scopus, EBSCOhost, and EconLit since their inception through May 2024. The search terms used consisted of three domains: *Increase AND Utilization AND Home Dialysis*. The search terms and search strategies used for each database are shown in Supplementary Table 1. Additionally, the reference lists of the included studies were examined to further identify relevant studies for the review.

Study selection

The study selection process was facilitated by the Covidence systematic review software (version 2, Veritas Health Innovation, Melbourne, VIC, Australia). Titles and abstracts of the identified studies were screened by one reviewer (all authors). Full texts of the studies were reviewed independently by two reviewers if the decision could not be made based on titles and abstracts (all authors).

Observational studies (i.e., case-control, cross-sectional, and cohort studies), quasi-experimental studies, and randomized controlled trials (RCT) were eligible for this review if they met all of the following criteria: 1) studies that included participants as non-dialysis dependent CKD or dialysis-requiring CKD, and 2) studies that assessed and reported the effect of public health intervention on increasing utilization or retention of HoD. Therapeutic interventions, such as the use of innovative dialysate, were deemed beyond the scope of the review and were excluded.

In this review, “home dialysis” is defined as any dialysis modality conducted at the patient’s house, including PD and HHD. Public health interventions in this review are defined as the interventions that are focused on individual, or system levels⁷⁸. The interventions focused on individual levels aim to change beliefs, attitudes, and knowledge about home dialysis in CKD patients such as education and shared decision-making. The interventions focused on system levels and aimed to change the organization, laws, and policy of home dialysis such as change in service provision (e.g., home visit by nurse, insert catheter by nephrologist), or change in payment system or policy.

Data extraction

After the study selection process was completed, the included studies then went through a data extraction process by a single reviewer using Microsoft Excel. During this process, data regarding the study characteristics, details of the intervention, study context, impact, costs of implementing the intervention, as well as the supporting and limiting factors to the success of the intervention were extracted. Later, the impact data extracted was then cross-checked by another reviewer (TA and PS).

Data analysis

The effects of interventions on the utilization and retention of home dialysis were summarized qualitatively by intervention types and outcomes. However, as PD is the predominant home dialysis modality, the term HoD in the studies that did not specify PD or HHD was assumed to refer to PD in our analysis.

ACKNOWLEDGEMENTS

We would like to extend our gratitude to all parties involved in the Commission: The Learning Committee, chaired by Prof. Emeritus Kriang Tungsanga. Prof. Vivekanand Jha; The National Health Security Office's Working Group, chaired by Prof. Kearkiat Praditpornsilpa; and the Secretariat team of both the Committee and the Working Group. Their invaluable guidance and feedback have been instrumental throughout the research.

In addition, we would like to thank Kinanti Khansa Chavarina of the Health Intervention and Technology Assessment Program (HITAP) for helping design the first stages of the study and for giving a detailed review of the manuscript. We would also like to thank Siriyakorn Khamkom of the Faculty of Pharmacy, Silpakorn University for her contribution to the title and abstract screening process.

AUTHOR CONTRIBUTIONS

N.Y., Y.T., S.D., and T.A. contributed to study's concept and design. N.G. and T.A. searched for the relevant studies. N.Y., Y.T., P.S., D.P., T.C., J.S., N.G., N.K., S.D. and T.A. screened studies for the titles, abstracts, and full texts and performed the data extraction. N.Y., Y.T., and T.A. analysed the data and interpreted the results. N.Y. and T.A. drafted the manuscript. Y.T., P.S., D.P., T.C., J.S., N.G., N.K., and S.D. critically revised the manuscript. All authors approved the final version of the manuscript.

FUNDING SOURCES

This study was funded by the Health Systems and Research Institute (HSRI), Thailand (grant number HSRI 67-067), and the National Science, Research and Innovation

Fund (NSRF) through the Program Management Unit for Human Resources & Institutional Development, Research and Innovation (B41G670025). The funders had no role in the design, data collection, analysis, interpretation, or writing of the report. The findings, interpretations, and conclusions expressed in this article are those of the authors and do not necessarily reflect the views of the funding agencies.

The Health Intervention and Technology Assessment Program (HITAP) Foundation in Thailand supports evidence-informed priority-setting and decision-making in healthcare and is funded by both national and international public agencies.

COMPETING INTEREST DECLARATION

No authors declare competing interests.

DATA AVAILABILITY

Not applicable. This scoping review did not apply meta-analysis and only qualitatively summarized the data presented in the primary studies.

REFERENCES

1. Hill, N. R. *et al.* Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis. *PLoS ONE* **11**, e0158765 (2016).
2. Chin, H. J. *et al.* Moderately decreased renal function negatively affects the health-related quality of life among the elderly Korean population: a population-based study. *Nephrology Dialysis Transplantation* **23**, 2810–2817 (2008).
3. Perlman, R. L. *et al.* Quality of life in Chronic Kidney Disease (CKD): A cross-sectional analysis in the Renal Research Institute-CKD study. *American Journal of Kidney Diseases* **45**, 658–666 (2005).
4. National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification. *Am J Kidney Dis* **39**, S1-266 (2002).
5. Kloss, K. *et al.* Factors influencing access to kidney transplantation: a research protocol of a qualitative study on stakeholders' perspectives. *BMJ Open* **9**, e032694 (2019).
6. Ogutmen, B. *et al.* Health-Related Quality of Life After Kidney Transplantation in Comparison Intermittent Hemodialysis, Peritoneal Dialysis, and Normal Controls. *Transplantation Proceedings* **38**, 419–421 (2006).
7. Yang, F., Liao, M., Wang, P., Yang, Z. & Liu, Y. The Cost-Effectiveness of Kidney Replacement Therapy Modalities: A Systematic Review of Full Economic Evaluations. *Appl Health Econ Health Policy* **19**, 163–180 (2021).
8. Mudiayi, D. *et al.* Global Estimates of Capacity for Kidney Transplantation in World Countries and Regions. *Transplantation* **106**, 1113–1122 (2022).
9. Chander, S. *et al.* Mortality and mode of dialysis: meta-analysis and systematic review. *BMC Nephrol* **25**, 1 (2024).
10. Ng, C. H., Ong, Z. H., Sran, H. K. & Wee, T. B. Comparison of cardiovascular mortality in hemodialysis versus peritoneal dialysis. *Int Urol Nephrol* **53**, 1363–1371 (2021).

11. Walker, R. C. *et al.* A Discrete Choice Study of Patient Preferences for Dialysis Modalities. *CJASN* **13**, 100–108 (2018).
12. Tong, A. *et al.* The Perspectives of Adults Living With Peritoneal Dialysis: Thematic Synthesis of Qualitative Studies. *American Journal of Kidney Diseases* **61**, 873–888 (2013).
13. Ferguson, T. W. *et al.* Cost-Utility of Dialysis in Canada: Hemodialysis, Peritoneal Dialysis, and Nondialysis Treatment of Kidney Failure. *Kidney Medicine* **3**, 20-30.e1 (2021).
14. Baboolal, K. *et al.* The cost of renal dialysis in a UK setting - a multicentre study. *Nephrology Dialysis Transplantation* **23**, 1982–1989 (2008).
15. Karopadi, A. N., Mason, G., Rettore, E. & Ronco, C. Cost of peritoneal dialysis and haemodialysis across the world. *Nephrology Dialysis Transplantation* **28**, 2553–2569 (2013).
16. Teerawattananon, Y., Mugford, M. & Tangcharoensathien, V. Economic Evaluation of Palliative Management versus Peritoneal Dialysis and Hemodialysis for End-Stage Renal Disease: Evidence for Coverage Decisions in Thailand. *Value in Health* **10**, 61–72 (2007).
17. Cho, Y. *et al.* Peritoneal Dialysis Use and Practice Patterns: An International Survey Study. *American Journal of Kidney Diseases* **77**, 315–325 (2021).
18. Abdulkarim, S., Shah, J., Twahir, A. & Sokwala, A. P. Eligibility and patient barriers to peritoneal dialysis in patients with advanced chronic kidney disease. *Perit Dial Int* **41**, 463–471 (2021).
19. Paudel, K. *et al.* Overcoming barriers and building a strong peritoneal dialysis programme – Experience from three South Asian countries. *Perit Dial Int* **41**, 480–483 (2021).

20. Chan, C. T. *et al.* Overcoming Barriers for Uptake and Continued Use of Home Dialysis: An NKF-KDOQI Conference Report. *American Journal of Kidney Diseases* **75**, 926–934 (2020).
21. Edalat-Nejad, M. & Abdi, F. Medical and socioeconomic barriers to peritoneal dialysis utilization in the central province of Iran. *Iran J Kidney Dis* **7**, 147–150 (2013).
22. Teerawattananon, Y. & Chavarina, K. K. Nature Medicine Commission on Dialysis Policy in Low- and Middle-Income Countries: From Policy to Pivotal Impact: Thailand’s Dialysis Reform Journey and Its Unexpected Consequences. (*submitted*).
23. Phannajit, J., Praditpornsilpa, K. & Tungsanga, K. A Promising Start, A Troubling End: The Fallout of Thailand’s 2022 Universal Renal Dialysis Policy. (*submitted*).
24. Castledine, C. I., Gilg, J. A., Rogers, C., Ben-Shlomo, Y. & Caskey, F. J. Renal centre characteristics and physician practice patterns associated with home dialysis use. *Nephrology Dialysis Transplantation* **28**, 2169–2180 (2013).
25. Kurella Tamura, M. *et al.* Educational programs improve the preparation for dialysis and survival of patients with chronic kidney disease. *Kidney International* **85**, 686–692 (2014).
26. Fortnum, D., Ludlow, M. & Morton, R. L. Renal unit characteristics and patient education practices that predict a high prevalence of home-based dialysis in Australia. *Nephrology* **19**, 587–593 (2014).
27. Chan, D. T., Faratro, R. & Chan, C. T. The impact of simulation-based teaching on home hemodialysis patient training. *Clin Kidney J* **8**, 594–598 (2015).
28. Prieto-Velasco, M., Quiros, P., Remon, C., & Spanish Group for the Implementation of a Shared Decision Making Process for RRT Choice with Patient Decision Aid Tools. The Concordance between Patients’ Renal Replacement Therapy Choice and Definitive Modality: Is It a Utopia? *PLoS ONE* **10**, e0138811 (2015).

29. de Maar, J. S., de Groot, M. A. J., Luik, P. T., Mui, K. W. & Hagen, E. C. GUIDE, a structured pre-dialysis programme that increases the use of home dialysis. *Clin Kidney J* **9**, 826–832 (2016).
30. Shukla, A. M. *et al.* Effects of A Comprehensive Predialysis Education Program on the Home Dialysis Therapies: A Retrospective Cohort Study. *Perit Dial Int* **37**, 542–547 (2017).
31. Dubin, R. & Rubinsky, A. A Digital Modality Decision Program for Patients With Advanced Chronic Kidney Disease. *JMIR Form Res* **3**, e12528 (2019).
32. Lee, C.-T. *et al.* Shared Decision Making Increases Living Kidney Transplantation and Peritoneal Dialysis. *Transplantation Proceedings* **51**, 1321–1324 (2019).
33. Shukla, A. M. *et al.* What do the US advanced kidney disease patients want? Comprehensive pre-ESRD Patient Education (CPE) and choice of dialysis modality. *PLoS ONE* **14**, e0215091 (2019).
34. Parapiboon, W., Pitsawong, W., Wongluechai, L., Thammavaranucupt, K. & Raegasint, L. Customized versus conventional video counseling for peritoneal dialysis decision-making in patients with stage 5 chronic kidney disease under a PD-first policy: a randomized controlled study. *Kidney Res Clin Pract* **39**, 451–459 (2020).
35. Imamura, Y. *et al.* Effect of multidisciplinary care of dialysis initiation for outpatients with chronic kidney disease. *Int Urol Nephrol* **53**, 1435–1444 (2021).
36. Shukla, A. M. *et al.* Utilization of CMS pre-ESRD Kidney Disease Education services and its associations with the home dialysis therapies. *Perit Dial Int* **41**, 453–462 (2021).
37. Mckeon, K. *et al.* Utilization of Home Dialysis and Permanent Vascular Access at Dialysis Initiation Following a Structured CKD Education Program. *Kidney Medicine* **4**, 100490 (2022).

38. Shah, N., Cole, A., McCarthy, K. & Baharani, J. A Quality Improvement Process to Increase and Sustain a Peritoneal Dialysis Programme in the United Kingdom. *Blood Purif* **51**, 1022–1030 (2022).
39. Blankenship, D. M. *et al.* Assessing the impact of transitional care units on dialysis patient outcomes: A multicenter, propensity score-matched analysis. *Hemodialysis International* **27**, 165–173 (2023).
40. Sakurada, T. *et al.* Effects of one-hour discussion on the choice of dialysis modality at the outpatient clinic: A retrospective cohort study using propensity score matching. *Ther Apher Dial* **27**, 442–451 (2023).
41. Asif, A. *et al.* Does Catheter Insertion by Nephrologists Improve Peritoneal Dialysis Utilization? A Multicenter Analysis. *Seminars in Dialysis* **18**, 157–160 (2005).
42. Lobbedez, T. *et al.* Assisted Peritoneal Dialysis. Experience in a French Renal Department. *Perit Dial Int* **26**, 671–676 (2006).
43. Oliver, M.-J. *et al.* Home care assistance and the utilization of peritoneal dialysis. *Kidney International* **71**, 673–678 (2007).
44. Jiang, Z. & Yu, X. Advancing the Use and Quality of Peritoneal Dialysis by Developing a Peritoneal Dialysis Satellite Center Program. *Perit Dial Int* **31**, 121–126 (2011).
45. Chen, Y.-R. *et al.* Effectiveness of multidisciplinary care for chronic kidney disease in Taiwan: a 3-year prospective cohort study. *Nephrology Dialysis Transplantation* **28**, 671–682 (2013).
46. Yu, Y. *et al.* Impact of Continuous Quality Improvement Initiatives on Clinical Outcomes in Peritoneal Dialysis. *Perit Dial Int* **34**, 43–48 (2014).
47. Blaauw, M. Use of Sharesource in Remote Patient Management in Peritoneal Dialysis: A UK Nurse's Perspective. in *Contributions to Nephrology* (eds. Ronco, C., Crepaldi, C. & Rosner, M. H.) vol. 197 73–83 (S. Karger AG, 2019).

48. Boyer, A. *et al.* Trends in assisted peritoneal dialysis over the last decade: a cohort study from the French Peritoneal Dialysis Registry. *Clinical Kidney Journal* **13**, 1003–1011 (2020).
49. Liu, A. Y. L. & Nikam, M. Nephrologists-led intervention increases utilisation of peritoneal dialysis in Singapore. *Ann Acad Med Singap* **50**, 919–921 (2021).
50. van Eck van der Sluijs, A. *et al.* Assisted peritoneal dialysis across Europe: Practice variation and factors associated with availability. *Perit Dial Int* **41**, 533–541 (2021).
51. Yao, Y.-H., Chen, C.-M., Chou, Y.-J. & Huang, N. Impact of time-varying center volume on technique failure and mortality in peritoneal dialysis. *Perit Dial Int* **41**, 569–577 (2021).
52. Quinn, R. R. *et al.* The impact of the Starting dialysis on Time, At home on the Right Therapy (START) project on the use of peritoneal dialysis. *Perit Dial Int* 08968608231225013 (2024) doi:10.1177/08968608231225013.
53. Kaiser, P. *et al.* A Virtual Multidisciplinary Care Program for Management of Advanced Chronic Kidney Disease: Matched Cohort Study. *J Med Internet Res* **22**, e17194 (2020).
54. Tombocon, O. *et al.* Home before Hospital \square : a whole of system re-design project to improve rates of home-based dialysis therapy: Experience and outcomes over 8 years. *International Journal for Quality in Health Care* **33**, mzab108 (2021).
55. Manns, B. J. *et al.* Multifaceted Intervention to Increase the Use of Home Dialysis: A Cluster Randomized Controlled Trial. *CJASN* **17**, 535–545 (2022).
56. Mendelssohn, D. C., Langlois, N. & Blake, P. G. Peritoneal Dialysis in Ontario: A Natural Experiment in Physician Reimbursement Methodology. *Perit Dial Int* **24**, 531–537 (2004).
57. Praditpornsilpa, K. *et al.* Prevalence Trend of Renal Replacement Therapy in Thailand: Impact of Health Economics Policy. **94**, (2011).

58. Hirth, R. A. *et al.* The Initial Impact of Medicare's New Prospective Payment System for Kidney Dialysis. *American Journal of Kidney Diseases* **62**, 662–669 (2013).
59. Lin, E. *et al.* Home Dialysis in the Prospective Payment System Era. *JASN* **28**, 2993–3004 (2017).
60. Sloan, C. E. *et al.* Trends in Peritoneal Dialysis Use in the United States after Medicare Payment Reform. *CJASN* **14**, 1763–1772 (2019).
61. Lin, E., Chertow, G. M., Bhattacharya, J. & Lakdawalla, D. Early Delays in Insurance Coverage and Long-term Use of Home-based Peritoneal Dialysis. *Medical Care* **58**, 632–642 (2020).
62. Sriravindrarajah, A. *et al.* Impact of supplemental private health insurance on dialysis and outcomes. *Internal Medicine Journal* **50**, 542–549 (2020).
63. Trachtenberg, A. J. *et al.* Association between change in physician remuneration and use of peritoneal dialysis: a population-based cohort analysis. *cmajo* **8**, E96–E104 (2020).
64. Ji, Y., Einav, L., Mahoney, N. & Finkelstein, A. Financial Incentives to Facilities and Clinicians Treating Patients With End-stage Kidney Disease and Use of Home Dialysis: A Randomized Clinical Trial. *JAMA Health Forum* **3**, e223503 (2022).
65. Chang, R.-E., Lin, S.-P. & Myrtle, R. C. Impact of dialysis reimbursement policies on care outcomes in peritoneal dialysis patients in Taiwan. *Health Policy* **132**, 104816 (2023).
66. Johansen, K. L. *et al.* Association of the End-Stage Renal Disease Treatment Choices Payment Model With Home Dialysis Use at Kidney Failure Onset From 2016 to 2022. *JAMA Netw Open* **6**, e230806 (2023).
67. Wuerth, D. B. *et al.* Patients' Descriptions of Specific Factors Leading to Modality Selection of Chronic Peritoneal Dialysis Or Hemodialysis. *Perit Dial Int* **22**, 184–190 (2002).

68. Quinn, R. R. *et al.* Selection Bias Explains Apparent Differential Mortality between Dialysis Modalities. *Journal of the American Society of Nephrology* **22**, 1534–1542 (2011).
69. Merkus, M. P. *et al.* Quality of life in patients on chronic dialysis: Self-assessment 3 months after the start of treatment. *American Journal of Kidney Diseases* **29**, 584–592 (1997).
70. Moreiras-Plaza, M. M. Where we come from and where we are going in terms of peritoneal dialysis: identifying barriers and strategies for the future. *Nefrologia* **34**, 756–767 (2014).
71. Asif, A. Peritoneal Dialysis Access □ Related Procedures by Nephrologists. *Seminars in Dialysis* **17**, 398–406 (2004).
72. Ash, S. R. Bedside peritoneoscopic peritoneal catheter placement of Tenckhoff and newer peritoneal catheters. *Adv Perit Dial* **14**, 75–79 (1998).
73. Gadallah, M. F. *et al.* Peritoneoscopic versus surgical placement of peritoneal dialysis catheters: a prospective randomized study on outcome. *Am J Kidney Dis* **33**, 118–122 (1999).
74. Povlsen, J. V. & Ivarsen, P. Assisted automated peritoneal dialysis (AAPD) for the functionally dependent and elderly patient. *Perit Dial Int* **25 Suppl 3**, S60-63 (2005).
75. Kao, T.-W. *et al.* Lifetime Costs for Peritoneal Dialysis and Hemodialysis in Patients in Taiwan. *Perit Dial Int* **33**, 671–678 (2013).
76. Hornberger, J. & Hirth, R. A. Financial Implications of Choice of Dialysis Type of the Revised Medicare Payment System: An Economic Analysis. *American Journal of Kidney Diseases* **60**, 280–287 (2012).

77. Michie, S., Van Stralen, M. M. & West, R. The behaviour change wheel: A new method for characterising and designing behaviour change interventions. *Implementation Sci* **6**, 42 (2011).
78. Litvak, E. *et al.* Making sense of what exactly public health does: a typology of public health interventions. *Can J Public Health* **111**, 65–71 (2020).
79. The White House. Advancing American Kidney Health Executive Order. *Federal Register* **84**, 33817–33819 (2019).
80. Tricco, A. C. *et al.* PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* **169**, 467–473 (2018).

Figure 1. PRISMA Diagram

medRxiv preprint doi: <https://doi.org/10.1101/2024.12.29.24319745>; this version posted December 30, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity.

It is made available under a [CC-BY-NC-ND 4.0 International license](https://creativecommons.org/licenses/by-nc-nd/4.0/) .




Table 1. Characteristics of included studies

First Author (Year)	Study Design	Country/Region	Study Setting	Intervention	Comparator	Intervention Lead	Target Population
<i>Education</i>							
Castledine (2013)	Retrospective Cohort	UK	National Level	Pre-dialysis education program using home visit	No intervention	Nurses and Existing Patients	Patients
				Pre-dialysis education program using group session			
				Pre-dialysis education program using Existing Patients			
				Pre-dialysis education program using review of modality			
				Pre-dialysis education program using video/DVD materials			
Tamura (2013)	Case-Control	USA	National Level	Kidney Early Evaluation Program (KEEP) for screening and education	No intervention	Healthcare Providers	Patients
Fortnum (2014)	Cross-sectional	Australia	National Level	Renal units offering more group sessions per year	No intervention	Healthcare Providers	Patients
Chan (2015)	Retrospective Cohort	Canada	Single Center	Simulation-based teaching	Conventional teaching	Nurses	Patients
Prieto-Velasco (2015)	Prospective Cohort	Spain	Multiple Centers	Education Process (EP) with Patient Decision Aid (PDA) tools	No intervention	Multidisciplinary Care Team	Patients
de Maar (2016)	Retrospective Cohort	Netherlands	Single Center	Pre-dialysis programme (GUIDE)	Historical control	Multidisciplinary Care Team	Patients
Shukla (2017)	Retrospective Cohort	USA	Single Center	Comprehensive pre-dialysis education program	United States Renal Data System	Multidisciplinary Care Team	Patients
Dubin (2019)	Prospective Cohort	USA	Multiple Centers	Digital Modality Decision Program	Historical control	Multidisciplinary Care Team	Patients
Lee (2019)	Prospective Cohort	Taiwan	Not Reported	Shared decision-making (SDM)	Historical control	Multidisciplinary Care Team	Patients
Shukla (2019)	Prospective Cohort	USA	Single Center	Comprehensive pre-dialysis education program	CKD care in USRDS data	Nurses and Nephrologists	Patients and Caregivers
					Historical control		
Parapiboon (2020)	RCT	Thailand	Single Center	Customized multimedia	Conventional multimedia	Dialysis Facilities	Patients
Imamura (2021)	Retrospective	Japan	Single	Multidisciplinary care (MDC)	No intervention	Multidisciplinary	Patients

First Author (Year)	Study Design	Country/Region	Study Setting	Intervention	Comparator	Intervention Lead	Target Population
	Cohort		Center			Care Team	
Shukla (2021)	Retrospective Cohort	USA	National Level	Kidney Disease Education (KDE)	No intervention	Government Healthcare Payers and Dialysis Facilities	Dialysis Facilities and Patients
McKeon (2022)	Retrospective Cohort	USA	Multiple Centers	A structured CKD Education Program	No intervention	Multidisciplinary Care Team	Patients
Shah (2022)	Prospective Cohort	UK	Single Center	Quality Improvement (QI) by training nurses and patients	Pre-QI Period	Healthcare Providers and Existing Patients	Nurses and Patients
Blankenship (2023)	Retrospective Cohort	USA	Multiple Centers	Transitional care units (TCUs) or dedicated care programs or dialysis orientation units	Historical control	Healthcare Providers	Patients
Sakurada (2023)	Retrospective Cohort	Japan	Single Center	Shared decision-making (SDM)	No intervention	Nephrologists and Nurses	Patients and Caregivers
Service Provision							
Asif (2005)	Retrospective Cohort	USA	Multiple Centers	PD catheter insertion by nephrologists	PD catheter insertion by surgeon	Nephrologists	Patients
Oliver (2007)	Retrospective Cohort	Canada	Multiple Centers	Home Plus Program	No intervention	Nurses	Patients
Jiang (2011)	Cross-sectional	China	Multiple Centers	PD satellite center program	Historical control	Physicians and Nurses	Physicians and Nurses
Chen (2012)	Prospective Cohort	Taiwan	Multiple Centers	Multidisciplinary care (MDC)	Usual care group	Multidisciplinary Care Team	Patients
Castledine (2013)	Retrospective Cohort	UK	National Level	Provision home visits to PD patients	No intervention	Nurses and Existing Patients	Patients
				PD catheter insertion by members of renal team	PD catheter insertion by surgeon		
				Provision same day hospital visits for PD patients	No intervention		
Yu (2014)	Retrospective Cohort	China	Single Center	Continuous quality improvement	Historical control	Multidisciplinary Care Team	Patients
Blaauw (2019)	Prospective Cohort	UK	Not Reported	Remote patient management (RPM) systems	Historical control	Nurses	Nurses and Patients
Boyer (2020)	Retrospective	France	National	Nurse-assisted PD	Historical control	Nurses	Dialysis

First Author (Year)	Study Design	Country/Region	Study Setting	Intervention	Comparator	Intervention Lead	Target Population
	Cohort		Level				Facilities and Patients
Liu (2021)	Retrospective Cohort	Singapore	Single Center	PD catheter insertion by nephrologists	Historical control	Nephrologists	Patients
van Eck van der Sluijs (2021)	Cross-sectional	Europe	Multiple Centers	Assisted PD program	No intervention	Nurses and Caregiver	Patients
Yao (2021)	Retrospective Cohort	Taiwan	National Level	PD center volume (26-42 incident patients per year)	PD center volume (1-12 incident patients per year)	Dialysis Facilities	Patients
Quinn (2024)	Prospective Cohort	Canada	Sub-National Level	At Home, on the Right Therapy (START) project	Historical control	Dialysis Facilities	Patients
Combined Education and Service Provision							
Kaiser (2020)	Prospective Cohort	USA	Single Center	Virtual Multidisciplinary Care Program	No intervention	Multidisciplinary Care Team	Patients
					Self-control (pre-education)		
Tombocon (2021)	Prospective Cohort	Australia	Multiple Centers	Quality Improvement through establishing treatment pathways that coordinates local home treatment, raise awareness of HoD, and develop flexible individualized treatment (Home before Hospital)	Historical control	Multidisciplinary Care Team	Patients
Manns (2022)	RCT	Canada	Sub-National Level	Multifaceted Interventions, including phone surveys from a knowledge translation broker, 1-year center-specific audit/feedback on home dialysis use, delivery of an educational package, and an academic detailing visit	No intervention	Nephrologists	Dialysis Facilities
Payment							
Mendelssohn (2004)	Case-Control	Canada	Sub-National Level	Equal physician reimbursement for all dialysis modalities	Historical control	Government Healthcare Payers	Physicians
Praditpornsilpa (2011)	Retrospective Cohort	Thailand	National Level	PD-first policy (2009)	Historical control (2007)	Government Healthcare Payers	Dialysis Facilities
Hirth (2013)	Retrospective	USA	National	Medicare Prospective Payment	Historical control	Government	Dialysis

First Author (Year)	Study Design	Country/Region	Study Setting	Intervention	Comparator	Intervention Lead	Target Population
	Cohort		Level	System (PPS)	(2007)	Healthcare Payers	Facilities
Lin (2017)	Retrospective Cohort	USA	National Level	Medicare Prospective Payment System (PPS)	Historical control (2007)	Government Healthcare Payers	Dialysis Facilities
				Add-on paying for home dialysis training (Medicare Parts A/B subgroup)	No intervention		
Sloan (2019)	Retrospective Cohort	USA	National Level	Medicare Prospective Payment System (PPS)	Historical control	Government Healthcare Payers	Dialysis Facilities
Lin (2020)	Retrospective Cohort	USA	National Level	PD catheter paid for by Medicare	No intervention	Government Healthcare Payers	Patients
Sriravindrarajah (2020)	Retrospective Cohort	Australia	Sub-National Level	Private health insurance (PHI)	No intervention	Patients	Patients
Trachtenberg (2020)	Retrospective Cohort	Canada	Sub-National Level	Equal nephrologist fee-for-service (FFS) for HD and PD	Salaried nephrologist	Government Healthcare Payers	Physicians
Ji (2022)	RCT	USA	National Level	End-Stage Renal Disease Treatment Choices (ETC) Payment Model	No intervention	Government Healthcare Payers	Dialysis Facilities
Chang (2023)	Retrospective Cohort	Taiwan	National Level	PD-encouraging reimbursement policy	Historical control	Government Healthcare Payers	Dialysis Facilities
Johansen (2023)	Retrospective Cohort	USA	National Level	End-Stage Renal Disease Treatment Choices (ETC) Payment Model	Historical control	Government Healthcare Payers	Dialysis Facilities

Table 2. Results of Included Studies

First Author (Year)	Intervention	Comparator	Outcome	Odds Ratio	Percent Change	P-value
Castledine (2013)	Pre-dialysis education program using home visit	No intervention	HoD utilization	1.39 (1.06 - 1.83)	-	0.02
	Pre-dialysis education program using group session		HoD utilization	1.21 (0.88 - 1.66)	-	0.3
	Pre-dialysis education program using Existing Patients		HoD utilization	0.98 (0.76 - 1.26)	-	0.9
	Pre-dialysis education program using review of modality		HoD utilization	0.93 (0.72 - 1.2)	-	0.6
	Pre-dialysis education program using video/DVD materials		HoD utilization	0.63 (0.46 - 0.86)	-	0.003
Tamura (2013)	Kidney Early Evaluation Program (KEEP) for screening and education	No intervention	PD initiation	1.68 (1.24 - 2.28)	-	-
Fortnum (2014)	Renal units offering more group sessions per year	No intervention	HoD utilization	1.013 (1.01 - 1.02)	-	0.008
Chan (2015)	Simulation-based teaching	Conventional teaching	PD retention		96.4% vs 100% (0.16 - 107.62)	0.54
Prieto-Velasco (2015)	Education Process (EP) with Patient Decision Aid (PDA) tools	No intervention	PD initiation	13.2 (5.20 - 33.54)	-	<0.001
de Maar (2016)	Pre-dialysis programme (GUIDE)	Historical control	HoD initiation	1.93 (0.79 - 4.72)	-	-
Shukla (2017)	Comprehensive pre-dialysis education program	United States Renal Data System	PD initiation	-	9% vs 55%	-
Dubin (2019)	Digital Modality Decision Program	Historical control	PD utilization	5.69 (1.51 - 21.42)	-	0.004
			HHD utilization	2.19 (0.36 - 13.22)	-	0.5
Lee (2019)	Shared decision-making (SDM)	Historical control	PD utilization	2.33 (1.47 - 3.69)	-	-
Shukla (2019)	Comprehensive pre-dialysis education program	CKD care in USRDS data	PD initiation	-	62% vs 8%	-
		Historical control	HoD utilization	-	12% vs 27%	<0.0001
Parapiboon (2020)	Customized multimedia	Conventional multimedia	PD initiation	1.16 (0.55 - 2.45)	-	0.86
Imamura (2021)	Multidisciplinary care (MDC)	No intervention	PD retention	2	-	0.012
			PD initiation	2.52 (1.04 - 6.11)	-	0.038

Shukla (2021)	Kidney Disease Education (KDE)	No intervention	HoD utilization	1.7 (1.52 - 1.90)	-	-
			HoD initiation	1.99 (1.66 - 2.39)	-	-
McKeon (2022)	A structured CKD Education Program	No intervention	HoD utilization	3.35 (2.93 - 3.83)	-	<0.001
			HoD initiation	4.34 (3.75 - 5.03)	-	<0.001
Shah (2022)	Quality Improvement (QI) by training nurses and patients	Pre-QI Period	PD utilization	3.13 (2.06 - 4.73)	-	<0.001
Blankenship (2023)	Transitional care units (TCUs) or dedicated care programs or dialysis orientation units	Historical control	PD utilization	-	2.8% vs 9.9%	<0.0001
			HHD utilization	-	7.3% vs 15.7%	<0.0001
Sakurada (2023)	Shared decision-making (SDM)	No intervention	PD initiation	4.81 (2.81 - 8.24)	-	<0.001
Service Provision						
Asif (2005)	PD catheter insertion by nephrologists	PD catheter insertion by surgeon	PD utilization	1.55 (1.25 - 1.93)	-	-
Oliver (2007)	Home Plus Program	No intervention	PD utilization	1.49 (0.73 - 3.04)	-	0.27
Jiang (2011)	PD satellite center program	Historical control	PD retention	1.84 (1.53 - 2.21)	-	0.01
Chen (2012)	Multidisciplinary care (MDC)	Usual care group	PD initiation	4.77 (1.36 - 16.68)	-	-
Castledine (2013)	Provision home visits to PD patients	No intervention	HoD utilization	1.63 (1.11 - 2.42)	-	0.01
	PD catheter insertion by members of renal team	PD catheter insertion by surgeon	PD utilization	1.1 (0.83 - 1.43)	-	0.5
	Provision same day hospital visits for PD patients	No intervention	HoD utilization	0.96 (0.58 - 1.60)	-	0.9
Yu (2014)	Continuous quality improvement	Historical control	PD retention	-	89.6% vs 95.6%	<0.001
Blaauw (2019)	Remote patient management (RPM) systems	Historical control	PD retention	-	37% vs 71%	-
Boyer (2020)	Nurse-assisted PD	Historical control	PD initiation	1.13 (1.06 - 1.21)	-	-
Liu (2021)	PD catheter insertion by nephrologists	Historical control	PD utilization	-	10-23% vs 25-29%	0.015
van Eck van der	Assisted PD program	No intervention	PD	2.81	-	<0.001

Sluijs (2021)			utilization	(1.77 - 4.47)		
			PD initiation	1.91 (1.21 - 3.01)	-	<0.001
Yao (2021)	PD center volume (26-42 incident patients per year)	PD center volume (1-12 incident patients per year)	PD retention	1.1 (0.91 - 1.33)	-	-
Quinn (2024)	At Home, on the Right Therapy (START) project	Historical control	PD initiation	-	MD = 5.4% (1.5 - 9.2)	-
Combined Education and Service Provision						
Kaiser (2020)	Virtual Multidisciplinary Care Program	No intervention	PD utilization	5.33 (0.47 - 60.80)	-	0.99
		Self-control (pre-education)	PD initiation	3.2 (0.91 - 11.27)	-	-
		No intervention	HHD utilization	0.9 (0.02 - 50.24)	-	0.99
		Self-control (pre-education)	HHD initiation	1.58 (0.24 - 10.52)	-	-
Tombocon (2021)	Quality Improvement through establishing treatment pathways that coordinates local home treatment, raise awareness of HoD, and develop flexible individualized treatment (Home before Hospital)	Historical control	HoD utilization	-	14.8% vs 35%	-
Manns (2022)	Multifaceted Interventions, including phone surveys from a knowledge translation broker, 1- year center-specific audit/feedback on home dialysis use, delivery of an educational package, and an academic detailing visit	No intervention	HoD utilization	1.16 (0.92 - 1.45)	-	0.21
			HoD initiation	1.31 (0.88 - 1.93)	-	0.17
Payment						
Mendelssohn (2004)	Equal physician reimbursement for all dialysis modalities	Historical control	PD utilization	-	19.7% vs 22.6%	-
Praditpornsilpa (2011)	PD-first policy (2009)	Historical control (2007)	PD utilization	3.47 (3.25 - 3.70)	-	-
			PD initiation	5.89 (5.32 - 6.52)	-	-
Hirth (2013)	Medicare Prospective Payment System (PPS)	Historical control (2007)	PD utilization	-	6.48% vs 6.96%	-
			HHD utilization	-	0.67% vs 1.44%	-
Lin (2017)	Medicare Prospective Payment System (PPS)	Historical control (2007)	HoD utilization	-	MD = 5.8% (4.3 - 6.9)	-

	Add-on paying for home dialysis training (Medicare Parts A/B subgroup)	No intervention	HoD utilization	-	MD = -0.2% (-1.0 - 0.5)	-
Sloan (2019)	Medicare Prospective Payment System (PPS)	Historical control	PD utilization	1.39 (1.37 - 1.41)	-	<0.001
			PD retention	1.09 (1.02 - 1.15)	-	0.004
			PD initiation	1.38 (1.36 - 1.40)	-	<0.001
Lin (2020)	PD catheter paid for by Medicare	No intervention	PD utilization	12 (9.60 - 15)	-	-
			PD initiation	81 (53.34 - 123)	-	-
Sriravindrarajah (2020)	Private health insurance (PHI)	No intervention	PD utilization	0.92 (0.76 - 1.11)	-	0.36
			PD initiation	0.81 (0.67 - 0.98)	-	0.03
			HHD utilization	1.38 (1.01 - 1.89)	-	0.04
Trachtenberg (2020)	Equal nephrologist fee-for-service (FFS) for HD and PD	Salaried nephrologist	PD utilization	1.52 (0.96 - 2.4)	-	-
Ji (2022)	End-Stage Renal Disease Treatment Choices (ETC) Payment Model	No intervention	HoD utilization	-	MD = 0.12% (-1.42 - 1.65)	0.89
Chang (2023)	PD-encouraging reimbursement policy	Historical control	PD utilization	1.28 (1.22 - 1.34)	-	0.029
			PD retention	0.89 (0.80 - 0.96)	-	0.029
Johansen (2023)	End-Stage Renal Disease Treatment Choices (ETC) Payment Model	Historical control	HoD utilization	-	MD = 1.07% (0.16 - 1.97)	-