

# Optimising covariate allocation at design stage using Fisher Information Matrix for Non-Linear Mixed Effects Models in pharmacometrics

Lucie Fayette<sup>1,2</sup>, Karl Brendel<sup>2</sup>, and France Mentré<sup>1</sup>

<sup>1</sup>Université Paris Cité, Inserm, IAME, F-75018 Paris, France

<sup>2</sup>Pharmacometrics, Ipsen Innovation, Les Ulis, France

lucie.fayette@inserm.fr

## Abstract

This work focuses on designing experiments for pharmacometrics studies using Non-Linear Mixed Effects Models including covariates to describe between-subject variability. Before collecting and modelling new clinical trial data, choosing an appropriate design is crucial. Assuming a known model with covariate effects and a joint distribution for covariates in the target population from previous clinical studies, we propose to optimise the allocation of covariates among the subjects to be included in the new trial. It aims achieving better overall parameter estimations and therefore increase the power of statistical tests on covariate effects to detect significance, and more importantly, clinical relevance or non-relevance of relationships. We suggested dividing the domain of continuous covariates into clinically meaningful intervals and optimised their proportions, along with the proportion of each category for the discrete covariates. We used the Fisher Information Matrix and developed a fast and deterministic computation method, leveraging Gaussian quadrature and copula modelling. The optimisation problem was formulated as a convex problem subject to linear constraints, allowing resolution using Projected Gradient Descent algorithm. Different scenarios for a pharmacokinetics model were explored. We showed the benefit of covariate optimisation in reducing the number of subjects needed to achieve desired power in covariate tests.

**Keywords:** Pharmacometrics; Optimal design, Fisher Information Matrix, Non-Linear Mixed Effects Models, Covariates, Sample size, Projected Gradient Descent

## Funding statement

This work was financed by a CIFRE agreement (Conventions Industrielles de Formation par la Recherche) of the ANRT (Association Nationale de la Recherche et de la Technologie). The CIFRE agreement is a partnership between a public laboratory and a company, here the UMR (Unité Mixte de Recherche) 1137 and Ipsen, respectively.

## Data availability statement

All the R scripts are available at the following link: <https://doi.org/10.5281/zenodo.14778034>

## 1 Introduction

In pharmacometrics longitudinal data collected during clinical trials are leveraged through Non-Linear Mixed Effect Models (NLMEM). In particular, NLMEM allow to quantify the relationships between covariates and parameters [1]. Indeed, covariates are often included in pharmacometrics models to describe predictable between subjects variability, thereby enhancing model fit and/or model-based predictions [2].

Before collecting and modelling the data, a key step is to chose an appropriate design. In NLMEM, a population design refers to the number of subjects and their allocation to elementary designs, an elementary design being a combination of values for the design variables, such as number of samples, sampling times, dosage regimen, etc. Usually, covariates are not considered at this step.

In its Population Pharmacokinetics (PK) Guidance for Industry 2022 [3], the FDA emphasises that "Simulations and optimal design methods can maximize the utility of population PK data collection and analyses" and stresses in particular that these methods can aim to estimate "major covariate effects of interest" with greater precision. Nevertheless, the method proposed for power assessment and sample size computation in this recommendation is clinical trial simulations (CTS), although it is computationally expensive and less exhaustive than optimal design strategies. The traditional theory of optimal experimental design, initially developed for non-linear models [4], have already been imported to the field of NLMEM and pharmacometrics [5, 6]. It relies on Fisher Information Matrix (FIM) computation, through linearization of the structural model, or stochastic simulations or Gaussian approximation [7–9]. Thus, before the start of a new study, given

a NLMEM and parameters value, one can compute the expected standard error (SE) of the model parameters and deduce whether enough information will be collected to meet the objectives of the study. Thereafter, design optimisation consist in finding the design of experiments that provides the most informative data for estimating model parameters while respecting some design constraints, such as number of subjects, number of sampling times or samplings times windows. These optimal design strategies require to assume a model and a vector of population parameters values. Methods to compute robust designs to the prior choice of parameter values or/and to the prior choice of model in the context of NLMEM have already been explored [10–13] but will not be considered here.

Various optimisation software especially dedicated to design optimisation in the context of clinical trial using NLMEM have been developed in the past 20 years. Among them we can cite *PFIM* [14], which was the first software tool proposed in 2001 for evaluating a design without using simulations, especially, the R package *PFIM 6.1* [15, 16] (CRAN release October 2024); PopED [17, 18] and NONMEM\$DESIGN [19].

Numerous studies have demonstrated the benefits of using this methodology to optimise sampling times but very few studies have addressed the question of which covariate values would give the most information.

Nevertheless, if a study aims to detect whether some covariate relationships are significant or relevant, this design optimisation procedure needs to account for covariates. In a previous work, we explored methods for handling covariates in FIM computation [20]. There are based on Monte-Carlo (MC) approximation, and propose to leverage either an historical covariate dataset, a known covariate distribution or a copula. Their accuracy in predicting uncertainty and power of significance and relevance tests was assessed in a simulation study [20] using a simple PK example and under various scenarios, including different sample sizes, sampling points, and IIV. In addition the FIM computation using a prior covariate dataset was applied to a Phase 3 example with a model including 27 covariate relationships and was found to accurately predict uncertainty even in this more complex setting. These methods have not yet been used for computing the FIM during design optimisation process, but it does not present any particular technical difficulties. In this work we propose to go a step further: assuming a covariate distribution within the target population, we propose, for a fixed design, NLMEM and parameter vector, to optimise the selection of subjects to be included in the future trial from this distribution.

Previous works have looked at how to account for covariates when allocating patients to different arms, and how an unbalanced design could improve efficiency and ethics, including model-based approaches (see for instance [21] for a review). However, these approaches do not use the longitudinal data framework nor NLMEM. In addition, they aim to optimised the design sequentially and seek, for each new patient entering the trial, to allocate them to the best possible arm in terms of efficacy and ethical considerations, while maintaining randomisation. Consequently, it does not correspond to our setting aiming to optimise the design prior to setting up the study. In [22], a new method based of the FIM for optimising design using NLMEM where covariates are design variables is proposed. However, the proposed methods require the FIM to be evaluated for a set of fixed covariate values, without taking into account the distribution of these covariates, so the combinatorial approach can result in a very large number of elementary designs to be evaluated and the optimisation result is discrete, making it difficult to manage continuous covariates. In addition, there are no recommendations on how to take into account multiple covariates that are correlated with each other, or on the practical management of covariate distributions.

To this end, we developed in this work a new method aiming to reduce the computational cost of the FIM computation, to handle continuous and correlated covariates and to optimise their distribution. Indeed, with optimisation purposes, the required number of FIM evaluation increases and run time becomes a challenge. Therefore, avoiding MC simulations is a lever to reduce this cost. As a consequence, we introduce a Gaussian quadrature [23] (GQ) based approach, thus faster and deterministic. We have coupled this approach with the use of copula to describe the distribution of covariates, as it is very flexible and if already available, it can be used without individual data. Regarding optimisation, a hurdle to be overcome is that optimising the distribution of continuous covariates is not trivial as it is unrealistic to recommend including a given number of patients matching an exact value of the covariate. Therefore, we propose to segment the continuous covariates into different intervals and to optimise the proportion of each of them among the trial population. This has the combined advantage to remains realistic in practice while defining discrete parameters to be optimised. Finally, we propose a method for the constrained optimisation of covariates using a projected gradient method. The FIM is computed over a finite set combining elementary designs and covariate distributions and each combination is optimally weighted, given the constraints, using the Projected Gradient Descent (PGD) algorithm [24]. Once covariate distribution is optimised, the sample size can be optimised to reach desired level of precision or desired power in statistical test on covariate effects .

All this methods have been implemented using the R language [25] in a working version of *PFIM 6*. Scripts are available in the Zenodo repository <https://doi.org/10.5281/zenodo.14778034>. Based on the previously used simple PK model, we developed a workflow to assess the number of nodes required in the quadrature for an accurate FIM computation. Because prior data are not always available at design step, we used public database from the National Health and Nutrition Examination Surveys (NHANES) database [26] to get prior covariate distributions. Thereafter we explored various scenarios using a simple PK model demonstrating the benefit of covariate distribution optimisation.

In Section 2, the notations and methods are detailed. In Section 3 we go through the example settings and results are presented in Section 4. Lastly global results are discussed in section 5.

## 2 Methods

In this section we detail notations and methods on study design and NLMEM. We expose the FIM computation accounting for covariate distribution, and especially we describe the segmentation of continuous covariates for optimisation purposes and how it is handle in the FIM computation. Then, we explain the FIM integration using copula and Gauss-Legendre Quadrature (GLQ). Lastly, we present optimisation of covariate distributions using PGD.

### 2.1 Notations on study designs and NLMEM

In the following, a population design is denoted  $\Xi = \{N, (\xi_1, \dots, \xi_N)\}$ , where  $N$  is the number of subjects and the  $\xi_i$  are their elementary designs.

The observation vector  $y_i$  for the  $i^{th}$  subject is modelled as given in equation (1) where  $f$  denotes the structural non-linear model and  $\theta_i = u(\mu, \beta, Z_i, \eta_i)$ , the  $p$ -vector of individual parameters. The latter are defined as a function  $u$ , depending on fixed effects, composed of the typical values denoted  $\mu$  and of the covariate effects  $\beta$ , and on individual random effects  $\eta_i$ , where  $\eta_i \sim \mathcal{N}(0, \Omega)$ . Individual parameters also depend on the vector of possibly transformed individual covariates  $Z_i$ .

$$y_{ij} = f(\xi_i, \theta_i) + g(\xi_i, \theta_i, \sigma)\varepsilon_{ij} \quad (1)$$

We denote  $Z_i = (Z_{Dis,i}, Z_{Cont,i})$  a covariate vector when considered as a random vector, and  $z_i = (z_{Dis,i}, z_{Cont,i})$  as a realisation of this random vector. The lower  $.Dis$  refers to the discrete sub-vector of the covariate vector, while  $.Cont$  refers to the continuous sub-vector.

In pharmacometrics modelling, parameters are generally log-normally distributed with additive covariate relationships on the log scale, i.e.  $\log \theta_i = \log \mu + \beta(z_i - z_{ref}) + \eta_i$ .

The residual error is modelled by the function  $g$ , depending on  $\xi_i$ ,  $\theta_i$  and some parameters  $\sigma$ , and by the random variable  $\varepsilon_i$ , with  $\varepsilon_i \sim \mathcal{N}(0, I_{n_i})$ .

The  $P$ -vector of population parameters is denoted  $\Psi = \{\mu, \beta, \lambda\}$ , where  $\lambda$  contains the variance parameters (elements of  $\Omega$  and of the residual error model  $\sigma$ ).

To assess the magnitude of a covariate effect, one can compute the ratio of change in primary parameters when covariate values change relative to a reference value  $z_{ref}$ . Detailed formulas are given in [20], and for a log-normally distributed parameter with an additive covariate relationship on the log scale, the ratio writes  $r_{l,c} = e^{\beta_{l,c}}$  if the covariate is binary, and the ratio associated to a given percentile  $PX$  of the covariate distribution if continuous writes  $r_{l,c(PX)} = e^{\beta_{l,c}(PX - P50)}$ , where  $P50$  denotes the median of the covariate distribution.

### 2.2 Elementary and population FIM

Given an elementary design  $\xi_i$ , a population parameters vector  $\Psi$ , and a covariate realisation  $z_i$  the expected elementary FIM,  $M_F(\xi_i, \Psi, z_i)$ , is defined as the covariance matrix of the Fisher score (equation (2)) where  $l(y_i; \Psi, z_i, \xi_i)$  is the likelihood of the vector of observations  $y_i$  for the population parameters  $\Psi$ , given the individual vector of covariates  $z_i$  and the elementary design  $\xi_i$  (equation (3)).

$$M_F(\xi_i, \Psi, z_i) = \mathbb{E} \left( \frac{\partial \log l(y_i; \Psi, z_i, \xi_i)}{\partial \Psi} \frac{\partial \log l(y_i; \Psi, z_i, \xi_i)}{\partial \Psi^T} \right) \quad (2)$$

$$l(y_i; \Psi, z_i, \xi_i) = \int p(y_i | \eta_i; \Psi, z_i, \xi_i) p(\eta_i; \Psi, z_i) d\eta_i \quad (3)$$

Knowing the distribution  $p_Z$  for the covariate vector, the expected FIM can be expressed as an expectation over the covariate vector as given in equation (4).

$$M_F(\xi_i, \Psi, p_Z) = \int M_F(\xi_i, \Psi, z_i) p_Z(z) dz \quad (4)$$

Assuming that the covariate distribution  $p_Z$  is the same for all the elementary design, the expected population FIM is the sum of the elementary FIM (equation (5)).

$$M_F(\Xi, \Psi, p_Z) = \sum_{i=1}^N M_F(\xi_i, \Psi, p_Z) \quad (5)$$

In the following, we consider that the  $N$  subjects have the same elementary design  $\xi$ , thus the expected population FIM is  $M_F(\Xi, \Psi, p_Z) = N M_F(\xi, \Psi, p_Z)$ .

### 2.2.1 Elementary FIM with discrete covariates only

In the case with only  $D$  discrete covariates, each with  $Q_d$ ,  $d = 1, \dots, D$  categories. The covariate vector follows a discrete distribution with  $n^{(QD)} = \prod_{d=1}^D Q_d$  possible values, denoted  $z_{Dis,d}$ ,  $d = 1, \dots, n^{(QD)}$ , each having the probability  $p_{Dis,d}$ ,  $d = 1, \dots, n^{(QD)}$ . Therefore, the elementary FIM for  $\xi$  writes as a weighted sum as given in equation (6).

$$M_F(\xi, \Psi, p_Z) = \sum_{d=1}^{n^{(QD)}} p_{Dis,d} M_F(\xi, \Psi, z_{Dis,d}) \quad (6)$$

### 2.2.2 Elementary FIM with discrete and continuous covariates

In case there are both non-independent discrete and continuous covariates, we denote  $p_{Dis}$  the distribution of the discrete sub-vector, and for each of its possible value  $z_{Dis,d}$ , we denote  $p_{Cont|z_{Dis,d}}$  the distribution of the continuous sub-vector conditionally to  $z_{Dis,d}$ . The elementary FIM for a given  $\xi$  is given in equation (7).

$$M_F(\xi, \Psi, p_Z) = \sum_{d=1}^{n^{(QD)}} p_{Dis,d} M_F(\xi, \Psi, z_{Dis,d}, p_{Cont|z_{Dis,d}}) \quad (7)$$

### 2.3 Segmentation of continuous covariates

We assume here that a priori distributions are known for the covariates, and our aim is to distribute the subjects in the study. To achieve this, we suggested to segment the continuous covariates into different intervals, chosen for their clinical meaning. We then balanced the allocation between the different continuous intervals and between the discrete covariate categories. The advantage is that it maintains the continuous nature of the covariate in the FIM calculus, while offering a discrete segmentation, which is useful both for reducing dimensionality during optimisation and for inclusion of subjects based on the optimisation results.

We take the general framework, with respectively  $D$  discrete and  $C$  continuous covariates. The domain of each continuous covariate, indexed by  $c$ , is segmented in  $n_c^{(I)}$  intervals denoted  $\mathcal{I}_j^{(c)}$ ,  $j = 1, \dots, n_c^{(I)}$ . Combining those continuous intervals leads to  $n^{(I)} = \prod_c n_c^{(I)}$  possible intervals for  $Z_{Cont}$ . Finally, there are  $L = n^{(QD)} \times n^{(I)}$  possible distributions for the covariate vector  $Z$ .

Conditionally to  $Z_{Dis} = z_{Dis,d}$ , the distribution of  $Z_{Cont} = (Z_1 \in \mathcal{I}_{j_{c_1}}^{(c_1)}, \dots, Z_C \in \mathcal{I}_{j_{c_C}}^{(c_C)})$  is denoted  $p_{Cont, |z_{Dis,d}, \mathcal{I}(j_{c_1}, \dots, j_{c_C})}$ . The proportion of subjects within the trial population and for which  $Z_{Dis,i} = z_{Dis,d}$  and  $Z_{Cont,i} \sim p_{Cont, |z_{Dis,d}, \mathcal{I}(j_{c_1}, \dots, j_{c_C})}$  is denoted:  $x_{|z_{Dis,d}, \mathcal{I}(j_{c_1}, \dots, j_{c_C})}$ .

This segmentation leads to the elementary FIM formula given in equation (8), which now depends on  $\mathbf{x}$ .

$$M_F(\xi, \Psi, p_Z(\mathbf{x})) = \sum_{d=1}^{n^{(QD)}} \sum_{i_1=1}^{n_{c_1}^{(I)}} \dots \sum_{i_{n_C}=1}^{n_{c_C}^{(I)}} x_{|z_{Dis,d}, \mathcal{I}(j_{c_1}, \dots, j_{c_C})} M_F(\xi, \Psi, z_{Dis,d}, p_{Cont, |z_{Dis,d}, \mathcal{I}(j_{c_1}, \dots, j_{c_C})}) \quad (8)$$

We denote  $\mathbf{x}$  the vector of size  $L$  containing the  $x_{|z_{Dis,d}, \mathcal{I}(j_{c_1}, \dots, j_{c_C})}$  and for a better clarity in the notations, we denote  $x_l$  the  $l^{th}$  element of  $\mathbf{x}$  and  $p_{Z,x_l}$  the associated distribution for  $Z$ . The overall distribution for  $Z$  is denoted  $p_Z(\mathbf{x})$ . These notations result in the FIM formula given in equation (9).

$$M_F(\xi, \Psi, p_Z(\mathbf{x})) = \sum_{l=1}^L x_l M_F(\xi, \Psi, p_{Z,x_l}) \quad (9)$$

Finally, the multivariate distribution of covariates within the trial is described as the weighting of  $L$  multivariate distributions, and it is these proportions that we are seeking to optimise.

### 2.4 FIM integration using copula and Gauss-Legendre Quadrature

We describe here how copula modelling and GLQ can be leverage to compute an elementary FIM  $M_F(\xi, \Psi, z_{Dis}, p_{Cont, |z_{Dis}})$ .

The first aspect of our method is to characterise the covariate distribution through copula modelling to account for correlations in FIM computation as already proposed [20].

It is well known that given a random vector  $(Z_1, Z_2, \dots, Z_C)$  which follows the distribution  $p_{Cont, |z_{Dis}}$  and has continuous marginals  $F_1, \dots, F_C$ , the random vector  $(U_1, U_2, \dots, U_C) = (F_1(Z_1), F_2(Z_2), \dots, F_C(Z_C))$  has marginals uniformly distributed on  $[0; 1]$ . By definition, the copula of  $(Z_1, Z_2, \dots, Z_C)$  is the cumulative distribution function (cdf)

of  $(U_1, U_2, \dots, U_C)$ . The Rosenblatt transform [27] denoted  $T_{Cont, |z_{Dis}}$  is defined in equation (10) and its inverse in equation (11).

$$T_{Cont, |z_{Dis}} : \begin{pmatrix} z_1 \\ \vdots \\ z_C \end{pmatrix} \mapsto \begin{pmatrix} \mathbb{P}(Z_1 \leq z_1) \\ \mathbb{P}(Z_2 \leq z_2 | Z_1 \leq z_1) \\ \vdots \\ \mathbb{P}(Z_C \leq z_C | Z_{C-1} \leq z_{C-1}, \dots, Z_1 \leq z_1) \end{pmatrix} = \begin{pmatrix} F_1(z_1) \\ F_2(z_2 | z_1) \\ \vdots \\ F_C(z_C | z_{C-1}, \dots, z_1) \end{pmatrix} \quad (10)$$

$$T_{Cont, |z_{Dis}}^{-1} : \begin{pmatrix} u_1 \\ \vdots \\ u_C \end{pmatrix} \mapsto \begin{pmatrix} F_1^{-1}(u_1) \\ F_2^{-1}(u_2 | u_1) \\ \vdots \\ F_C^{-1}(u_C | u_{C-1}, \dots, u_1) \end{pmatrix} \quad (11)$$

Therefore, applying the inverse Rosenblatt transform to a vector of random uniform variables return a vector that has the same distribution as  $(Z_1, Z_2, \dots, Z_C)$ .

Consequently,  $M_F(\xi, \Psi, z_{Dis}, p_{Cont, |z_{Dis}})$  can be expressed using  $T_{Cont, |z_{Dis}}^{-1}$  as given in equation (12).

$$M_F(\xi, \Psi, z_{Dis}, p_{Cont, |z_{Dis}}) = \int_{[0;1]^C} M_F(\xi, \Psi, z_{Dis}, T_{Cont, |z_{Dis}}^{-1}(u_1, \dots, u_C)) du_1 \dots du_C \quad (12)$$

The second aspect is to approximate this integral through a GQ [23]. The latter consists in approximating an integral by a weighted sum of the function to be integrated, evaluated at a number  $n^{(Q)}$  of given points in the integration domain, called quadrature nodes. The advantage of GQ integration over the MC is that, since the nodes are chosen to represent the distribution harmoniously, equivalent accuracy is achieved with fewer GQ nodes than with random draws for MC [28].

In our case the integration domain for each of the  $C$  integrals is  $[0; 1]$ . Therefore, we apply  $C$  GLQ with a substitution, as GLQ allows to integrate over the interval  $[-1, 1]$ . Denoting  $v_q$  and  $w_q$  the  $q^{th}$  node and its weight in GLQ, the substitutions are:  $v'_q = \frac{1}{2}(1 + v_q)$  and  $w'_q = \frac{w_q}{2}$ .

Finally,  $M_F(\xi, \Psi, z_{Dis}, p_{Cont, |z_{Dis}})$  can be expressed as given in equation (13).

$$M_F(\xi, \Psi, z_{Dis}, p_{Cont, |z_{Dis}}) \approx \sum_{q_1=1}^{n^{(Q)}} \dots \sum_{q_C=1}^{n^{(Q)}} w'_{q_1} \dots w'_{q_C} M_F(\xi, \Psi, z_{Dis}, T^{-1}(v'_{q_1}, \dots, v'_{q_C})) \quad (13)$$

This methodology can thereafter be applied to compute each of the  $L$  matrices that appears in equation (9).

## 2.5 Optimisation of covariate distributions

We describe here the proposed algorithm to optimise the proportions  $x_l$  of the  $L$  distributions of covariates within the population.

### 2.5.1 Optimisation criteria

In order to compare the FIM between different designs, a criteria mapping the matrix to a scalar needs to be defined. Among various optimality criterion used in theory of optimal design (see for instance [4]), the D-criterion is a popular one and is widely used in the field of pharmacometrics. It uses the determinant of the FIM, as given in equation (14) and maximizing it ensures that the parameter estimates are as precise as possible. Indeed, a larger determinant of the FIM corresponds to a smaller volume of the confidence ellipsoid for the parameter estimates, indicating higher precision.

$$\phi_D(\Xi, \Psi, p_Z(\mathbf{x})) = \left( \det(M_F(\Xi, \Psi, p_Z(\mathbf{x}))) \right)^{\frac{1}{P}} \quad (14)$$

The comparison of covariate repartition  $\mathbf{x}$  against a reference  $\mathbf{x}_{ref}$  for the D-criterion is done by calculating the D-efficiency  $E_D$  (equation (15)).

$$E_D(\Xi, \Psi, p_Z(\mathbf{x})_{ref}, \mathbf{x}) = \frac{\phi_D(\Xi, \Psi, p_Z(\mathbf{x}))}{\phi_D(\Xi, \Psi, p_Z(\mathbf{x})_{ref})} \quad (15)$$

### 2.5.2 Optimising covariate proportions using Projected Gradient Descent

Our aim is to maximize the D-criterion, for a fixed design  $\Xi$  by balancing between the different covariate distributions weighted by the  $x_l$ , and given some constraints on these proportions.

**Linear constraints** First, as  $\mathbf{x}$  represents a vector of proportions, it should sum to 1 and each  $x_l$  has to be in  $[0; 1]$ . Additional constraints can be set on the  $x_l$ , for instance to make sure that a given interval for a continuous covariate does not represent more than a desired proportion and not less than another proportion. Therefore there are both equality and inequality constraints, and the optimisation problem is given in equation (16). We denote  $\mathcal{Q}$  the constraint space.

$$\begin{aligned} \min \quad & -\phi_D(\Xi, \Psi, p_Z(\mathbf{x})) \\ \text{s.t.} \quad & E\mathbf{x} = F \\ & G\mathbf{x} \geq H \end{aligned} \quad (16)$$

**Projected Gradient Descent** As the D-criterion is a convex function, the PGD [24] can be used to solve the problem given in equation (16).

Starting from an initial point  $\mathbf{x}_0 \in \mathcal{Q}$ , PGD iterates the equation given in (17), where  $\alpha_k \in [0; \infty]$  is the gradient step-size at the  $k^{\text{th}}$  iteration,  $\nabla\phi_D$  the gradient of  $\phi_D$  with respect to  $\mathbf{x}$  and  $\mathcal{P}_{\mathcal{Q}}$  is the projection on the constraint space. This projection is also the solution of a minimization problem, given in equation (18).

$$\mathbf{x}_{k+1} = \mathcal{P}_{\mathcal{Q}}\left(\mathbf{x}_k + \alpha_k \nabla\phi_D(\Xi, \Psi, p_Z(\mathbf{x}_k))\right) \quad (17)$$

$$\mathcal{P}_{\mathcal{Q}}(\mathbf{x}) = \min \|\mathbf{x} - \mathbf{y}\| \quad \text{s.t.} \quad E\mathbf{y} = F, G\mathbf{y} \geq H \quad \iff \quad \mathcal{P}_{\mathcal{Q}}(\mathbf{x}) = \min \frac{1}{2}\mathbf{y}^\top \mathbf{y} - \mathbf{x}^\top \mathbf{y} \quad \text{s.t.} \quad E\mathbf{y} = F, G\mathbf{y} \geq H \quad (18)$$

PGD iterates until a stopping condition is met. The detailed computation for  $\nabla\phi_D(\Xi, \Psi, p_Z, \mathbf{x}_k)$  when all the subjects have the same elementary design  $\xi$  is given in [29] and the result is recall in equation (19), where  $\text{tr}$  denotes the trace operator.

$$\frac{\partial\phi_D(\Xi, \Psi, p_Z(\mathbf{x}))}{\partial x_l} = \frac{N}{P} \text{tr} \left( M_F(\Xi, \Psi, p_Z(\mathbf{x}))^{-1} M_F(\xi, \Psi, p_{Z, x_l}) \right) \det \left( M_F(\Xi, \Psi, p_Z(\mathbf{x})) \right)^{1/P} \quad (19)$$

## 2.6 Optimisation of sample size

After having optimised the proportion of each covariate distribution, we can compute the sample size needed,  $NSN$ , all other things being equal, to reach desired confidence level on parameter estimates or on power of statistical tests on covariate effect.

Denoting  $SE(\Xi, \Psi, p_Z(\mathbf{x}))_\beta$  the standard error on a parameter  $\beta \in \Psi$  for a design  $\Xi$ , population vector  $\Psi$  and a covariate distribution defined by  $p_Z$  and  $\mathbf{x}$ , the number of subjects needed  $NSN$  to reach a desired level  $SE^*$  on this parameter is given in equation (20).

$$NSN(\Xi, \Psi, p_Z(\mathbf{x}))_\beta = N \left( \frac{SE(\Xi, \Psi, p_Z(\mathbf{x}))_\beta}{SE^*} \right)^2 \quad (20)$$

From this, we can calculate the  $NSN$  required to achieve the desired power for significance tests on covariates [20, 30], as well as relevance tests on covariate relationships [20] for a given equivalence interval  $[R_{inf}; R_{sup}]$ . The non-relevance of a covariate effect is assessed by a TOST procedure at level  $\alpha$ , with the null hypothesis is  $H_0$ : "the covariate effect is relevant", i.e.  $r_{l,c} \notin [R_{inf}; R_{sup}]$  while the alternative hypothesis is  $H_1$ : "the covariate is not relevant", i.e.  $r_{l,c} \in [R_{inf}; R_{sup}]$ .

This two sided null hypothesis can be split into two, respectively  $H_{0,inf}$  and  $H_{0,sup}$ :

$$\begin{cases} H_{0,inf} : r_{l,c} \leq R_{inf} \quad \text{or} \quad H_{0,sup} : r_{l,c} \geq R_{sup} \\ H_1 : r_{l,c} > R_{inf} \quad \text{and} \quad r_{l,c} < R_{sup} \end{cases}$$

Thus  $H_0$  is rejected if both  $H_{0,inf}$  and  $H_{0,sup}$  are rejected.

For log-normally distributed parameter with additive covariate relationships on the log-scale,  $r_{l,c} = e^{\beta_{l,c}(z - z_{ref})}$  and at level  $1 - \alpha$ , the null hypothesis is rejected if  $\widehat{\beta}_{l,c} + q_{1-\alpha} SE_{\beta_{l,c}} > B_{inf}$  and if  $\widehat{\beta}_{l,c} - q_{1-\alpha} SE_{\beta_{l,c}} < B_{sup}$  where depending on the sign of  $z - z_{ref}$ ,  $B_{sup}$  and  $B_{inf}$  equal  $\frac{\log R_{sup}}{z - z_{ref}}$  or  $\frac{\log R_{inf}}{z - z_{ref}}$ . The power is given by the equation (21), which holds if  $2q_{1-\alpha} < \frac{B_{sup} - B_{inf}}{SE_{\beta_{l,c}}}$ .

$$P_{nonrelev} = \Phi \left( -q_{1-\alpha} + \frac{B_{sup} - \beta_{l,c}}{SE_{\beta_{l,c}}} \right) - \Phi \left( q_{1-\alpha} + \frac{B_{inf} - \beta_{l,c}}{SE_{\beta_{l,c}}} \right) \quad (21)$$

The  $NSN$  to reach a desired non-relevance power  $P^*$  is found using a round finder approach to solve equation (22).

$$P^* = \Phi \left( -q_{1-\alpha} + \sqrt{\frac{NSN}{N}} \frac{B_{sup} - \beta_{l,c}}{SE_{\beta_{l,c}}} \right) - \Phi \left( q_{1-\alpha} + \sqrt{\frac{NSN}{N}} \frac{B_{inf} - \beta_{l,c}}{SE_{\beta_{l,c}}} \right) \quad (22)$$

## 2.7 Implementation

All these methods have been implementing using the R language [25]. For FIM evaluation an extension of the package *PFIM6*, based on previous work [20] have been done. All scripts are available online at NEWZENODO. The quadrature nodes were computed using the function *gauss.quad()* from the package *statmod* [31]. For PGD, the projection on the constraint space was made using the function *lse()* from the package *limSolve* [32].

## 3 Evaluation example using a simple PK model

In this section we explored an example to demonstrate the benefit of covariate distribution optimisation in reducing the sample size needed to draw conclusions about the relevance of covariate relationships.

### 3.1 Settings

#### 3.1.1 Model

The PK model was a one compartment model with IV bolus and linear elimination,  $f(t) = \frac{Dose}{V} e^{-\frac{Cl}{V}t}$ , with two parameters: the clearance ( $Cl$ ) and the volume of distribution ( $V$ ). The residual error was modelled as a combined error model:  $g = a + bf$  and exponential random effects without correlation were assumed for  $V$  and  $Cl$ ,  $\eta_{V,i} \sim \mathcal{N}(0, \omega_V^2)$  and  $\eta_{Cl,i} \sim \mathcal{N}(0, \omega_{Cl}^2)$ . All covariate effects were modelled as additive on the log-scale:  $\log \theta_i = \log \theta_{pop} + \beta_{\theta,Z} Z_i + \eta_{\theta,i}$ .

Firstly, three univariate scenarios were considered, with the model including either an effect of the Sex ( $SEX$ ) on  $V$ , an effect of the Body Mass Index ( $BMI$ ) on  $V$ , or an effect of the creatinine clearance ( $CLCR$ ) on  $Cl$ . Both  $BMI$  and  $CLCR$  were first normalised and log transformed, applying respectively  $\log\left(\frac{BMI_i}{BMI_{pop}}\right)$  and  $\log\left(\frac{CLCR_i}{CLCR_{pop}}\right)$ , where  $BMI_{pop}$  and  $CLCR_{pop}$  respectively denoted the average  $BMI$  and  $CLCR$  among Male individuals with Normal RF and Healthy Weight for  $BMI$ . These scenarios are hereafter referred to as ‘ $SEX$  only’, ‘ $BMI$  only’ and ‘ $CLCR$  only’. Then, the three effects were taken into account simultaneously, and this scenario is referred to as the ‘Three covariates’ scenario.

#### 3.1.2 Covariate distribution specification

Because covariate data may not be available at design step, we used the public database from the National Health and Nutrition Examination Surveys (NHANES) database [26] to get prior covariate distributions. Covariate data from 2009 to 2020 were extracted, keeping only subjects between 18 and 80 years old, with  $BMI$  higher or equal to  $18.5 \text{ kg/m}^2$  and for whom there were no missing data for Age, Body Weight (BW),  $BMI$ , Height, Creatinine, and  $SEX$ . The body weight was adjusted for ideal body weight computed with Robinson’s formula [33] and the Cockcroft and Gault equation with adjusted body weight [34] was then used to estimate the creatinine clearance,  $CLCR$ . 29 463 covariate vectors were selected in the NHANES database and Table 1 provides the summary statistics of these covariates.

The  $BMI$  was segmented into three usual intervals, namely Obesity if  $BMI \geq 30 \text{ kg/m}^2$ , Overweight if  $25 \leq BMI < 30 \text{ kg/m}^2$  and Healthy Weight if  $18.5 \leq BMI < 25 \text{ kg/m}^2$ . The  $CLCR$  was segmented into the four usual intervals to define the renal function (RF), namely Normal RF if  $CLCR \geq 90 \text{ mL/min}$ , Mild RF if  $60 \leq CLCR < 90 \text{ mL/min}$ , Moderate RF if  $30 \leq CLCR < 60 \text{ mL/min}$  and Severe RF if  $CLCR < 30 \text{ mL/min}$ . Of note,  $BMI_{pop} = 22.9 \text{ kg/m}^2$  and  $CLCR_{pop} = 112.2 \text{ mL/min}$ .

While there was a good balance between Male and Female and between the different  $BMI$  intervals, Severe RF was poorly represented with only 1.2% of the data.

Combining  $SEX$ ,  $BMI$  intervals and  $CLCR$  intervals, there were  $2 \times 4 \times 3 = 24$  possible covariate combinations. Nevertheless, those including Severe RF were poorly represented in the dataset (e.g. only 40 subjects are Male, with Severe RF and Obesity). Therefore, Severe RF individuals were pooled into two classes depending on their  $SEX$  and without taking into account their  $BMI$  status. This pooling was carried out to ensure the adequacy of copula fits. The detailed repartition between the resulted 20 covariate combinations is given in the first panel of Figure 7.

#### 3.1.3 Parameter values

The PK parameter values are given in Table 2, such as ratios of change caused by being a Female on  $V$ , being Obese on  $V$  and having Severe RF on  $Cl$ . With ratios of 0.70, i.e. outside of  $[0.80; 1.25]$ , both the effect of being a Female on  $V$  and having Severe RF on  $Cl$  were tested for relevance; while  $r_{V,BMI}(\text{Obesity})$  was within the equivalence interval, thus the effect of being Obese on  $V$  was tested for non-relevance. It should be noted that  $r_{V,BMI}(\text{Obesity})$  and  $r_{Cl,CLCR}(\text{Severe RF})$  were computed for the threshold values for belonging to these intervals, namely for  $30 \text{ kg/m}^2$  and  $30 \text{ mL/min}$ .

Table 1: Covariate data summary from the NHANES database, 2009-2020

Continuous covariates		
Covariate	mean (sd)	median [min; max]
<i>BW</i> (kg)	82.4 (21.8)	78.9 [37.5; 254.3]
<i>BMI</i> (kg/m <sup>2</sup> )	29.5 (7.0)	28.2 [18.5; 92.3]
Age (years)	48.3 (18.3)	48 [18; 80]
<i>CLCR</i> (ml/min)	99.0 (35.6)	97.3 [4.1; 342.9]
Discrete covariates		
Covariate	Category Name	N (%)
<i>SEX</i>	Male	14368 (48.8)
	Female	15095 (51.2)
Continuous segmentation		
Condition on covariate	Interval Name	N (%)
$18.5 \leq BMI < 25 \text{ kg/m}^2$	Healthy Weight	8234 (27.9)
$25 \leq BMI < 30 \text{ kg/m}^2$	Overweight	9635 (32.7)
$BMI \geq 30 \text{ kg/m}^2$	Obesity	11594 (39.4)
$CLCR \geq 90 \text{ mL/min}$	Normal RF	17221 (58.4)
$60 \leq CLCR < 90 \text{ mL/min}$	Mild RF	8343 (28.3)
$30 \leq CLCR < 60 \text{ mL/min}$	Moderate RF	3540 (12.0)
$CLCR < 30 \text{ mL/min}$	Severe RF	359 (1.2)

*BW*: refers to Body Weight, *BMI*: Body Mass Index, *CLCR*: Creatinine Clearance

Table 2: Population PK parameter values used in the evaluation example

Parameter	$\mu_{Cl}$ (L/h)	$\mu_V$ (L)	$\omega_{Cl}$	$\omega_V$	$a$ (mg/L)	$b$	$\beta_{V,SEX}$	$\beta_{V,BMI}$	$\beta_{Cl,CLCR}$
Value	0.43	7.13	0.30	0.24	2.39	0.08	-0.35	0.50	0.27
Ratio	$r_{V,SEX}$	$r_{V,BMI}$ (Obesity)	$r_{Cl,CLCR}$ (Severe RF)						
Value	0.70	1.14	0.70						

$\mu_{\cdot}$ : refers to the typical value,  $\beta_{\cdot}$ : covariate parameter,  $\omega$ : standard deviation of the random effect,  $a$  and  $b$ , parameters of the residual error model  
 $r_{l,c}$ : ratio on the parameter  $l$  for the covariate  $c$

### 3.1.4 Design

The considered design includes  $N = 24$  subjects receiving 250 mg of drug at time 0, and for each of them, three samples are collected at 1, 4 and 12 hours post-dose. The concentration curves  $f(t)$  accounting for univariate covariate effects and with parameters given in Table 2 are shown on Figure 1. Because being a Female reduces the volume, the typical concentration curve for Female is above the Male one. Because the higher the *BMI* the higher the volume, the typical area for concentration curve in individual with Healthy Weight is above the one for Overweight, which is above the one for Obesity. Regarding *CLCR* status, the lower the *CLCR*, the lower the typical curve, since renal impairment decreases *CLCR*.

## 3.2 FIM evaluation using copula and quadrature

### 3.2.1 Copula fitting

In the example with *SEX* only (i.e. without continuous covariate), the population FIM was computed for each *SEX* and the total FIM was obtained as the weighted sum, each FIM weighted by the proportion of the corresponding *SEX* in the database. With *BMI* only, three copula were fitted: one for each of the three datasets corresponding to *BMI* intervals. In the same way, in the example with *CLCR* only, four copula were fitted: one for each of the four datasets corresponding to *CLCR* intervals.

For the example with three covariates, a copula was fitted for each of the 20 covariate combination on the associated subset of data.

All copula were fitted as vine copula, using the R package *rvinecopulib* 0.6.3.1.1 [35]. Copula fit evaluation was conducted through a simulation-based strategy as described in [36] using 100 replicates. The relative error (RE) between summary






Figure 1: Evolution of the concentration  $f(t)$  according to the fixed effects in the three univariate scenarios: the first panel corresponds to an effect of  $SEX$  on  $V$ , the second panel to an effect of  $BMI$  on  $V$  and the third panel to an effect of  $CLCR$  on  $Cl$ . For the second and third panels, the black lines correspond to the limits of the intervals of the continuous covariates.

metrics  $M_{sim}$  in copula-simulated populations and  $M_{obs}$  in the NHANES dataset was computed as  $\frac{M_{sim} - M_{obs}}{M_{obs}}$ . Summary metrics were the mean, the standard deviation, the median and the 5<sup>th</sup> and 95<sup>th</sup> percentiles. Overlaps between simulated and true distributions were also computed such as the simulated and observed correlations between  $BMI$  and  $CLCR$ . Diagnostics plots were explored using the R package *pmxcopula* [37] with minor edits.

### 3.2.2 Quadrature settings

According to the Gauss-Legendre Quadrature rule, the approximation using  $n^{(Q)}$  nodes is exact for integrands that are polynomials of degree  $2n^{(Q)} - 1$  or less. Nevertheless, in our case, the integrand is the composition of the FIM and the inverse Rosenblatt transform of the copula describing the continuous covariate distribution, therefore, in a general setting, it is not a polynomial function. Consequently, the quadrature remains an approximation and increasing  $n^{(Q)}$  increases the accuracy.

To determine the number of nodes required in the quadrature for an acceptable accuracy in FIM evaluation, we computed the FIM for  $n^{(Q)}$  between 1 and 25, and looked for stabilization in the D-criterion. The same number of nodes was used for all the continuous covariates and for each copula. As a benchmark, we also computed the FIM using copula with Monte Carlo (MC) simulations. This computation was performed using  $n^{(MC)} = 10, 100, 500, 1000,$  and  $2000$  Monte Carlo samples drawn from each copula. Additionally, since it is a stochastic method, the process was repeated  $S = 100$  times to assess the uncertainty of the computed FIM. The target value was chosen as the average  $\phi_D$  obtained with 1000, and 2000 Monte-Carlo samples, and the RE on the D-criterion was computed with respect to this value. In the same time, we computed the run time as a function of  $n^{(Q)}$ , and the average run time in MC procedure as a function of the number of samples.

This comparison was performed for both  $BMI$  only and  $CLCR$  only, and because of computation already done, optimisation was then carried out with the higher number of nodes, i.e.  $n^{(Q)} = 25$ . Because of reasonable run time,  $n^{(Q)} = 25$  was also used for the Three covariate scenario and MC approach using  $n^{(MC)} = 500$  was also explored as a reference.

## 3.3 Covariate distribution optimisation

In all scenarios, covariate distribution optimisation was performed using PGD. The step-size  $\alpha$  was constant and set to 0.001 and the stopping condition was that the objective function should not change by more than  $\epsilon = 10^{-7}$  between two iterations.

In univariate cases, optimisation was done for various values of the covariate effects, while the other PK parameter values were the same as those given in Table 2.

Different constraints were applied depending on the scenario.

### 3.3.1 SEX only

The optimal balance between Male and female was explored for  $\beta_{V,Sex} \in [-1.20; 0.00]$  with a 0.05 step, with no other constraint than that the proportions should be within  $[0; 1]$  and sum to 1.

### 3.3.2 BMI only

Optimisation between the three *BMI* intervals was explored for  $\beta_{V,BMI} \in [0.00; 1.00]$  with a 0.05 step, and for two optimisation settings. First without any other constraints than proportion definition. Then, optimisation among the three *BMI* intervals was performed with the additional constraint that each interval should represent at least 5% of the population. This setting is hereafter referred to as 'Lower constraints'.

### 3.3.3 CLCR only

Optimisation between the four *CLCR* intervals was explored for  $\beta_{Cl,CLCR} \in [0.00; 1.00]$  with a 0.05 step. First, the two same settings as *BMI* only were explored. In addition, because Severe RF subjects are rare in the population, a third optimisation scenario was explored by adding to the lower limit constraint an upper limit set at 10% for Severe RF in the trial population. This setting is hereafter referred to as 'Lower and Upper constraints'.

### 3.3.4 Three covariates

For the case with the three covariates, the three same optimisation settings as for *CLCR* only were explored.

## 3.4 Optimised distributions evaluation

The initial distribution between the *SEX*, *CLCR* and *BMI* intervals corresponds to the observed proportions in the selected subset from the NHANES database.

For each optimised distribution, the D-Efficiency relative to the design with the initial distribution was computed, in addition with the *NSN* to conclude with a 80% power on the statistical significance for each covariate effects, and the *NSN* to conclude with a 80% power on the relevance (or non relevance) of relationship for the effect of being a Female on *V*, of having Obesity on *V* and of having Severe RF on *Cl*.

## 4 Results

### 4.1 Copula fitting

The performance of copula model was evaluated for each covariate combination.

#### 4.1.1 BMI only

For the scenario with *BMI* only, mean, standard deviation, median and 5<sup>th</sup> and 95<sup>th</sup> percentiles of the simulated population agreed with the observed one, with a small RE within  $\pm 2.5\%$  (Appendix A.1.1, Figure A.1.1). Visual predicted checks (VPC) were also in accordance with the observed percentiles (Appendix A.1.1, Figure A.1.2).

#### 4.1.2 CLCR only

Similarly, with *CLCR* only, RE on summary statistics was satisfactory, within  $\pm 5\%$  for all the intervals (Appendix A.1.2, Figure A.1.3) and VPC were also in accordance with the observed percentiles (Appendix A.1.2, Figure A.1.4).

#### 4.1.3 Three covariates

As shown in Appendix A.1.3, Figure A.1.5, RE was within  $\pm 10\%$  for all the summary statistics for almost all the intervals of *BMI* and *CLCR*. Indeed the RE on the 5<sup>th</sup> or 95<sup>th</sup> percentiles of the simulations were a little higher for *sd* on *BMI* for Male, with Obesity and Moderate RF; for *sd* on *BMI*, *P5%* and *P50%* on *CLCR* for both Male and Female with Moderate RF. The simulated correlations from the copula models were very similar to observed correlations between *BMI* and *CLCR* (Appendix A.1.3, Figure A.1.7) and overlap between simulated and observed joint distribution for *BMI* and *CLCR* was always above 80%, except for Female with Severe RF, for which the overlap was above 80% in only in three quarters of the simulations (Appendix A.1.3, Figure A.1.6). Moreover, the VPC given in Figure 2 show a good match between simulated and observed level lines, for all combinations.

Overall, copula fits were judged to be very satisfactory.




Figure 2: Three Covariates - Visual predictive checks based on the contours of the bivariate density between  $BMI$  and  $CLCR$ . For each of the 20 covariate combination, the virtual population was simulated 100 times from the copula and the 99% prediction intervals of percentile contours of the joint distribution was derived and compared to the contours observed in the NHANES database.

The ribbon areas correspond to the 99% prediction interval of 5<sup>th</sup> (●), 50<sup>th</sup> (●) and 95<sup>th</sup> (●) percentile contours of the joint distribution of  $BMI$  and  $CLCR$ . The lines corresponds to the observed 5<sup>th</sup> (—), 50<sup>th</sup> (—) and 95<sup>th</sup> (—) percentile contours.

## 4.2 FIM evaluation using quadrature

### 4.2.1 $BMI$ only

As shown in Figure 3, quadrature method was much faster than MC. Indeed, with 25 nodes, the FIM computation was perform in 1.7 seconds, whereas even with only 100 MC samples, the average run time was 6.4 seconds, and it increased to 63.8 and 126.8 seconds for respectively 1000 and 2000 MC samples.

Regarding the accuracy in  $\phi_D$  computation, the first observation is that MC integration led to a non-negligible uncertainty when the number of MC samples  $n^{(MC)}$  is low (Figure 3 and Figure A.2.1 given in Appendix A.2.1). In our example, the uncertainty is however very similar between 500, 1000 and 2000 MC sample, suggesting that 500 MC sample are acceptable for the FIM computation. For the quadrature approach, the RE to the target, defined as the average  $\phi_D$  over the repetitions with 1000 or 2000 MC sample, was below 1% starting from 4 quadrature nodes and higher (Figure 3).




Figure 3: *BMI* only - Run time for  $\phi_D$  computation (top) and relative error ( $RE$ ) in  $\phi_D$  computation (bottom), with FIM integration using on the left copula and MC and on the right copula and GLQ. The target value was the average  $\phi_D$  obtained with  $n^{(MC)} = 1000$  and  $n^{(MC)} = 2000$ .

The green line corresponds to 0. The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. The diamond corresponds to the mean.

$\phi_D$ : refers to the D-criterion, FIM: Fisher Information Matrix MC: Monte-Carlo, GLQ: Gauss-Legendre Quadrature

#### 4.2.2 CLCR only

Once again, GQ method was much faster than MC with a run time of 2.3 seconds with 25 nodes whereas even with only 100 MC samples, the average run time was 8.6 seconds, and it increased to 86.0 and 170.4 seconds for respectively 1000 and 2000 MC samples (Figure A.2.3 given in Appendix A.2.2).

The uncertainty in  $\phi_D$  computation with the MC approach was also very similar between 500, 1000 and 2000 MC sample, suggesting that 500 MC sample are acceptable for this FIM computation (Figure A.2.2 given in Appendix A.2.2).

With GQ, the relative error to the target was below 1% starting from 3 quadrature nodes and higher (Figure A.2.3 given in Appendix A.2.2).

#### 4.2.3 Three Covariates

In this scenario, as the number of covariate combinations was higher, the run time was longer for  $n^{(Q)} = 25$ , 533.0 seconds than the average run time with 500 MC samples: 412.7 seconds ( $sd = 2.1$ ). Both methods gave very similar results with an average D-criterion of 62.5 with 500 MC samples ( $sd = 0.2$ ) and 62.5 with  $n^{(Q)} = 25$  (Figure A.2.4 given in Appendix A.2.3).

### 4.3 Covariate distribution optimisation

#### 4.3.1 SEX only

Optimal balance between Male and Female for the D-criteria and as a function of  $\beta_{V,SEX}$  is given in Figure 4. The optimal distribution did not vary a lot and remained close to the 50-50 balance. For  $\beta_{V,SEX} = 0$ , the optimal balance was perfect equilibrium between Male and Female. When  $\beta_{V,SEX}$  decreased, the optimal weight for Female increased up to 67.8% for  $\beta_{V,SEX} = -1.20$ . For the value used in the scenario with the three covariates, i.e.  $\beta_{V,SEX} = -0.35$ , the optimal Female proportion in this univariate case was 62.0%.

In Figure 4 are given the D-criterion, the D-Efficiency relative to the initial design, the  $NSN$  to conclude on the significance of  $\beta_{V,SEX}$ , and the  $NSN$  to conclude on the relevance/non-relevance (when appropriate) of  $\beta_{V,SEX}$  are given as a function of  $\beta_{V,SEX}$  for both the initial design (51% of Females) and the Optimal design. In this context, the benefit of using the optimal covariate distribution was very small. As expected,  $NSN_{signif}(\beta_{V,SEX})$  and  $NSN_{relev}(V, SEX, (Female))$  decreased when  $\beta_{V,SEX}$  get further from 0. For the value used in the scenario with the three covariates, the relative D-Efficiency was quite low: only 1.01. The optimal design increased slightly the uncertainty on  $\beta_{V,SEX}$  therefore  $NSN_{signif}(\beta_{V,SEX})$  which initially was 28 became 30, and similarly  $NSN_{relev}(V, SEX, (Female))$  has been increased from 165 to 176.




Figure 4: *SEX* only - Optimal proportions for the covariate distributions for the D-criterion (top),  $\phi_D$ , D-Efficiency and *NSN* to reach 80% power (bottom), as a function of  $\beta_{V,SEX}$

The purple vertical line corresponds to the value used in the Three covariates scenario. The grey area corresponds to the non-relevance interval.  $\phi_D$ : refers to the D-criterion, *NSN*: Number of subjects needed

### 4.3.2 BMI only

The optimal distribution between the three *BMI* intervals as a function of  $\beta_{V,BMI}$  is given in Figure 5. With no constraints on the representation of all the statuses, we can see that the balance was only between the extremes: when  $\beta_{V,BMI} = 0$ , the optimal was having 53.7% of Obese and 46.3% of Healthy Weight. When the *BMI* effect increased, the optimal proportion of Obese individual decreased, being 36.0% for  $\beta_{V,BMI} = 1$ . With the lower constraint of having at least 5% of each interval, the optimal proportion of Overweight was always the minimum of 5%. With this lower constraint, the proportion of Obesity decreased less as the *BMI* effect increased (-2.2% when  $\beta_{V,BMI} = 0$  and -1.7% when  $\beta_{V,BMI} = 1$ ).

For the value used in the scenario with the three covariates, i.e.  $\beta_{V,BMI} = 0.5$ , the optimal Obesity proportion without constraint was 39.7% and with the Lower constraints, 38.0%.

The higher  $\beta_{V,BMI}$ , the more there was benefit in using optimal distribution, no matter the constraint, in terms of overall estimation as the D-Efficiency increased with  $\beta_{V,BMI}$ , from 1.05 when  $\beta_{V,BMI} = 0$  to respectively 1.15 and 1.14 when  $\beta_{V,BMI} = 1$  without and with lower constraints (Figure 5). Moreover, the addition of the lower constraints, which makes the distribution more realistic in terms of clinical requirements, was still very valuable, as the  $\phi_D$  curve for the lower constraints was much closer to the  $\phi_D$  curve without constraints than to the curve with the initial distribution. In the same way, the *NSN* to reach 80% in statistical tests with the lower constraints were closer to those without constraint than to those with the initial distribution. For high effect, the *NSN* to conclude of the significance of  $\beta_{V,BMI}$  was all ready low (18 for  $\beta_{V,BMI} = 1$ ) and decreased with optimal distribution (respectively 14 and 15 without and with constraints). Bellow  $\beta_{V,BMI} = \frac{\log(1.25)}{\log(\frac{30}{BMI_{pop}})}$ , the ratio of change caused by the Obesity on *V* was bellow 1.25, and obviously the *NSN* needed to conclude to the non-relevance of this effect decreased with  $\beta_{V,BMI}$ , and was minimal in 0. With the initial distribution,  $NSN_{nonrelev}(V, BMI, (Obesity)) = 28$ , and it decreased to 20 and 21 without and with constraints.

For the value used in the scenario with the three covariates,  $NSN_{signif}(\beta_{V,BMI})$  was respectively 70, 52 and 55 for the




Figure 5: *BMI* only - Optimal proportions for the covariate distributions for the D-criterion (top),  $\phi_D$ , D-Efficiency and *NSN* to reach 80% power (bottom), as a function of  $\beta_{V,BMI}$

The purple vertical line corresponds to the value used in the three covariates scenario. The grey area corresponds to the relevance interval.  $\phi_D$ : refers to the D-criterion, *NSN*: Number of subjects needed

initial, without constraint and with lower constraints distributions, while the  $NSN_{nonrelev}(V, BMI, (Obesity))$  was 130, 97, 101.

### 4.3.3 CLCR only

With *CLCR* only, a similar pattern as for *BMI*, with allocation to the extreme statuses was observed without constraint (Figure 6). Thus, for  $\beta_{Cl,CLCR} = 0$ , the optimal distribution was having 52.2% of Severe RF and 47.8% of Normal RF. When the *CLCR* effect increased, the optimal proportion of Severe individual decreased, first slowly until  $\beta_{Cl,CLCR} = 0.85$  (26.7%) then fell to 4.2% for  $\beta_{Cl,CLCR} = 1$ . In the same time, while the proportion of Moderate RF was null for  $\beta_{Cl,CLCR} = 0.85$ , it increased up to 19.3% for  $\beta_{Cl,CLCR} = 1$ . In other words, when the effect was stronger, less extreme small *CLCR* values were sufficient and better than the smaller ones.

When adding the lower constraint of having at least 5% of each *CLCR* interval, until  $\beta_{Cl,CLCR} = 0.85$ , 5% was the optimal for both Mild and Moderate RF. With a greater effect and as previously, the optimal proportion of Moderate increased up to 15.1 for  $\beta_{Cl,CLCR} = 1$ .

With the additional upper constraint of 10% for the Severe RF, there was a shift from Severe to Moderate, while Mild remained at the lower limit of 5%.

For the value used in the scenario with the three covariates, i.e.  $\beta_{Cl,CLCR} = 0.27$ , the optimal distribution Normal - Mild - Moderate - Severe were respectively 65.0 - 0 - 0 35.0 %, 58.6 - 5.0 - 5.0 - 31.4 % and 69.4 - 5 - 15.6 - 10.0 % without constraint, with lower constraints, and with lower and upper constraints.

As in the scenario with *BMI* only, adding the lower constraint on the proportions did not deteriorate too much the performance of the optimal distribution in terms of  $\phi_D$ , D-efficiency and *NSN*. Constraining Severe RF being below 10% reduced the information and therefore increased the *NSN* but was still valuable compared to the initial distribution. The greater difference in D-Efficiency when using an optimal distribution compared to the initial one was observed for a null




Figure 6: *CLCR* only - Optimal proportions for the covariate distributions for the D-criterion (top),  $\phi_D$ , D-Efficiency and *NSN* to reach 80% power (bottom), as a function of  $\beta_{CI,CLCR}$

The purple vertical line corresponds to the value used in the Three covariates scenario. The grey area corresponds to the non-relevance interval.  $\phi_D$ : refers to the D-criterion, *NSN*: Number of subjects needed

effect; because it was in this case than the optimal proportion of Severe RF was the higher while there were only 1.2% Severe in the initial distribution. Above,  $\beta_{CI,CLCR} = \frac{\log(1.25)}{\log(\frac{30}{CLCR_{pop}})}$ , the ratio of change caused by the Severe RF on *CLCR* was above 1.25, and logically the *NSN* needed to conclude to the relevance of this effect decrease with the increasing in  $\beta_{CI,CLCR}$ .

For the value used in the scenario with the three covariates,  $NSN_{signif}(\beta_{CI,CLCR})$  was respectively 121, 32, 35 and 62 for the initial, without constraint, with lower constraints and with lower and upper constraints distributions, while the  $NSN_{relev}(CI, CLCR, (Severe))$  was respectively 684, 179, 194 and 348. Therefore, the use of the more constrained optimal distribution still made it possible to reduce by almost 50% the *NSN* to conclude on the relevance of the effect of having a Severe RF on *Cl*.

#### 4.3.4 Three Covariates

The optimal distribution between the 20 covariate combinations in the scenario with three covariates are given in Figure 7 for the initial distribution and for the three optimisation settings. The first observation is that these distributions were very sparse in the sense that many categories had a weight of 0.

Because being a Female reduces *V* while *V* increases with the *BMI*, in optimal distributions, Female were more associated with Healthy Weight while Male were associated with Obesity. Indeed, this balance allows to maximise the difference in the theoretical concentration curves, higher for Healthy Female than for Obese Male. In the same way, because having a Severe RF decreases *Cl* and therefore slows the elimination and leads to higher concentration curves, in optimal distribution, only Female Severe RF were present. We thus find the same tendency to distribute between extremes as in the univariate analysis.

Marginal distributions are given in Appendix A.3 in Table 4. Regarding the *SEX*, no matter the constraint settings, the optimal balance is close to 66.3% Female, not so far from the 62.0% observed in the univariate case for *SEX* only.






Figure 7: Three Covariates - Optimal proportions for the covariate distributions for the D-criterion for the different sets of constraints

For the *BMI* intervals, even without constraint, there were Overweight in the optimal distribution because of the pool of the *BMI* statuses in the Severe RF combinations. Nonetheless, the proportions of Obesity were close to the univariate scenario: 42.0% without constraint and 39.7% in univariate for the optimisation without constraint, 40.6% and 38.0% with lower constraints, and finally 43.0% with additional upper constraint on Severe RF.

Once again for the *CLCR* intervals, when there were no constraint, Mild and Moderate RF were absent from the optimal distribution, and the balance Normal - Severe was 68.7 - 31.3% therefore close to 65.0 - 35.0 % from the univariate case. With lower constraints, Mild and Moderate were still to their minimum and with the additional upper constraint, Severe RF was replaced by Moderate RF.

Regarding performances, having both lower and upper constraints remained close to the two other optimisation settings in terms of D-Efficiency and still allowed to reduce the *NSN*. Indeed the higher *NSN* to conclude on the significance of the three covariates effects, on the relevance of the effect of being a Female on *V* and of the effect of having Severe RF on *Cl*, and on the non-relevance of having Obesity on *V* was 596 with the initial design and fell to respectively 197 for both without constraint and with lower constraint optimisation and to 229 with the additional upper bound on Severe. Thus, even the more constrained optimisation allowed to reduce the *NSN* by more than 60%.

Table 3: Three Covariates -  $\phi_D$ , D-Efficiency and *NSN* to reach 80% power in covariate testing with the initial and optimal covariate distributions for the D-criterion for the different sets of constraints

Name	Initial	Optimal designs		
	Design	Without constraint	Lower constraints	Lower and Upper constraints
$\phi_D$	62.47	77.1	76.11	74.56
D-Efficiency	1	1.23	1.22	1.19
$NSN_{signif}(\beta_{V,SEX})$	28	33	33	33
$NSN_{signif}(\beta_{V,BMI})$	69	<b>57</b>	<b>58</b>	<b>52</b>
$NSN_{signif}(\beta_{Cl,CLCR})$	<b>106</b>	26	28	41
$NSN_{relev}(V,SEX)$	167	<b>197</b>	<b>197</b>	196
$NSN_{nonrelev}(V,BMI (Obesity))$	129	106	108	97
$NSN_{relev}(Cl,CLCR (Severe RF))$	<b>596</b>	142	153	<b>229</b>

**in bold:** the higher *NSN* across the three covariate relationships, for respectively the significance tests and the (non-)relevance tests, for each design.  $\phi_D$ : refers to the D-criterion, *NSN*: Number of subjects needed

## 5 Discussion

From an application point of view, we proposed in this work to optimise the allocation of the covariates among the patients to be including in a future study. The main objective was to improve the estimation, but we were particularly interested in the effects of the covariates and the power of the statistical tests relating to them, as well as the *NSN* to conclude on their relevance with a sufficient degree of confidence. For relevance assessment, we used the equivalence region [0.80;1.25] which is the one used in bioequivalence studies to define whether a dosing adjustment is required but alternative methods have

been proposed to chose equivalence regions appropriate to a specific context [2]. This optimisation was carried out between categories for discrete covariates, while for continuous ones, we introduced a segmentation of their domains into intervals and optimised the proportion between them. In terms of methods, we have introduced a new way of integrating the FIM for the covariate distribution by leveraging copula modelling and GLQ, and we solved the optimisation problem using the PGD algorithm. All these methods have been implemented in R based on the PFIM 6.1 package, and the source code is available online at the following link: <https://doi.org/10.5281/zenodo.14778034>.

We applied the proposed workflow on a PK example with one discrete covariate and two continuous ones. First, univariate analysis showed classical and intuitive results, namely an equal distribution between the more extreme classes and a disappearance of intermediate values in cases where there are no constraints. Constraints were proposed to better reflect real life conditions for patients' inclusion in clinical studies. Even with these constraints, we showed that in the scenarios with the three covariate effects, the optimal distribution with the most realistic constraints allowed to reduce by more than 60% the *NSN* to conclude on the clinical relevance (or non-relevance) of the relationships. The challenge in implementing this optimal distribution is the need to include 23 females with Severe RF, while with the initial distribution, it was just 5 subjects without restriction on the *SEX*. Nevertheless, as the reduction in the sample size is otherwise significant, it may be worth concentrating on this difficult inclusion.

We have shown that using GLQ greatly reduces the computation time compared with MC, while providing good accuracy in FIM computation. However, the choice of the quadrature settings has not been examined in depth. Indeed, we compared the results depending on the number of quadrature nodes with the ones obtained with MC simulations; but using this approach in practice negates all the benefits of run time reduction. In practice, a suggestion can be to increase progressively the number of nodes and just look for a stabilization of the D-criterion without reference. In addition, we have seen that the computation time remains quite low even with a high number of nodes, in our case 25. Nonetheless, this holds for three covariates but if the number of continuous covariates  $C$  increases, due to the quadrature of each of them, the number of FIM evaluations  $(n(Q))^C$  increases exponentially. This difficulty could be countered, for example, by refining the number of nodes required for each distribution rather than using the same number for all the covariates, or by using adaptive methods that require fewer nodes (see for instance a review in [38]). However, this point should be qualified by the fact that, in practice, it seems unlikely that the number of covariates to be optimised will be very large. Moreover, it would still be faster than CTS, which requires simulation and then estimation on a large number of data sets.

We note that we did not explore the PGD hyper-parameters such as the step size nor some of performance-improving variants because the need was not felt in our situation, but it can be an option in more complex cases. However, in the current version of this algorithm, only linear constraints can be used for an easy projection on the constraint space, while in practice some non-linear ones may be useful (e.g., to impose the power of a test or a SE level on a parameter). To address this, a penalty term could be added to the objective function or another class of algorithms could be explored. Regarding the optimisation criteria, we only used the D-criterion while others exist such as the DS-criterion which allows focus on a specific subset of the parameters (see for instance [13] for an application). In our case, the covariates effects could have been the only parameters of interest. This change of criterion is not straightforward in the sense that its derivative has to be calculated, but apart from this calculation, it presents no particular difficulty.

In this work, we have assumed the same elementary design for all the subjects, but the computations can easily be extended to the case where there are several. It would therefore certainly be more beneficial to adapt the elementary design to the covariate combination. Then, an interesting path may be to jointly optimize the elementary designs and the covariates. If discrete optimization is performed for elementary designs, it can be combined with the covariates and the global process remains the same, only the number of FIM evaluations increases. We can easily imagine that this joint optimisation would enable us to improve the designs, for example in our case, for subjects with poor elimination, having a higher final observation time would guarantee a better estimation. On the other hand, for continuous optimization, the extension is harder. In both cases, it will first be necessary to consider the practical and ethical constraints and discuss the actual implementation with experts.

In our example we used a covariate database and the goodness of fit diagnosis suggested that a large amount of data is required but copulas could have been used directly if available, which is one of their great interests in pharmaceutical applications as they can be made public without threatening anonymity. In addition, the optimization approach can be coupled with another method for FIM computation if preferred.

Regarding FIM computation, we used a linearization approach but methods avoiding this approximation have already been explored using MC and GQ [39] [40]; and future work could extend them to include the integration of the covariate distribution.

Furthermore, we need to point out the possible difficulties in implementing these optimal design distributions. For example, it seems implausible not to include patients because they are not in the most extreme interval, when inclusion processes are already long and costly. The spirit of this optimisation would therefore be rather to aim towards the optimal in practice and above all to be aware of the conditions on the distribution of covariates and on the number of subjects that would be needed to reach a conclusion.

Last but not least, this approach assumes that the model and its parameters are known, including the effects of the

covariates, whereas the trial to be designed aims to better estimate them, and is not robust to misspecifications. In the univariate example, we computed optimal allocation and  $NSN$  for a range of  $\beta$  and this could be extended into a robust approach as already explored in the NLMEM context [10–13]. Moreover, this methodology could be used to design a Phase 3 study based on Phase 2 results; therefore, the SE from the covariate effects estimates based on the Phase 2 data may be used to specify a prior distribution on the  $\beta$ . This absence of robustness also holds for the covariate distribution assumed for FIM computation and covariate optimisation. In this case, Adaptive Designs and especially Two-stage approaches [10, 29, 41] could be a good avenue to explore for refining the distribution of covariates halfway through inclusion.

## References

1. Mould, D. R. & Upton, R. Basic concepts in population modeling, simulation, and model-based drug development. *CPT: Pharmacometrics & Systems Pharmacology* **1**, 1–14 (2012).
2. Sanghavi, K. *et al.* Covariate modeling in pharmacometrics: General points for consideration. *CPT: Pharmacometrics & Systems Pharmacology* **13**, 710–728 (2024).
3. *FDA Guidance for Industry Population Pharmacokinetics* <https://www.fda.gov/media/128793/download>. 2022.
4. Atkinson, A., Donev, A. & Tobias, R. *Optimum experimental designs, with SAS* (OUP Oxford, 2007).
5. Mentré, F. *et al.* Current use and developments needed for optimal design in pharmacometrics: a study performed among DDMoRe’s European Federation of Pharmaceutical Industries and Associations Members. *CPT: Pharmacometrics & Systems Pharmacology* **2**, 1–2 (2013).
6. Strömberg, E. A., Nyberg, J. & Hooker, A. C. The effect of Fisher information matrix approximation methods in population optimal design calculations. *Journal of Pharmacokinetics and Pharmacodynamics* **43**, 609–619 (2016).
7. Mentre, F., Mallet, A. & Baccar, D. Optimal design in random-effects regression models. *Biometrika* **84**, 429–442 (1997).
8. Retout, S., Mentré, F. & Bruno, R. Fisher information matrix for non-linear mixed-effects models: evaluation and application for optimal design of enoxaparin population pharmacokinetics. *Statistics in Medicine* **21**, 2623–2639 (2002).
9. Bazzoli, C., Retout, S. & Mentré, F. Fisher information matrix for nonlinear mixed effects multiple response models: evaluation of the appropriateness of the first order linearization using a pharmacokinetic/pharmacodynamic model. *Statistics in Medicine* **28**, 1940–1956 (2009).
10. Dodds, M. G., Hooker, A. C. & Vicini, P. Robust population pharmacokinetic experiment design. *Journal of Pharmacokinetics and Pharmacodynamics* **32**, 33–64 (2005).
11. Foo, L. K., McGree, J., Eccleston, J. & Duffull, S. Comparison of robust criteria for D-optimal designs. *Journal of Biopharmaceutical Statistics* **22**, 1193–1205 (2012).
12. Loingeville, F., Nguyen, T. T., Riviere, M.-K. & Mentré, F. Robust designs in longitudinal studies accounting for parameter and model uncertainties—application to count data. *Journal of Biopharmaceutical Statistics* **30**, 31–45 (2020).
13. Seurat, J., Nguyen, T. T. & Mentré, F. Robust designs accounting for model uncertainty in longitudinal studies with binary outcomes. *Statistical Methods in Medical Research* **29**, 934–952 (2020).
14. Bazzoli, C., Retout, S. & Mentré, F. Design evaluation and optimisation in multiple response nonlinear mixed effect models: PFIM 3.0. *Computer Methods and Programs in Biomedicine* **98**, 55–65 (2010).
15. Leroux, R., Seurat, J., Fayette, L., Bach, N. T. & Mentré, F. Design evaluation and optimisation in nonlinear mixed effects models with the R package PFIM 6.0. *PAGE* (2023).
16. Mentré, F., Leroux, R., Seurat, J. & Fayette, L. *PFIM: Population Fisher Information Matrix* R package version 6.1 (2024). <https://CRAN.R-project.org/package=PFIM>.
17. Foracchia, M., Hooker, A. C., Vicini, P. & Ruggeri, A. POPED, a software for optimal experiment design in population kinetics. *Computer Methods and Programs in Biomedicine* **74** (2004).
18. Nyberg, J. *et al.* PopED: An extended, parallelized, nonlinear mixed effects models optimal design tool. *Computer Methods and Programs in Biomedicine* **108** (2012).
19. Bauer, R. J., Hooker, A. C. & Mentre, F. Tutorial for \$ DESIGN in NONMEM: Clinical trial evaluation and optimization. *CPT: Pharmacometrics & Systems Pharmacology* **10**, 1452–1465 (2021).
20. Fayette, L., Brendel, K. & Mentré, F. Using Fisher Information Matrix to predict uncertainty in covariate effects and power to detect their relevance in Non-Linear Mixed Effect Models in pharmacometrics. *medRxiv*, 2024–10 (2024).
21. Rosenberger, W. F. & Sverdlov, O. Handling Covariates in the Design of Clinical Trials. *Statistical Science* **23**. <http://dx.doi.org/10.1214/08-STS269> (Aug. 2008).

22. Bogacka, B., Latif, M. A., Gilmour, S. G. & Youdim, K. Optimum designs for non-linear mixed effects models in the presence of covariates. *Biometrics* **73**, 927–937 (2017).
23. Golub, G. H. & Welsch, J. H. Calculation of Gauss quadrature rules. *Mathematics of Computation* **23**, 221–230 (1969).
24. Calamai, P. H. & Moré, J. J. Projected gradient methods for linearly constrained problems. *Mathematical Programming* **39**, 93–116 (1987).
25. R Core Team. *R: A Language and Environment for Statistical Computing* R Foundation for Statistical Computing (Vienna, Austria, 2021). <https://www.R-project.org/>.
26. Centers for Disease Control and Prevention (CDC). National Center for Health Statistics (NCHS). *National Health and Nutrition Examination Survey Data*. Hyattsville, MD: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention <https://wwwn.cdc.gov/nchs/nhanes/default.aspx>. 2009-2020.
27. Rosenblatt, M. Remarks on a multivariate transformation. *The Annals of Mathematical Statistics* **23**, 470–472 (1952).
28. James, F. Monte Carlo theory and practice. *Reports on Progress in Physics* **43**, 1145 (1980).
29. Fayette, L., Leroux, R., Mentré, F. & Seurat, J. Robust and adaptive two-stage designs in nonlinear mixed effect models. *The AAPS Journal* **25**, 71 (2023).
30. Retout, S., Comets, E., Samson, A. & Mentré, F. Design in nonlinear mixed effects models: optimization using the Fedorov–Wynn algorithm and power of the Wald test for binary covariates. *Statistics in Medicine* **26**, 5162–5179 (2007).
31. Giner, G. & Smyth, G. K. statmod: probability calculations for the inverse Gaussian distribution. *R Journal* **8**, 339–351 (2016).
32. Karline Soetaert, Karel Van den Meersche & Dick van Oevelen. *limSolve: Solving Linear Inverse Models* R package 1.5.1 (2009).
33. Robinson, J. D., Lupkiewicz, S. M., Palenik, L., Lopez, L. M. & Ariet, M. Determination of ideal body weight for drug dosage calculations. *American Journal of Hospital Pharmacy* **40**, 1016–1019 (1983).
34. Chin, P., Florkowski, C. & Begg, E. The performances of the Cockcroft-Gault, modification of diet in renal disease study and chronic kidney disease epidemiology collaboration equations in predicting gentamicin clearance. *Annals of Clinical Biochemistry* **50**, 546–557 (2013).
35. Nagler, T. & Vatter, T. rvinecopulib: High performance algorithms for vine copula modeling. *R package version 0.5 5*. <https://CRAN.R-project.org/package=rvinecopulib> (2021).
36. Guo, Y., Guo, T., Knibbe, C. A., Zwep, L. B. & van Hasselt, J. Generation of realistic virtual adult populations using a model-based copula approach. *Journal of Pharmacokinetics and Pharmacodynamics*, 1–12 (2024).
37. Guo, Y., Guo, T., van Hasselt, J. G. C. & Zwep, L. B. pmxcopula - R package for copula-based covariate simulation. *PAGE* (2024).
38. Malcolm, M. A. & Simpson, R. B. Local versus global strategies for adaptive quadrature. *ACM Transactions on Mathematical Software (TOMS)* **1**, 129–146 (1975).
39. Riviere, M.-K., Ueckert, S. & Mentré, F. An MCMC method for the evaluation of the Fisher information matrix for non-linear mixed effect models. *Biostatistics* **17**, 737–750 (2016).
40. Ueckert, S. & Mentré, F. A new method for evaluation of the Fisher information matrix for discrete mixed effect models using Monte Carlo sampling and adaptive Gaussian quadrature. *Computational Statistics & Data Analysis* **111**, 203–219 (2017).
41. Lestini, G., Dumont, C. & Mentré, F. Influence of the size of cohorts in adaptive design for nonlinear mixed effects models: an evaluation by simulation for a pharmacokinetic and pharmacodynamic model for a biomarker in oncology. *Pharmaceutical Research* **32**, 3159–3169 (2015).

## A Appendix

### A.1 Copula diagnostics

#### A.1.1 BMI only




Figure A.1.1: *BMI* only - Relative error (*RE*) of distribution metrics (mean, standard deviation and percentiles) of *BMI* for the three intervals, as compared to the statistics of the NHANES database. For each *BMI* interval, the virtual population was simulated 100 times from the copula.

The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. *sd* refers to the standard deviation, and *P5%*, *P50%* and *P95%* to the 5<sup>th</sup>, 50<sup>th</sup> and 95<sup>th</sup> percentiles respectively.




Figure A.1.2: *BMI* only - Visual predictive checks based on the percentile of the *BMI* distribution. The virtual population was simulated 100 times from the copula and the 99% prediction intervals of the percentiles the distribution was derived and compared to the percentiles observed in the NHANES database.

The ribbon areas correspond to the 99% prediction interval of 5<sup>th</sup> (●), 50<sup>th</sup> (●) and 95<sup>th</sup> (●) percentiles of the *BMI* distribution. The lines correspond to the observed 5<sup>th</sup> (—), 50<sup>th</sup> (—) and 95<sup>th</sup> (—) percentiles of the *CLCR* distribution.

#### A.1.2 CLCR only




Figure A.1.3: *CLCR* only - Relative error (*RE*) of distribution metrics (mean, standard deviation and percentiles) of *CLCR* for the four intervals, as compared to the statistics of the NHANES database. For each *CLCR* interval, the virtual population was simulated 100 times from the copula.

The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. *sd* refers to the standard deviation, and *P5%*, *P50%* and *P95%* to the 5<sup>th</sup>, 50<sup>th</sup> and 95<sup>th</sup> percentiles respectively.




Figure A.1.4: *CLCR* only - Visual predictive checks based on the percentile of the *CLCR* distribution. The virtual population was simulated 100 times from the copula and the 99% prediction intervals of the percentiles the distribution was derived and compared to the percentiles observed in the NHANES database.

The ribbon areas correspond to the 99% prediction interval of 5<sup>th</sup> (●), 50<sup>th</sup> (●) and 95<sup>th</sup> (●) percentiles of the *CLCR* distribution. The lines correspond to the observed 5<sup>th</sup> (—), 50<sup>th</sup> (—) and 95<sup>th</sup> (—) percentiles of the *CLCR* distribution.

### A.1.3 Three covariates




Figure A.1.5: Three Covariates - Relative error ( $RE$ ) of marginal metrics (mean, standard deviation and percentiles) of continuous covariates for the 20 covariate combinations, as compared to the statistics of the NHANES database. For each combination, the virtual population was simulated 100 times from the copula.

The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles.  $sd$  refers to the standard deviation, and  $P5\%$ ,  $P50\%$  and  $P95\%$  to the 5<sup>th</sup>, 50<sup>th</sup> and 95<sup>th</sup> percentiles respectively.




Figure A.1.6: Three Covariates - Overlap metric of 95<sup>th</sup> density contours of virtual population relative simulated 100 times from the copula to the NHANES database.

Overlap metric of 95<sup>th</sup> density contours of virtual population relative to observed population. Virtual population was simulated 100 times. The gray dashed line indicates 85% overlap percentage.




Figure A.1.7: Three Covariates - Correlations of  $\log BMI$  and  $\log CLCR$  pair in the NHANES database (red diamond) and in virtual population relative simulated 100 times from the copula (boxplot).

The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. sd refers to the standard deviation, and P5%, P50% and P95% to the 5<sup>th</sup>, 50<sup>th</sup> and 95<sup>th</sup> percentiles respectively.

## A.2 FIM integration: MC vs GQ

### A.2.1 BMI only






Figure A.2.1: *BMI* only -  $\phi_D$  computation with FIM integration using copula and MC (left) and copula and GLQ (right).

The green line corresponds to the average  $\phi_D$  obtained with  $n^{(MC)} = 1000$  and  $n^{(MC)} = 2000$ . The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. The diamond corresponds to the mean.

$\phi_D$ : refers to the D-criterion, FIM: Fisher Information Matrix MC: Monte-Carlo, GLQ: Gauss-Legendre Quadrature

## A.2.2 *CLCR* only




Figure A.2.2: *CLCR* only -  $\phi_D$  computation with FIM integration using copula and MC (left) and copula and GLQ (right).

The green line corresponds to the average  $\phi_D$  obtained with  $n^{(MC)} = 1000$  and  $n^{(MC)} = 2000$ . The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. The diamond corresponds to the mean.

$\phi_D$ : refers to the D-criterion, FIM: Fisher Information Matrix MC: Monte-Carlo, GLQ: Gauss-Legendre Quadrature




Figure A.2.3: *CLCR* only - Run time for  $\phi_D$  computation (top) and relative error ( $RE$ ) in  $\phi_D$  computation (bottom), with FIM integration using on the left copula and MC and on the right copula and GLQ. The target value was the average  $\phi_D$  obtained with  $n^{(MC)} = 1000$  and  $n^{(MC)} = 2000$ .

The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. The diamond corresponds to the mean.  $\phi_D$ : refers to the D-criterion, FIM: Fisher Information Matrix MC: Monte-Carlo, GLQ: Gauss-Legendre Quadrature

### A.2.3 Three Covariates




Figure A.2.4: Three covariates -  $\phi_D$  with FIM integration using copula and MC (left) and copula and GLQ (right).

The boxplot displays the median, the 25<sup>th</sup> and 75<sup>th</sup> percentiles, while the whiskers are 5<sup>th</sup> and 95<sup>th</sup> percentiles. The diamond corresponds to the mean.  $\phi_D$ : refers to the D-criterion, FIM: Fisher Information Matrix MC: Monte-Carlo, GLQ: Gauss-Legendre Quadrature

### A.3 Optimisation results: marginal distributions

Table 4: Three Covariates - Marginal optimal proportions for the covariate distributions for the D-criterion for the different sets of constraints

Name	Interval	Initial Design	Optimal designs		
			Without constraint	Lower constraints	Lower and Upper constraints
<i>SEX</i>	Male	0.488	0.337	0.335	0.337
	Female	0.512	0.663	0.665	0.663
<i>BMI</i>	Healthy Weight	0.279	0.463	0.441	0.483
	Overweight	0.327	0.117	0.154	0.087
	Obesity	0.393	0.420	0.406	0.430
<i>CLCR</i>	Normal RF	0.584	0.687	0.622	0.617
	Mild RF	0.283	0.000	0.050	0.050
	Moderate RF	0.120	0.000	0.050	0.233
	Severe RF	0.012	0.313	0.278	0.100