

1 Quantification of Optical Coherence
2 Tomography Angiography in Age and Age-
3 related Macular Degeneration using Vessel
4 Density analysis

5

6 **Ehsan Vaghefi, PhD^{1,2}, Sophie Hill, MD³, Hannah M Kersten, PhD¹, David**
7 **Squirrell, FRANZCO³**

8 1- School of Optometry and Vision Science, University of Auckland, New
9 Zealand

10 2- Auckland Bioengineering Institute, University of Auckland, New Zealand

11 3- Department of Ophthalmology, University of Auckland, New Zealand

12

13 * Corresponding Author

14 Dr Ehsan Vaghefi

15 85 Park Rd, Grafton

16 School of Optometry and Vision Science

17 University of Auckland, Auckland

18 New Zealand

19 E: e.vaghefi@auckland.ac.nz

20 P: +64 9 9231036

21 This work was funded by Faculty Research Development fund, The University of

22 Auckland.

23 None of the authors of this manuscript has a proprietary interest to declare.

24 **Keywords:** Age-related Macular Degeneration, Optical Coherence Tomography

25 Angiography, Retinal Blood Flow, Vessel Density

26

27 Word Count: 3768

28

29 **Brief Summary**

30 In this manuscript, we have studied the potential of retinal vessel density as

31 measured by optical coherence tomography angiography (OCT-A), as a biomarker

32 for detection of high-risk of developing exudative age-related macular degeneration

33 (AMD).

34

35 **Abstract**

36 **Purpose**

37 To determine whether vessel density (VD) as measured by optical coherence tomography
38 angiography (OCT-A) provide insights into retinal and choriocapillaris vascular changes with
39 ageing and intermediate dry age related macular degeneration (AMD).

40 **Methods**

41 Seventy-five participants were recruited into three cohorts; young healthy (YH) group, old healthy
42 (OH) and those at high-risk for exudative AMD. Raw OCT and OCT-A data from TOPCON DRI
43 OCT Triton were exported using Topcon IMAGENET 6.0 software, and 3D datasets were
44 analysed to determine retinal thickness and vessel density.

45 **Results**

46 Central macular thickness measurements revealed a trend of overall retinal thinning with
47 increasing age. VD through the full thickness of the retina was highest in ETDRS sector 4 (the
48 inferior macula) in all the cohorts. Mean VD was significantly higher in the deep capillary plexus
49 than the superficial capillary plexus in all ETDRS sectors in all cohorts but there was no
50 significant difference noted between groups. Choriocapillaris VD was significantly lower in all
51 ETDRS sectors in the in the AMD group compared with the YH and the OH groups.

52 **Conclusions**

53 Retinal vessel density maps, derived from the retinal plexi are not reliable biomarkers for
54 assessing the ageing macular. Our non-proprietary analysis of the vascular density of the
55 choriocapillaris revealed a significant drop off of VD with age and disease but further work is
56 required to corroborate this finding. If repeatable, choriocapillaris VD may provide a non-invasive
57 biomarker of healthy ageing and disease.

58 **Introduction**

59 Currently we lack a reliable, non-invasive vascular biomarker to monitor both the “healthy”
60 ageing and disease progression of age related macular degeneration (AMD). Optical
61 coherence tomography angiography (OCT-A) represents a novel, non-invasive, dye-less
62 retinal vascular imaging technique that can be rapidly acquired during a clinical consultation.
63 OCT-A is a progression of OCT, using repeated B-scans to detect motion contrast¹. Based
64 on the assumption that this represents the movement of erythrocytes travelling through the
65 retinal blood vessels, the retinal vasculature can therefore be visualised.² Previous studies
66 have shown the value of this imaging technique in the evaluation of glaucoma, diabetic
67 retinopathy (DR), retinal vein occlusions and AMD³.

68 The laminar structure of the retina lends itself to segmentation analysis. Histologically, the
69 retina has four retinal capillary plexuses, within the macular only three of these are
70 considered; the superficial, deep and intermediate capillary plexuses^{4, 5}. When the macular
71 vasculature is imaged using OCT-A the vascular layers are segmented automatically by the
72 software. This results in the merging of the intermediate with the deep capillary plexus³. The
73 regions of interest identified on the OCT-A therefore include the superficial capillary plexus
74 (SCP) and the deep capillary plexus (DCP)³. The SCP consists of the vasculature within the
75 retinal nerve fibre layer (RNFL) and ganglion cell layer (GCL). The DCP represents the
76 vascular plexuses present at two locations: These plexuses are at the border of the inner
77 nuclear layer and the outer plexiform layer, and the border of the inner plexiform layer and
78 inner nuclear layer⁶.

79 OCT-A images can be subjectively appraised for the presence of disease or post processed
80 to produce quantitative data that can be evaluated objectively. A number of studies^{7, 8} have
81 used perfusion indices such as vessel density and flow index⁹ as a method of quantitatively
82 analysing OCT-A images. Vessel density is defined as the “percentage area occupied by
83 vessels in the segmented area”⁹. Flow index is defined as “the average decorrelation values
84 in the segmented area”⁹. It has been proposed that these indices could be of value in

85 monitoring disease progression in AMD and DR^{7, 10, 11}. When measuring vessel density, both
86 vessel length and diameter need to be considered. Poor perfusion can also result in vessel
87 dilation, therefore vessel density may not give complete information about the retinal
88 vascular status¹².

89 There are a wide range of OCT-A devices available, each using a variety of algorithms to
90 capture retinal images. These include OCT-based optical micro-angiography (OMAG); split-
91 spectrum amplitude decorrelation angiography (SSADA); OCT angiography ratio analysis
92 (OCT-ARA); speckle variance; phase variance; and correlation mapping¹³. Specific OCT-A
93 models such as the AngioVue (Optovue, Inc., Fremont, CA, USA) have inbuilt software for
94 perfusion indices calculation¹⁴. However, the majority of studies have applied post image
95 analysis in a variety of methods after the images are exported from the OCT-A instrument².

96 This heterogeneity in the image capture and analysis means that data calculated using
97 different algorithms are not directly comparable, due to systemic difference and poor
98 agreement². Therefore, external normative databases are unreliable and longitudinal
99 monitoring of disease using vessel density requires scans to be performed using the same
100 instrument, scan location and algorithm².

101 The aims of this study are to compare the macular vessel density values for; young healthy,
102 old healthy, and patients with intermediate dry age related macular degeneration at high risk
103 of progression to exudative AMD, and to determine if these perfusion indices could usefully
104 serve as a biomarker for both “healthy” ageing and the identification of disease risk.

105

106

107

108

109

110 **Materials and Methods**

111 Seventy-five participants were recruited through Auckland Optometry Clinic and Milford Eye
112 Clinic, Auckland, New Zealand. All participants provided written informed consent prior to
113 imaging. Ethics approval (#018241) from the University of Auckland Human Participants Ethics
114 Committee was obtained for this study. This research adhered to the tenets of the Declaration of
115 Helsinki.

116 Participants were divided into three groups, young healthy (YH) group, old healthy (OH) group
117 and the high-risk intermediate AMD group. Twenty participants were recruited into the YH group,
118 twenty-one were recruited into the OH group, and thirty-four recruited into the AMD group. A
119 comprehensive eye exam was conducted on each participant prior to the OCT and OCT-A scans
120 in order to determine high contrast BCVA and the ocular health status of the fundus. Patients
121 with any posterior eye disease that could potentially affect the choroidal or retinal vasculature
122 including, but not limited to glaucoma, polypoidal choroidal vasculopathy, DR, hypertensive
123 retinopathy, and high myopia ($\geq 6D$), were excluded from the study. Patients with any history of
124 neurological disorders were also excluded. The Young Healthy group consisted entirely of
125 individuals between the ages of 20 and 26 and a best corrected visual acuity of $\geq 6/9$ in the eye
126 under test. The “Old Healthy” group consisted of individuals over the age of 55 years and who
127 had and a best corrected visual acuity of $\geq 6/9$ in the eye under test. The “AMD group” consisted
128 entirely of patients with high risk intermediate dry AMD. This being diagnosed if the individual
129 had at least two of the following risk factors; reticular pseudodrusen, established neovascular
130 AMD in the fellow eye, confluent soft drusen with accompanying changes within the retinal
131 pigment epithelium. In order to ensure that all patients in the “AMD cohort” could maintain
132 fixation during OCT-A imaging, only those patients with a best-corrected visual acuity (BCVA) of
133 6/15 or better were enrolled. The mean age of the participants in the YH group, OH and AMD
134 participants were 23 ± 3 , 65 ± 10 and 75 ± 8 years respectively. Only one eye of each patient
135 was used for the analysis, and if the patient had both eyes scanned, the OCT-A scan that was
136 the better quality (assessed subjectively by the clinical grader) of the two was used. Mean best-
137 corrected visual acuity for the YH, OH and AMD group were 6/6, 6/9, and 6/12 respectively.

138 The ocular health of all participants was assessed at Auckland Optometry Clinic, by a registered
139 optometrist, prior to enrolment in the study. The macular status of patients enrolled into the AMD
140 group was assessed separately by an experienced retinal specialist (DS).

141

142 **Data collection protocol**

143 Participants were dilated with 1.0% tropicamide if the pupils were deemed to be too small for
144 adequate OCT scans. Intraocular pressures were measured before and after dilation by iCare
145 tonometry. OCT/OCT-A scans, fundus photography and clinical measurements were all
146 performed at the University of Auckland Optometry Clinic, in a single session.

147

148 **SS-OCT-A device and scanning protocol**

149 The swept source OCT-A device (Topcon DRI OCT Triton, Topcon Corporation, Japan) was
150 used to obtain the OCT and OCT-A scans. A macular line and 3x3mm² OCT, and 3x3mm² OCT-
151 A scan were performed on each participant. All scans were centered on the fovea, and retinal
152 layers were identified using the IMAGENET 6.0 automated layer detection tool so that the
153 superficial (SCP) and deep (DCP) capillary plexuses could be evaluated. The quality of the
154 generated layers were checked manually. Scan quality was evaluated at the time of acquisition,
155 and repeated if required.

156

157 **Quantitative analysis of OCT-A Images**

158 Raw OCT and OCT-A data were exported using Topcon IMAGEnet 6.0 software. Since OCT-A
159 images are only qualitative, we chose to normalize the OCT-A datasets, prior to further analysis.
160 These datasets were then correlated with vessel density in each retinal sector. OCT-A 3D
161 datasets were analysed in three different ways (outlined below), to investigate the retinal and
162 choroidal thickness and vessel density (VD).

163

164 **1. ETDRS-specific**

165 The nine Early Treatment Diabetic Retinopathy Study (ETDRS) sectors were computationally
166 generated and centred manually by the clinical examiner, on the fovea (using enface OCT-A) for

167 each participant. In some instances, the macular region was not in the centre of the enface
168 $3 \times 3 \text{mm}^2$ OCT-A scan. Hence, after manual adjustment of the ETDRS central positioning on the
169 fovea, the outer ETDRS regions (i.e. 6-9) could have been positioned outside of the imaging
170 area. Therefore, in this study in which we were interested in assessing the role vessel density may
171 play as a potential biomarker for ageing and disease of the central macular, we only used
172 ETDRS regions 1 to 5 for consistency in our analysis. In this process, the OCT-A signal is
173 extracted and saved for each ETDRS region, superimposed on every OCT-A enface layer. In
174 other words, all the $3 \times 3 \text{mm}^2$ OCT-A layers through the retinal thickness are combined to create
175 an OCT-A volume dataset, 920 of them per $3 \times 3 \text{mm}^2$ scan, using the Topcon DRI OCT. Finally,
176 the ETDRS regions are manually applied to each of the 920 enface OCT-A images and OCT-A
177 normalized data per region are extracted (see Figure 1). This method has been recently
178 published¹⁵. The averaged normalized vessel density (VD) of the full thickness retina and
179 choriocapillaris of each ETDRS region was then extracted from every participant for further
180 analysis. Normalisation was performed by converting the raw data to percentage of VD in all
181 ETDRS regions, throughout the thickness of the retina.

182

183 **2. SCP, DCP and Choriocapillaris**

184 Using the same ETDRS regions as above, the VD was extracted from layer 1 and 2 for the SCP
185 calculation, layers 3 and 4 for the DCP and layer 8 for the choriocapillaris. Averaged normalized
186 VD from the SCP, DCP and choriocapillaris was then extracted for each participant, in the three
187 groups.

188

189 **3. Retinal and choroidal thickness**

190 In addition to OCT-A data, we also recorded the foveal thickness and subfoveal choroidal
191 thickness from the OCT B-scans.

192

193 **Statistical analysis**

194 MATLAB programming software and custom-written code was used to import and analyse the
195 data, as detailed above. Statistical analysis was performed using MATLAB Statistics Toolbox.

196 Due to unequal sample sizes between the sample groups, we performed the two sample non-
197 parametric Kolmogorov and Smirnov test, which is one of the most general nonparametric
198 methods for comparing two samples. The Kolmogorov and Smirnov method assumes that the
199 data in both groups follow Gaussian distributions. Significant differences between groups were
200 defined as $D < 0.565$ for the Kolmogorov and Smirnov test.
201

202 **Results**

203

204 **Mean age, central macular thickness, and foveal avascular zone diameter.**

205 Seventy-five participants were included in this study of which 20 were in the Young Healthy
206 group (23 ± 3 years old), 21 in the Old Healthy group, (65 ± 10 years old) and 34 in the AMD
207 group (75 ± 8 years old). OCT data was segmented by IMAGEnet 6.0 software, and the mean
208 central macular thickness measurements for the three participating cohorts was extracted: YH
209 217.4 ± 9.28 μm , OH 209.8 ± 6.13 μm , AMD 174.2 ± 20.3 μm . Horizontal diameter of the foveal
210 avascular zone (FAZ) was also measured from each OCT and mean FAZ measurements were
211 calculated: YH 672.5 ± 73.4 μm , OH 577.7 ± 98.8 μm , AMD 639 ± 126.2 μm .

212 **Vessel density measurements: Full retinal thickness.**

ETDRS Sectors	Foveal % (1)	Superior % (2)	Nasal % (3)	Inferior % (4)	Temporal % (5)
Young Healthy n=20	7.5 ± 6.6	18.1 ± 7.3	16.0 ± 7.0	20.5 ± 8.3	18.2 ± 8.1
Old	14.7 ± 9.8	21.1 ± 8.3	20.8 ± 9.9	21.5 ± 11	21.1 ± 9.6

213 *Table 1: Mean Vessel Density from full retinal thickness per ETDRS Sector, for each group.*

Healthy n=21					
AMD n=34	29.6±19.8	48.5±31.4	43.8±31.9	47.7±27.9	48.6±24.8

ETDRS Sectors	Foveal % (1)	Superior % (2)	Nasal % (3)	Inferior % (4)	Temporal % (5)
--------------------------------	---------------------	-----------------------	--------------------	-----------------------	-----------------------

214

215 Table 1 shows the mean VD throughout the **full retinal thickness**, for each ETDRS sector. In
 216 all groups the highest VD was found in sector 4, the inferior macula. Lowest vessel density in all
 217 groups was in sector 1 the foveal/parafoveal region. A statistical difference between the vessel
 218 density in Sector 1 was found between all the groups, with the vessel density being higher in the
 219 AMD cohort (29.6±19.8) compared to the Young Healthy (7.5±6.6) and Old healthy groups
 220 (14.7±9.68). Analysis of the vessel density of ETDRS sectors 2-5, revealed that there was a
 221 statistical difference in the VD between the AMD group and both the YH and OH groups; but not
 222 between the YH and OH groups.

223 **Vessel density measurements: SCP Segment.**

224 *Table 2: Mean Vessel Density in superficial capillary plexus (SCP) defined in sectors based on the*
 225 *ETDRS chart (Foveal, Superior, Nasal, Inferior, and Temporal).*

Young Healthy		27.0±12.9	62.3±23.1	60.0±20.1	64.8±21.5	59.9±21.2
ETDRS n=20 Sectors	Foveal % (1)		Superior % (2)	Nasal % (3)	Inferior % (4)	Temporal % (5)
Old						
Young Healthy		35.9±20.3	57.1±19.9	56.3±22.6	56.6 ± 22.1	54.1±24.8
Healthy n=21	19.8±3.3		25.0±4.7	24.5±4.2	27.8±4.8	25.5±4.8
AMD n=20		43.8±31.6	55.0±44	53.9±41.1	62.0±50.1	55.6±40.6
Old n=34						
Healthy n=21	21.3±3.9		25.9±3.5	25.7 ± 4.1	26.7.±4.9	24.6.±5.0

226

227 Table 2 shows the mean VD of the SCP segment within each ETDRS sector in each group.

228 Although, there was a trend towards the mean vessel density derived from the SCP in all ETDRS

229 sectors being higher in the AMD cohort compared to the 2 healthy cohorts, no significant

230 difference was found in the VD's recorded in any of the 5 ETDRS sectors between the 3 cohorts.

231 **Vessel density measurements: DCP Segment.**

232 *Table 3: Mean Vessel Density in deep capillary plexus (DCP) defined in sectors based on the*
 233 *ETDRS chart (Foveal, Superior, Nasal, Inferior, and Temporal).*

AMD	25.4 ±7.9	30.4±10.5	30.7±11.1	32.5±11.0	32.0±8.5
ETDRS	Foveal ratio	Superior ratio	Nasal ratio	Inferior ratio	Temporal ratio

234

235 Table 3 shows the mean VD of the DCP segment within each ETDRS sector for all groups.

236 The mean VD was found to be significantly higher (<0.01) in the DCP than the SCP in all ETDRS
237 sectors and in all groups. Although, there was a trend towards the mean vessel density derived
238 from the DCP in ETDRS sector 1 being higher in the AMD cohort compared to the 2 “healthy”
239 cohorts, the difference was not statistically significant. There was a trend for the mean vessel
240 density derived from the DCP in the remaining ETDRS sectors; 2-5, to be higher in the YH cohort
241 compared to the other 2 cohorts, this difference was not statistically significant.

242

243 **Vessel density measurements: Choriocapillaris Segment.**

244

245 *Table 4: Mean Vessel Density in the choriocapillaris, based on the ETDRS chart (Foveal, Superior,*
246 *Nasal, Inferior, and Temporal)*

S Sector s	(1)	(2)	(3)	(4)	(5)
Young Health y n=20	97 ± 10.5	86.2 ± 10.7	90.8 ± 11	103.8 ± 9.8	103 ± 11.3
Old Health y n=21	97 ± 4.5	85.5 ± 8.5	83.9 ± 9.5	86.3 ± 10.5	80.1 ± 12.8
AMD n=34	71.4 ± 24.3	68.1 ± 23.8	69.7 ± 24.4	68.4 ± 22.7	73.1 ± 19.4

247

248 Table 4 shows the mean VD of the Choriocapillaris segment within each ETDRS sector for all
 249 groups. In contrast to the measurements recorded in the retinal segments, statistically
 250 significantly lower mean VD's were recorded in the choriocapillaris segment layer in the AMD
 251 group across all ETDRS sectors compared to the two healthy groups. There was no significant
 252 difference in the VD's measured in each of the ETDRS sectors between YH and the OH groups.

253 **Discussion**

254 Although a number of changes within the ageing macular have previously been reported, a
255 reliable, non-invasive biomarker of both “healthy” macular ageing and disease risk remains
256 elusive. Retinal thickness, both at the fovea and extra-foveal regions, has previously been shown
257 to differ significantly between individuals with early AMD and age-matched controls with the
258 retina being thinner in individuals with disease¹⁶. The results of our study also revealed that there
259 was a trend for the overall retinal thickness to thin with age but this difference was not
260 statistically significant between cohorts. It is also recognised that the choroid tends to thin with
261 age; with the nasal area being thinnest, and the subfoveal region being thickest¹⁷. In addition to
262 thinning with “normal ageing,” Choroidal thinning has also been demonstrated in individuals with
263 early AMD compared to age matched controls^{16, 18-20} however, the significance of this
264 observation is controversial as both the extent and the pattern of thinning is significant in some²⁰
265 but not all studies^{16,18,19}. The histological evidence regarding choroidal thickness is also
266 contradictory with some studies showing reduction in late AMD²¹ and others showing no
267 change¹⁹; moreover, the changes in the vasculature of the choriocapillaris in early AMD that
268 have been reported from histological studies^{19,21} appear to be independent of choroidal
269 thickness.

270 Similarly inconclusive observations have been reported with respect to the width of the Foveal
271 avascular zone with age. Whilst some studies report no difference with age^{22, 23} others have
272 found that the FAZ enlarges with increasing age^{24, 25}. Where reported, the FA at the level of the
273 SCP has been found to be significantly smaller in >60 year olds compared to younger study
274 groups¹⁴, but the significance of this difference is unknown and furthermore no difference was
275 observed when the FAZ at the level of the DCP was analysed¹⁴. Recently, a study studying the
276 FAZ in patients with intermediate non-exudative AMD found no significant difference in the width
277 of the FAZ between patients with non-exudative AMD and healthy controls²⁶. We likewise found
278 there to be no statistically significant difference in the diameter of the FAZ between any of the 3
279 cohorts studied.

280 Where objective measures of macular anatomy have failed, OCTA may provide an opportunity to
281 develop a reliable, non-invasive vascular biomarker of the ageing macular. The proprietary

282 OCTA software now has a vessel density function which, in brief uses the proportion of bright
283 pixels to the proportion of dark pixels to derive a measure of vascular density. In the first instance
284 we reviewed the mean vessel density percentage throughout the **full** retinal thickness, for each
285 ETDRS sector. We then used the **segmentation** function of the enface OCTA to explore the
286 relationship of the vessel densities within each of the ETDRS sectors 1-5 in each of the 3
287 vascular zones of the central macular; (SCP, DCP, Choriocapillaris) across the 3 cohorts; YH,
288 OH and AMD.

289 As one might expect the VD's derived from the relatively avascular ETDRS sector 1 were, in the
290 **full** thickness maps, significantly reduced compared to the VD's derived from ETDRS sectors 2-5
291 in the YH, OH, and AMD cohorts. However, and curiously, the VD recorded in the AMD cohort
292 was significantly higher than in the two "Healthy" cohorts, and whilst there was a trend for the
293 VD's derived from the YH cohort to be reduced compared to the OH cohort, this difference was
294 not significant. Aside the marked difference between the AMD cohort and the other two "healthy"
295 cohorts, no other clinically relevant or statistically significant patterns were noted in the VD's
296 derived from the full thickness retinal maps.

297 Overall, and when the retinal vascular complexes were analysed as separate segments, the
298 highest vessel densities were found in sectors 2 and 4; the superior and inferior macula. Again
299 the VDs derived from ETDRS sector 1 was greater in the AMD cohort compared to the two
300 "Healthy" cohorts. Whilst there was a trend for the VD's derived from the OH cohort to be lower
301 compared to the YH cohort, this difference was not significant. The spatial variation in VD's that
302 we observed; a higher mean VD in the inferior and superior EDTRS sectors and a lower mean
303 VD in the temporal and nasal sectors, in both the deep and superficial capillary plexi, is
304 consistent with published data from previous studies.^{14, 22, 23} Like others^{14, 27}, we also observed
305 that in all ETDRS sectors and in all 3 study groups, the mean VD derived from the DCP was
306 significantly higher than that derived from the SCP.

307 The association between age and vessel density remains unclear, with reports of statistically
308 significant lower mean vessel density in SCP, DCP and full thickness of retina in each ETDRS
309 sector in patients >60 years old compared to younger cohorts;¹⁴, however, others who have
310 studied this have found no such correlation²². On the whole, our results do not further our

311 understanding of VD changes with age, being similarly inconclusive. On the basis of these data,
312 one has to conclude that without further work, VD's derived from the retinal segments are not a
313 reliable biomarker for "healthy" ageing.

314 One consistent feature of the VD's derived from the retinal segments in the present study was
315 the finding that the VD's recorded from ETDRS sector 1, and indeed in many other ETDRS
316 sectors in the AMD cohort, were as a group consistently higher than those obtained from the
317 "healthy" cohorts. One possible explanation for this surprising result is the technical challenges of
318 obtaining high quality OCT-A scans from individuals with even modest visual impairment as
319 accurate central fixation during the OCT-A capture can be challenging. Many of the patients in
320 our AMD cohort struggled to maintain fixation during the image acquisition stage, despite the
321 majority recording a Snellen acuity of 6/12. . Increased acquisition time and the resultant
322 compromise in image quality means there is a higher likelihood of projection and banding
323 artefacts.²⁸ Motion artefacts can also lead to the duplication of inner retinal vascular structures
324 ²⁹.As vessel density is derived from the percentage of the area occupied by bright pixels in a
325 segmented area, errors such as vessel duplication and banding artefacts could be incorrectly
326 interpreted during computational analysis as increased vessel density within the image. Such an
327 error would be magnified in ETDRS sector 1 where the effects of these banding and motion
328 artefacts would be more pronounced in what should be avascular tissue.

329 Although the proprietary OCTA software to assess VD is designed to measure the inner retinal
330 plexi, a vascular bed which has a very different architecture to the choriocapillaris³⁰ we were
331 curious to know whether the principle of VD analysis could be used to assess the vascularity of
332 the choriocapillaris. We therefore extracted the raw data, and then used a manual analysis
333 method previously published in the literature¹⁵, to assess the "Vascular density" of the
334 choriocapillaris segment. Even allowing for the influence of banding and motion artefacts,
335 issues which should increase VD measurements, we observed a significant reduction in the VD
336 in both the AMD and OH cohorts compared to the YH cohort with the reduction in VD being most
337 marked in the AMD cohort (Table 4). This contrasts sharply with the data derived from the retinal
338 plexi which broadly showed no difference between groups. Although interesting, one has to
339 interpret these findings with caution. It is possible that the observed reduction in VD is simply a

340 result of increased projection artefact and as VD has, to the best of our knowledge, not
341 previously been used to assess the choriocapillaris, it is not possible to corroborate these
342 findings. However, a global reduction in choriocapillaris density, associated with drusen on OCT-
343 A in dry AMD has been demonstrated previously.⁶ It is also recognised that the inner choroid and
344 the choriocapillaris disproportionately thin with age and disease³¹. It is feasible that a thinning
345 choriocapillaris would be less vascular and other groups have hypothesised that choroidal
346 perfusion changes on the OCT-A, may predict disease progression⁶. Debate about whether this
347 is related to actual flow impairment⁶ or projection artefact “masking” the choriocapillaris^{29, 32} due
348 to large drusen or pigment epithelial detachments attenuating the OCT-A signal is still ongoing.
349 Without further corroborative work, the importance of our finding that there is a progressive
350 reduction in choriocapillaris VD in age and disease is difficult to assess. Furthermore, because
351 our old healthy group and AMD cohorts are not aged matched, it is not possible to comment on
352 whether any observed reduction is simply an ageing phenomenon or a marker of disease.
353 Nevertheless, if reproducible, such a measure has the potential to provide a non-invasive
354 biomarker of both “healthy” macular ageing and disease risk. Further limitations of this study
355 include a small sample size, which might reduce the generalisability of our findings and the
356 interpretation of the statistical significance of our data. Furthermore, the Kolmogorov and
357 Smirnov statistical test assumes the Gaussian distribution of the tested data.

358

359 **Conclusion**

360

361 OCT-A mean retinal vessel density results for normal subjects are in line with the existing
362 literature. Challenges of image acquisition and the significant influence of image artefact on post-
363 acquisition analysis of patients with mild visual loss due to intermediate age related macular
364 degeneration further limits the value of VD measurements in this patient group. Traditional
365 measures of vessel density, derived from the retinal plexi are therefore unreliable biomarkers for
366 both “healthy” ageing and quantifying risk of developing AMD. Our non-proprietary analysis of
367 the vascular density of the choriocapillaris revealed a significant drop off of VD with age and

368 disease but further work is required to corroborate this finding. If repeatable choriocapillaris VD
369 may provide a non-invasive biomarker of healthy ageing and disease.

370

371 **Acknowledgments**

372 The authors wish to acknowledge the assistance by Mr Diego Sonderegger, from Device
373 Technologies®.

374

375 **References**


- 376 1. Spaide RF, Fujimoto JG, Waheed NK, Sadda SR, Staurengi G. Optical coherence
377 tomography angiography. *Prog Retin Eye Res* 2018;64:1-55.
- 378 2. Rabiolo A, Gelormini F, Sacconi R, et al. Comparison of methods to quantify macular
379 and peripapillary vessel density in optical coherence tomography angiography. *PLoS One*
380 2018;13:e0205773.
- 381 3. Spaide RF, Klancnik JM, Jr., Cooney MJ. Retinal vascular layers imaged by
382 fluorescein angiography and optical coherence tomography angiography. *JAMA Ophthalmol*
383 2015;133:45-50.
- 384 4. Tan PE, Yu PK, Balaratnasingam C, et al. Quantitative confocal imaging of the retinal
385 microvasculature in the human retina. *Invest Ophthalmol Vis Sci* 2012;53:5728-5736.
- 386 5. Gariano RF, Iruela-Arispe ML, Hendrickson AE. Vascular development in primate
387 retina: comparison of lamellar plexus formation in monkey and human. *Invest Ophthalmol Vis*
388 *Sci* 1994;35:3442-3455.
- 389 6. de Carlo TE, Romano A, Waheed NK, Duker JS. A review of optical coherence
390 tomography angiography (OCTA). *Int J Retina Vitreous* 2015;1:5.
- 391 7. Reif R, Qin J, An L, Zhi Z, Dziennis S, Wang R. Quantifying optical
392 microangiography images obtained from a spectral domain optical coherence tomography
393 system. *Int J Biomed Imaging* 2012;2012:509783.
- 394 8. Jia Y, Bailey ST, Hwang TS, et al. Quantitative optical coherence tomography
395 angiography of vascular abnormalities in the living human eye. *Proc Natl Acad Sci U S A*
396 2015;112:E2395-2402.
- 397 9. Chalam KV, Sambhav K. Optical Coherence Tomography Angiography in Retinal
398 Diseases. *J Ophthalmic Vis Res* 2016;11:84-92.
- 399 10. Pichi F, Nucci P, Baynes K, Lowder CY, Srivastava SK. Sustained-release
400 dexamethasone intravitreal implant in juvenile idiopathic arthritis-related uveitis. *Int*
401 *Ophthalmol* 2017;37:221-228.
- 402 11. Agemy SA, Sripsema NK, Shah CM, et al. Retinal Vascular Perfusion Density
403 Mapping Using Optical Coherence Tomography Angiography in Normals and Diabetic
404 Retinopathy Patients. *Retina* 2015;35:2353-2363.
- 405 12. Chu Z, Lin J, Gao C, et al. Quantitative assessment of the retinal microvasculature
406 using optical coherence tomography angiography. *J Biomed Opt* 2016;21:66008.
- 407 13. Zhang A ZQ, Chen CL, Wang RK. Methods and algorithms for optical coherence
408 tomography-based angiography: a review and comparison. *J Biomed Opt* 2015;20:100901.
- 409 14. Coscas F, Sellam A, Glacet-Bernard A, et al. Normative Data for Vascular Density in
410 Superficial and Deep Capillary Plexuses of Healthy Adults Assessed by Optical Coherence
411 Tomography Angiography. *Invest Ophthalmol Vis Sci* 2016;57:OCT211-223.
- 412 15. Hirano T, Chanwimol K, Weichsel J, Tepelus T, Sadda S. Distinct Retinal Capillary
413 Plexuses in Normal Eyes as Observed in Optical Coherence Tomography Angiography Axial
414 Profile Analysis. *Sci Rep* 2018;8:9380.
- 415 16. Wood A, Binns A, Margrain T, et al. Retinal and choroidal thickness in early age-
416 related macular degeneration. *Am J Ophthalmol* 2011;152:1030-1038 e1032.
- 417 17. MANJUNATH V TM, FUJIMOTO JG, DUKER JS. Choroidal Thickness in Normal
418 Eyes Measured Using Cirrus HD Optical Coherence Tomography. *Am J Ophthalmol*
419 2010;150:325-329.
- 420 18. Chung SE, Kang SW, Lee JH, Kim YT. Choroidal thickness in polypoidal choroidal
421 vasculopathy and exudative age-related macular degeneration. *Ophthalmology* 2011;118:840-
422 845.

- 423 19. Ramrattan RS, van der Schaft TL, Mooy CM, de Bruijn WC, Mulder PG, de Jong PT.
424 Morphometric analysis of Bruch's membrane, the choriocapillaris, and the choroid in aging.
425 *Invest Ophthalmol Vis Sci* 1994;35:2857-2864.
- 426 20. Sigler EJ, Randolph JC. Comparison of macular choroidal thickness among patients
427 older than age 65 with early atrophic age-related macular degeneration and normals. *Invest*
428 *Ophthalmol Vis Sci* 2013;54:6307-6313.
- 429 21. Sarks SH. Ageing and degeneration in the macular region: a clinico-pathological
430 study. *Br J Ophthalmol* 1976;60:324-341.
- 431 22. Rosen RK, B; Mo, S; Geyman, L; Phillips, E; Carroll, J; Weitz, R; Chui, TYP. Age-
432 Related Variations in Foveal Avascular Zone Geometry and Vessel Density - an Optical
433 Coherence Tomography Angiography (OCTA) study. *Invest Ophthalmol Vis Sci* September
434 2016;57:5501.
- 435 23. Gadde SG, Anegondi N, Bhanushali D, et al. Quantification of Vessel Density in
436 Retinal Optical Coherence Tomography Angiography Images Using Local Fractal
437 Dimension. *Invest Ophthalmol Vis Sci* 2016;57:246-252.
- 438 24. Iafe NA, Phasukkijwatana N, Chen X, Sarraf D. Retinal Capillary Density and Foveal
439 Avascular Zone Area Are Age-Dependent: Quantitative Analysis Using Optical Coherence
440 Tomography Angiography. *Invest Ophthalmol Vis Sci* 2016;57:5780-5787.
- 441 25. Falavarjani KG, Shenazandi H, Naseri D, et al. Foveal Avascular Zone and Vessel
442 Density in Healthy Subjects: An Optical Coherence Tomography Angiography Study. *J*
443 *Ophthalmic Vis Res* 2018;13:260-265.
- 444 26. Stavrev V, Sivkova N, Koleva-Georgieva D. Quantitative Assessment of Foveal
445 Avascular Zone in Patients with Early and Intermediate Nonexudative Age-Related Macular
446 Degeneration Using Optical Coherence Tomography-Angiography %J Open Journal of
447 Ophthalmology. 2018;Vol.08No.03:7.
- 448 27. Shahlaee A, Samara WA, Hsu J, et al. In Vivo Assessment of Macular Vascular
449 Density in Healthy Human Eyes Using Optical Coherence Tomography Angiography. *Am J*
450 *Ophthalmol* 2016;165:39-46.
- 451 28. Al-Sheikh M, Ghasemi Falavarjani K, Akil H, Sadda SR. Impact of image quality on
452 OCT angiography based quantitative measurements. *Int J Retina Vitreous* 2017;3:13.
- 453 29. Chen FK, Viljoen RD, Bukowska DM. Classification of image artefacts in optical
454 coherence tomography angiography of the choroid in macular diseases. *Clin Exp Ophthalmol*
455 2016;44:388-399.
- 456 30. Spaide RF. Choriocapillaris flow features follow a power law distribution:
457 implications for characterization and mechanisms of disease progression. *American journal*
458 *of ophthalmology* 2016;170:58-67.
- 459 31. Esmaeelpour M, Kajib V, Zabihian B, et al. Choroidal Haller's and Sattler's layer
460 thickness measurement using 3-dimensional 1060-nm optical coherence tomography. *PLoS*
461 *one* 2014;9:e99690.
- 462 32. Spaide RF, Fujimoto JG, Waheed NK. Image Artifacts in Optical Coherence
463 Tomography Angiography. *Retina* 2015;35:2163-2180.
- 464

465

466 **Figure Legend**

467 **Figure 1: Retinal layers were automatically identified by IMAGENET 6 software (A) the SCP and**
468 **DCP were then isolated from layers 1 and 2, and 3 and 4 respectively. ETDRS regions 1-9 are**
469 **superimposed onto the enface OCT-A images and manually adjusted so that sector 1 was**
470 **centred on the fovea. In this study we selected sectors 1-5 (B).**

A

NFL: Nerve Fiber Layer
GCL: Ganglion Cell Layer
IPL: Inner Plexiform Layer
INL: Inner Nuclear Layer
OPL: Outer Plexiform Layer

ONL: Outer Nuclear Layer
IS: Inner Segment
OS: Outer Segment
RPE: Retinal Pigment Epithelium

B