

Time to death and its Predictors among Adult TB/HIV Co-Infected Patients in Mizan Tepi University Teaching Hospital, South West Ethiopia

Wondimagegn Wondimu^{1*}: wonde1983.ww@gmail.com, Lamessa Dube²:
dubelamessa@yahoo.com, Teshome Kabeta²: tka1204@gmail.com

¹ Mizan Tepi University, College of Health Science, Department of Public Health, Mizan Aman,
Ethiopia

² Jimma University, Faculty of Public Health, Department of Epidemiology, Jimma, Ethiopia

* Corresponding author

Abstract

Background: Tuberculosis (TB) and Human Immuno Deficiency Virus (HIV) co-infection represents a complex pathogenic scenario with synergistic effect and leads to about 300,000 HIV-associated TB deaths in the world in 2017. Despite this burden of death, time to death and its predictors among TB-HIV co-infected patient was not adequately studied; and the existing evidences are inconsistent. Therefore, this study was aimed to determine time to death and identify its predictors among adult TB/HIV co-infected patients.

Method: Retrospective cohort study was conducted by reviewing registers of randomly selected 364 TB/HIV co-infected patients enrolled in health care from July 2, 2007 up to July 1, 2017 at Mizan Tepi University Teaching Hospital. The hospital was located in Bench Maji Zone, South West Ethiopia. Data were collected from March 1 through 31, 2018, entered to Epi data 3.1 and exported to SPSS version 21. Each patient was followed from date of TB treatment initiation till death, loss to follow up and treatment completed. On the other hand, events other than death were considered as censored. After checking the proportional hazard model assumption, Cox-regression was used to identify the predictors. In bivariable analyses, $P\text{-value} \leq 0.25$ was used to identify candidate variables for multivariable analysis. The 95% CI of hazard ratio (HR) with respective $P\text{-value} < 0.05$ was used to declare significance in the final model.

Result: All the 364 patients were followed for 1,654 person months. There were 83 (22.8%) deaths and most 38 (45.8%) were occurring within the first two months of anti-TB treatment initiation. The overall incidence rate and median survival time were 5.02 per 100 person months (95% CI: 4.05, 6.22) and 10 months respectively. Statistically significant better survival was observed among patients: with $CD4 \geq 200$ cells/mm³ ($P < 0.001$), who had a history of cotrimoxazole preventive therapy (CPT) use ($P < 0.001$), who disclose their HIV status ($P < 0.001$) and with working functional status ($P < 0.001$). Not using CPT (adjusted hazard ratio [AHR] = 1.72; $P = 0.023$), bedridden functional status (AHR = 2.55; $P = 0.007$), not disclosing HIV status (AHR = 4.03; $P < 0.001$) and $CD4 < 200$ cells/mm³ (AHR = 6.05; $P < 0.001$) were predictors of time to death among TB/HIV co-infected patients.

Conclusion: The median survival time was 10 months and poor survival was associated with low CD4 count, not using CPT, not disclosing HIV status and having bedridden functional status. Close monitoring of bedridden and low CD4 count patients, prompt CPT initiation and encouraging HIV status disclosure are recommended.

Key words: TB/HIV co-infection, South West Ethiopia, time to death, MTUTH

Introduction

TB/HIV co-infection has dangerous synergy which leads to highly complicated features of both diseases, including epidemiologic profile, pathogenesis, clinical presentation, treatment, and prevention methods (1). It has also social, economic and political crisis as consequences. The most common consequences are stigma, high susceptibility to other comorbidities like mental disorders, cost of care, increased number of orphans and decreased productivity due to long time illness and absence from work place (2,3). In most of the situations, this co-infection affects the productive segment of the population and increases the crisis by many folds, especially for low income countries like Ethiopia (4).

For people living with HIV/AIDS (PLWHA), the risk of developing TB is estimated to be between 17-23 times greater than those without HIV infection in 2017. Globally, the proportion of HIV-positive patients who died during TB treatment was 11%, in the same year (5,6).

TB was the leading cause of death among people with HIV and there were about 300,000 people who died from HIV-associated TB in 2017 in which the majorities of deaths (84%) occurred in Africa (6).

Ethiopia is among the 30 high TB/HIV burden countries and among TB patients with known HIV status, which is about 11%. It was estimated that there were about 3600 deaths due to TB-HIV co-infection in 2017 (5,7). On the top of knowledge gap, the problems of lay beliefs, misconceptions and prejudices regarding TB/HIV co-infection are other devastating issues in Ethiopia; since this all might contribute to delay from care and in turn for poor survival among co-infected patients (8–10).

Currently, there is no published study in the study area on time to death and its predictors among TB/HIV co-infected patients; although it is immediate near to Gambella region, which is among the regions with higher TB/HIV co-infection in Ethiopia and from where also a significant number of TB/HIV patients come to Mizan-Tepi University Teaching Hospital (MTUTH).

In addition, studies conducted in North West and South West part of Ethiopia identified inconsistent predictors (11,12). This study was aimed to determine time to death and its predictors among TB-HIV co-infected adult patients in MTUTH.

Methods

Study Design and Setting

A retrospective cohort study was conducted among TB/HIV co-infected patients enrolled from July 2, 2007 to July 1, 2017 at Mizan Tepi University Teaching Hospital. Each patient was followed from date of TB treatment initiation till death, loss to follow up and treatment completed. On the other hand, events other than death were considered as censored.

The hospital is in Mizan-Aman town, which is 593 km to South West of Addis Ababa, capital of Ethiopia. The hospital started offering free ART service in 2006. About 5946 PLWHA were enrolled till January 2019 and among these 1830 were active. Assessment for TB was being done routinely for every HIV patient. Similarly, if their test results become positive, all TB patients were supported to know their HIV status and linked to ART clinic.

Study Population and Sampling Techniques

All adult TB/HIV co-infected patients, who received care from MTUTH from July 2, 2007 to July 1, 2017 were included. Incomplete registries, transfer-out and death in the same day of anti-TB treatment initiation were excluded.

The sample size was determined using the online calculator for test time-to-event data for the Cox proportional hazard (13). The CD4 count ≤ 50 is known predictor of mortality and used to classify the groups into two (i.e. subjects with CD4 count ≤ 50 cells/mm³ and subjects with CD4 count > 50 cells/mm³) (4,14). Using the parameters (power=80%, type one error rate=5%, hazard ratio=2.31, overall probability of event=13% and proportion of sample in group of subjects with CD4 count ≤ 50 cells/mm³=38.7%) from previous study (15), the sample size became 364. Final samples were selected by simple random sampling technique after listing medical registration number of all 670 eligible records as a sampling frame.

Data Collection Technique and Data Quality Control

Data were collected using a data extraction tool adapted from national ART and anti TB treatment standard registries. Data sources were standard registries of ART and anti-TB treatments, electronic format, patient medical record (card) and intake forms. Two trained nurses each from the ART and TB clinics of the hospital collected the data. Trained supervisor supervises all the

collection process. Both the supervisor and principal investigator daily checked the completeness and consistency of checklists during data collection. Socio demographic, clinical, behavioural and health service-related base line data were collected from the sources.

Data entry, processing and analysis

Data were entered to Epi data 3.1 and exported to SPSS version 21 for analysis. Exploration of the data was done to check the missing values, outliers, and proportionality of hazards over time. Continuous variables were described using means and medians with respective 95% CI; while categorical variables were described using counts, percentages and tables. The median survival time, survival probability and the incidence rate were determined. The association between the independent variables and outcome variable was assessed by Cox-proportional hazard model and hazard ratio (HR) was used as a measure of association. Survival status of TB/HIV co-infected patients was presented by life table.

All variables with p-value ≤ 0.25 were candidates for the final model. By using backward stepwise likely hood ratio technique and p-value < 0.05 the final significant predictors were identified.

Ethical consideration

The study proposal was approved by an institutional review board (IRB) of Jimma University. Support letter was obtained from IRB of Jimma University to MTUTH administrator's office. Informed consent was obtained from the concerning administrative body of the hospital. Confidentiality was assured strictly. In addition, patient identifiers (medical registration number and ART unique number) were replaced by a new identification number during data collection and the name of the patients was not used.

Results

Characteristics of cohort

Among eligible 670 patients, 364 were included to the study. The mean age (\pm SD) of the participants was 30.52(\pm 9.03). From the total participants, 211(58%) were females, 150(41.2%) were married and 133(36.5%) had attended secondary school. **(Table 1)**

Table 1- Base line socio demographic characteristics of study subjects in MTUTH, South West Ethiopia, 2007-2017

Variable	Variable category	Frequency	Percent
Age	15-24	91	25
	25-34	160	44
	35-44	81	22.2
	\geq 45	32	8.8
Sex	Male	153	42
	Female	211	58
Educational status	No education	70	19.2
	Primary (grade 1-8)	80	22
	Secondary (grade 9-12)	133	36.5
	Tertiary (college/university)	81	22.3
Marital status	Never married	140	38.4
	Married	150	41.2
	Divorced	37	10.2
	Widowed	37	10.2
Occupation	Gov't employed	48	13.2
	Private organization employed	31	8.5
	Jobless	44	12.1
	Student	46	12.6
	Farmer	35	9.6
	Commercial sex worker	42	11.5

	Merchant	83	22.8
	Housewife	35	9.6
Place of residence	Within catchment area	261	71.7
	Out of catchment area	103	28.3
Having HIV-positive family member (n=362) *	No	100	27.6
	Yes	262	72.4

* Two participants had no records on either they had HIV positive family member or not.

Participants with baseline CD4 count at least 200 cells/mm were 240 (65.9%). Significant portion of study participants had one or two opportunistic infections other than TB, 232(63.7%). Cotrimoxazole was used as preventive therapy among 283 (77.7%) of the participants. (**Table 2**)

Table 2- Clinical characteristics of study participants in MTUTH, South West Ethiopia, 2007-2017

Variables	Category of variables	Frequency	Percent
CD4 count	<200	124	34.1
	≥200	240	65.9
ART regimen	First line	343	94.2
	Second line	21	5.8
Use of CPT	Yes	283	77.7
	No	81	22.3
OIs other than TB	No at all	105	28.8
	One or two	232	63.7
	At least three	27	7.4
ART failure history	Yes	21	5.8
	No	343	94.2
Functional status	Working	139	38.2
	Ambulatory	149	40.9
	Bedridden	76	20.9
WHO clinical stages	One	60	16.5

Two	51	14
Three	220	60.4
Four	33	9.1

ART-anti retroviral therapy, CPT-cotrimoxazole preventive therapy, CD4- Cluster of differentiation four, TB-tuberculosis, OIs-opportunistic infections, BMI-body mass index, WHO-world health organization

Survival status and its predictors

Three hundred sixty-four participants were followed for total of 1,654 person months. There were 83 (22.8%) deaths and 41(11.3%) loss to follow up. From the total deaths, 38 (45.8%) occurred within the first two months of anti-TB treatment. The overall incidence rate of death was 5.02 per 100 person months (95% CI: 4.05, 6.22). The median survival time was at around the 10th month. Survival at 2, 6 and 12 months after initiating TB treatment were 82.02%, 73% and 45% respectively. (Table 3)

Table 3- Overall life table of TB/HIV co-infected patients in MTUTH, South West Ethiopia, 2007-2017

Time in months	Number at risk	Number of events	Proportion of events	Proportion Surviving	Cumulative Proportion Surviving
0	354.5	38	0.11	0.89	0.89
2	295	24	0.08	0.92	0.82
4	177	11	0.06	0.94	0.77
6	52.5	3	0.06	0.94	0.73
8	16.5	5	0.3	0.70	0.51
10	9.5	1	0.11	0.89	0.45
12	8	0	0	1	0.45
14	8	0	0	1	0.45
16	6	1	0.17	0.83	0.38
18	2	0	0	1	0.38
20	0.5	0	0	1	0.38

The significant predictors of time to death for TB/HIV co-infected patients were; not using CPT (AHR=1.72; 95% CI: 1.08, 2.74), bedridden functional status (AHR=2.55; 95% CI: 1.29, 5.06), not disclosing HIV status (AHR=4.03; 95% CI: 2.21, 7.35) and CD4 count of less than 200 cells/mm³. (**Table 4**)

Table 4- Multivariable cox- regression analysis for predictors of time to death among TB/HIV co-infected patients in MTUTH, South West Ethiopia, 2007-2017

Variables	Category of Variables	AHR	95% CI		P value
Sex	Male*	1			
	Female	1.27	0.63	2.55	0.508
Educational status	No education*	1			
	Primary	1.44	0.63	3.25	0.386
	Secondary	1.05	0.45	2.45	0.913
	Tertiary	1.32	0.37	4.71	0.669
Marital status	Never married	0.85	0.45	1.61	0.623
	Married	0.87	0.40	1.91	0.732
	Divorced	2.16	0.93	5.04	0.073
	Widowed*	1			
Occupation	Gov't employed*	1			
	Private organization employed	0.52	0.12	2.27	0.387
	Jobless	0.34	0.10	1.36	0.137
	Student	0.43	0.12	1.51	0.188
	Farmer	0.68	0.19	2.40	0.551
	Commercial sex worker	1.10	0.39	3.09	0.861
	Merchant	1.08	0.38	3.08	0.884
	Housewife	0.92	0.25	3.38	0.906

CPT use	Yes*	1			
	No	1.72	1.08	2.74	0.023**
WHO clinical stages	One*	1			
	Two	1.05	0.31	3.59	0.935
	Three	1.23	0.59	2.69	0.555
	Four	2.02	0.75	5.43	0.162
Functional status	Working*	1			
	Ambulatory	0.93	0.41	2.11	0.861
	Bedridden	2.55	1.29	5.06	0.007**
TB registration group	New*	1			
	Previously treated	0.88	0.52	1.49	0.634
Disclosure of status	Yes*	1			
	No	4.03	2.21	7.35	0.000**
BMI	<18.5	0.84	0.24	2.94	0.787
	18.5-24.9	0.52	0.16	1.66	0.267
	25-29.9	0.72	0.19	2.73	0.632
	>29.9*	1			
Number of OIs other than TB	No at all*	1			
	One or two	1.37	0.62	3.02	0.439
	At least three	1.06	0.40	2.83	0.902
CD4 count in cells/mm³	<200	6.05	2.83	12.95	0.000**
	≥200*	1			

* Reference category, ** statistically significant at $p < 0.05$, CI-confidence interval, AHR-adjusted hazard ratio, CPT-cotrimoxazole preventive therapy, CD4- Cluster of differentiation four, TB-tuberculosis, OIs-opportunistic infections, BMI-body mass index, WHO-world health organization

Discussion

This study revealed that there were 22.8% deaths among TB/HIV co-infected patients during TB treatment and this finding is comparable to the studies done in Jimma and Bahirdar towns where there were 20.2% and 22% deaths respectively. The overall incidence rate was 5.02 per 100 person months, which is comparable to a study done in Bahirdar in which there was 4.09 per 100 person months incidence rate during TB treatment (12,16). The comparability of death rates in these all studies is suggestive for Ethiopia's rank of being among TB/HIV high burden countries in the world and the need of future high attention and priority particularly for the timely achievement of ending both epidemics (7,17).

The majority of the deaths, 38 (45.8%) occurred within the two months of anti-TB treatment initiation and this is fairly comparable with a study conducted in Malaysia where 40% of deaths occurred in the same period of TB treatment (18). This is suggestive of the need of appropriate management of drug interactions and overlapping toxicities of anti TB and ART drugs, immune reconstitution inflammatory syndrome (IRIS) and other OIs which were factors highly associated with early deaths in TB/HIV co-infected patients (19–21).

The survival at 2, 6 and 12 months after initiating TB treatment were 82.02%, 73% and 45% respectively. This is lower than the Malaysian finding where there were survival of 90.7%, 82.8% and 78.8% respectively at 2, 6 and 12 months after initiating TB treatment (18). This might be due to the pioneering efforts in Malaysia to combat TB/HIV co-infection; which is evidenced by its early achievement of the first and third 90s of UNAIDS' 90-90-90 target (22).

Patients who didn't use CPT were nearly two times (AHR=1.72) at higher risk of death and in agreement with this, study conducted in Bahirdar revealed that not taking CPT was significantly associated with mortality among TB/HIV co-infected patients during TB treatment (12). Severe bacterial infections and malaria, which can be successfully prevented by CPT might be attributable to higher risk of death among CPT non users (14,23,24).

Bedridden patients were two and a half times (AHR=2.55) risky to die than those who had working functional status. This is similar to findings of studies conducted in Jimma and Ambo (11,25,26). This might be due to the delayed presentation of bedridden patients to care after developing

numerous infections and complications which were in turn associated with poor survival (10,18).

The bidirectional and synergistic effect of TB and HIV infection has profound impact on the survival of TB/HIV co-infected patients. Latent or active TB activates CD4 cells to be captured by HIV and HIV depletes CD4 cells, which in turn accelerates the progression of latent TB to active TB disease (1,27). CD4 count of less than two hundred cells per cubic millimeter was significantly associated with poor survival of TB/HIV co-infected patients in this study (AHR=6.05). This is supported by previous findings from different parts of the world including Ethiopia. Although the current estimation is consistent with the findings of Bahirdar and Malaysia, it is higher compared to that of a study conducted in Brazil (12,15,18). This might be due to the categorization CD4 count into small intervals in a previous study.

Failure to disclose status puts TB/HIV co-infected patients and their families at higher risk of death. It prevents other at risk family members from being tested and patients who do not disclose their status are highly likely to be lost to follow up, as they cannot explain their ongoing need to visit health institutions (28). In the current study, patients who didn't disclose their HIV status to anybody were about four times (AHR= 4.03) at higher risk of death than those who disclosed their status. This is in line with the finding of the study done in Jinka, Ethiopia (29). This might be due to patients who didn't disclose their status would not be comfortable to take their medications appropriately, couldn't get important family supports and wouldn't visit health institutions as expected; which might contribute to the unwanted outcome like ART failure and poor survival (30). One of the main reasons for non-disclosure of the status is fear of stigmatization; and this needs a sustainable awareness creation in the community to bring positive attitudes toward TB/HIV co-infected patients and alleviate lay beliefs and misconceptions (8,9).

Limitation of the study

Selection bias might be introduced; during the exclusion of incomplete registers and subjects with lost cards.

Conclusions

This study demonstrated short survival time, among TB/HIV co-infected patients during TB treatment, compared to previous findings. A high proportion of deaths occurred within the first two months of anti-TB initiation. In addition, it identified nonuse of CPT, bedridden functional status, nondisclosure of HIV status and CD4 count less than 200 cells/mm³ as predictors of time to death among TB/HIV co-infected patients. Close monitoring of bedridden and low CD4 count patients, prompt CPT initiation and encouraging HIV status disclosure are recommended.

List of abbreviations

AHR	Adjusted Hazard Ratio
AIDS	Acquired Immuno Deficiency Syndrome
AOR	Adjusted Odds Ratio
ART	Anti-Retro Viral Therapy
BMI	Body Mass Index
CD4	Cluster of Differentiation 4
CI	Confidence Interval
CPT	Cotrimoxazole Preventive Therapy
HIV	Human Immuno Deficiency Virus
HR	Hazard Ratio
IRB	Institutional Review Board
OIs	Opportunistic Infections

PLWHA	Peoples Living With HIV/IDS
TB	Tuberculosis
UNAIDS	The joint United Nations Programme on HIV/AIDS
WHO	World Health Organization

Acknowledgements

We would like to acknowledge the administrative bodies of Mizan Tepi University teaching hospital who allowed us to do this study on their institution and staffs of ART and TB clinics who participated in the collection of data and supervision of its overall processes.

Author Contributions

Conceptualization: Wondimagegn Wondimu

Formal analysis: Wondimagegn Wondimu, Teshome Kabeta

Methodology: Wondimagegn Wondimu, Teshome Kabeta, Lamessa Dube

Validation: Wondimagegn Wondimu, Teshome Kabeta, Lamessa Dube

Writing original draft: Wondimagegn Wondimu

Writing review & editing: Wondimagegn Wondimu, Teshome Kabeta, Lamessa Dube

Competing interests

The authors have declared that no competing interests exist.

References

1. Bruchfeld J, Correia-neves M, Ka'llenius G. Tuberculosis and HIV Coinfection. *Cold Spring Harb Perspect Med.* 2015;5(a017871).
2. Deribew A, Tesfaye M, Hailmichael Y, Apers L, Abebe G, Duchateau L, et al. Common mental disorders in TB / HIV co-infected patients in Ethiopia. *BMC Infect Dis.* 2010;10(21):1–8.
3. Deribew A, Hailemichael Y, Tesfaye M, Desalegn D, Wogi A, Daba S. The synergy between TB and HIV co-infection on perceived stigma in Ethiopia. *BMC research notes.* 2010;3(249):2–5.
4. UNAIDS. Global AIDS Monitoring [Internet]. 2016. Available from: <http://www.aidsdatahub.org/global-aids-monitoring-2017-indicators-monitoring-2016-united-nations-political-declaration-hiv-an-0>
5. WHO. Global tuberculosis report [Internet]. France; 2018. Available from: https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=2&ved=2ahUKEwiJy_Dy9efgAhUM6KQKHaurBiYQFjABegQIAhAC&url=http%3A%2F%2Fwww.unaids.org%2Fsites%2Fdefault%2Ffiles%2Fmedia_asset%2Funaid-data-2018_en.pdf&usg=AOvVaw1AkNS54cNJvDxrLgpO5qOe
6. WHO. HIV-Associated Tuberculosis, Achievements in 2017 [Internet]. 2018. Available from: https://www.who.int/tb/areas-of-work/tb-hiv/tbhiv_factsheet.pdf?ua=1
7. EPHI, WHO F. Ethiopia: Service Availability and Readiness Assessment 2016: Summary Report [Internet]. Addis Ababa; 2016. Available from: <https://www.washinhc.org/documents/Final-SARA-Report-Jan-2017.pdf>
8. Gebremariam MK, Bjune GA, Frich JC. Lay beliefs of TB and TB / HIV co-infection in Addis Ababa , Ethiopia : a qualitative study. *BMC Res Notes.* 2011;4(277):2–6.
9. Deribew A, Abebe G, Apers L, Jira C, Tesfaye M, Shifa J, et al. Prejudice and misconceptions about tuberculosis and HIV in rural and urban communities in Ethiopia : a challenge for the TB / HIV control program. *BMC Public Health.* 2010;10(400):1–10.
10. Gesesew H, Tsehaineh B, Massa D, Tesfay A, Kahsay H, Mwanri L. The prevalence and associated factors for delayed presentation for HIV care among tuberculosis / HIV co-infected patients in Southwest Ethiopia : a retrospective observational cohort. *Infect Dis Poverty* [Internet]. 2016;5(96):1–10. Available from: <http://dx.doi.org/10.1186/s40249-016-0193-y>
11. Gesesew H, Tsehayneh B, Massa D, Gebremedhin A, Kahsay H, Mwanri L. Predictors of

mortality in a cohort of tuberculosis / HIV co-infected patients in Southwest Ethiopia. *Infect Dis Poverty*. 2016;5(109):1–9.

12. Sileshi B, Deyessa N, Girma B, Melese M, Suarez P. Predictors of mortality among TB-HIV Co-infected patients being treated for tuberculosis in Northwest Ethiopia : a retrospective cohort study. *BMC Infect Dis*. 2013;13(297):1–10.

13. Chow S, Shao J WH. *Sample Size Calculations in Clinical Research* [Internet]. Second. 2008. 277 p. Available from: <http://powerandsamplesize.com/Calculators/Test-Time-To-Event-Data/Cox-PH-2-Sided-Equality>

14. WHO. Consolidated guidelines on the use of antiretroviral drugs for treating and preventing HIV infection: Recommended for a public health approach [Internet]. Second. France; 2016. Available from: <http://www.who.int/hiv/pub/arv/arv-2016/en/>

15. Otavio R, Velasque L, Ribeiro SR, Cardoso SW, Monteiro L, Marins S, et al. Mortality in patients with HIV-1 and tuberculosis co-infection in Rio de Janeiro , Brazil - associated factors and causes of death. *BMC Infect Dis*. 2017;17(373):1–10.

16. Abrha H, Tsehayneh B, Massa D, Tesfay A, Kahsay H. Survival Experience and its Predictors among TB / HIV Co-infected Patients in Southwest Ethiopia. *Epidemiol*. 2015;5(3):3–9.

17. WHO. Global Tuberculosis report [Internet]. France; 2017. Available from: http://www.who.int/tb/publications/global_report/en/

18. Ismail I, Bulgiba A. Predictors of Death during Tuberculosis Treatment in TB / HIV Co-Infected Patients in Malaysia. *PLoS One*. 2013;8(8):1–7.

19. Aluoch JA. Tuberculosis / HIV Co-infection. *East Afr Med J*. 2013;89(12):68–74.

20. Swaminathan S, Padmapriyadarsini C, Narendran G. HIV-Associated Tuberculosis : Clinical Update. *Clin Infect Dis*. 2010;50(10):1377–1386.

21. Worodria W, Massinga-loembe M, Mazakpwe D, Luzinda K, Menten J, Leth F Van, et al. Incidence and Predictors of Mortality and the Effect of Tuberculosis Immune Reconstitution Inflammatory Syndrome in a Cohort of TB / HIV Patients Commencing Antiretroviral Therapy. *J Acquir Immune Defic Syndr*. 2011;58(1):32–7.

22. UNAIDS. Global AIDS update, Ending AIDS, Progress towards the 90-90-90 targets [Internet]. 2017. Available from: http://www.unaids.org/en/resources/documents/2017/20170720_Global_AIDS_update_2017

23. Nunn AJ, Mwaba P, Chintu C, Mwinga A, Darbyshire JH, Zumla A. Role of co-trimoxazole prophylaxis in reducing mortality randomised clinical trial. *BMJ*. 2008;337(a257):2–8.
24. Hasse B, Walker AS, Fehr J, Furrer H, Hoffmann M, Battegay M, et al. Co-Trimoxazole Prophylaxis Is Associated with Reduced Risk of Incident Tuberculosis in Participants in the Swiss HIV Cohort Study. *Antimicrob Agents Chemother*. 2014;58(4):2363–8.
25. Hailu Refera EW. Survival of HIV-TB co-infected adult patients under ART in Ambo Referral Hospital, Ethiopia. *Ethiop J Heal Dev*. 2013;27(2).
26. Gesesew H, Tsehaineh B, Massa D, Tesfay A, Kahsay H, Mwanri L. The role of social determinants on tuberculosis / HIV co - infection mortality in southwest Ethiopia : a retrospective cohort study. *BMC Res Notes*. 2016;9(89):4–11.
27. Sullivan ZA, Wong EB, Ndung T, Kasprowicz VO, Bishai WR. Latent and Active Tuberculosis Infection Increase Immune Activation in Individuals Co-Infected with HIV. *EBioMedicine*. 2015;2(4):334–40.
28. PEPFAR Ethiopia. Ethiopia Country / Regional Operational Plan (COP / ROP) 2017 Strategic Direction Summary [Internet]. Addis Ababa; 2017. Available from: <https://www.pepfar.gov/documents/organization/272012.pdf>
29. Erdaw T, Ameni G. Survival and predictors of mortality among human immunodeficiency virus patients on anti-retroviral treatment at Jinka Hospital , South Omo , Ethiopia : a six years retrospective cohort study. *Epidemiol Health*. 2016;38(e2016049):1–10.
30. Yimer YT, Yalew AW. Magnitude and Predictors of Anti-Retroviral Treatment (ART) Failure in Private Health Facilities in Addis Ababa , Ethiopia. *PLoS One*. 2015;10(5).