

1 Efficient Prediction of Vitamin B Deficiencies via
2 Machine-learning Using Routine Blood Test Results in Patients
3 With Intense Psychiatric Episode

4
5 Hidetaka Tamune^{1)2)3)5)*}, Jumpei Ukita³⁾⁴⁾⁵⁾, Yu Hamamoto¹⁾²⁾, Hiroko Tanaka¹⁾²⁾, Kenji
6 Narushima¹⁾, Naoki Yamamoto¹⁾

7 1) Department of Neuropsychiatry, Tokyo Metropolitan Tama Medical Center, Tokyo,
8 Japan

9 2) Department of Neuropsychiatry, Graduate School of Medicine, The University of
10 Tokyo, Tokyo, Japan

11 3) Mental Health Research Course, Faculty of Medicine, The University of Tokyo,
12 Tokyo, Japan

13 4) Department of Physiology, Graduate School of Medicine, The University of Tokyo,
14 Tokyo, Japan

15 5) H. Tamune and JU contributed equally to this work

16

17 * Correspondence:

18 Hidetaka Tamune, M.D., tamune-tyk@umin.ac.jp

19

20 Abstract: 294 words

21 Main text: 1878 words + 4 Tables + 4 Figures + 24 references

22 Keywords: Machine Learning; Random Forest Classifier; Vitamin B Deficiency; Folic
23 Acid; Early Diagnosis; Decision support techniques or decision making.

24

25 **Abstract**

26 **Background:** Vitamin B deficiency is common worldwide and may lead to psychiatric
27 symptoms; however, vitamin B deficiency epidemiology in patients with intense
28 psychiatric episode has rarely been examined. Moreover, vitamin deficiency testing is
29 costly and time-consuming. It hampered to effectively rule out vitamin
30 deficiency-induced intense psychiatric symptoms. In this study, we aimed to clarify the
31 epidemiology of these deficiencies and efficiently predict them using machine-learning
32 models from patient characteristics and routine blood test results that can be obtained
33 within one hour.

34 **Methods:** We reviewed 497 consecutive patients deemed to be at imminent risk of
35 seriously harming themselves or others over 2 years. Machine-learning models were
36 trained to predict each deficiency from age, sex, and 29 routine blood test results.

37 **Results:** We found that 112 (22.5%), 80 (16.1%), and 72 (14.5%) patients had vitamin
38 B₁, vitamin B₁₂, and folate (vitamin B₉) deficiency, respectively. Also, the
39 machine-learning models well generalized to predict the deficiency in the future unseen
40 data; areas under the receiver operating characteristic curves for the validation dataset
41 (i.e. dataset not used for training the models) were 0.716, 0.599, and 0.796, respectively.
42 The Gini importance of these vitamins provided further evidence of a relationship
43 between these vitamins and the complete blood count, while also indicating a hitherto
44 rarely considered, potential association between these vitamins and alkaline phosphatase
45 (ALP) or thyroid stimulating hormone (TSH).

46 **Discussion:** This study demonstrates that machine-learning can efficiently predict some
47 vitamin deficiencies in patients with active psychiatric symptoms, based on the largest
48 cohort to date with intense psychiatric episode. The prediction method may expedite
49 risk stratification and clinical decision-making regarding whether replacement therapy
50 should be prescribed. Further research includes validating its external generalizability in
51 other clinical situations and clarify whether interventions based on this method can

52 improve patient care and cost-effectiveness.

53

54 **1. Introduction**

55 Vitamin B deficiency is common worldwide and may lead to psychiatric
56 symptoms¹⁻⁴. For example, meta-analyses have shown that patients with schizophrenia
57 or first-episode psychosis have lower folate (vitamin B₉) levels than their healthy
58 counterparts^{4,5}. Moreover, vitamin therapy can effectively alleviate symptoms in a
59 subgroup of patients with schizophrenia^{3,6-8}. However, the epidemiology of vitamin B
60 deficiency in patients with active mental symptoms requiring immediate hospitalization
61 has rarely been examined.

62 In a psychiatric emergency, psychiatrists should promptly distinguish treatable
63 patients with altered mental status due to a physical disease from patients with an
64 authentic mental disorder (international statistical classification of diseases and related
65 health problems-10, ICD-10 code: F2-9). However, vitamin deficiency testing is very
66 costly (around 60 dollars for each measurement of vitamin B₁ (vitB₁), vitamin B₁₂
67 (vitB₁₂), or folate in the U.S.; 15–25 dollars for each test in Japan) and usually requires
68 at least two days. Therefore, an efficient, cost-effective method of predicting vitamin B
69 deficiency is needed.

70 Although several studies have applied machine-learning to the prediction of
71 diagnosis or treatment outcomes⁹⁻¹¹, no study using machine-learning has focused on
72 vitamin B deficiencies. We herein explore whether vitB₁, vitB₁₂, and folate deficiencies
73 can be predicted using a machine-learning classifier from patient characteristics and
74 routine blood test results obtained within one hour based on a large cohort of patients
75 requiring urgent psychiatric hospitalization.

76

77 **2. Methods**

78 **2.1. Medical chart review**

79 We reviewed consecutive patients admitted to the Department of
80 Neuropsychiatry at Tokyo Metropolitan Tama Medical Center between September 2015
81 and August 2017 under the urgent involuntary hospitalization law, which requires the
82 immediate psychiatric hospitalization of patients at imminent risk of seriously harming
83 themselves or others. The necessity of hospitalization was judged by designated mental
84 health specialists. The patient characteristics, ICD-10 codes, and laboratory data were
85 gathered retrospectively.

86 Since the reference ranges for vitB₁, vitB₁₂, and folate are 70–180 nmol/L
87 (30–77 ng/mL), 180–914 ng/L, and > 4.0 µg/L, respectively¹², a deficiency of the
88 nutrients was defined as < 30 ng/mL, < 180 ng/L, and < 4.0 µg/L, respectively, unless
89 otherwise stated.

90

91 **2.2. Random forest classifier and statistics**

92 A random forest classifier was trained to predict the deficiency of each
93 substance from age, sex, and 29 routine blood variables (described in the Result section
94 with values). The random forest classifier was trained using the dataset populated in the
95 period from September 2015 to December 2016 (the “Training set”). First, we
96 optimized the hyperparameters of the classifier by selecting the best combination of
97 hyperparameters that maximized the “5-fold cross validation” accuracy, among many
98 combinations within appropriate ranges. The cross-validation accuracy was computed as
99 follows; in one session, the classifier was trained using 80% of the training set and
100 evaluated on the withheld 20% of the training set. This session was performed five
101 times so that every data would be withheld once. The accuracies were finally averaged
102 across sessions to yield the cross-validation accuracy. By incorporating this process, the
103 classifier was generalized to unseen data (Graphical method is shown in **Figure 1**).

104 Using the optimized hyperparameters, the classifiers were then validated using
105 data collected from January 2017 through August 2017 (the “Validation set”). We report
106 the classification performance on the validation set in the results section unless
107 otherwise stated. We quantified the sensitivity, specificity, and accuracy (defined as the
108 average of the sensitivity and the specificity on the optimal operating point) using
109 receiver operating characteristic curves (ROCs). We also quantified the 95% confidence
110 interval of the accuracy using 1000-times bootstrapping.

111 When investigating the Gini importance and the partial dependency¹³, we
112 retrained the classifiers using all datasets. All data analyses were performed using
113 Python (2.7.10) with the Scikit-learn package (0.19.0) and R (3.4.2) with the edarf
114 package (1.1.1).

115

116 **2.3. Ethical considerations**

117 Informed consent was obtained from participants using an opt-out form on the
118 website. The study protocol was approved by the Research Ethics Committee, Tokyo
119 Metropolitan Tama Medical Center (Approval number: 28-8). The study complied with
120 the Declaration of Helsinki and the STROBE statement.

121

122 **3. Results**

123 **3.1. Eligible patients**

124 During the 2-year study period, 497 consecutive patients (496 were Asian)
125 were enrolled. The mean age (standard deviation, SD) was 42.3 (\pm 15.4) years, and 228
126 patients (45.9%) were women. F2 (Schizophrenia, schizotypal, delusional, and other
127 non-mood psychotic disorders) was diagnosed in over 60% of the patients. The ICD-10
128 codes of the patients and the number of deficiencies at several cut-off values for vitB₁,
129 vitB₁₂, and folate are shown in **Table 1**. According to the predefined cut-off values¹²,
130 112 (22.5%), 80 (16.1%), and 72 (14.5%) patients exhibited a deficiency of vitB₁ (<30
131 ng/mL), vitB₁₂ (<180 ng/L), and folate (<4.0 μ g/L), respectively. Vitamin B deficiencies
132 in sub-groups are shown in **Table 2**. A summary of the full dataset is shown in **Table 3**.
133 Detailed information (sub-datasets) is shown in **Supplementary Table 1, 2, and 3**
134 online. Histograms of vitB₁, vitB₁₂, and folate values are shown in **Figure 2 A-C**.

135

136 **3.2. Prediction via machine-learning using routine blood test results**

137 A random forest classifier was trained to predict the deficiency of each
138 substance from patient characteristics and routine blood test results. The classifier was
139 trained using the dataset gathered in the period from September 2015 to December 2016
140 (the “Training set”, n = 373), which was then validated from January 2017 through
141 August 2017 (the “Validation set”, n = 124).

142 The area under the ROCs (AUCs) for the validation set were 0.716, 0.599, and
143 0.796, for vitB₁, vitB₁₂, and folate, respectively (**Figure 2 D-F** and **Table 4**). With some
144 operative points on the ROC, the sensitivity, specificity and accuracy for the validation
145 set were calculated (**Table 4**. See also **Supplementary Table 4** for training set and
146 **Supplementary Table 5** for different operating points).

147 When the prediction performances were compared between the classifiers
148 trained using the dataset from the F2 population and the classifiers trained using the

149 dataset from the other population, the AUC was not statistically different (DeLong's
150 test), except in the case of vitB₁ (see **Supplementary Table 6**).

151 **Figure 3** shows the Gini importance (a–c) and partial dependency plots (d–f)
152 for the eight most important variables for each substance. The results provided further
153 evidence of a relationship between the vitamin B levels and complete blood count while
154 also indicating the hitherto rarely considered, potential association between these
155 vitamins and alkaline phosphatase (ALP) or thyroid stimulating hormone (TSH).

156

157 **3.3. Robustness verification**

158 We verified the robustness of the results by two independent means. First, we
159 used different cut-off values to define the deficiency^{14–16}. Although the AUC for the
160 validation set, shown in **Supplementary Table 7**, tended to be higher when strict
161 cut-off values were used, the obtained AUCs were not statistically significant ($p > 0.05$,
162 DeLong's test with Bonferroni correction).

163 Second, we trained and evaluated random forest classifiers using a dataset split
164 in a different way; the classifier was trained using the dataset collected in the period
165 from the 31st of January, 2016 to August 2017, which was then validated with data
166 gathered from September 2015 to the 31st of January, 2016. Note that the sample sizes
167 of the training and validation sets were equal to those in the original setting. The AUCs
168 for the validation set were 0.771, 0.621, and 0.745 for vitB₁, vitB₁₂, and folate,
169 respectively; none were statistically different from the AUC trained using the original
170 setting (DeLong's test), further demonstrating the robustness of the performance.

171

172 **4. Discussion**

173 **4.1. Relevance of the present study**

174 Based on the largest cohort to date of patients at imminent risk of seriously
175 harming themselves or others, this study indicated that deficiency of certain vitamins
176 can be predicted in an efficient manner via machine-learning using routine blood test
177 results. Given the large number of patients with vitamin B deficiencies, empirical
178 therapy might be acceptable; however, risk stratification is preferred for personalized
179 medicine and shared decision-making. The prediction method presented here may
180 expedite clinical decision-making as to whether vitamins should be prescribed to a
181 patient (Graphical abstract is shown in **Figure 4**).

182 Remarkably, the AUC for folate deficiency was 0.796. Folate features the
183 potential to maintain neuronal integrity and is one of the homocysteine-reducing
184 B-vitamins⁵; homocysteine has been linked to the etiology of schizophrenia¹⁷, and
185 vitamin B supplements have been reported to reduce psychiatric symptoms significantly
186 in patients with schizophrenia⁷. As our study does not present longitudinal results, an
187 intervention effect of folate supplementation in the cohort remains to be clarified.

188

189 **4.2. Trade-off of interpretability and generalizability using machine-learning**

190 Compared to the AUC of folate, AUCs of vitB₁ and vitB₁₂ were relatively low.
191 Using other parameters that were not incorporated into this model or using other models
192 including deep neural networks might increase the accuracy of prediction.

193 However, interpretability and completeness of machine-learning classifiers are
194 subject to trade-off¹⁷. Although completeness and generalizability are desirable,
195 interpretability is also indispensable, especially in clinical settings, since it provides
196 meaningful and trustworthy findings for clinical physicians as well as new biological
197 insights¹⁸. In this study we chose random forest classifiers since they provide expressive
198 and interpretable data, with sufficient accuracy.

199

200 **4.3. Biological mechanism suggestion**

201 Using the random forest classifiers, as shown in **Figure 3**, we identified several
202 items related to complete blood count as top hits. Notably, our classifier was blind to
203 any biological knowledge, including the well-established association between anemia
204 and vitamin B deficiency, including folate¹⁹. The results provide further evidence of a
205 relationship between vitamin B levels and the complete blood count and support the use
206 of machine-learning to investigate novel, underlying biological mechanisms²⁰.

207 ALP and its metabolites indicate the vitamin B₆ status²¹; low vitB₁₂ is
208 potentially associated with low ALP²². More generally, ALP may have a close and
209 complicated relationship with the overall vitamin B group. Autoimmune disorders,
210 especially thyroid disease, are commonly associated with pernicious anaemia²³, but
211 there has been no established hypothesis regarding the causal relationships between
212 thyroid disease and vitamin B deficiencies. The potential association between the levels
213 of these vitamins and ALP or TSH awaits further study, both investigations of
214 populations and basic research²⁴.

215

216 **4.4. Limitations**

217 This study is subject to several limitations. First, the findings of this
218 single-center retrospective study may have limited generalizability. Second, the patients'
219 long-term prognosis was not investigated due to administrative restrictions; the extent to
220 which this method can expedite clinical decision-making is therefore unclear. Further,
221 we did not investigate the relationship between serological values and the need for
222 intervention. The lack of data for vitamin B deficiency in the Japanese general
223 population hampered the comparison between the experimental cohort and their
224 counterparts who lacked psychiatric symptoms. Establishing appropriate reference
225 values and an assessment method requires further investigation. Finally, we did not

226 assess the predictive value of other nutritional impairments, including vitamin B₆ and
227 homocysteine deficiency, which were previously shown to have a close link with
228 psychiatric symptoms^{3,5}; however, our study provides fundamental data on nutritional
229 impairment based on the largest cohort of patients with intense psychiatric episode ever
230 assembled for this purpose and presents a potential framework for predicting nutritional
231 impairment using machine-learning.

232

233 **4.5. Conclusion**

234 The present report is, to the best of our knowledge, the first to demonstrate that
235 machine-learning can efficiently predict nutritional impairment. Further research is
236 needed to validate the external generalizability of the findings in other clinical situations
237 and clarify whether interventions based on this method can improve patient care and
238 cost-effectiveness.

239

240 **5. Contribution to the Field Statement**

241 Vitamin B deficiency is common worldwide and may lead to psychiatric
242 symptoms; however, vitamin B deficiency epidemiology in patients with intense
243 psychiatric symptoms has rarely been examined. Moreover, vitamin deficiency testing is
244 costly and time-consuming. Based on the largest cohort to date of patients at imminent
245 risk of seriously harming themselves or others, this study demonstrated that the
246 deficiency of certain vitamins can be predicted in an efficient manner via
247 machine-learning models from patient characteristics and routine blood test results
248 obtained within one hour.

249 In detail, among the 497 patients investigated (over 60% was diagnosed with
250 schizophrenia or related psychotic disorders), 22.5%, 16.1%, and 14.5% patients had a
251 deficiency of vitamin B₁, B₁₂, and folate, respectively, by direct measurement. Also, the
252 machine-learning models well generalized to predict the deficiency in unseen datasets;
253 areas under the receiver operating characteristic curves for the validation dataset were
254 0.716, 0.599, and 0.796, respectively. The prediction method presented in this study
255 may expedite risk stratification and clinical decision-making regarding whether
256 replacement therapy should be prescribed. The results also provided further evidence for
257 a well-known relationship between these vitamins and the complete blood count and
258 supported the application of machine-learning to investigate novel, underlying
259 biological mechanisms.

260

261 **6. Acknowledgements**

262 We thank Mr. James Robert Valera for his assistance in editing this manuscript
263 and all the staff for their care of the patients and their contributions to this study.

264

265 **7. Author Contributions Statement**

266 H. Tamune has full access to all data and takes responsibility for the integrity of
267 the data. H. Tamune, JU, KN, and NY conceived the study. H. Tamune, YH, and H.
268 Tanaka collected the data. JU performed the statistical analyses. H. Tamune and JU
269 drafted the first version of the manuscript. All authors critically revised the manuscript
270 for intellectual content and approved the final version.

271

272 **8. Data Availability Statements**

273 The datasets and source code utilized in the current study are available from the
274 corresponding author upon reasonable request.

275

276 **9. Conflict of Interest Statement**

277 The authors declare no conflict of interest, except for a scholarship grant
278 awarded to JU from Takeda Science Foundation and Masayoshi Son Foundation.

279

280

281 **References**

- 282 1. Harper C. Thiamine (vitamin B1) deficiency and associated brain damage is still
283 common throughout the world and prevention is simple and safe! *Eur J Neurol.* (2006)
284 13: 1078–1082.
- 285 2. Reynolds E. Vitamin B12, folic acid, and the nervous system. *Lancet Neurol.* (2006)
286 5: 949–960.
- 287 3. Arai M, Yuzawa H, Nohara I, Ohnishi T, Obata N, Iwayama Y *et al.* Enhanced
288 carbonyl stress in a subpopulation of schizophrenia. *Arch Gen Psychiatry* (2010) 67:
289 589–597.
- 290 4. Cao B, Wang DF, Xu MY, Liu YQ, Yan LL, Wang JY *et al.* Lower folate levels in
291 schizophrenia: A meta-analysis. *Psychiatry Res.* (2016) 245: 1–7.
- 292 5. Firth J, Carney R, Stubbs B, Teasdale SB, Vancampfort D, Ward PB, *et al.* Nutritional
293 deficiencies and clinical correlates in first-episode psychosis: A systematic review and
294 meta-analysis. *Schizophr Bull.* (2018) 44: 1275–1292.
- 295 6. Levine J, Stahl Z, Sela BA, Ruderman V, Shumaico O, Babushkin I *et al.*
296 Homocysteine-reducing strategies improve symptoms in chronic schizophrenic patients
297 with hyperhomocysteinemia. *Biol Psychiatry* (2006) 60: 265–269.
- 298 7. Firth J, Stubbs B, Sarris J, Rosenbaum S, Teasdale S, Berk M *et al.* The effects of
299 vitamin and mineral supplementation on symptoms of schizophrenia: A systematic
300 review and meta-analysis. *Psychol Med.* (2017) 47: 1515–1527.
- 301 8. Itokawa M, Miyashita M, Arai M, Dan T, Takahashi K, Tokunaga T *et al.*
302 Pyridoxamine: A novel treatment for schizophrenia with enhanced carbonyl stress.
303 *Psychiatry Clin Neurosci.* (2018) 72: 35–44.
- 304 9. Koutsouleris N, Kahn RS, Chekroud AM, Leucht S, Falkai P, Wobrock T *et al.*
305 Multisite prediction of 4-week and 52-week treatment outcomes in patients with
306 first-episode psychosis: A machine learning approach. *Lancet Psychiatry* (2016) 3:
307 935–946.

- 308 10. Mechelli A, Lin A, Wood S, McGorry P, Amminger P, Tognin S *et al.* Using clinical
309 information to make individualized prognostic predictions in people at ultra-high risk
310 for psychosis. *Schizophr Res.* (2017) 184: 32-38.
- 311 11. Vieira S, Pinaya WH, Mechelli A. Using deep learning to investigate the
312 neuroimaging correlates of psychiatric and neurological disorders: Methods and
313 applications. *Neurosci Biobehav Rev.* (2017) 74: 58–75.
- 314 12. Mayo Foundation for Medical Education and Research, Rochester Test Catalog.
315 <https://www.mayomedicallaboratories.com/test-catalog/> (2018).
- 316 13. Friedman, JH. Greedy function approximation: A gradient boosting machine. *Ann.*
317 *Stat.* (2001) 29: 1189–1232.
- 318 14. Sasaki T, Yukizane T, Atsuta H, Ishikawa H, Yoshiike T, Takeuchi T *et al.* A case of
319 thiamine deficiency with psychotic symptoms: Blood concentration of thiamine and
320 response to therapy. *Seishin Shinkeigaku Zasshi* (2010) 112: 97–110.
- 321 15. Clarke R, Refsum H, Birks J, Evans JG, Johnston C, Sherliker P *et al.* Screening for
322 vitamin B-12 and folate deficiency in older persons. *Am J Clin Nutr.* (2003) 77:
323 1241–1247.
- 324 16. Goff DC, Bottiglieri T, Arning E, Shih V, Freudenreich O, Evins AE *et al.* Folate,
325 homocysteine, and negative symptoms in schizophrenia. *Am J Psychiatry* (2004) 161:
326 1705–1708.
- 327 17. Muntjewerff JW, Kahn RS, Blom HJ, den Heijer M. Homocysteine,
328 methylenetetrahydrofolate reductase and risk of schizophrenia: A meta-analysis. *Mol*
329 *Psychiatry* (2006) 11: 143–149.
- 330 18. Gilpin LH, Bau D, Yuan BZ, Bajwa A, Specter M, Kagal L. Explaining
331 explanations: An overview of interpretability of machine learning. In *2018 IEEE 5th*
332 *International Conference on Data Science and Advanced Analytics (DSAA)*: 80–89.
- 333 19. Evans TC, Jehle D. The red blood cell distribution width. *J Emerg Med.* (1991) 9:
334 71–74.

- 335 20. So HC, Chau CK, Chiu WT, Ho KS, Lo CP, Yim SH *et al.* Analysis of genome-wide
336 association data highlights candidates for drug repositioning in psychiatry. *Nat Neurosci.*
337 (2017) 20: 1342–1349.
- 338 21. Ueland PM, Ulvik A, Rios-Avila L, Middtun Ø, Gregory JF. Direct and functional
339 biomarkers of vitamin B6 status. *Annu Rev Nutr.* (2015) 35: 33–70.
- 340 22. Carmel R, Lau KH, Baylink DJ, Saxena S, Singer FR. Cobalamin and
341 osteoblast-specific proteins. *N Engl J Med.* (1977) 319: 70–75.
- 342 23. Stabler SP. Vitamin B12 deficiency. *N Engl J Med.* (2013) 368: 149–160.
- 343 24. Zheng Y, Cantley LC. Toward a better understanding of folate metabolism in health
344 and disease. *J Exp Med.* (2018) 216: 253–266.
- 345

346 **Supporting Material List**

347 Supplementary Table 1 (related to Table 1). Divided patient distribution data (n = 497)

348 Supplementary Table 2 (related to Table 2). Divided data of vitamin B deficiencies in
349 sub-groups

350 Supplementary Table 3 (related to Table 3). Divided dataset of age, sex, and 29
351 parameters

352 Supplementary Table 4 (related to Table 4). Summary of sensitivity, specificity, and
353 accuracy for the training set

354 Supplementary Table 5 (related to Table 4). Sensitivities and specificities at other
355 operating points

356 Supplementary Table 6. Subgroup analyses

357 Supplementary Table 7. AUC with different cut-off values

358

359 **Legends**

360 **Figure 1: Graphical illustration of method of machine-learning**

361

362 **Figure 2: Histogram and ROC curves of each vitamin B value**

363 (A-C) The histograms for vitamin B₁, vitamin B₁₂, and folate (vitamin B₉).

364 Their medians (1st–3rd quartile) are 35 (30–42) ng/mL, 285 (206–431) ng/L, and 7.2

365 (4.9–10.8) µg/L, respectively.

366 (D-F) ROC curves for vitamin B₁, vitamin B₁₂, and folate. Operating points

367 used in **Table 4** and **Supplementary Table 5** are depicted in blue.

368 Abbreviations: Vit B₁, vitamin B₁; Vit B₁₂, vitamin B₁₂.

369

370 **Figure 3: Gini importance and partial dependence plots of vitamin B deficiencies**

371 The Gini importance (A-C) and partial dependency plots of the probability of

372 deficiency (D-F) are shown for the eight most important variables for vitamin B₁,

373 vitamin B₁₂, and folate (vitamin B₉). Combined with these, this machine-learning

374 classifier without hypothesis also provided further evidence of a relationship between

375 vitamin B levels and the complete blood count while also indicating a potential

376 association between these vitamins and alkaline phosphatase (ALP) or

377 thyroid-stimulating hormone (TSH).

378 Abbreviations: Vit B₁, vitamin B₁; Vit B₁₂, vitamin B₁₂; Hb, hemoglobin; Hct,

379 hematocrit; WBC, white blood cell count; CK, creatine kinase; RDW.CV, red blood cell

380 distribution width-coefficient variation; Plt, platelet; ALT, alanine transaminase; Lym,

381 lymphocyte fraction; Cre, creatinine; Neu, neutrocyte fraction; γGTP,

382 γ-glutamyltransferase; MCV, mean corpuscular volume; glu, plasma glucose.

383

384 **Figure 4: Graphical abstract**

385

386 **Table 1. Patient distribution data (n = 497)**

387

	ICD-10 code										VitB ₁			VitB ₁₂			Folate		
	F0	F1	F2	F3	F4	F5	F6	F7	F8	F9	<20	<28	<30*	<150	<180*	<200	<3.0	<4.0*	<5.0
N	28	21	300	58	16	0	29	20	24	1	15	81	112	37	80	107	29	72	134
%	5.6	4.2	60.4	11.7	3.2	0.0	5.8	4.0	4.8	0.2	3.0	16.3	22.5	7.4	16.1	21.5	5.8	14.5	27.0

388

389 Asterisks show the predefined cut-off values for vitamin B₁, vitamin B₁₂, and folate
 390 (vitamin B₉) based on a reference¹²; different cut-off values based on other papers¹⁴⁻¹⁶
 391 are also presented for further investigation.

392

393 ICD-10 codes. F0, Mental disorders due to known physiological conditions; F1, Mental
 394 and behavioral disorders due to psychoactive substance use; F2, Schizophrenia,
 395 schizotypal, delusional, and other non-mood psychotic disorders; F3, Mood disorders;
 396 F4, Anxiety, dissociative, stress-related, somatoform, and other non-psychotic mental
 397 disorders; F5, Behavioral syndromes associated with physiological disturbances and
 398 physical factors; F6, Disorders of adult personality and behavior; F7, Intellectual
 399 disabilities; F8, Pervasive and specific developmental disorders; F9, Behavioral and
 400 emotional disorders with onset usually occurring in childhood and adolescence.

401

402 **Table 2. Vitamin B deficiencies in sub-groups**

403

	F0	F1	F2	F3	F4	F6	F7	F8	F9
vitB ₁ < 30	9 (32%)	4 (19%)	70 (23%)	11 (19%)	3 (19%)	7 (24%)	5 (25%)	3 (13%)	0
vitB ₁₂ < 180	5 (18%)	4 (19%)	53 (18%)	7 (12%)	3 (19%)	1 (3%)	4 (20%)	3 (13%)	0
Folate < 4.0	5 (18%)	7 (33%)	38 (13%)	6(10%)	5 (31%)	3 (10%)	4 (20%)	4 (17%)	0

404

405 Abbreviations; see **Table 1**.

406

407

408 **Table 3. Summary of full dataset of age, sex, and 29 parameters for**
 409 **machine-learning**

410

Parameters	Units	Mean	SD	UN	mg/dL	12.9	6.7
Age	years	42.3	15.4	Cre	mg/dL	0.7	0.2
Sex	Woman	n = 228		T.bil	mg/dL	0.7	0.4
WBC	$\times 10^3/\mu\text{L}$	8.2	2.8	Na	mmol/L	139	3
Hb	g/dL	13.7	1.7	Cl	mmol/L	105	4
Hct	%	40.3	4.5	K	mmol/L	3.7	0.4
MCV	fL	89	6.6	cor.Ca	mg/dL	9.1	0.5
Plt	$\times 10^4/\mu\text{L}$	24.9	6.3	CK	IU/L	514	1230
RDW.CV	%	13.5	1.3	AST	IU/L	31	34
Neu	%	70	11	ALT	IU/L	27	24
Lym	%	23	10	LDH	IU/L	239	91
Mono	%	6	2	ALP	IU/L	224	81
Eo	%	1	2	γGTP	IU/L	37	63
Baso	%	0	0	Glu	mg/dL	112	40
TP	g/dL	7.2	0.6	CRP	mg/dL	0.4	0.9
Alb	g/dL	4.4	0.4	TSH	$\mu\text{IU/mL}$	1.7	2.4

411

412 Two patients lacked age data (no photo ID was available), and one patient lacked
 413 biochemistry data (inappropriate sample processing). For machine-learning, the missing
 414 values were replaced using the mean.

415

416 Abbreviations: WBC, white blood cell count; Hb, hemoglobin; Hct, hematocrit; MCV,
 417 mean corpuscular volume; RDW.CV, red blood cell distribution width-coefficient
 418 variation; Plt, platelet; Neu, neutrocyte fraction; Lym, lymphocyte fraction; Mono,
 419 monocyte fraction; Eo, eosinocyte fraction; Baso, basocyte fraction; TP, total protein;
 420 Alb, albumin; UN, urea nitrogen; Cre, creatinine; T.bil, total bilirubin; Na, sodium; Cl,

421 chloride; K, potassium; cor.Ca, corrected calcium; CK, creatine kinase; AST, aspartate
422 transaminase; ALT, alanine transaminase; LDH, lactate dehydrogenase; ALP, alkaline
423 phosphatase; γ GTP, γ -glutamyltransferase; Glu, plasma glucose; CRP, C-reactive
424 protein; TSH, thyroid-stimulating hormone.
425

426 **Table 4. Summary of AUC, sensitivity, specificity, and accuracy for the validation**
427 **set**

428

	vitB ₁	vitB ₁₂	Folate
AUC	0.716	0.599	0.796
Sensitivity	0.594	0.316	0.667
Specificity	0.783	0.943	0.917
Accuracy	0.688 [0.597–0.787]	0.629 [0.523–0.746]	0.792 [0.665–0.909]

429

430 Generalization performance of the classifiers was evaluated using AUC of the validation
431 set. Sensitivity, specificity, and accuracy of the classification at the optimal operating
432 points that maximized accuracy on the receiver operating characteristic curve of the
433 validation set are also shown (see also **Figure 2 D-F**). Accuracy was defined as the
434 average of the sensitivity and specificity. Square brackets indicate the 95% CI. Note that
435 the 95% CI of each accuracy does not include 0.5, which demonstrates statistical
436 significance. For further information, see **Figure 2** and **Supplementary Table 5**.

437

438 Abbreviations: AUC, area under the receiver operating characteristic curve; CI,
439 confidence interval.

440




Figure 1




Figure 2




Figure 3




Figure 4