

## Quaternary Prevention: Is this Concept Relevant to Public Health? A Bibliometric and Descriptive Content Analysis.

Miguel Andino Depallens<sup>1</sup>  
Jane Mary de Medeiros Guimarães<sup>1</sup>  
Naomar Almeida Filho<sup>2</sup>

<sup>1</sup> Federal University of the South of Bahia (UFSB)

<sup>2</sup> Federal University of Bahia (UFBA)

### Abstract

**Objective** To measure and map research output on Quaternary Prevention (P4) and outline research trends; to assess the papers content, mainly regarding methods and subjects approached in order to contribute to the improvement of global knowledge about P4 and to evaluate its relevance for public health.

**Design** Bibliometric and descriptive content analysis.

**Articles reviewed** Scientific articles about P4 recorded in Pubmed, LILACS, Scielo or CINAHL published until August 2018, with correspondent full articles available in Portuguese, English, Spanish, German or French.

**Main outcome measures** Year of publication, first authors' name and nationality, journals' name, country and ranking, publication language, used methods and main reported subjects.

**Results** 65 articles were included, published in 33 journals of 16 countries between 2003 and 2018 with a peak of publications in 2015. The first authors came from 17 different countries, 23% of them were Brazilian and Uruguay was the leading nation according to the scientific production per capita. 40% of all the selected articles were in English, 32% in Portuguese, 26% in Spanish. 28% of the papers were published in Q1 or Q2 journals. The research outputs on P4 begun first in the South of Europe, went to South America and then expanded worldwide. 88% of the articles were bibliographic research and 38% of all focused on specific examples of medical overuse (including several screening tests).

**Conclusions** Quaternary prevention represents an ethical and valid approach to prevent occurrence of iatrogenic events and to achieve equal and fair access to health services. Conceptual, geographical and linguistic elements, as well as WONCA conferences and type of healthcare systems in the authors' country were fundamental factors that affected research output. The quality and quantity of available studies is still limited, therefore further investigations are recommended to assess the effective impact of P4 on public health.

Key-words: “Quaternary Prevention”; “Medical Overuse”; “Public Health”; “Overdiagnosis”; “Overmedicalization”.

## Introduction

The general notion of prevention has been instrumental for health care provision since early years in Ancient Greece. The simple idea that human beings should be protected from the influence of factors potentially harmful to their health indeed guided modern forms of medical practice (clinical, pharmacological and surgical interventions) and public health (sanitation, vaccination etc.) oriented by scientific criteria. These developments took place initially in Western Europe, during the Enlightenment era, resulting in the movements of Hygiene, Medical Police and Social Medicine.<sup>1</sup> The institutionalization and spread of preventive practices and measures in Medicine and Public Health were consolidated during the first half of the XXth century, largely due to Leavell & Clark’s ternary model of types or levels of prevention: primary, secondary and tertiary.<sup>2</sup>

The idea that medical practice in all manifestations could be fully described as different modalities or levels of preventive action have only prevailed particularly since World War II. However, despite strong efforts towards integrality and equity in health care, such as the WHO’s Health for All campaign, only late in past century criticisms and conceptual advances were brought about concerning this subject-matter.<sup>3</sup> Among them, a definition of “quaternary prevention” regarding iatrogenic effects, as proposed the Belgian family physician Marc Jamouille (1986)<sup>4</sup>, was an add-on to the theory of levels of prevention. This novel conceptual and practical approach may have enormous potential to foster better-quality, socially referenced, more humane health care systems and practices, needed especially in these days of widespread budget cuts and exclusionary policies in public health services, all over the world.

The main objective of this paper is to report a bibliometric and descriptive content analysis of research output on the concept of Quaternary Prevention (P4) published until August of 2018, and to prospectively correlate research trends. To achieve such goals, first, we briefly provide some theoretical and methodological background, and context information on the historical development of prevention approaches. Second, we present methods, describe the criteria for considering the papers and point the types of data that were collected, analyzed and classified, in order to measure and map scientific production as well as to assess the papers content. Third, we expose the results, focusing on: year of publication, authors’ and journals’ features, methods used in the selected studies and main topics approached. And fourth, we outline and discuss some trends towards research on Quaternary Prevention, emphasizing the main methods and subjects treated.

We hope to contribute to improve the global knowledge of P4 in order to further the understanding of this recent and relevant concept, highlighting authors, countries and journals involved in the research and publications on this topic and to classify the selected articles. In further papers, we intend to report a critical synthesis and in-depth analyses of the selected articles in a qualitative systematic review.

## Quaternary Prevention: Conceptual Background

The term iatrogenesis comes from the Greek language: “iatro” (medicine) and “genesis” (creation); it can be defined as medical harm or adverse effect. Iatrogenic events always existed in historical times. The Hippocratic oath (450-350 B. C. E.) confirms this statement and the awareness of the Ancient Greece physicians of the potential iatrogenic outcomes that could result from their practice. Across the centuries, the beneficence and non-maleficence principles – *primum non nocere* (first do no harm) – became one of the cornerstones of current bioethics.<sup>5</sup>

Over the course of the next 20 centuries, there was little increase in the effectiveness of healing therapies, remained not much effective. However, in those times, a few researchers produced very relevant public health knowledge and tangible strategies became available in the field of disease prevention. In this perspective, James Lind (1716–1794) established that the ingestion of some specific nutrients, like vitamin C, could prevent diseases, scurvy in this case. In a corresponding period, public health and prevention of disease became a priority for a few bellicose states that needed strong soldiers and workers. Johan Peter Frank (1745-1821) in Germany, implemented massive health policies to tackle infectious outbreaks related to hospitals, military medicine, litter sanitation, food and housing. Then, Jenner (1749-1823), Pasteur (1822-1895) and Koch (1843-1910) discovered the pathogenic role of some microorganisms and developed effective vaccines to prevent the related diseases (e. g.: smallpox, rabies). And the basics of modern epidemiology were established during the cholera outbreak in London (1849) by John Snow (1813–1858), who discovered the pathogenic role of infected water, mapped the cases of cholera and identified the contaminated spread of the sources, enabling the prevention of its dissemination.<sup>6</sup>

Spectacular individual and biochemical treatments followed upon the discovery of insulin therapy, by Banting (1922) and his team that enabled to save young diabetic end-stage patients from certain death.<sup>7</sup> A few years later, in 1928, Fleming (1881-1955) made another fabulous finding and discovered the penicillin that open the period of curative therapies. Around 1940, Sir Archibald Cochrane (1909-1988) developed a new and very significant concept: the evidence-based medicine (EBM). In this way, the randomized clinical trial (RCT) came to be considered a revolutionary scientific method, which enable the comparison of two different therapies.<sup>8</sup> Up to the present, RCTs are used worldwide to guide most of clinical decision-making.

At the end of the 1980s, influenced by Illich’s book about iatrogenesis<sup>9</sup>, the Belgian family physician Marc Jamoulle revised<sup>10</sup> the prevention levels of Leavell and Clark<sup>11</sup>, integrating a public health approach into individual clinical practice. His classification was designed according to two specific dimensions: one defined by the patient’s feeling (illness) and the other by the clinical assessment (disease)<sup>12</sup>. We put hereinafter the definition of the levels of prevention together with the initial approach of Jamoulle:

1. Primary prevention (P1) acts on healthy individual (absence of illness) to avoid the incidence of a specific disease (absence of disease), for example, through counseling about smoking cessation to prevent lung cancer and chronic obstructive pulmonary disease (COPD);
2. Secondary prevention (P2) aims to detect early a potentially severe disease in an asymptomatic population (absence of illness), increasing the probability of cure by

identifying and treating an initial stage of the detected disease (presence of disease), for example through a screening method (e. g.: annual eye funduscopy to detect diabetic retinopathy).

3. Tertiary prevention (P3) proposes to reduce the impacts of any disease on the quality of life (illness and disease present) through medical treatment or rehabilitation, for example through an earlier neurorehabilitation therapy after a disabling stroke.
4. Quaternary prevention (P4, first definition) aims to prevent medical overuse in situations where the patient feels ill (illness present), but the physician does not link the symptoms to any biological disease (absence of disease). An example of P4 could be the use of a “watchful waiting” strategy (watch and follow-up) when a young healthy patient without any cardiovascular risk or symptom worries about his cholesterol level.

We can define overmedicalization or medical overuse as a medical intervention that brings more harm than benefit.<sup>13</sup> In this situation and if a drug was prescribed to the young patient, it would be considered a typical case of overmedicalization. There are plenty of medical overuse/P4 situations and we will see some in further topics.

*Figure 1*

**Quaternary Prevention in all the levels of prevention according to the patient’s feeling and the physician’s assessment (adapted from Jamoulle<sup>14</sup>).**

In 2000, Barbara Starfield reported data showing that iatrogenic events represented the third cause of death in the United States.<sup>15</sup> Moreover, several authors were researching medical overuse and P4, denouncing and attempting to prevent iatrogenic procedures.<sup>16, 17, 18, 19</sup> In 2003, the concept of P4 was finally acknowledged by the scientific community and the World Organization of National Colleges, Academies and Academic Associations of General Practitioners/Family Physicians (WONCA) International Classification Committee. The following definition has been accepted: “Action taken to identify a patient or a population at risk of overmedicalization, to protect them from new medical invasions and suggest to them interventions, which are ethically acceptable”.<sup>20</sup> This new definition has an enhanced scope and is supported by the nonmaleficence bioethical principle. It aims to prevent all kinds of medical overuse without considering the features of the clinical situation, neither the patient’s feeling nor the physician’s assessment. Figure 1 may help in understanding both the initial and actual definition, and how P4 can be applied according to the specificities of some clinical situations.

As well as the prevention of medical overuse and iatrogenesis, P4 also aims to significantly lower some related specific costs of overall health expenditure responding to one of the actual worldwide governments’ main objectives.<sup>21, 22</sup> Consequently, P4 could allow the expansion of global health coverage and confirms its relevance for public health.

## **Methods**

To achieve an integral assessment of research output on P4, two complementary methods were used: a quantitative bibliometric approach and a qualitative content analysis. Bibliometrics is a methodology which focuses on measuring and evaluating research trends.<sup>23, 24</sup> Topics of bibliometric studies on adjacent topics are: medicalization during pregnancy,<sup>25</sup> overdiagnosis,<sup>26</sup> which emphasized breast cancer screening. Also, we found two more papers about “unnecessary procedures”.<sup>27, 28</sup> However, these studies focused on specific examples of overmedicalization or had significantly different objectives (e.g.: focusing the impact of Cochrane Reviews on clinical decision); none of them approached P4. The criteria of the Cochrane Handbook<sup>29</sup> have been used to build the search strategy and the papers’ selection process. “Evidence from qualitative studies can play an important role in adding value to systematic reviews for policy, practice and consumer decision-making”,<sup>30</sup> therefore, complementing the bibliometric quantitative approach, a content analysis has been achieved according to the Standards for Reporting Qualitative Research (SRQR)<sup>31</sup> and constitutes the second part of the present article.

#### *Criteria for considering articles for this review*

We included all scientific articles, published up until August, 7<sup>th</sup>, 2018, from the research databases, available in Portuguese, English, Spanish, German or French, and using the WICC (2003) definition’s meaning when it mentioned P4. We excluded articles that were commentary, letter or editorial as well as documents that were not articles, such as: master or doctorate thesis, teaching material, videos, among others, as well as all other records that did not fit our inclusion criteria.

#### *Data and outcomes*

The following data were collected: title; name of the author(s) and their nationality; year of publication; name of the journal and country of its headquarters; publication’s language; methods; abstracts.

#### *Search methods for identification of articles*

We did the search using the following databases: Pubmed, LILACS, Scielo and CINAHL. There were no existing records about P4 in the Cochrane Library; for this reason, this renowned database was not included. P4 is a relatively recent concept and thus there is not much literature about it, therefore we decided to use the widest possible Search Builder: using the conjunction “OR” and, when available, coupled with a Medical Subject Headings (MeSH). In Pubmed, Scielo and CINAHL, there was no pre-existing MeSH about Quaternary Prevention; therefore, we used the following Search Builder: “Quaternary Prevention” OR “Prevención Cuaternaria” OR “Prevenção Quaternária“. In LILACS, there was already a descriptor (*Descritores em Ciências da Saúde: DECS*) - “Quaternary Prevention” - that corresponds to a

MeSH in the Pubmed database. So, we expanded our Search Builder in LILACS as follows: (tw:("prevenção quaternária" OR "quaternary prevention" OR "prevención cuaternária")) OR (mh:(prevenção quaternária)). To illustrate the selection process of the papers, we used Review Manager 5.3® and designed a flow diagram (figure 2).

### *Data collection, analysis and extraction*

The searches were exported directly from the databases to Zotero®, which accelerated the transfer of information, the removal of duplicates and facilitated the achievement of the initial bibliographic list with a specific tool. Results from electronic databases searches were reviewed to check the inclusion and exclusion criteria. The abstracts were read, the papers classified and the extracted was digitalized in Excel® tables. The articles that were not freely available have been accessed through the CAPES (*Coordenação de Aperfeiçoamento de Pessoal de Nível Superior*) website.<sup>32</sup> We read the abstracts of the selected articles to check the meaning of P4 and, if necessary, excluded the ones which used a very different definition. Finally, we searched for the journal rank in two acknowledged databases: one European – Scimago<sup>33</sup> – and one South American institution – Qualis/CAPES.<sup>34</sup> The journal ranks were entered into the Excel® tables, together with previously collected data.

### *Bibliometric analysis*

We conducted the bibliometric analysis, building dynamic cross tables and outlining some trends according to the following data: year of publication; language of publication; name of the first authors and nationality; Publications per capita (Number of publications per first author's country / number of residents in million); The data of each country's population in 2019 has been obtained calculating an estimation stem from the *United Nations population estimates and projections*<sup>35</sup>; name of the journal and country of its headquarters; journal ranking (2017 Scimago: quartile Q1-Q5); journal ranking (2013-2016 Qualis/CAPES: A1-C).

### *Content analysis*

Focusing on the title and abstract, we categorized each paper according to its method, such as: bibliographic research, qualitative research, quantitative research, case report, among others. Articles were also classified under one of the following categories:

1. Epistemological and conceptual research focusing on P4 general concept and/or the overmedicalization process;
2. Studies addressing bioethical issues related to P4, discussing in particular the non-maleficence and the justice principles;
3. Reviews that consider empirical evidence on medical overuse and iatrogenesis, highlighting examples of overmedicalization (except screening);

4. Screening also represents a type of potential iatrogenesis, so we separated it from the previous category because it is one of the main focus of P4;
5. Investigations that outline P4 implementation strategies and/or aim to divulge the concept.

We identified 245 records in the electronic databases: 152 in LILACS, 58 in Pubmed, 25 in Scielo and 10 in CINAHL. 121 duplicates were removed, and the remaining 124 records left were screened. We excluded one record, because we did not find it even after sending an email to the journal and the author (Gofrit, 2000); 16 because they were not scientific articles (master or doctorate thesis, teaching material, MeSH, second professional opinion or videos), four full-articles were in other languages (three in Serbo-Croat and one in Corean) and 22 were commentaries, letters or editorials. After these steps, we read the 71 abstracts left and excluded 6 more articles (Gadelha, 1992; McColl and al., 2017; Moses, Mawby, Phillips, 2013; Trova and al., 2015; Weinstein, 2001) as they used a different definition of P4, defining it like a physical or social rehabilitation, which did not match with the meaning of the definition accepted by the WICC (2003). Finally, 65 articles were included in the study (Figure 2).

*Figure 2*

#### **Flow chart of the articles selection**

### **Results**

#### **Bibliometric analysis**

The 65 selected articles were published between 2003 and 2018. Until 2011, there was less than one publication a year. Starting from 2012, we observe a slow increase in the scientific output until 2014; 2015 represents a peak with 23 publications (35% of the total number). After 2015 (2016-2018), the publications returns to their basal numbers, similar to the 2012-2014 period (Figure 3).

*Figure 3*

#### **Number of papers according to the year of publication (n=65).**

#### *Authors*

The first authors came from 17 different countries in Europe, South and North America and Asia. Brazilian authors were the leaders of 15 papers (23%), the Spanish were in the second position with 11 published articles (17%) followed by the Portuguese with 8 publications (12%) (Table 1). To consider the population size of each countries, we calculated the scientific production per capita. Consequently, we reached another ranking in favour of the smaller countries: Uruguay was first (1,18), Portugal (0,80) moved up to the second place and there were three countries with a very close index in the third place: Belgium (0,36), Switzerland (0,24) and Spain (0,23); Brazil (0,07) fell to the 7<sup>th</sup> position.

Table 1

	till 2011 (included)	2012	2013	2014	2015	2016	2017	2018	Total	Prod. p/capita (in millions)
Brazil	1	2		4	3	2	2	1	15	0,07
Spain	6	1	1	1	2				11	0,23
Portugal		2	1	1	3			1	8	0,80
Argentina					3		2		5	0,11
Belgium				1	3				4	0,36
Uruguay					2	1	1		4	1,18
India					1	2			3	0,00
France					1			1	2	0,03
South Korea					1	1			2	0,04
Switzerland				1	1				2	0,24
Peru						1	1		2	0,06
Canada					2				2	0,05
Germany							1		1	0,01
Vietnam					1				1	0,01
Norway						1			1	0,2
Bolivia			1						1	0,09
Italy				1					1	0,02
Total	7	5	4	8	23	8	7	3	65	0,22

**Number of articles according to first author's nationality and year of publication (n=65) and production per capita.**

Until 2011, six out of seven articles were written by Spanish authors and, in particular, three by Juan Gérvas (2003, 2006, 2006) - he wrote another article in 2012. Gérvas is a family physician. He is a retired professor from the University of Valladolid (Spain) and the Johns Hopkins University (United States), committed researcher who published more than 400 articles in scientific journals, and has been a member of the WICC since 1986. He is one of the most active proponents of P4 and wrote one of the most significant books about this topic, *Sano y salvo* ("Healthy and Safe", 2012). Then, Norman and Tesser, promoted the concept in South America through their first article about P4 (2009). Both are Brazilian family physician, professors and researchers. We noted that until 2012, there were only Spanish, Brazilian and Portuguese authors who wrote about P4. Tesser, professor of the Federal University of Santa Catarina and of the University of Coimbra, and Jamoulle, the "father" of P4, family physician and researcher, are the most established authors on the subject of P4. Both had six articles included here. Beyond Europe and South America, 11 out of 17 (65%) authors came from Latin-speaking countries, we can see that Asian authors from South Korea, India and Vietnam, as much as North America, with Canadian authors also took part in this process (map 1).

Map 1

**Map of the expansion of the publications about P4 according to first author's nationality.**

## Languages







Korea													
Spain	4	5				2		4	1	2	4	11	
Switzerland			1			1			1	1		2	
Uruguay						4			3	1		4	
Vietnam						1				1		1	
<b>Total</b>	<b>4</b>	<b>14</b>	<b>8</b>	<b>6</b>	<b>1</b>	<b>32</b>	<b>0</b>	<b>1</b>	<b>0</b>	<b>2</b>	<b>3</b>	<b>24</b>	<b>65</b>

**Number of articles according to the journal rank (Scimago Quartile and Qualis/CAPES) and first author's nationality**

### *Methods of the selected papers*

The vast majority of the articles (88%) used a bibliographic research as their main method. There were also 5% of papers that used case reports and the same ratio used qualitative research (e. g. interviewing family physician about P4); quantitative research was quite rare with only one article (1%).

### *Main categories*

According to the distribution of the main approached topics, 16 articles (25%) were graded in the conceptual category “general considerations about P4 and overmedicalization process”. We classified many of the discussed aspects as: theoretical and conceptual aspects (levels of prevention, EBM, distinction between preventive approach and clinical care), medicalization drivers and consequences (conflict of interest, disease mongering, medicalization of risk factors, overdiagnosis and overtreatment) and protective factors (PHC attributes: longitudinality and role of gate-keeper; communication skills: doctor-patient relationship, patient-based medicine; Bioethics: non-maleficence and justice principles; and patient-safety strategies.

Only four articles (6%) were graded in “Bioethics”, approaching many topics, such as: the definition of health; the humanization of the relationship doctor-patient; the clinical context of uncertainty; political, economic and social determinants in health; autonomy, beneficence, non-maleficence and justice principles; intellectual freedom and responsibility; epistemology of medicine.

We classified 32 papers (49% of all) under the category “specific examples of overmedicalization and iatrogenesis”, including “screening”. 31% belong to “specific iatrogenic situations”, like chronic diseases and polypharmacy (e. g.: diabetes, genetic syndrome), birth, mental health and others (odontology, geriatrics, pathology) and 18% belong to the “screening” category (general aspects, cervical, breast and prostate cancers, genetic testing and neonatal screening).

And 13 articles (20%) were categorized as belonging to the subject “implementation and divulgation strategies of P4”: four of them developed the humanization of health assistance and the major role of the communication skills, such as the patient-centered medicine, to reduce iatrogenic events. The other articles discussed medical prescription, strategies to ensure patient's safety, applicability of “Choosing Wisely” campaigns, manifestoes in favour of P4 and

defending a medicine without conflict of interests, P4 teaching and search for P4 in electronic databases.

## Discussion

To understand the timeline of research output on P4, we need to remember the substantial publications of 2000 and 2002 about iatrogenesis, such as: “Is US health really the best in the world?”,<sup>36</sup> “The arrogance of preventive medicine”<sup>37</sup> and “Too much medicine?”.<sup>38</sup> Moreover, Gérvas had already published a paper in this respect in 1997, reporting the inefficiency of statin treatment in primary prevention of ischemic cardiopathy.<sup>39</sup> In this this context, the WICC accepted the definition of P4 in 2003, acknowledging its relevance and encouraging future publications. It is exactly the same year that Gérvas published the first article included in our review.<sup>40</sup> Afterwards, the scientific output increased slowly and progressively until its peak in 2015 which was probably linked to the disclosure, in February 2015: P4 would be one of the main subjects of the 21<sup>st</sup> WONCA (World Organization of Family Doctors) World Conference 2016 in Rio de Janeiro. Consequently, this reputed event encouraged many international authors to undergo research and publish on the subject of P4 (figure 3). Furthermore, the same year, Jamouille led a specific edition of the RBMFC about P4 and the Iberoamerican Family and Community Medicine Conference took place in Uruguay, also emphasized quaternary prevention. Those events led to a significant increase in the publications in English, together with a geographical expansion of the first authors countries. After 2015, the scientific output declined and came back to its baseline.

Until 2012, the first authors came from Spain, Portugal and Brazil, countries that have universal public health system. These systems, by definition, are very concerned about equity, which is one of the P4 main objectives; it can be achieved totally or partially, through the reduction of unnecessary medical procedures, which, indirectly, promotes equity and favors the access to health services. We also demonstrated that Latin-speaking countries first authors seem to be more acquainted with the concept and published more about it; therefore the same countries occupied the three first positions of the publications ranking. However, when we assessed the number of papers per capita, Uruguay was first. It is a very small country of 3,47 million people that lead the WONCA group on P4, which also has a universal public health system and borders the South of Brazil and is greatly influenced by its larger neighbour of 212 million of inhabitants.<sup>41</sup>

The results reported a clear predominance of Brazilian journals and we suggest at least six reasons to explain it. Firstly, the Brazilian authors, Norman and Tesser,<sup>42</sup> were precursors of the concept in Latin America since 2009, which promoted the submission of articles in Brazilian journals. Secondly, because Brazil is one of the biggest universal healthcare systems in the world and suffer from insufficient funding, It requires a cost-effective management that can be achieved, at least partly, with P4 implementation policies, tackling medical overuse. Thirdly, because Brazilian journals accept articles in many languages, drawing international authors. Fourthly, as we wrote above, P4 was one of the main subjects in the World WONCA Conference of Rio de Janeiro in 2016, and it encouraged the Brazilian journals and authors to

publish on this subject. Fifthly, the municipality of Rio de Janeiro, reformed completely and enhanced its primary health care (PHC) system in 2009-2012,<sup>43</sup> improving the access of the population to health services, emphasizing universality and equity principles. And sixthly, the Brazilian government launched a massive program in 2013 to strengthen its national PHC, the “More Doctors Program” (*Programa Mais Médicos*) based on the principles of universality and equity, engaging more than 18.000 national and foreign physicians, favouring the access of 63 million of Brazilians to the health system.<sup>44</sup>

Most of the papers were published in non-classified journals. However, a significant number (18 articles, 28%) also published in Q1 or Q2 journals. Some readers could deduct that the majority of the papers were published in low quality journals. However, according to Seglen (1997):

- Journals' impact factors are determined by technicalities unrelated to the scientific quality of their articles;
- Journal impact factors depend on the research field: high impact factors are likely in journals covering large areas of basic research with a rapidly expanding but short lived literature that use many references per article.<sup>45</sup>

In this regard, some determinant factors, like the language of publication, can strongly impact the number of citations and, consequently, affect the journal impact factor and the article's visibility. Therefore, if P4 is published in non-classified journals, it can be one more factor that impairs its disclosure and, in this way, its expansion. Another aspect can possibly explain the publications in low impact factor journals: publication costs. Thus, most of the selected studies have been realized in low or middle-income countries, without any sponsorship and probably could not afford high cost of publication in high impact factor journals. Nevertheless, according to the four publications in Q1 journals, we note four characteristics that could have promoted them in the best classified journals. First, the geographical situation promoted the publications of European authors in European Journals. Second, all of them used the English language that constitutes the main official language of the European Union. Third, we also observed that three of the four first authors were from Latin-speaking countries (France, Italy and Portugal) that confirmed the approximation between such countries and the concept of P4. And fourth, the economic aspect could have facilitated authors from high-income countries to pay the expensive publication costs of high impact journals.

Assessing the quality of the selected papers has been a quite difficult and sensitive task, especially because most of the articles were literature reviews and that type of study rarely uses a structured methodology; we confirm this trend in our study. In addition, while the papers frequently brought a broad conceptual understanding, including multidimensional analysis, they usually supported P4 thesis and seldom critically developed the potential weaknesses of the concept.

P4 is a complex concept based on historical, theoretical and scientific fundamentals that involves several areas of knowledge, suggesting the necessity of an exhaustive expertise in these areas (e. g.: politics, economics, sociology, law, public health management, history and epistemology of medicine, medical teaching, professional practice, EBM, bioethics, psychology,

communication skills, among others). The previous understanding of these basics and their interconnections is essential and enable the development of effective interventions to tackle unnecessary medical procedures and their iatrogenic effects. Therefore, the category “general considerations about P4 and overmedicalization process” is more than necessary to be developed. The overmedicalization process is described as an effect of biased studies, conflict of interest and disease mongering strategies, involving mainly the pharmaceutical industry but also some physicians.<sup>46</sup> In this regard, considerable advertising campaigns enhanced this phenomenon; it lead the population to a more concerned condition about their own health and raised the expectations according to the health care system, even when healthy. Concerning the definition of P4, the authors wrote it generally just after the first three levels of prevention of Leavell and Clark.

Quaternary prevention constitutes a counter-hegemonic concept, which approaches modern and occidental medicine being based on a critical and multidimensional perspective. Moreover, actual biomedical research, regardless of its funding,<sup>47, 48</sup> focuses on high-technology studies<sup>49</sup> and does not prioritize medical overuse issues. Therefore, we understand why the vast majority of the included papers were bibliographic research and low cost studies. Bibliographic studies were also necessary to create the robust theoretic base of such a recent and interdisciplinary concept. Qualitative studies are essential to promote the identification and analysis of key elements involved in the medical overuse process, for example regarding the perception and practice of the health actors. In this way, these points are precious because they can lead to the elaboration of effective strategies, in order to achieve significant changes in health practice and impact medical overuse. However, only a small number of qualitative studies were found, probably because they were very time-consuming and of high cost. Quantitative research was also very scarce, probably because it is generally used to evidence the efficacy of a medical intervention and rarely to establish its harm. However, the only quantitative included study<sup>50</sup> highlighted significant data about overscreening and underscreening for cervical cancer in a specific population. This type of study is brief, low-cost and very relevant, because able to reveal precise quantitative estimations about overmedicalization procedures, including the expenditure that can be saved if a P4 approach would be implemented. Case-report studies can also be used to improve theoretical aspects on P4 but have less potential to contribute to elaborate concrete strategy to impact overmedicalization because they stem from the analysis of a singular and practical example.

In a context where medicine is inducing much medical overuse, leading to an outbreak of iatrogenic events<sup>51</sup> and is, by definition, contrary to the principles of medicine; the papers approached this issue through an ethical perspective. This point represents the heart of the P4 problematic and implies to consider the risk and benefits of each clinical decision, integrating a constant critical reflection on beneficence and non-maleficence principle into health practice. A broad theoretical knowledge, accurate competence to assess critically the available evidences and a global understanding of each clinical case are required to apply such type of practice. The second point focuses on the autonomy principle, since the patient always takes part in the clinical decision-making. In some situations, its role can be more significant, especially in case of shared decision. Besides, plenty of other factors are also able to affect the patient's opinion, for example: an advertisement which minimizes the risks of a medical intervention (e. g. PSA

screening<sup>52</sup>) and does not mention the prevalence of severe iatrogenic damages. This situation illustrates well a possibility of bias in the free and informed patient decision-making concerning a single screening test. In this case, the role of P4 is to identify such situations and be able to build strategies to face or prevent them.

Last but not least, P4 aims to tackle medical overuse, saving funding and allowing some necessary medical interventions which could not be realized because of insufficient funding. This last example highlights the role of P4 advocating the justice principle. In this category, the authors considered some conceptual aspects as: knowledge (epistemology) and ethics (non-maleficence and beneficence principles); analyzed specific bioethical challenges in the case of “incidentalomas”;<sup>53</sup> outlined schemes, remembering the importance of cross-sectoral engagement, including political, economic and social dimensions, to recover the principles of a human-centered medicine,<sup>54</sup> others wrote also about the practitioners’ responsibility.<sup>55</sup> And one of the few criticisms found on the concept, suggested that P4 “needs to be theoretically clarified and more widespread among - and dialogued with - health professionals of various specialties”<sup>56</sup>, and cannot be a medical practice only, but has to become an interdisciplinary practice.

The papers classified in the “specific iatrogenic examples” category addressed some specific population groups, for example pregnant women or mental health users. The authors generally criticized the medicalization of healthy people or physiological situations, emphasizing the frequent iatrogenic results. We develop briefly here the examples of pregnancy and mental health. Pregnancy is a physiologic stage of life, however, health professionals usually handled it as a disease, especially the labour. Souza and Pileggi-Castro (2014) emphasized that the:

Use of health technologies favors reduction in maternal morbidity and mortality, but hyper-medicalization – or the excessive and unnecessary use of health technology in care during pregnancy and childbirth – also represent risks for women, fetuses and newborns.<sup>57</sup>

Tesser and Norman (2015) reported the excess of episiotomy, cesarean section and routine use of oxytocine in Brasil. They also suggest PHC-centered P4 strategies, as the: elaboration of labor plans (personalized care itinerary), inclusion of trained family physicians to assist low-risk labour/childbirth and empowerment of the social movements which fight for “humanizing” it.<sup>58</sup> In this regard, mental health represents a knowledge area whose epistemology, practice and results are frequently questioned.<sup>59</sup> The selected papers of this topic point in the same direction: criticizing the treatment of healthy people, as well as the biological reductionist theories that are used to explain mental health problems, ruling out key elements like the path of life and other social dimensions. Another interesting example concerns the elderly people group. For multiple reasons, it constitutes a vulnerable group to iatrogenesis, because of usual: polymorbidity and polypharmacy, lower rate of metabolism (e. g.: renal clearance), cerebral and other functions (slower reflexion and understanding, difficulty of locomotion, hearing loss, etc.), shorter life expectancy, among others. In all these specific groups or situations, one frequent outlined strategy is “not to do”, using a “wait and see” approach, and sometimes even “desprescribing” suppressing prescriptions of medications and heavy treatments that are more harmful than beneficial to health. And to do so, the professionals need to have a good knowledge on EBM, so

they can critically approach the available evidence, and adapt it to every individual situations. The other articles about odontology, pathology, chronic diseases developed similar reflections, reporting discrepancies between the excess of medical procedures, the cost-effectiveness and the non-maleficence principle.

Screening belongs to the second level of prevention and aims to detect asymptomatic diseases and treat them early. Consequently, screening procedures have to be realized, in the cases that the risks/benefits balance is unequivocally favorable to the patient. This differs completely from curative situations, where the patient has a disease and suffers; which allows a thinner margin between risks and benefits, because if we do not act, the clinical outcomes are much worse than if we do.<sup>60</sup> Therefore, research on the harm of screening has to be enhanced and, maybe, connected with other type of iatrogenesis that can affect healthy people like primary prevention.

We will discuss further now the publications focused on the category “implementation and divulgation strategies of P4”. The authors proposed strategies intrinsically related to the drivers of medical overuse focusing on the main actors: physicians, patients, pharmaceutical industry, politicians as well as the means of communication between them. Improving the communication abilities of the physicians, for example, through a patient-centered approach<sup>61</sup> was one of the most P4 suggested strategies. This approach involves: empathy, the construction of a trustworthy therapeutic relationship, exploring the illness, identifying the patient’s stress-factors and shared decision-making. All these communication skills evidenced effectiveness to improve mental health, reducing the overuse of health system and improving most of the symptoms. In this way, we understand that patient-centered medicine is one key element that could tackle medical overuse and reduce unnecessary medical procedures. Another significant and complementary perspective focuses particularly on the clinicians and invites them to enhance their critical skills to assess clinical guidelines, encouraging them, when possible, to use a watchful waiting approach and consequently avoiding overmedicalization. One of the selected papers<sup>62</sup> mentioned the “Choosing Wisely” (CW) campaign, launched by the American Board of Internal Medicine in 2012<sup>63</sup> that created and disclosed a list of five unnecessary medical procedures which had to be avoided and encouraged other medical societies to do the same. Moreover, this type of CW strategy showed evidences of cost-effectiveness in multiple studies.<sup>64</sup> Another suggested and pertinent strategy was the claim to reach a medicine without conflict of interests that biases most scientific studies and impacts strongly the guidelines. Consequently, it highly affects the clinical decision-making. To conclude this topic, we highlight an important aspect affecting the divulgation of the concept of P4. It is currently not well represented in the online search databases; and one of main causes is probably the fact that most of the databases does not classify it as a MeSH, or as a relevant keyword. Therefore, in order to boost significantly the promotion of P4 scientific identification and production, P4 needs to become a MeSH in the electronic databases.

#### Strengths and weaknesses of the study

As we report in the introduction, the present study is the first bibliometric analysis and Descriptive Content Analysis on P4. While some bibliometric studies about adjacent topics like medicalization,<sup>65</sup> overdiagnosis<sup>66</sup> or “unnecessary procedures”<sup>67</sup> does already exist, their



objectives focused on some very specific clinical situations, such as: pregnancy or mammography screening. The last one focused on the impact of the Cochrane Reviews on clinical decision. For these reasons, we chose not to make any comparison between these publications and our study.

In addition, we employed a systematic methodology and used internationally acknowledged databases with minimal restriction criteria. Such sensitive approach positively supports the reliability of our results. However, in spite of our wide search methods, several key articles escaped our study and were not selected; some of them, probably because of the non-inclusion of P4 in the MeSH or keywords in the major databases, such as: one of Jamouille (1986)<sup>68</sup> and two articles of the renowned Barbara Starfield, Gérvas and Heath.<sup>69, 70</sup> Although these papers could have helped to promote the concept of P4, we chose not to include them for two reasons. First, because we would to highlight the existing impairment of P4 search in the selected databases. And second, their inclusion could invalidate our search method. Moreover, according to language criteria, another four articles had to be excluded because they were published in Serbo-Croat or Korean.

Furthermore, the selected papers had an heterogeneous number of authors and with a concern on a strong methodology, we decided to consider only the first authors data to perform the statistical results. This decision probably resulted in an overall reduction of the authors nationalities.

We knew that the journal ranking did not represent the most significant factor to assess the publications and Nature published about it in 2016:

It effectively undervalues papers in disciplines that are slow-burning or have lower characteristic citation rates. Being an arithmetic mean, it gives disproportionate significance to a few very highly cited papers, and it falsely implies that papers with only a few citations are relatively unimportant.<sup>71</sup>

However, we decided to include the journal impact factor in our bibliometric analysis, because it is one of the few tools that can assess and, maybe, roughly explain the visibility – and not the quality – of some scientific studies in the world. Consequently, it could also help to understand and highlight journals and countries inequalities according to research output.

According to the method used in the selected articles, most of them were bibliographic studies. On the one hand, these types of studies provide interesting informations with conceptual or historical aspects that help to establish a robust background. But on the other hand, the studies were frequently non-systematized and aimed to support the concept without seeking a critical approach, thus weakening the content of those articles.

Several articles approached more than one category making it difficult to classify them under a single one. And with a concern on a global view on research output, we decided to classify each included paper under only one category according to the main article's objective. So, as a qualitative classification, any bias according to the coder subjective decision could occur.

Implication for practice

Quaternary Prevention represents a relevant concept for public health because it promotes the identification and prevention of potential iatrogenic situations and, consequently, can reduce overall iatrogenic mortality and health costs. In this respect, the authors suggest concrete implementation policies through: cross-sectoral collaboration (political, legal, social, educational and sanitary, among others), P4 teaching in the health graduation and post-graduation programs, improving the critical approach of clinical studies and guidelines, practical changes in the medical prescription and appropriate use of the communication skills by the physicians.

<b>WHAT IS ALREADY KNOWN ON THIS TOPIC</b>
<ul style="list-style-type: none"><li>• The concept of Quaternary Prevention (P4) has been designed initially at the end of the 1980s, when the Belgian family physician Jamouille revised the prevention levels of Leavell and Clark according to the patient's perception (illness) and the physician's assessment (disease), integrating a public health approach into individual clinical practice.</li><li>• P4 is a recent theoretical concept for public health because it aims to tackle the outspread of unnecessary medical procedures and iatrogenic events; and consequently, can be able to lower global cost expenditures.</li></ul>
<b>WHAT THIS STUDY ADDS</b>
<ul style="list-style-type: none"><li>• Conceptual, geographical and linguistic aspects impacted the scientific production on P4, as well as the WONCA conferences and the model of healthcare systems in the authors' country.</li><li>• The Brazilian were the first authors who most published (absolute number of publications) and the Uruguayan lead the production per capita.</li><li>• Most of the papers used a bibliographic research method and focused on the causes and examples of overmedicalization, just as strategies to implement P4.</li><li>• The quality and quantity of available studies is still limited and further investigations are recommended to assess the effective impact of P4 on public health.</li></ul>

#### Implications for research

As we see above, Quaternary Prevention constitutes a critical approach on modern medicine and exposes most of its limitations. In this regard, P4 represents a counter-hegemonic concept and, consequently, research on it is clearly underfunded. While we observed a significant increase in the scientific production in 2015, the publications decreased again after this "golden year". In this context, we consider that research to further the categories developed in this study is required. According to our analysis, three purposes have to be prioritized: the detection of overmedicalization situations (bibliographic investigative research), the identification of factors that lead to these situations and the development of P4 strategies (bibliographic research and qualitative studies) and testing the developed P4 strategies (quantitative studies).

For practical reasons and to provide effective results, we suggest that future research should focus on:

- Quantitative studies on situations of medical overuse, evaluating the iatrogenic outcomes of medical intervention (e.g.: primary and secondary prevention interventions) on previously healthy people;
- Studies assessing the impact of P4 strategies (e.g.: communication skills teaching, critical approach on EBM, disclosure of the physician's conflict of interests) on clinical outcomes (mortality, iatrogenic events, cost-effectiveness, patient and physician's perceptions), and;
- Studies and interventions of health promotion (e.g.: physical activity, nutritional, stress management, social inclusion measures) and/or health education (e.g.: population education about P4), without medical interventions and analyzing the outcomes (e.g.: mortality, cost-effectiveness, patient and physician's perceptions);

In addition, the total number of articles remains low if compared to other search terms like "medical overuse" which have up to 100 times more results than "quaternary prevention" in Pubmed. We understand that a linguistic and conceptual gap could explain this consistent difference and consider that some strategies could be able to transcend it, involving linguistic, conceptual, information and communication technologies dimensions, respectively:

- To insist on the English language to promote P4 publications worldwide;
- To disclose the concept and clarify possible misunderstanding regarding the initial and actual acknowledged definition of the concept;
- To come closer to other adjacent and more "famous" concepts, such as "medical overuse", "overdiagnosis" or even "overtreatment" and put them in the abstracts and as keywords;
- To publish more often in Q1 or Q2 reviews and, when possible, addressing quantitative studies that support P4 (e. g.: cost-effectiveness and mortality reduction);
- To claim the inclusion of P4 as a MeSH in the international databases.

## **Conclusion**

The present bibliometric and content analysis highlighted key elements that affected the research output on P4, such as: conceptual, geographical, linguistic, WONCA conferences and model of healthcare systems in the authors' country. Most of the included papers used a bibliographic research method and approached quite equally the causes, the examples of overmedicalization and the implementation strategies. Quaternary prevention represents an ethical combat towards fairness and equity of access to health services. The concept and its respective scientific output are relevant for public health, however we consider that there is still a limited quality and quantity of available studies on the topic and further studies are recommended, especially in order to assess its effectiveness to prevent the outbreak of iatrogenic events and to lower global health expenditure.

- <sup>1</sup> Rosen G. **Uma história da saúde pública**. São Paulo: Hucitec, editora da Universidade Estadual Paulista; Rio de Janeiro: Associação Brasileira de Pós-Graduação em Saúde Coletiva; 1994.
- <sup>2</sup> Leavell HD, Clark EG, editors. **Preventive medicine for the doctor in his community: an epidemiologic approach**. 3rd ed. New York: McGraw-Hill;1960:20-21.
- <sup>3</sup> Arouca SAS. **O dilema preventivista: contribuição para a compreensão e crítica da medicina preventiva** [dissertation on the Internet]. Campinas: Universidade Estadual de Campinas; 1975. [cited 2019 Sep 14]. Available from: <https://teses.icict.fiocruz.br/pdf/aroucaass.pdf>.
- <sup>4</sup> Jamouille M. **Information et informatisation en médecine générale**. In: Les informagiciens. Namur: Presses Universitaires de Namur;1986:193-209.
- <sup>5</sup> Drabkin IE. Medical education in Ancient Greece and Rome. *J Med Edu*. 1957;32:286-296.
- <sup>6</sup> Magner LN. **A History of Medicine**. Boca Raton: Taylor & Francis Group; 2005.
- <sup>7</sup> Bliss M. **The Discovery of Insulin**. The Canadian Encyclopedia, Toronto [Internet]. 2015 August 19 [cited 2019 Sep 15]; Available from: <https://www.thecanadianencyclopedia.ca/en/article/the-discovery-of-insulin>.
- <sup>8</sup> Cochrane AL. **Sickness in Salonica: my first, worst, and most successful clinical trial**. *Br Med J (Clin Res Ed)*. 1984;289(6460):1726–1727.
- <sup>9</sup> Illich I. Medical nemesis. **The expropriation of health**. New York: Pantheon Books; 1976.
- <sup>10</sup> Jamouille M. **Information et informatisation en médecine générale**. In: Les informagiciens. Namur: Presses Universitaires de Namur;1986:193-209.
- <sup>11</sup> Leavell HD, Clark EG, editors. **Preventive medicine for the doctor in his community: an epidemiologic approach**. 3rd ed. New York: McGraw-Hill;1960:20-21.
- <sup>12</sup> Kleinman A. **The illness narratives: suffering, healing & the human condition**. New York: Basic Books, 1988.
- <sup>13</sup> Gérvas J, Pérez Fernández MP. **Genética y prevención cuaternaria. El ejemplo de la hemochromatosis**. *Aten Primaria* 2003;32(3):158-62.
- <sup>14</sup> Jamouille M. Prevenção quaternária: primeiro não causar dano. *Rev Bras Med Fam Comum*. 2015;10(35):1-3. Available from: [https://doi.org/10.5712/rbmfc10\(35\)1064](https://doi.org/10.5712/rbmfc10(35)1064).
- <sup>15</sup> Starfield B. **Is US health really the best in the world?** *JAMA*. 2000;284:483-5.
- <sup>16</sup> Moynihan, R, Cassels A. **Selling sickness: How the World's Biggest Pharmaceutical Companies Are Turning Us All into Patients**. Vancouver: Greystone Books; 2005.
- <sup>17</sup> Norman AH, Tesser CD. **Prevenção quaternária na atenção primária à saúde: uma necessidade do Sistema Único de Saúde**. *Cad. Saúde Pública*, Rio de Janeiro [Internet]. 2009 Sep [cited 2019 Sep 28];25(9):2012-2020. Available from: [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0102-311X2009000900015&lng=en&nrm=iso](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0102-311X2009000900015&lng=en&nrm=iso).
- <sup>18</sup> Gérvas J, Pérez Fernández MP. **São e salvo: e livre de intervenções médicas desnecessárias** [electronic version] / Juan Gérvas, Mercedes Pérez Fernández; tradução: Marcela Ceci Dohms, Gustavo de Araújo Porto Landsberg ; revisão técnica: Gustavo Diniz Ferreira Gusso. Porto Alegre: Artmed; 2016.
- <sup>19</sup> Gøtzsche PC. **Deadly Medicines and Organised Crime: How Big Pharma Has Corrupted Healthcare**. 1. ed. London: Radcliffe Publishing Ltd; 2013.
- <sup>20</sup> WONCA International Classification Committee. **WONCA International Dictionary for General/Family Practice**. Niels Bentzen. 2003:110-115. Available from: <http://www.ph3c.org/PH3C/docs/27/000092/0000052.pdf>
- <sup>21</sup> World Health Organization. **The World Health Report. Health Systems Financing: The Path to Universal Coverage**. WHO Library Cataloguing-in-Publication Data. 2010. Available from:

[https://apps.who.int/iris/bitstream/handle/10665/44371/9789241564021\\_eng.pdf;jsessionid=D6EBB74DE86879B4E635B8390689BA06?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/44371/9789241564021_eng.pdf;jsessionid=D6EBB74DE86879B4E635B8390689BA06?sequence=1).

<sup>22</sup> World Health Organization. **Global Report. New Perspectives on Global Health Spending for Universal Health Coverage.** WHO Library Cataloguing-in-Publication Data. 2018. Available from:

<https://apps.who.int/iris/bitstream/handle/10665/259632/WHO-HIS-HGF-HFWWorkingPaper-17.10-eng.pdf;jsessionid=50C803BE192E622506C80967364023B0?sequence=1>.

<sup>23</sup> Narin F, Olivastro D, Stevens K. **Bibliometrics theory, practice and problems.** Eval Rev. 1994;18:65–76.

<sup>24</sup> Pendlebury DA. **White Paper using Bibliometrics in Evaluating Research.** Thomson Reuters. 2010. Available from: <http://thomsonreuters.com/web-of-science/>.

<sup>25</sup> Marcellus L. **Bibliometric and Textual Analysis of Historical Patterns in Maternal-Infant Health and Nursing Issues in The Canadian Nurse Journal, 1905-2015.** Can J Nurs Res. 2018 Dec 21:844562118804119. doi: 10.1177/0844562118804119.

<sup>26</sup> Rasmussen K, Jørgensen KJ, Gøtzsche PC. **Citations of scientific results and conflicts of interest: the case of mammography screening.** BMJ Evidence-Based Medicine 2013;18:83-89.

<sup>27</sup> Bunn F, Trivedi D, Alderson P, Hamilton L, Martin A, Pinkney E, et al. **The impact of Cochrane Reviews: a mixed method evaluation of outputs from Cochrane review groups supported by the NIHR.** Health Technol Assess 2015;19(28):1-99, v-vi.

<sup>28</sup> Bunn F, Trivedi D, Alderson P, Hamilton L, Martin A, Iliffe S. **The impact of Cochrane Reviews: a mixed method evaluation of outputs from Cochrane review groups supported by the NIHR.** Syst Rev. 2014 Oct 27;3:125.

<sup>29</sup> Higgins JPT, Green S (editors). **Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0.** The Cochrane Collaboration, 2011. Available from: [www.cochrane-handbook.org](http://www.cochrane-handbook.org).

<sup>30</sup> Noyes J, Popay J, Pearson A, Hannes K, Booth A. **Chapter 20: Qualitative research and Cochrane reviews.** In: Higgins JPT, Green S (editors), Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0. The Cochrane Collaboration, 2011. Available from [www.handbook.cochrane.org](http://www.handbook.cochrane.org).

<sup>31</sup> O'Brien BC, Harris IB, Beckman TJ, Reed DA, Cook DA. **Standards for Reporting Qualitative Research: A Synthesis of Recommendations.** Academic Medicine. September 2014;89(9):1245-1251.

<sup>32</sup> Coordenação de Aperfeiçoamento de Pessoal de Nível Superior [Portal]. [cited 2019 Sep 10]. Available from: [www.periodicos.capes.gov.br](http://www.periodicos.capes.gov.br).

<sup>33</sup> SCImago, (n.d.). SJR — SCImago Journal & Country Rank [Portal]. [cited 2019 Sep 10]. Available from: <https://www.scimagojr.com/journalrank.php>.

<sup>34</sup> Plataforma Sucupira [Internet]. Qualis Periódicos [cited 2019 Aug 17]. Available from: <https://sucupira.capes.gov.br/sucupira/public/consultas/coleta/veiculoPublicacaoQualis/listaConsultaGeralPeriodicos.jsf>.

<sup>35</sup> United Nations Population Division [Internet]. Probabilistic Projections [cited 2019 Aug 14]. Available from:

[https://population.un.org/wpp/DVD/Files/2\\_Indicators%20\(Probabilistic%20Projections\)/UN\\_PP\\_P2017\\_Output\\_PopTot.xls](https://population.un.org/wpp/DVD/Files/2_Indicators%20(Probabilistic%20Projections)/UN_PP_P2017_Output_PopTot.xls).


<sup>36</sup> Starfield B. **Is US health really the best in the world?** JAMA. 2000;284:483-5.

<sup>37</sup> Sackett DL. **The arrogance of preventive medicine.** CMAJ. 2002;167(4):363–364.

<sup>38</sup> Moynihan R., Smith R. **Too much medicine?** BMJ (Clinical research ed.), 2002;324(7342):859–860.


- <sup>39</sup> Gérvas J, Pérez Fernández M. **Las hiperlipemias y la prevención primaria de la cardiopatía isquémica.** Med Clin (Barc) 1997; 109: 549-552.
- <sup>40</sup> Gérvas J, Pérez Fernández MP. Genética y prevención cuaternaria. El ejemplo de la hemochromatosis. Aten Primaria 2003;32(3):158-62.
- <sup>41</sup> World Population Review. Available from: <http://worldpopulationreview.com/> . 2019 [Cited in 2019 Sep 16]
- <sup>42</sup> Norman AH, Tesser CD. **Prevenção quaternária na atenção primária à saúde: uma necessidade do Sistema Único de Saúde.** Cad. Saúde Pública, Rio de Janeiro [Internet]. 2009 Sep [cited 2019 Aug 19];25(9):2012-2020. Available from: [http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0102-311X2009000900015&lng=en&nrm=iso](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0102-311X2009000900015&lng=en&nrm=iso).
- <sup>43</sup> Organização Pan-Americana da Saúde. **Reforma da Atenção Primária à Saúde na cidade do Rio de Janeiro – avaliação dos três anos de Clínicas da Família. Pesquisa avaliativa sobre aspectos de implantação, estrutura, processo e resultados das Clínicas da Família na cidade do Rio de Janeiro.** Porto Alegre, RS: OPAS; 2013.
- <sup>44</sup> Brazil. Ministério da Saúde. Secretaria de Gestão do Trabalho e da Educação na Saúde: Departamento de Planejamento e Regulação da Provisão de Profissionais de Saúde. Programa Mais Médicos [recurso eletrônico] / Ministério da Saúde, Secretaria de Gestão do Trabalho e da Educação na Saúde – Brasília: Ministério da Saúde; 2017.
- <sup>45</sup> Seglen PO. **Why the impact factor of journals should not be used for evaluating research.** BMJ. 1997 Feb 15; 314(7079): 498–502.
- <sup>46</sup> Gøtzsche PC. **Deadly Medicines and Organised Crime: How Big Pharma Has Corrupted Healthcare.** 1. ed. London: Radcliffe Publishing Ltd; 2013.
- <sup>47</sup> Dorsey ER, de Roulet J, Thompson JP, et al. **Funding of US Biomedical Research, 2003-2008.** JAMA. 2010;303(2):137–143. doi: 10.1001/jama.2009.1987.
- <sup>48</sup> "2 Sources of Funding for Biomedical Research." Institute of Medicine. 2004. Strategies to Leverage Research Funding: Guiding DOD's Peer Reviewed Medical Research Programs. Washington, DC: The National Academies Press. doi: 10.17226/11089.
- <sup>49</sup> Owens B. **Mapping biomedical research in the USA.** The Lancet. 2014 Jul 5;384(9937):11–4.
- <sup>50</sup> Cardozo Rufo V, Fernández Saavedra ML. **Tamizaje de câncer de cuello de útero en usuarias de la Policlínica Villa Ferré (Uruguay). Una mirada desde la prevención cuaternaria.** Archiv med fam gen (En línea). 2017 Nov;14(2):7–13.
- <sup>51</sup> Starfield B. **Is US health really the best in the world?** JAMA. 2000; 284: 483-5.
- <sup>52</sup> Modesto AAD, Lima RLB de, D'Angelis AC, Augusto DK. **Um novembro não tão azul: debatendo rastreamento de câncer de próstata e saúde do homem.** Interface (Botucatu, Online). 2018 Mar;22(64):251–62.
- <sup>53</sup> Mariño MA. **Incidentalomas: concept, relevance and challenges for medical practice.** Rev bras med fam comunidade. 2015 Jun;10(35):1–9.
- <sup>54</sup> La Valle R. **Sobre la definición de salud.** Archiv med fam gen (En línea). 2017 Nov;14(2):27–31.
- <sup>55</sup> Widmer D. **Philosophical roots of Quaternary Prevention.** Rev bras med fam comunidade. 2015 Jun;10(35):1–8.
- <sup>56</sup> Oliveira CC, Reis A. **Questões epistemológicas e bioéticas da prevenção quaternária.** Physis (Rio J). 2012;22(4):1485.
- <sup>57</sup> Souza JP, Pileggi-Castro C. **On labor and childbirth: the importance of quaternary prevention.** Cad Saude Publica. 2014 Aug;30 Suppl 1:S1-3.


- <sup>58</sup> Tesser CD, Knobel R, Andrezzo HF de A, Diniz SG. **Violência obstétrica e prevenção quaternária: o que é e o que fazer.** Rev bras med fam comunidade. 2015 jun;10(35). Available from: <https://www.rbmf.org.br/rbmfc/article/view/1013/716>.
- <sup>59</sup> Whitaker R. **Anatomy of an Epidemic.** New York: Crown Publishers. 2010.
- <sup>60</sup> Gérvas J, Pérez Fernández M. **Limits to the power of medicine to define disease and risk factor, and quaternary prevention.** Gac Sanit. 2006 Dec;20 Suppl 3:66–71.
- <sup>61</sup> Stewart M et al. **Patient-centered medicine: transforming the clinical method.** California: Sage Publications, Inc; 1995.
- <sup>62</sup> Cuba Fuentes MS, Zegarra Zamalloa CO, Reichert S, Gill D. **Attitudes, perceptions and awareness concerning quaternary prevention among family doctors working in the Social Security System, Peru: a cross-sectional descriptive study.** Medwave. 2016 Apr 27;16(3):e6433.
- <sup>63</sup> Choosing Wisely Organization. History [Internet]. Philadelphia: Choosing Wisely Organization; [Cited in 2019 Sep 26] Available from: <https://www.choosingwisely.org/our-mission/history/>.
- <sup>64</sup> Choosing Wisely Organization. History [Internet]. Philadelphia: Choosing Wisely Organization; [Cited in 2019 Sep 10] Available from: <https://www.choosingwisely.org/getting-started/resource-library/does-choosing-wisely-work/>
- <sup>65</sup> Marcellus L. **Bibliometric and Textual Analysis of Historical Patterns in Maternal-Infant Health and Nursing Issues in The Canadian Nurse Journal, 1905-2015.** Can J Nurs Res. 2018 Dec 21:844562118804119. doi: 10.1177/0844562118804119.
- <sup>66</sup> Rasmussen K, Jørgensen KJ, Gøtzsche PC. **Citations of scientific results and conflicts of interest: the case of mammography screening.** BMJ Evidence-Based Medicine 2013;18:83-89.
- <sup>67</sup> Bunn F, Trivedi D, Alderson P, Hamilton L, Martin A, Pinkney E, et al. **The impact of Cochrane Reviews: a mixed method evaluation of outputs from Cochrane review groups supported by the NIHR.** Health Technol Assess 2015;19(28)
- <sup>68</sup> Jamouille M. **Information et informatisation en médecine générale.** In: Les informagiciens. Namur: Presses Universitaires de Namur;1986:193-209.
- <sup>69</sup> Starfield B, Hyde J, Gérvas J, et al. **The concept of prevention: a good idea gone astray?** Journal of Epidemiology & Community Health 2008;62:580-583.
- <sup>70</sup> Gérvas J, Starfield B, Heath I. **Is clinical prevention better than cure?** The Lancet. 2008 Dec 6;372(9654):1997–9.
- <sup>71</sup> [No authors listed]. **Time to remodel the journal impact factor.** Nature. 2016 Jul 28;535(7613):466. doi: 10.1038/535466a.





## Articles per Year of Publication (n=65)





# Quaternary Prevention (P4)

## PHYSICIAN'S ASSESSMENT

		Disease	
		ABSENT	PRESENT
PATIENT'S FEELING	Illness ABSENT	<p><b>P4 in Primary Prevention:</b></p> <ul style="list-style-type: none"><li>Prevent adverse effects (e.g.: statins, vaccines).</li></ul> <p><b>P4 in Secondary Prevention (majority of healthy asymptomatic people):</b></p> <ul style="list-style-type: none"><li>Prevent false-positive tests (e.g.: breast cancer, prostate cancer).</li><li>Prevent unnecessary exams (e.g.: too frequent cervical cancer screening)</li></ul>	<p><b>P4 in Secondary Prevention (minority of diseased patient):</b></p> <ul style="list-style-type: none"><li>Prevent false-negative, overdiagnosis and overtreatment (e.g.: breast cancer, prostate cancer).</li><li>Prevent unnecessary exams before surgery in patient well diagnosed but without other comorbidity (XR, ECG).</li></ul>
	PRESENT	<p><b>P4 in Quaternary Prevention (initial definition)</b></p> <ul style="list-style-type: none"><li>Exclude red flags;</li><li>Prevent prescription of unnecessary medications if the patient doesn't fill the diagnostic criteria (e.g.: antidepressives, anxiolytics);</li><li>Prevent the requests of unnecessary tests in patient without red flag (e.g.: brain or spine MRI/CT).</li></ul>	<p><b>P4 in Tertiary Prevention</b></p> <ul style="list-style-type: none"><li><b>Prevent excessive treatment (e.g.: in incurable disease);</b></li><li><b>Prevent adverse effects attributable to the interactions between comorbidities, polypharmacy, unnecessary treatments and exams;</b></li></ul>

P4