

Epidemiological Risk Factors Associated with Death and Severe Disease in Patients Suffering From COVID-19: A Comprehensive Systematic Review and Meta-analysis

Kunchok Dorjee, MBBS, PhD, MPH¹ and Hyunju Kim, PhD, MPH²

¹School of Medicine Division of Infectious Diseases, Center for TB Research, Johns Hopkins University, Baltimore, USA and

²Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, USA

Running head: *COVID-19 Disease Progression: A Meta-analysis*

Keywords: coronavirus disease-19, COVID-19, severe acute respiratory syndrome coronavirus-2, SARS-CoV-2

Word Count: Abstract: 248; Text: 3418

Correspondence to:

Kunchok Dorjee, MBBS, PhD, MPH
Johns Hopkins University School of Medicine
Division of Infectious Diseases
Center for Tuberculosis Research
1503 E Jefferson St, Room 114
Baltimore, MD 21205
Ph: 410-502-7135
kdorjee1@jhmi.edu

Abstract

Introduction: Progression of COVID-19 to severe disease and death is insufficiently understood.

Objective: Summarize the prevalence adverse outcomes, risk factors, and association of risk factors with adverse outcomes in COVID-19 patients.

Methods: We searched Medline, Embase and Web of Science for case-series and observational studies of hospitalized COVID-19 patients through May 22, 2020. Data were analyzed by fixed-effects meta-analysis, using Shore's adjusted confidence intervals to address heterogeneity.

Results: Forty-four studies comprising 20594 hospitalized patients met inclusion criteria; 12591 from the US-Europe and 7885 from China. Pooled prevalence of death [% (95% CI)] was 18% (15-22%). Of those that died, 76% were aged ≥ 60 years, 68% were males, and 63%, 38%, and 29% had hypertension, diabetes and heart disease, respectively. The case fatality risk [% (95% CI)] were 62% (48-78) for heart disease, 51% (36-71) for COPD, and 42% (34-50) for age ≥ 60 years and 49% (33-71) for chronic kidney disease (CKD). Summary relative risk (sRR) of death were higher for age ≥ 60 years [sRR=3.8; 95% CI: 2.9-4.8; n=12 studies], males [1.3; 1.2-1.5; 17], smoking history [1.9; 1.1-3.3; n=6], COPD [2.0; 1.6-2.4; n=9], hypertension [1.8; 1.7-2.0; n=14], diabetes [1.5; 1.4-1.7; n=16], heart disease [2.0; 1.7-2.4; 16] and CKD [2.0; 1.3-3.1; 8]. The

overall prevalence of hypertension (55%), diabetes (31%) and heart disease (16%) among COVID-19 patients in the US were substantially higher than the general US population.

Conclusions: Public health screening for COVID-19 can be prioritized based on risk-groups. A higher prevalence of cardiovascular risk factors in COVID-19 patients can suggest increased risk of SARS-CoV-2 acquisition in the population.

Introduction: Coronavirus disease-19 (COVID-19) caused by severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) that first emerged in Wuhan, China in late December 2019 has spread with such rapidity and efficiency that in less than six months, it has caused more than 8.5 million cases and 450,000 deaths globally.¹ Driven by an urgency to solve the crisis, studies are being published at an unprecedented pace. However, across the publications, prevalence of death, severe disease and their association with epidemiological risk factors have greatly varied^{2,3}, with studies showing conflicting results for association of key risk factors such as sex,⁴⁻⁸ smoking,⁹⁻¹² hypertension^{5,6,8,13,14} and diabetes^{5,6,8,13,14} with COVID-19 disease severity and death. Whether or how cardiovascular risk factors, especially prior hypertension, diabetes and heart disease are associated with acquisition of SARS-CoV-2 and progression to severe disease or death is being debated and not understood well.¹⁵⁻¹⁷ Meta-analyses conducted so far on prevalence of epidemiological risk factors and association with disease progression were mostly based on studies from China^{16,18,19} and many of the analyses on prevalence estimates included studies focused on critically ill patients,^{9,18,20,21} which can overestimate the prevalence and affect generalizability of results. To our knowledge, none of the analyses were restricted to hospitalized COVID-19 patients that indeed provides an efficient sampling frame to investigate disease progression in relation to risk factors.

Therefore, we undertook a comprehensive systematic review and meta-analysis to investigate the association between key epidemiological factors—age, gender, smoking, hypertension, diabetes, heart disease, chronic obstructive pulmonary disease (COPD), and chronic kidney

disease (CKD)—and progression to death in patients hospitalized due to COVID-19. We additionally compared the 1) the prevalence of risk factors and death in the US-Europe with that of China; 2) the prevalence of co-morbidities at baseline with the general population prevalence, and 3) prevalence of cardiovascular disease, chronic kidney disease and COPD at baseline with corresponding organ injuries (acute cardiac injury, acute lung injury, and acute kidney injury) during hospital admission.

Methods:

Literature search, study selection and data abstraction

We searched Medline, Embase, Web of Science and the WHO COVID-19 database to identify studies published through May 22, 2020 that investigated the risk of severe disease or death in hospitalized patients with confirmed COVID-19 disease. We used search terms, ‘coronavirus disease 19’, ‘COVID-19’, ‘severe acute respiratory syndrome coronavirus 2’ and ‘SARS-CoV-2’. We started the search on March 18, 2020 with biweekly search thereafter and final search on May 22, 2020. We included case series and observational studies that described the prevalence of death or severe disease in adult population stratified by risk factors: age, sex, hypertension, diabetes, heart disease, COPD, CKD and CLD. We excluded studies that exclusively focused on pregnant women, children, and elderly patients. We excluded studies that exclusively studied critically ill patients from calculation of prevalence of death but included them for calculating the association of risk factors with death.

Risk factors and outcomes

Primary outcome was prevalence of death and association of risk factors with death. We extracted data on death as recorded in the publications. We measured prevalence of severe disease and association with risk factors as secondary outcome. We defined outcome as severe disease for any of 1) the study classified COVID-19 disease as severe or critical, 2) intensive care unit (ICU) admission, 3) acute respiratory distress syndrome, or 4) mechanical ventilation. Severe disease was defined by studies as respiratory rate ≥ 30 per minute, oxygen saturation $\leq 93\%$, and $\text{PaO}_2/\text{FiO}_2 < 300$ and/or lung infiltrates $> 50\%$ within 24-48 hours.³ Critical illness was defined as respiratory failure, shock and/or multiple organ dysfunction or failure.³ Heart disease as a pre-existing condition was broadly defined by most studies as 'cardiovascular disease' (CVD). Additional outcomes were acute cardiac and kidney injury in the hospitalized patients that were defined as such by the studies.

Statistical analysis

We calculated and reported summary estimates from fixed-effects models²². We assessed heterogeneity across studies using Cochran's Q-test (χ^2 p value < 0.10)²³ and I^2 statistics ($I^2 > 30\%$)²⁴. In the presence of heterogeneity, we adjusted the 95% confidence intervals for between-study heterogeneity using the method described by Shore et al²⁵. We have presented the results from random effects meta-analysis as well. The meta-analysis was performed in Microsoft[®] Excel 2020 (Microsoft Corporation, Redmond, WA). We analyzed publication bias using funnel plots, and Begg's and Egger's tests. Quality of each study was assessed using the Newcastle-Ottawa assessment scales using the PRISMA guidelines. We calculated the following

as a part of our analyses: 1) prevalence of severe disease or death, 2) prevalence of risk factors, and 3) relative risk for the association of age, sex, and comorbidities with outcome. When not reported or when unadjusted odds ratio was presented, we calculated the relative risk (95% confidence interval) using the frequencies provided. Adjusted estimates were used where available. Case fatality risk (and case severity risk) for a specific risk factor was calculated as number of deaths (or severe disease) in patients with a risk factor out of all patients possessing the risk factor.

Results

Study characteristics

Initial search yielded 13919 citations. Articles were then filtered (**Figure 1**). We identified 275 articles for full text review, of which 44 studies met inclusion criteria (**Table 1**).^{4,5,7,8,13,14,26-61} The studies were conducted in: China (n=31), USA (n=8), Italy (n=2), UK (n=1), Iran (n=1) and Singapore (n=1). Two studies were prospective, one cross sectional, and remaining retrospective in nature.

Population and demographics

There were 20594 total COVID-19 hospitalized patients that included 12591 patients from the US and Europe (>90% from the US), and 7885 patients from China. Median age was 57 years [IQR: 54-60 years; $I^2=62\%$; n=36 studies]; 65 years [IQR: 61-70 years; $I^2=0\%$; n=9] for the US and Europe; and 54 years [IQR: 51-58 years; $I^2=0\%$; n=26] for China. Forty-three percent [95% CI: 38-48%; $I^2=96\%$; n=22] were aged ≥ 60 (**Table 2**); 46% [95% CI: 43-50%; $I^2=88\%$; n=4] for US and

Europe; and 31% [95% CI: 25-38%; $I^2=95\%$; $n=18$] for China. Fifty-eight percent [(95% CI: 56-60%); $I^2=78\%$; $n=43$] were males.

Prevalence of Death and Severe Disease

We calculated a pooled prevalence of death of 18% [95% CI: 15-22%; $I^2=92\%$; $n=26$], ranging from 1% to 28% across the studies, and a pooled prevalence of severe disease of 25% [95% CI: 21-31%; $I^2=96\%$; $n=36$] (**Table 2**). The prevalence of death was 21% [95% CI: 18-24%; $I^2=78\%$; $n=9$] in the US and Europe, and 13% [95% CI: 9-18%; $I^2=93\%$; $n=17$] in China. Prevalence of severe disease was 18% [95% CI: 13-24%; $I^2=97\%$; $n=10$] for US and Europe, and 33% [95% CI: 27-41%; $I^2=95\%$; $n=25$] for China. The measures for severe disease are described in

Supplemental table 1.

Pooled Prevalence of Risk Factors and Association with Death or Severe Disease

Age and Sex (Tables 2 and Figure 2)

Median age for people who died was 67 years [IQR: 63-71; $I^2=74\%$; $n=18$] and who had severe disease was 60 years [IQR: 58-63; $I^2=47\%$; $n=26$]. Seventy-six percent [95% CI: 71-83; $I^2=76\%$; $n=7$] of the deaths were in people aged ≥ 60 years and 68% [95% CI: 63-74; $n=13$] were in males. The CFR (95% CI) was 42% (34-50%) for age ≥ 60 years and 30% (23-38%) for males.

Patients aged ≥ 60 years [summary relative risk (sRR): 3.77; 95% CI: 2.94-4.82; $I^2=73\%$; $n=12$] and males [sRR: 1.34; 95% CI: 1.23-1.54; $I^2=19\%$; $n=17$] had higher risk of death. The risk of severe disease was similarly higher (**Supplemental table 1**).

Smoking and COPD (Table 2 and Figure 2)

The pooled prevalence for any history of smoking in the patients was 13% [95% CI: 11-16%; $I^2=22%$; $n=12$]. For patients with smoking history, the CFR was 22% (95% CI: 11-42%) and CSR was 42% (95% CI: 33-53). Compared to never smokers, patients with smoking history had higher relative risk of death [sRR: 1.87; 95% CI: 1.05-3.33; $I^2=80%$; $n=6$] and severe COVID-19 disease [sRR: 1.33; 95% CI: 1.16-1.54; $I^2=42%$; $n=15$]. The prevalence of COPD was 8% [95% CI: 6-10%; $I^2=93%$; $n=27$]. Patients with COPD had a CFR of 51% (95% CI: 36-71%), CSR of 49% (95% CI: 42-56%), a sRR of death of 2.80 [95% CI: 1.69-4.66; $I^2=82%$; $n=9$].

Hypertension (Tables 2 and Figure 3)

The pooled prevalence of hypertension in the COVID-19 patients was 47% [95% CI: 41-54% $I^2=98%$; $n=34$], with a CFR in hypertensive patients of 24% (95% CI: 16-36%) and a CSR of 53% (95% CI: 44-64%). Of the COVID-19 patients that died, 63% [95% CI: 57-70%; $I^2=98%$; $n=13$] were having hypertension. Hypertensives had higher relative risk of death [sRR: 1.84; 95% CI: 1.66-2.03; $I^2=34%$; $n=14$] and severe disease [sRR: 1.62; 95% CI: 1.37-1.92; $I^2=79%$; $n=23$] compared to non-hypertensives.

Diabetes (Tables 2 and Figure 3)

The pooled prevalence of diabetes was 26% [95% CI: 22-30%; $I^2=96%$; $n=36$] with a CFR of 22% (95% CI: 16-32%) and CSR of 46% (95% CI: 40-52%) in the diabetics. Of the COVID-19 patients that died, 38% [95% CI: 32-44%; $I^2=75%$; $n=13$] were diabetics. Diabetics had higher relative risk

of death [sRR: 1.51; 95% CI: 1.37-1.68; $I^2=0\%$; n=16] and severe disease [sRR: 1.50; 95% CI: 1.32-1.71; $I^2=65\%$; n=25] compared to non-diabetics.

Cardiovascular Disease (Tables 2 and Figure 3)

The pooled prevalence of CVD was 14% [95% CI: 12-17%; $I^2=95\%$; n=35] with a CFR of 62% (95% CI: 48-78%) and CSR of 56% (95% CI: 47-68%) among cardiac patients. Of all the patients that died, 29% [95% CI: 21-41%; $I^2=81\%$; n=12] had heart disease. Cardiac patients had higher relative risk of death [sRR: 1.99; 95% CI: 1.72-2.38; $I^2=33\%$; n=16] and severe disease [sRR: 1.67; 95% CI: 1.42-1.96; $I^2=83\%$; n=20] compared to patients without cardiac disease.

Chronic Kidney Disease (Tables 2 and Figure 3)

The pooled prevalence of CKD was 7% [95% CI: 5-10%; $I^2=93\%$; n=26] with a CFR of 49% (95% CI: 33-71%) and CSR of 34% (95% CI: 32-36%) in CKD patients. CKD was present in 23% [95% CI: 13-39%; $I^2=62\%$; n=7] of all COVID-19 patients that died. CKD patients had higher relative risk of death [sRR: 2.17; 95% CI: 1.30-3.13; $I^2=75\%$; n=8] and severe disease [sRR: 1.67; 95% CI: 1.30-2.16; $I^2=90\%$; n=14] compared to non-CKD patients.

COVID Related Organ System Injury (Figure 3)

To understand how pre-existing health conditions may be correlated with the risk of specific organ injury, we calculated the pooled prevalence of acute injury to lung, heart and kidney for studies that reported prevalence of both the pre-existing condition(s) and corresponding organ injury. Pooled across 12 studies,^{8,14,26,30,36-42,50} the prevalence of COPD at baseline was 6% (95%

CI: 4-11%) and that of ARDS was 48% (31-74%). The pooled prevalence of baseline CVD (n=11 studies) was 11% (95% CI: 9-12%) and that of acute cardiac injury (ACI) during hospitalization was 21% (95% CI: 16-29%).^{4,14,26,30,31,35,37,38,40,50,53} The prevalence of CKD (n=8 studies) was 5% (95% CI: 3-6%) and that of acute kidney injury (AKI) was 20% (95% CI: 15-26%).^{4,14,26,30,36,37,43,50}

Sensitivity and Sub-group Analyses

The relative risk of death in the patients was homogeneously elevated for hypertension, diabetes, COPD and heart disease. We carried out sensitivity analyses by excluding outliers. For the risk of death for hypertension and smoking history, we removed the study by Yao et al⁵³ which showed significantly higher risk compared to other studies; the results for both hypertension (sRR=2.02; 95% CI: 1.70-2.38) and smoking (sRR:1.59; 95% CI: 1.01-2.49) remained significant. Guan et al.¹³ had published a second study with additional patients and reported adjusted estimates for COPD, diabetes and hypertension. We used the adjusted risk estimates for the analyses. For the risk of death with other risk factors (CVD & CKD) for Guan et al., we conducted sensitivity analyses by using the counts only from the original study.³⁶ The results [sRR (95% CI)] were similar as: CVD=1.99 (1.69-2.33) and CKD=1.90 (1.27-2.86). We observed variation in the prevalence of risk factors and death based on region. Upon sub-group analysis, we noted significantly higher prevalence of risk factors among COVID-19 patients in the US and Europe than in China (**Figure 3. D**). Pooled prevalence of co-morbidities between US-Europe and China differed as follows: **1) for US-Europe:** HTN=54% (95% CI: 49-61%); diabetes=31% (95% CI: 28-34); CVD=17% (12-22%); smoking history=15% (11-21%); COPD=9% (6-13%) and **2)**

For China: HTN=23% (21-27%); diabetes=11% (10-13%); CVD=9% (7-11%); smoking history=11% (9-13%); and COPD=3% (2-4%).

Publication Bias

We observed asymmetry in the funnel plot for studies that reported prevalence of death in COVID-19 patients (Egger's test $p=0.007$ and Begg's test $p=0.004$) (**Supplemental Figure 1**). On further analysis, the plot remained asymmetrical when restricted to studies from China (Egger's $p=0.060$ and Begg's $p=0.053$) but was symmetrical for studies from US-Europe (Egger's $p=0.462$ and Begg's $p=0.210$). We observed symmetrical funnel plots with no bias for pooled prevalence severe disease or for analyses of association of risk factors with death or severe disease.

Discussion

We carried out a comprehensive systematic review and meta-analysis including 20594 hospitalized patients from 44 studies, 12591 from the US and Europe, and 7885 from China to investigate the prevalence and risk factors for death and severe disease in COVID-19 patients. We calculated a prevalence of death of 18%–21% for US and Europe, and 13% for China—and a prevalence of severe disease of 25%–18% for US and Europe, and 33% for China. In the COVID patients who died, risk factors were distributed as: age ≥ 60 years: 76%; males: 68%; hypertension: 63%; diabetes: 38%; heart disease: 29%; CKD: 23%; smoking history: 27%; and COPD: 12%.

In comparison with the overall prevalence of death of 18% for all COVID-19 hospitalized patients, the CFR was higher for patients with comorbidities: heart disease (62%), COPD (51%), CKD (49%), CLD (41%), hypertension (24%), diabetes (22%), and smoking history (22%). CFR was also higher for age \geq 60 years (42%) and for males (30%). The elevation in the risk of death was statistically significant for age \geq 60 (sRR=3.8; 95% CI: 2.9-4.8), male sex 1.3 (95% CI: 1.2-1.5), smoking history (sRR=1.9; 95% CI: 1.1-3.3), COPD (sRR=2.0; 95% CI: 1.6-2.4), heart disease (sRR=2.0; 95% CI: 1.7-2.4), CKD (sRR=2.0; 95% CI: 1.3-3.1), hypertension (sRR=1.8; 95% CI: 1.7-2.0), and diabetes (sRR=1.5; 95% CI: 1.4-1.7). All of the risk factors we analyzed were positively associated with progression to severe disease as well (**Table 2 & 3**). The results suggest that older age, male sex and the co-morbidities increase the risk of progression to severe disease and death in COVID-19 patients.

While we are unsure of the reason behind significant tests for publication bias for prevalence of death in studies from China, the funnel plot showed several studies with lower prevalence of death. This may suggest under reporting of death for the several studies from China. While the lower median age as well as lower prevalence of co-morbidities for COVID-19 patients in China can explain the lower prevalence of death, this should not lead to a statistical bias in publication as observed. The lower prevalence of death is incommensurate with the higher prevalence of severe disease observed for the studies from China. China's official death toll from COVID-19 was under-reported initially that was updated on April 17, 2020.⁶²

There is ongoing debate regarding predisposition by cardiovascular disease—hypertension, diabetes, heart disease—toward increasing the risk of acquisition of SARS-CoV-2.⁶³⁻⁶⁵ On one hand, Angiotensin converting enzyme 2 (ACE2)—by blocking the renin angiotensin aldosterone system (RAAS) and decreasing or countering the vasoconstrictive, proinflammatory and profibrotic properties of Angiotensin-II through catalysis of ANG-II to Ang-(1-7)—exerts cardiovascular protective effects and can prevent acute lung injury from SARS-CoV-2.⁶³⁻⁶⁵ However, on the other hand, it is unclear if angiotensin converting enzyme inhibitors (ACEI) or angiotensin receptor blockers (ARB)—used for treatment of hypertension, heart disease and diabetes—increases susceptibility toward SARS-CoV-2 infection through potential upregulation of ACE2, the host protein used by SARS-CoV-2 as a co-receptor to translocate intracellularly.⁶³⁻⁶⁶ In this context, it would be reasonable to posit that a substantially higher prevalence of cardiovascular comorbidities in the hospitalized patients compared to the prevalence in the general population may suggest elevated risk of acquisition of SARS-CoV-2 for patients with cardiovascular risk factors. To this end, we found that the prevalence of hypertension (55%), diabetes (31%) and heart disease (16%) in the hospitalized COVID-19 patients in the US were substantially higher than the corresponding prevalence of hypertension (29%),⁶⁷ diabetes (13%)⁶⁸ and heart disease (9%)¹⁵ in the general US population (**Figure 3 E**), suggesting either a tendency for SARS-CoV-2 to more effectively establish infection in cardiovascular patients or to cause more severe disease among them prompting hospitalization. The prevalence of smoking history (15%), COPD (9%) and CKD (8%) in the COVID-19 patients in the US was similar to or lower than the population prevalence of smoking (14%),⁶⁹ COPD (7%),⁷⁰ and CKD (15%)⁷¹ in the country. In the prevalence of hypertension (23%), diabetes (11%), CVD (9%), smoking (11%)

and COPD (3%) in the hospitalized patients in China (**Figure 3 D**) were all lower as compared to the US. The prevalence of these comorbidities in the COVID-19 patients was actually corresponding to the prevalence of hypertension (23%),¹⁷ diabetes (15%)⁷², and CVD (21%)⁷³ in the general population of China. A previous meta-analysis also noted this observation.¹⁸ Notably, the prevalence of smoking (11%) and CVD (9%) in the hospitalized COVID-19 patients in China are inexplicably lower than the prevalence of smoking (15%) and CVD (16%) among hospitalized patients in the US, despite the prevalence of smoking (47% in Chinese males)⁷⁴ and CVD (21%)⁷³ in the general Chinese population being significantly higher than that of the US. As such, we are unsure whether the lower prevalence of comorbidities noted for the COVID-19 patients in China are representative of the true prevalence. There was a such great urgency and a race to publish data in the early phase of the outbreak. As such, there exists the possibility of substantial under-recording of data on covariables.

We assessed if patients with specific co-morbidities at baseline had higher risk of specific organ injury from SARS-CoV-2 during hospitalization. While the available data did not allow direct assessment of this relation, we compared the prevalence of comorbidities with the prevalence of corresponding organ system injury for studies that reported both baseline comorbidity and corresponding organ injury. We observed that the risk of acute lung injury (measured as ARDS) (48%), ACI (21%), and AKI (20%) were substantially higher the baseline prevalence of COPD (6%), heart disease (11%) and CKD (5%), respectively (**Figure 3 F**). The higher prevalence of acute organ injury than the prevalence of baseline comorbidity simply indicates that ARDS, ACI

and AKI were also occurring in patients who did not have a corresponding comorbidity at baseline.

A limitation is that most studies simply reported frequencies of risk factor and did not present adjusted measures for disease severity or death. As such, the risk ratio we calculated from the frequencies are largely unadjusted estimates. Future studies could additionally present, at the least, age and sex adjusted measures for association of risk of comorbidities with death or severe disease. Many studies reported odds ratio for the measure of association between pre-existing conditions and risk of severe disease or death. Odds ratio poorly approximates risk ratio when the disease prevalence is high at baseline. For example, Zhou et al.¹⁴ calculated an odds ratio of 5.4 (95% CI: 0.96-30.4) for risk of death from COPD in COVID-19 patients whereas the risk ratio we calculated from the frequencies presented is RR=2.47; 95% CI: 1.34-4.55. Prevalence of severe disease or death in COVID-19 patients was high in several studies. Similarly, several meta-analyses calculated odds ratios instead of risk ratios to summarize the risk of disease severity or death in association with risk factors such as smoking, diabetes, hypertension and cardiovascular disease,^{10,11,16} often to be interpreted by media and even by researchers as a measure of relative risk. Lack of rigor in research design, analysis and interpretation could generate inconsistent and ungeneralizable results across studies leading to controversy and confusion around serious public health issues such as that existing for association (or not) of smoking with COVID-19 disease acquisition, severity or death.

As publications evolve at a pace that could be overwhelming for researchers and practitioners, we attempted to present a meaningful summary and inference for association of risk factors with death or severe disease from literatures published globally and an epidemiological framework for the risk of infection by SARS-CoV-2 based on presence of cardiovascular risk factors. This analysis can inform public health measures for COVID-19 screening and prevention, risk stratification and management of patients in clinical practice, analysis and presentation strategies for research data and inspire etiological investigations.

Research in Context

Evidence before this study: COVID-19 is an ongoing pandemic. Prevalence of death as a result of COVID-19 has greatly varied across regions. Epidemiological risk factors for progression of COVID-19 to severe disease and death, and the risk of acquisition of SARS-CoV-2, the causal agent for COVID-19, based on presence of pre-existing conditions are insufficiently understood. Particularly, there exist controversy and gap in knowledge on how pre-existing cardiovascular disease including hypertension, diabetes, and heart disease and use of angiotensin converting enzyme inhibitors and angiotensin receptor blockers influence the risk of disease acquisition as well as disease progression.

Added value of this study: Meta-analysis of 44 studies including 20285 COVID-19 patients revealed case fatality risk of 62% for heart disease, 51% for COPD, 49% for CKD, 24% for hypertension, 22% for diabetes, and 22% for smoking history. Of all patients that died, 76%

were aged ≥ 60 years, 68% were males, 63% had hypertension and 38% had diabetes. Age ≥ 60 , male sex, and pre-existing conditions were significantly associated with death and severe disease. Also, the prevalence of cardiovascular risk factors—hypertension, diabetes, and heart disease—were substantially higher in COVID-19 patients than their prevalence in the general US population.

Implications of all the available evidence: COVID-19 patients who are older, males and have pre-existing health conditions are at higher risk of progression to severe disease and death. The comparative prevalence data suggest people with cardiovascular risk factors may be at higher risk of acquisition of SARS-CoV-2 infection.

Funding: Dr. Dorjee is supported by grants from private philanthropists; the Johns Hopkins Alliance for Healthier World (Grant # 80045453); National Institute of Allergy and Infectious Diseases of the National Institute of Health (Grant # K01AI148583); the United Nations STOP TB PARTNERSHIP TB REACH (Grant # 134126); and the Pittsfield Anti-tuberculosis Association (PATA).

Table 1. Characteristics of studies in COVID-19 hospitalized patients to investigate association of risk factors with death or severe disease

Author, Year (Journal)	Country	Region	Study Period	Study Design	Sample size	Epidemiological Risk Factors	Outcomes	Measure of Association
Aggarwal S et al., 2020 (Diagnosis)	USA	Des Moines	3-1-2020 to 4-4-2020	Retrospective	16	Age, sex, smoking, substance use, obesity, HTN, DM, CVD, COPD, CKD, Cancer	Prevalence of death and primary end point (death, shock, or ICU admission). Association of risk factors with primary end point	Unadjusted RR (calculated)
CDC (MMWR)	USA	National	2-12-2020 to 3-28-2020	Retrospective	5285	Age, Current Smoker, DM, CVD, COPD, CKD, CLD	Prevalence of ICU admission. Association of risk factors with severe disease (ICU admission).	Unadjusted RR (calculated)
Chen G et al., 2020 (Journal of Clinical Investigation)	China	Wuhan	December 2019 to 01-27-2020	Retrospective	21	Age, sex, Huanan sea food market exposure, HTN, DM	Prevalence of severe disease. Comparison of moderate and severe cases based on risk factors.	Unadjusted RR (calculated)
Chen J et al., 2020 (Journal of Infection)	China	Shanghai	1-20-2020 to 2-6-2020	Retrospective	249	Age, sex	Prevalence of ICU admission. Association of age and sex with ICU admission.	Adjusted OR reported for age and sex
Chen Q et al., 2020 (Infection)	China	Zhejiang province	1-1-2020 to 3-11-2020	Retrospective	145	Age, sex, smoking, exposure history, BMI, HTN, DM, COPD, CKD, Solid tumor, Heart disease, HIV infection	Prevalence of severe disease. Association of risk factors with severe disease.	Unadjusted RR (calculated)
Chen T et al., 2020 (BMJ)	China	Wuhan, Hubei	1-13-2020 to 2-28-2020	Retrospective	274	Age, sex, sea food market exposure, contact history, smoking (current and former), HTN, DM, CVD, CHF, heart failure, cancer, HBV, HIV, CKD	Association of risk factors with death compared death and recovered group. Presence of hospitalized patients excluded from study	Unadjusted RR (calculated)
Cummings MJ et al., 2020 (The Lancet)	USA	New York City	3-2-2020 to 4-1-2020	Prospective	257	Age, sex, race, BMI, HTN, DM, chronic cardiac disease (CHD and CHF), CKD, smoking history, COPD or ILD, cancer, HIV infection, liver cirrhosis	Association of risk factors with death	Adjusted HR
Deng Y et al., 2020 (Chin Med J)	China	Wuhan	1-1-2020 to 2-21-2020	Retrospective	116 out of 964	Age, sex, HTN, DM, Heart Disease, Cancer	Association of risk factors with death compared death and recovered group. Presence of hospitalized patients excluded from study	Unadjusted RR (calculated)
Du R-H et al., 2020 (ERJ)	China	Wuhan, Hubei	1-25-2020 to 2-7-2020	Retrospective	179	Age, sex, HTN, DM, CVD, TB, cancer, CKD or CLD	Prevalence of death. Association of risk factors with death.	Adjusted OR for age≥65 and CVD. Unadjusted RR (calculated) for remaining variables.
Feng Y et al., 2020 (AJRCCM)	China	Wuhan, Shanghai, Anhui province	1-1-2020 to 2-15-2020	Retrospective	476	Age, age groups, sex, Wuhan exposure, smoking, alcohol, HTN, anti-hypertensives, CVD, DM, cancer, COPD, CKD	Prevalence of death. Association of risk factors with severe disease	Adjusted HR for HTN, CVD, DM. Unadjusted RR (calculated) for other variables.
Ferguson J et al., 2020 (EID)	USA	Northern California	03-13-2020 to 04-11-2020	Retrospective	72	Sex, race, smoking, HTN, DM, CKD, Heart Disease, COPD	Prevalence of ICU admission. Association of risk factors with severe disease (ICU admission).	Unadjusted RR (calculated)
Gold J et al, 2020	USA	Georgia	3-1-2020 to	Retrospective	305	Age, sex, race, HTN, DM, Heart Disease,	Prevalence of patient characteristics, death, and	Unadjusted RR

medRxiv preprint doi: <https://doi.org/10.1101/2020.06.19.20195483>; this version posted June 23, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

(MMWR)			3-30-2020			COPD, CKD, Cancer	ICU	(calculated)
Goyal P et al. 2020 (NEJM)	USA	New York City	3-3-2020 to 3-27-2020	Retrospective	393	Age, sex, race, smoking, HTN, DM, COPD, Heart Disease, Asthma	Prevalence of severe disease (mechanical ventilation). Association of risk factors with severe disease.	Unadjusted RR (calculated)
Guan et al., 2020 (NEJM)	China	National	12-11-2019 to 01-31-2020	Retrospective	1099	Age, sex, smoking (never, current, former), exposure to transmission source, HTN, DM, CHD, CKD, COPD, Cancer, HBV, cerebrovascular disease, immunodeficiency	Prevalence of death, composite outcome, ((Death/MV/ICU) and severe disease). Association of risk factors with severe disease and composite outcome.	Unadjusted RR (calculated)
Guan Wei-Jie, 2020 (ERJ)	China	Nationwide	12-11-2019 to 1-31-2020	Retrospective	1590	Age, sex, smoking, CKD, COPD, HTN, DM, CVD, Cancer, HBV	Prevalence of patient characteristic death and composite outcome (Death, ICU, MV)	Adjusted HR
Hu L et al., 2020 (CID)	China	Wuhan	1-8-2020 to 2-20-2020	Retrospective	323	Age, sex, current smoker, HTN, DM, CVD, COPD, CKD, CLD, Cancer	Prevalence of severe (severe and critical) disease. Association of risk factors with disease severity.	Unadjusted RR (calculated)
Huang C et al., 2020 (The Lancet)	China	Wuhan	12-16-2020 to 1-2-2020	Prospective	41	Age, sex, Huanan seafood market exposure, current smoking, HTN, DM, CKD, COPD, CVD, Cancer, CLD	Association of risk factors with severe disease (ICU care)	Unadjusted RR (calculated)
Inciardi R et al., 2020 (Eur Heart J)	Italy	Lombardy	3-4-2020 to 3-25-2020	Retrospective	99	Sex, smoking, HTN, DM, coronary artery disease, COPD, CKD, cancer	Prevalence of death. Association of risk factors with death.	Unadjusted RR (calculated)
Javanian M et al., 2020 (Rom J Intern Med)	Iran	Mazandaran province	2-25-2020 to 3-12-2020	Retrospective	100	Age, sex, HTN, DM, CVD, CKD, cancer, CLD	Prevalence of death. Association of risk factors with death.	Unadjusted RR (calculated)
Liu W et al. 2020 (Chin Med J)	China (Wuhan, Hubei)	Wuhan	12-30-2019 to 01-15-2020	Retrospective	78	Age, sex, history of smoking, exposure to Huanan seafood market, hypertension, diabetes, COPD, Cancer	Compared progression group and nonprogression group. Progression group defined as progression to severe or critical disease or death.	Unadjusted RR (calculated)
Li X et al., 2020 (J Allergy Clin Immunol)	China (Wuhan, Hubei)	Wuhan, Hubei	1-26-2020 to 2-5-2020	Retrospective	548	Age, sex, smoking, HTN, DM, Heart Disease, CKD, Cancer, COPD	Prevalence of death and severe disease. Association of risk factors with severe disease.	Unadjusted RR (calculated)
Nowak B et al., 2020 (Pol Arch Intern Med)	Poland	Warsaw	3-16-2020 to 4-7-2020	Retrospective	169	Sex, smoking, HTN, DM, coronary artery disease, COPD, CKD, AKI, cancer	Prevalence of death. Association of risk factors with death.	Unadjusted RR (calculated)
Palaiodimos L et al., 2020 (Metabolism)	USA	New York	3-9-2020 to 3-22-2020	Retrospective	200	Age, sex, race, smoking, HTN, DM, coronary artery disease, COPD, CKD, cancer	Prevalence of death. Association of risk factors with death.	Adjusted OR (provided by the study)
Richardson S et al., 2020 (JAMA)	USA	New York	3-1-2020 to 4-4-2020	Retrospective	5700	Age, sex, race, smoking, HTN, DM, COPD, asthma, coronary artery disease, kidney disease, Liver disease (Cirrhosis, hep B, Hep C, obesity, cancer	Prevalence of ICU admission and death. Association of risk factors with death.	Unadjusted RR (calculated)

Preprint JMIR Publications | doi:10.19182/jmip.2020.06.19.013543 | This version posted June 20, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted JMIR Publications a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.

Shi Y et al., 2020 (Crit Care)	China	Zhejiang province	Not specified to 02-11-2020	Retrospective	487	Age, sex, smoking, HTN, DM, CKD, CVD, CLD, cancer	Prevalence of and Association of risk factors with severe disease	Unadjusted RR (calculated)
Sun L et al., 2020 (Journal of Medical Virology)	China	Beijing	1-20-2020 to 2-15-2020	Retrospective	55	Age, sex, exposure, HTN, DM, CVD, Lung Disease, CKD, CLD	Prevalence of severe disease. Association of risk factors with severe disease.	Unadjusted RR (calculated)
Tian S et al., 2020 (Journal of Infection)	China	Beijing	1-20-2020 to 2-10-2020	Retrospective	262	Age, sex, contact history, exposure to Wuhan.	Prevalence of death. Association of severe disease with risk factors.	Unadjusted RR (calculated)
Tomlins J et al., 2020 (Journal of Infection)	UK	Bristol	3-10-2020 to 3-30-2020	Retrospective	95	Age, sex, HTN, DM, COPD, CVD, cancer, renal disease, gastrointestinal disease, neurological disease, immunocompromise	Prevalence of death. Association of risk factors with death.	Unadjusted RR (calculated)
Wan S et al., 2020 (Journal of Medical Virology)	China	Northeast Chongqing	1-23-2020 to 2-8-2020	Retrospective	135	Age, sex, smoking, CKD, COPD, HTN, DM, CVD, Cancer, CLD, exposure, travel history	Prevalence of severe disease. Association of risk factors with severe disease.	Unadjusted RR (calculated)
Wang D et al., 2020 (JAMA)	China	Wuhan	1-1-2020 to 1-28-2020	Retrospective	138	Age, sex, Huanan Seafood Market Exposure, HTN, DM, CVD, COPD, Cancer, CKD, CLD, HIV	Prevalence of death and ICU admission; Association of risk factors with severe disease (ICU care)	Unadjusted RR (calculated)
Wang R et al., 2020 (Internal Journal of Infectious Diseases)	China	Fuyang	1-20-2020 to 02-09-2020	Retrospective	125	Age, sex, CVD, Cancer	Prevalence of critical disease. Association of age, sex, and smoking with critical disease	Unadjusted RR (calculated)
Wang Z et al., 2020 (CID)	China	Wuhan	1-16-2020 to 01-29-2020	Retrospective	69	Age, sex, HTN, DM, CVD, COPD, Cancer, HBV, Asthma	Prevalence of death and severe disease (SpO2<90%); Association of risk factors with severe disease	Unadjusted RR (calculated)
Wu C et al., 2020 (JAMA Intern Med)	China	Wuhan	12-15-2019 to 01-26-2020	Retrospective	201	Age, sex, HTN, DM, CVD, CKD, Chronic Lung Disease, Cancer, CLD, Sea Food Market Exposure.	Prevalence of ARDS, ICU admission and death. Association of risk factors with severe disease (ARDS) and death	Unadjusted HR reported for age≥65, sex, HTN and DM. Unadjusted RR (calculated) for other variables
Yang X et al., 2020 (Lancet Respir Med)	China	Wuhan	12/24/2019 to 1-26-2020	Retrospective	52	Age, sex, exposure, COPD, diabetes, chronic cardiac disease, smoking, malnutrition	Association of risk factors with death	Unadjusted RR (calculated)
Yao Q et al., 2020 (Pol Arch Intern)	China	Huanggang city, Hubei	1-30-2020 to 2-11-2020	Retrospective	108	Age, sex, smoking, HTN, DM, CVD, CLD, cancer	Prevalence of severe disease, and death; Association of risk factors with severe disease and death	Unadjusted RR (calculated)
Young BE et al., 2020 (JAMA)	Singapore	Singapore	1-23-2020 to 2-3-2020	Retrospective	18	Age, sex	Prevalence of severe disease (receiving supplemental O2). Association of severe disease with age and sex.	Unadjusted RR (calculated)
Yu T et al., 2020	China	Guangdo	Jan 2020 to	Cross-	95	Age, sex, current smoker	Prevalence of ARDS. Association of age, sex, and	Unadjusted RR

(Clinical Therapeutics)		ng	Feb 2020	sectional			smoking with ARDS.	(calculated)
Yu X et al., 2020 (Transboundary and Emerging Diseases)	China	Shanghai	Up to 2-19-2020	Retrospective	333	Age, sex, BMI, smoking, alcohol, exposure, HTN, DM, CVD	Prevalence of death and severe disease (Severe/critical pneumonia). Association of risk factors with severe disease.	Adjusted OR for age group, sex, CVD, DM, HTN. Unadjusted RR (calculated) for remaining variables.
Zhang G et al., 2020 (BMC Respiratory Research)	China	Wuhan	1-16-2020 to 2-25-2020	Retrospective	95	Age, sex	Prevalence of severe disease, composite end point, and death. Association of risk factors with severe disease.	Unadjusted RR (calculated)
Zhang JJ et al., 2020 (Allergy)	China	Wuhan	1-16-2020 to 2-3-2020	Retrospective	140	Age, sex, current smoker, past smoker, exposure history, HTN, DM, CVD, COPD, CKD, CLD, PTB,	Prevalence of severe disease. Association of risk factors with severe disease (ICU admission).	Unadjusted RR (calculated)
Zhao X-Y et al., 2020 (BMC Inf Dis)	China	Hubei (Non Wuhan)	1-16-2020 to 2-10-2020	Retrospective	91	Age, sex, DM, COPD, Cancer, Kidney disease	Prevalence of death. Association of risk factors with severe disease	Unadjusted RR (calculated)
Zheng S et al., 2020 (BMJ)	China	Zhejiang province	1-19-2020 to 2-15-2020	Retrospective	96	Age, sex, HTN, DM, CVD, lung disease, Liver disease, renal disease, malignancy, viral Load, immunocompromise	Prevalence of death and severe disease. Association of risk factors with severe disease.	Unadjusted RR (calculated)
Zheng Y et al., 2020 (Pharmacological Research)	China	Shiyan, Hubei	1-16-2020 to 2-4-2020	Retrospective	73	Age, sex, exposure, smoking history, DM, CVD	Prevalence of severe (severe/ critical) disease. Association of smoking and diabetes with severe disease.	Unadjusted RR (calculated)
Zhou F et al., 2020 (The Lancet)	China	Wuhan	12-29-2019 To 01-31-2020	Retrospective	191	Age, sex, current smoking, exposure history, HTN, DM, CVD, COPD, Cancer, CKD	Prevalence of severe disease (ICU admission) and death; Association of risk factors with death	Adjusted OR for age and CHD. Unadjusted RR (calculated) for other variables.

HTN: hypertension; DM: diabetes mellitus; CVD: cardiovascular disease; COPD: chronic obstructive pulmonary disease; CHD: coronary heart disease; CKD: chronic kidney disease; OR: odds ratio; RR: relative risk;

Table 2. Pooled prevalence of death and severe disease stratified by epidemiological risk factors in COVID-19 patients hospitalized between December 2019-May 2020


medRxiv preprint doi: <https://doi.org/10.1101/2020.06.19.20105588>; this version posted June 20, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY 4.0 International license.

Risk group	Overall prevalence of risk across studies		Pooled Prevalence of Death (Case Fatality Risk) and Risk Factor			Summary Relative Risk of Death			
	No. of studies	Pooled prevalence of risk factor, % (95% CI)	No. of studies	* Case fatality risk (Pooled prevalence of death in risk group), % (95% CI)	#Prevalence of risk group in dead, % (95% CI)	No. of studies	Fixed Effects Summary relative risk; 95% CI (Shore adjusted)	Random Effects# sRR; (95% CI)	Heterogeneity I^2 ; χ^2 ; p value
Age ≥ 60 years	22	43 (38-48)	8	42 (34-50)	76 (71-82)	12	3.77 (2.94-4.82)	1.66 (1.19-2.27)	73%; 41; p<0.01
Male	43	58 (56-60)	16	30 (23-38)	68 (63-74)	17	1.34 (1.20-1.50)	1.39 (1.05-1.83)	19%; 16; p=0.20
Smoking history	22	13 (11-16)	5	22 (12-38)	27 (18-41)	6	1.87 (1.05-3.33)	1.89 (1.05-3.44)	80%; 25; p<0.01
Current smoker	13	8 (6-12)	3	21 (8-56)	21 (13-37)	4	2.20 (1.16-4.16)	2.51 (1.16-5.86)	78%; 14; p<0.01
COPD	27	8 (6-10)	8	51 (36-71)	12 (7-21)	9	1.97 (1.59-2.43)	homogeneous	0%; 3.55; p=0.83
Hypertension	34	47 (41-54)	13	24 (16-36)	63 (57-70)	14	1.84 (1.66-2.03)	homogeneous	34%; 20; p=0.10
Diabetes	36	26 (23-30)	13	22 (16-32)	38 (32-44)	16	1.51 (1.37-1.68)	homogeneous	0%; 10; p=0.68
Cardiovascular disease	35	14 (12-17)	12	62 (48-78)	29 (21-41)	16	1.99 (1.72-2.38)	homogeneous	33% (22; p=0.10)
Chronic kidney disease	26	7 (5-10)	7	49 (33-71)	23 (13-39)	8	2.02 (1.30-3.13)	2.14 (1.30-3.51)	75%; 28; p<0.01

* Case fatality risk of represent total number of people that died in the specific risk group divided by total population in the risk group.

Prevalence of risk group in dead represent total number of people having the risk group divided by total population that died.

medRxiv preprint doi: <https://doi.org/10.1101/2020.06.19.20135418>; this version posted June 20, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.



medRxiv preprint doi: <https://doi.org/10.1101/2020.06.19.20135483>; this version posted June 20, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a [CC-BY-NC-ND 4.0 International license](#).

Figure 2. Prevalence of death in COVID-19 patients and association with epidemiological risk factors






Figure 3. Risk of death in COVID-19 patients associated with pre-existing conditions, and prevalence of comorbidities and organ injury


A. Risk of death in COVID-19 patients with hypertension




B. Risk of death in COVID-19 patients with diabetes




C. Risk of death in COVID-19 patients with heart disease




D. Prevalence of comorbidities in hospitalized COVID-19 patients in the US-Europe and China



E. Pooled prevalence of co-morbidities in COVID-19 patients in the US and corresponding prevalence in the US population



F. Pre-existing condition and corresponding organ injury in COVID-19 patients



HTN: hypertension; CVD: cardiovascular disease; COPD: chronic obstructive pulmonary disease; CKD: chronic kidney disease; AKI: acute kidney injury; ACI: acute cardiac injury; ARDS: acute respiratory distress syndrome

medRxiv preprint doi: <https://doi.org/10.1101/2020.06.19.20136443>; this version posted June 20, 2020. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC-ND 4.0 International license.

Reference:

1. Johns Hopkins University. COVID-19 Dashboard by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). <https://coronavirus.jhu.edu/map.html>. Published 2020. Accessed June 4, 2020.
2. Garg S, Kim L, Whitaker M, et al. Hospitalization Rates and Characteristics of Patients Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 - COVID-NET, 14 States, March 1-30, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(15):458-464.
3. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72,314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA.* 2020;10.1001/jama.2020.2648.
4. Li X, Xu S, Yu M, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol.* 2020.
5. Palaiodimos L, Kokkinidis DG, Li W, et al. Severe obesity, increasing age and male sex are independently associated with worse in-hospital outcomes, and higher in-hospital mortality, in a cohort of patients with COVID-19 in the Bronx, New York. *Metabolism.* 2020;108:154262.
6. Cummings MJ, Baldwin MR, Abrams D, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet.* 2020.
7. Du RH, Liang LR, Yang CQ, et al. Predictors of Mortality for Patients with COVID-19 Pneumonia Caused by SARS-CoV-2: A Prospective Cohort Study. *Eur Respir J.* 2020.
8. Wu C, Chen X, Cai Y, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA internal medicine.* 2020.
9. Alqahtani JS, Oyelade T, Aldhahir AM, et al. Prevalence, Severity and Mortality associated with COPD and Smoking in patients with COVID-19: A Rapid Systematic Review and Meta-Analysis. *PLoS One.* 2020;15(5):e0233147.
10. Lippi G, Henry BM. Active smoking is not associated with severity of coronavirus disease 2019 (COVID-19). *European Journal of Internal Medicine.* 2020.
11. Patanavanich R, Glantz SA. Smoking is Associated with COVID-19 Progression: A Meta-Analysis. *Nicotine Tob Res.* 2020.
12. Rossato M, Russo L, Mazzocut S, Di Vincenzo A, Fioretto P, Vettor R. Current Smoking is Not Associated with COVID-19. *The European Respiratory Journal.* 2020.
13. Guan W-J, Liang W-H, Zhao Y, et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: A Nationwide Analysis. *The European respiratory journal.* 2020.
14. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet.* 2020.
15. Benjamin EJ, Muntner P, Alonso A, et al. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association. *Circulation.* 2019;139(10):e56-e528.
16. Li X, Guan B, Su T, et al. Impact of cardiovascular disease and cardiac injury on in-hospital mortality in patients with COVID-19: a systematic review and meta-analysis. *Heart.* 2020.
17. Wang Z, Chen Z, Zhang L, et al. Status of Hypertension in China: Results From the China Hypertension Survey, 2012-2015. *Circulation.* 2018;137(22):2344-2356.
18. Yang J, Zheng Y, Gou X, et al. Prevalence of comorbidities in the novel Wuhan coronavirus (COVID-19) infection: a systematic review and meta-analysis. *International Journal of Infectious Diseases.* 2020.
19. Zheng Z, Peng F, Xu B, et al. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. *J Infect.* 2020.

20. Li L-Q, Huang T, Wang Y-Q, et al. 2019 novel coronavirus patients' clinical characteristics, discharge rate and fatality rate of meta-analysis. *Journal of medical virology*. 2020;10.1002/jmv.25757-25710.21002/jmv.25757.
21. Rodriguez-Morales AJ, Cardona-Ospina JA, Gutierrez-Ocampo E, et al. Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel medicine and infectious disease*. 2020:101623-101623.
22. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials*. 1986;7(3):177-188.
23. Cochran WG. The Combination of Estimates From Different Experiments. *Biometrics*. 1954;10(1):101-129.
24. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. 2002;21(11):1539-1558.
25. Shore RE, Gardner MJ, Pannett B. Ethylene oxide: an assessment of the epidemiological evidence on carcinogenicity. *Br J Ind Med*. 1993;50(11):971-997.
26. Aggarwal S, Garcia-Telles N, Aggarwal G, Lavie C, Lippi G, Henry BM. Clinical features, laboratory characteristics, and outcomes of patients hospitalized with coronavirus disease 2019 (COVID-19): Early report from the United States. *Diagnosis*. 2020;7(2):91-96.
27. Chen G, Wu D, Guo W, et al. Clinical and immunologic features in severe and moderate Coronavirus Disease 2019. *The Journal of clinical investigation*. 2020.
28. Chen J, Qi T, Liu L, et al. Clinical progression of patients with COVID-19 in Shanghai, China. *Journal of Infection*. 2020.
29. Chen Q, Zheng Z, Zhang C, et al. Clinical characteristics of 145 patients with corona virus disease 2019 (COVID-19) in Taizhou, Zhejiang, China. *Infection*. 2020.
30. Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ*. 2020;368:m1091-m1091.
31. Deng Y, Liu W, Liu K, et al. Clinical characteristics of fatal and recovered cases of coronavirus disease 2019 (COVID-19) in Wuhan, China: a retrospective study. *Chinese Medical Journal*. 9000;Publish Ahead of Print.
32. Feng Y, Ling Y, Bai T, et al. COVID-19 with Different Severity: A Multi-center Study of Clinical Features. *Am J Respir Crit Care Med*. 2020.
33. Ferguson J, Rosser JI, Quintero O, et al. Characteristics and Outcomes of Coronavirus Disease Patients under Nonsurge Conditions, Northern California, USA, March-April 2020. *Emerg Infect Dis*. 2020;26(8).
34. Gold JAW, Wong KK, Szablewski CM, et al. Characteristics and Clinical Outcomes of Adult Patients Hospitalized with COVID-19 - Georgia, March 2020. *MMWR Morb Mortal Wkly Rep*. 2020;69(18):545-550.
35. Goyal P, Choi JJ, Pinheiro LC, et al. Clinical Characteristics of Covid-19 in New York City. *N Engl J Med*. 2020.
36. Guan W-j, Ni Z-y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*. 2020.
37. Hu L, Chen S, Fu Y, et al. Risk Factors Associated with Clinical Outcomes in 323 COVID-19 Hospitalized Patients in Wuhan, China. *Clinical Infectious Diseases*.
38. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020.
39. Inciardi RM, Adamo M, Lupi L, et al. Characteristics and outcomes of patients hospitalized for COVID-19 and cardiac disease in Northern Italy. *Eur Heart J*. 2020;41(19):1821-1829.
40. Javanian M, Bayani M, Shokri M, et al. Clinical and laboratory findings from patients with COVID-19 pneumonia in Babol North of Iran: a retrospective cohort study. *Rom J Intern Med*. 2020.
41. Liu W, Tao Z-W, Lei W, et al. Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chin Med J (Engl)*. 2020:10.1097/CM1099.0000000000000775.

42. Nowak B, Szymanski P, Pankowski I, et al. Clinical characteristics and short-term outcomes of patients with coronavirus disease 2019: a retrospective single-center experience of a designated hospital in Poland. *Pol Arch Intern Med.* 2020;130(5):407-411.
43. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA.* 2020.
44. Shi Y, Yu X, Zhao H, Wang H, Zhao R, Sheng J. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care.* 2020;24(1):108.
45. Sun L, Shen L, Fan J, et al. Clinical Features of Patients with Coronavirus Disease 2019 (COVID-19) from a Designated Hospital in Beijing, China. *Journal of medical virology.* 2020.
46. Team CC-R. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 - United States, February 12-March 28, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(13):382-386.
47. Tian S, Hu N, Lou J, et al. Characteristics of COVID-19 infection in Beijing. *J Infect.* 2020;80(4):401-406.
48. Tomlins J, Hamilton F, Gunning S, Sheehy C, Moran E, MacGowan A. Clinical features of 95 sequential hospitalised patients with novel coronavirus 2019 disease (COVID-19), the first UK cohort. *Journal of Infection.* 2020.
49. Wan S, Xiang Y, Fang W, et al. Clinical Features and Treatment of COVID-19 Patients in Northeast Chongqing. *Journal of medical virology.* 2020.
50. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA.* 2020:10.1001/jama.2020.1585.
51. Wang R, Pan M, Zhang X, et al. Epidemiological and clinical features of 125 Hospitalized Patients with COVID-19 in Fuyang, Anhui, China. *Int J Infect Dis.* 2020.
52. Wang Z, Yang B, Li Q, Wen L, Zhang R. Clinical Features of 69 Cases with Coronavirus Disease 2019 in Wuhan, China. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America.* 2020.
53. Yao Q, Wang P, Wang X, et al. Retrospective study of risk factors for severe SARS-Cov-2 infections in hospitalized adult patients. *Pol Arch Intern Med.* 2020.
54. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA.* 2020.
55. Yu T, Cai S, Zheng Z, et al. Association Between Clinical Manifestations and Prognosis in Patients with COVID-19. *Clinical Therapeutics.* 2020.
56. Yu X, Sun X, Cui P, et al. Epidemiological and Clinical Characteristics of 333 Confirmed Cases with Coronavirus Disease 2019 in Shanghai, China. *Transboundary and Emerging Diseases.*n/a(n/a).
57. Zhang G, Zhang J, Wang B, Zhu X, Wang Q, Qiu S. Analysis of clinical characteristics and laboratory findings of 95 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a retrospective analysis. *Respiratory Research.* 2020;21(1):74-74.
58. Zhang J-J, Dong X, Cao Y-Y, et al. Clinical characteristics of 140 patients infected by SARS-CoV-2 in Wuhan, China. *Allergy.* 2020:10.1111/all.14238.
59. Zhao X-Y, Xu X-X, Yin H-S, et al. Clinical characteristics of patients with 2019 coronavirus disease in a non-Wuhan area of Hubei Province, China: a retrospective study. *BMC Infectious Diseases.* 2020;20.
60. Zheng S, Fan J, Yu F, et al. Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January-March 2020: retrospective cohort study. *BMJ.* 2020;369:m1443.
61. Zheng Y, Xiong C, Liu Y, et al. Epidemiological and Clinical Characteristics Analysis of COVID-19 in the Surrounding Areas of Wuhan, Hubei Province in 2020. *Pharmacological Research.* 2020:104821.
62. Qin A. China Raises Coronavirus Death Toll by 50% in Wuhan. *The New York Times.* April 17, 2020, 2020.
63. Akhmerov A, Marban E. COVID-19 and the Heart. *Circ Res.* 2020;126(10):1443-1455.
64. Clerkin KJ, Fried JA, Raikhelkar J, et al. COVID-19 and Cardiovascular Disease. *Circulation.* 2020;141(20):1648-1655.

65. South AM, Diz DI, Chappell MC. COVID-19, ACE2, and the cardiovascular consequences. *Am J Physiol Heart Circ Physiol*. 2020;318(5):H1084-H1090.
66. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med*. 2020;8(4):e21.
67. Fryar CD, Ostchega Y, Hales CM, Zhang G, Kruszon-Moran D. Hypertension Prevalence and Control Among Adults: United States, 2015-2016. *NCHS Data Brief*. 2017(289):1-8.
68. United States Centers for Disease Control and Prevention. *National Diabetes Statistics Report*. Atlanta, GA2020.
69. Creamer MR, Wang TW, Babb S, et al. Tobacco Product Use and Cessation Indicators Among Adults - United States, 2018. *MMWR Morb Mortal Wkly Rep*. 2019;68(45):1013-1019.
70. Biener AI, Decker SL, Rohde F. Prevalence and Treatment of Chronic Obstructive Pulmonary Disease (COPD) in the United States. *JAMA*. 2019;322(7):602.
71. United States Centers for Disease Control and Prevention. Chronic Kidney Disease in the United States, 2019. https://www.cdc.gov/kidneydisease/pdf/2019_National-Chronic-Kidney-Disease-Fact-Sheet.pdf. Published 2019. Accessed June 7, 2020, 2020.
72. Hu C, Jia W. Diabetes in China: Epidemiology and Genetic Risk Factors and Their Clinical Utility in Personalized Medication. *Diabetes*. 2018;67(1):3-11.
73. Ma LY, Chen WW, Gao RL, et al. China cardiovascular diseases report 2018: an updated summary. *J Geriatr Cardiol*. 2020;17(1):1-8.
74. Wang M, Luo X, Xu S, et al. Trends in smoking prevalence and implication for chronic diseases in China: serial national cross-sectional surveys from 2003 to 2013. *Lancet Respir Med*. 2019;7(1):35-45.