What variables can better predict the number of infections and deaths worldwide by SARS-CoV-2? Variation through time

García de Alcañíz, J. G.; Romero-López, J.; Martínez-Esteban, R. P.; López-Rodas, V. and Costas, E.*

Genetics, Faculty of Veterinary Medicine, Complutense University, 28040 Madrid, Spain *Corresponding author: ecostas@ucm.es

ABSTRACT

Using data from 50 very different countries (which represent nearly 70% of world's population) and by means of a regression analysis, we studied the predictive power of different variables (mobility, air pollution, health & research, economic and social & geographic indicators) over the number of infected and dead by SARS-CoV-2. We also studied if the predictive power of these variables changed during a 4 months period (March, April, May and June). We approached data in two different ways, cumulative data and non-cumulative data.

The number of deaths by Covid-19 can always be predicted with great accuracy from the number of infected, regardless of the characteristics of the country.

Inbound tourism emerged as the variable that best predicts the number of infected (and, consequently, the number of deaths) happening in the different countries. Electricity consumption and air pollution of a country (CO₂ emissions, nitrous oxide and methane) are also capable of predicting, with great precision, the number of infections and deaths from Covid-19. Characteristics such as the area and population of a country can also predict, although to a lesser extent, the number of infected and dead. All predictive variables remained significant through time.

In contrast, a series of variables, which in principle would seem to have a greater influence on the evolution of Covid-19 (hospital bed density, Physicians per 1000 people, Researches in R & D, urban population...), turned out to have very little - or none- predictive power.

Our results explain why countries that opted for travel restrictions and social withdrawal policies at a very early stage of the pandemic outbreak, obtained better results. Preventive policies proved to be the key, rather than having large hospital and medical resources.

Competing Interest Statement

The authors have declared no competing interest.

Keywords: SARS-Cov-2, COVID-19, Regression Analysis, Tourism, Air Pollution, NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice. Urban Population, Hospital Bed density

INTRODUCTION

COVID-19 is a global pandemic caused by SARS-CoV-2 virus^{1,2}. The virus rapidly spread and is happening at the same time all around the world, to different people living in different countries (with different mobility, pollution, health, economic, geographic and social characteristics). Infection and death counts vary widely not only from country to country but also through time. In some countries the virus is in the remission phase, with chances of new outbreaks, while in others is yet uncontrolled.

The main feature of COVID-19 pandemic is the lack of knowledge, SARS-CoV-2 is a new coronavirus and little is known yet, despite the effort of scientists around the world. Under this setting, regression analysis is a very useful tool to find out which variables can predict number of infected and dead, and which variables cannot. Consequently, numerous studies have been performed since the outbreak of the pandemic using regression analysis 3-7, most of them focus on the clinical aspects of the pandemic and its implications on health care settings ^{5,8-13}, such as cardiac injury, kidney disease, respiratory conditions, diabetes, comorbidities 8,9,14-19 gender and underlying comorbidities ²⁰, severe COVID-19 condition and risk factors ^{21–24}, number of CD8 +T cells as risk factors for the duration of SARS-CoV-2 viral positivity ²⁵ ²⁶, clinical symptoms as smell loss ²⁷, smoking habits ²⁸. And not as many, on other aspects like looking into, for example, temperature as an affecting parameter to transmission rates^{3,29-31}. Some have used GIS modelling to understand what is happening with SARS CoV-2 in this pandemic ³². Some others try to unveil the relevance of temperature, humidity, climate, air quality, socioeconomic factors, demographics...^{3,4,38–40,5,30,31,33–37}.

The relationship of COVID-19 with socioeconomic variables has also been investigated. For example, Mollalo et al. (2020) have performed a study in the U.S.A. compiling 35 environmental, socioeconomic, topographic and demographic variables and create a geodatabase to explain the spatial variability of the COVID incidence ³⁷. GIS-based spatial modelling studies have also been done. ³² Guha et al (2020) studied community and socioeconomic characteristics of people in the United States and its possible associations with COVID-19 cases and deaths ⁶, Whittle et al (2020) studied socioeconomic predictors across neighbourhoods in New York ⁷, Pirouz et al (2020) mixed climate and urban parameters for three regions in Italy ⁴

In this study we used a regression analysis between the number of COVID-19 cases and the number of deaths, the dependent variables; and a series of independent variables, such as Mobility, Air Pollution, Health & Research, Economic, Geographical and Social Indicators, to estimate the predictive value of those independent variables and if it changed with time. We also analysed how the predictive value of these variables changed through May and June but using non-cumulative data.

Interestingly, variables related to mobility and air pollution better explain number of infected and dead from COVID-19 than other variables that estimate countries health system's quality or economic strength.

MATERIALS AND METHODS

We carried out a regression analysis, trying to unveil which characteristics make a country more vulnerable to Covid19.

The chosen dependent variables correspond to two of the most significant and worrisome characteristics of any outbreak: total number of infected people and total death toll. Data was collected from the World Health Organisation (WHO) Coronavirus Disease (COVID-19) Dashboard⁴¹ (retrieved from https://covid19.who.int the 21st of: March, April, May and June).

This study focuses on the predictive ability -or not- of the chosen variables, regarding infected and death counts. Calculations were performed using data in two different ways: cumulative count and non-cumulative count. Cumulative count data implies that figures from one period include previous ones. Non-cumulative count represent exclusively figures from the period of time studied (i.e. May period starts on the 21st of April and ends on the 21st May; June period starts on the 21st of May and ends on the 21st of June).

We first analysed how number of infected predicts number of deaths and its evolution through time. If there was a significant improvement in treatment protocols, its predictive value should change.

The predictive variables were selected searching for relations that may explain the differences within countries. Mobility easiness also implies an easy way to spread pathogens, we selected Inbound Tourism as an international mobility indicator. One Health speaks about Human Health, Animal Health and Environmental Health: we selected Air Pollution indicators such as CO₂, NO₂ and methane emissions, as they may show an influence on the disease incidence. Health

indicators (e.g. Hospital Bed Density -beds per 1000- or Physicians per 1000 people, Researches in R & D). Economic indicators as different outcomes could be influenced by the countries' wealth. Social and geographic indicators to study any pattern that explained countries differences (e.g. population, surface area, urban population...).

All data was obtained via internet repositories open to public access, such as The World Health Organization, The Central Intelligence Agency or The World Bank. The complete list of variables, as well as the link to each data set is presented in Table 1.

Table 1 List of predictive variables and data source

Predictive Variable	Source
	International Mobility Indicators
Inbound Tourism	https://www.unwto.org/es/country-profile-inbound-tourism
	Air Pollution Indicators
Electricity Consumption	https://www.cia.gov/library/publications/the-world-factbook/fields/253rank.html
Carbon dioxide emissions from consumption of energy	https://www.cia.gov/library/publications/the-world-factbook/fields/274rank.html
Nitrous oxide emissions (thousand metric tons of CO ₂ equivalent)	https://data.worldbank.org/indicator/EN.ATM.NOXE.KT.C
Methane emissions (kT of CO ₂ equivalent)	https://data.worldbank.org/indicator/EN.ATM.METH.KT.C
	Health and Research Indicators
Hospital Bed Density (beds per 1000 people)	https://www.cia.gov/library/publications/the-world-factbook/fields/360.html
Physicians (per 1,000 people)	https://data.worldbank.org/indicator/SH.MED.PHYS.ZS
Researches in R & D (per million people)	https://data.worldbank.org/indicator/SP.POP.SCIE.RD.P6

Economic Indicators						
Unemployment rate	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/220rank.html					
Inflation rate (consumer prices)	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/229rank.html					
Government expenditure on education, total (% of GDP)	https://data.worldbank.org/indicator/SE.XPD.TOTL.GD.ZS					
Military expenditures	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/330rank.html					

	Social and Demographic Indicators							
Population:	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/335rank.html							
Area:	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/279rank.html							
Urban Population	https://data.worldbank.org/indicator/SP.URB.TOTL.IN.ZS?vie w=chart							
Median age	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/343rank.html							
Population ages 65 and above	https://data.worldbank.org/indicator/SP.POP.65UP.TO.ZS							
Proportion of seats held by women in national parliaments	https://data.worldbank.org/indicator/SG.GEN.PARL.ZS							
Population Growth rate	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/344rank.html							
Infant Mortality Rate	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/354rank.html							
Life expectancy at birth	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/355rank.html							
Obesity - Adult prevalence rate	https://www.cia.gov/library/publications/resources/the-world-factbook/fields/367rank.html							

The territories for this study were the first 50 countries (which represent nearly 70% of world's population) with more inbound tourism (last data available from 2018), according to the World Tourism Organization's webpage⁴² (https://www.unwto.org/country-profile-inbound-tourism). In these countries, very different to each other but with lots of people entering from everywhere, the likelihood of a quick SARS-CoV-2 arrival was very high.

Regression analysis allows to establish a relation between two variables, a dependent variable y_i and an independent variable x_i . This linear regression follows the model:

$$y_i = x_i \beta + \alpha + \epsilon_i$$

and was performed using the statistic package StatPlus^{:mac} (AnalystSoft Inc.)

RESULTS & DISCUSSION

Before any further analysis we studied the relation between the number of dead by SARS-COV-2 (y_i = dependent variable) and the number of infected people (x_i = predictive variable). Among all the studied parameters, these had the highest significant regression (p<0.00001) consistently through the period studied (Cumulative March: R²=0.80; C. April: R²=0.76; C. May: R²=0.99; C. June: R²=0.85).

The coefficient of determination (R²) is the proportion of the variance in the dependent variable that is predictable from the independent variable (e.g. in C. May 99% of the deaths were explained by the number of infected people).

Furthermore, when non-cumulative data was used results were as significant and surprising as before (Non-Cumulative May: R²=0.85; N-C. June: R²=0.85). We were expecting some change as health care systems learnt how to fight the virus. This may appear as an obvious conclusion, but it gains relevance when you consider that other parameters studied (i.e. health and research indicators, country richness...) had no significance, this is in line with Walach et al (2020) findings, that none of the variables related to Health or other population parameters were predictive⁴³. No matter what country you choose (rich, poor, with a good or not so good health system, etc.) the number of infected people is going to rule the number of deaths. This drives the idea that what really matters to fight COVID-19 pandemic is to prevent infections among their citizens rather than better health care systems or richness or development. This may be the reason behind why poorer countries with less developed health care systems did so well fighting this pandemic outbreak (e.g. Viet Nam).

We present variables divided into clearly predictive and not predictive. As interesting is to know which ones can predict as it is to know the ones that do not predict at all.

1. Variables that are clearly predictive of the covid-19 pandemic (statistically significant linear regression p < 0.05):

Inbound tourism was the second variable with greater significance. It remains significant with the number of infected people during the cumulative four-month period studied (C. March: p<0.00001, R²=0.34; C. April: p<0.00001, R²=0.42; C. May: p<0.00006, R²=0.29; C. June: p<0.0003 R²=0,17) and with non-cumulative data (N-C May: p<0.005, R²=0.15; N-C June: p<0.0003 R²=0.17). Our results are concordant with the results obtained by Aldibasi et al (2020)⁴⁴. Daon et al (2020) suggest that an additional prominent reason for the especially wide and rapid spread of COVID-19 is likely to be the current high prevalence of international travel⁴⁵. According to Papadopoulus et al. (2020) early international travel restrictions may help control the pandemic⁴⁶, and Walch et al (2020) found that border closure had the potential of preventing cases⁴³. For Chinazzi et al. (2020) most of the imported cases outside China have likely originated from air travel⁴⁷ and Linka et al. suggest that the spread of COVID-19 in Europe closely followed

air travel patterns and that the severe travel restrictions implemented there resulted in substantial decreases in the disease's spread⁴⁸. Kubota et al (2020) found that relative frequency of foreign visitors per population was positively correlated⁴⁹ and Coelho et al. (2020) emphasized the role of the air transportation network in this epidemic⁵⁰. Gross et al studying spatial dynamics of the COVID-19 in China found a strong correlation between the number of infected individuals in each province studied and the population migration from Hubei to those provinces⁵¹ with a slight decay along time, which agrees with our results although remaining highly significant. Tuli et al (2020) show in their model that countries with higher air passenger traffic have higher likelihood of getting higher level of infections⁵². Lockdown-type measures had the largest effect on limiting viral transmission, followed by complete travel ban⁵³

Inbound tourism also has a significant association with the other dependent variable, number of dead people and stays significant with both cumulative (C. March: p<0.0003, R²=0.24; C. April: p<0.00001, R²=0.63; C. May: p<0.00001, R²=0.46; C. June: p=0.00001, R²=0.35) and non-cumulative data (N-C May: p<0.0001, R²=0.27; N-C June: p<0.00001, R²=0.35).

This means that the number of people arriving to a country is linked to number of infected or dead by SARS-CoV-2. This is concordant with the results obtained by Anzai et al.⁵⁴. Merler et al. conducted a study to see the effect of travel restrictions on the spread of the virus, and they found that the travel quarantine of Wuhan delayed the overall epidemic progression by only 3 to 5 days in mainland China but had a more marked effect on the international scale, where case importations were reduced by nearly 80% until mid-February ⁵⁵.

Other results that we estimated relevant was the strong regression found between infected and dead by COVID-19 and independent variables that talk about air pollution.

Electricity consumption had high statistical significance with both dependent variables, number of infected people and number of deaths, and were statistically significant with cumulative data and non-cumulative data. Other environmental indicators (CO₂ emissions, NO₂ emissions and methane emissions) also showed high predictive power since they had statistical significance (with p<0.001 values) Countries with more emissions and more energy usage had higher counts of infected or dead people (Results are shown on Table 2 and Table 3).

Although Travaglio et al (2020) worked only with data from England their findings are similar to ours since they conclude that the levels of some air pollutants are linked to COVID-19 cases and morbidity⁴⁰.

Similarly, Liu et al. (2020) studied the association of NO₂ atmospheric level over the spread of the COVID-19 in Chinese cities, suggesting that ambient NO₂ may contribute to the spread of COVID-19⁵. Yao et al. (2020) although studying the effect of other pollution indicators (PM_{2.5} and PM₁₀) found that COVID-19 held higher death rates with increasing concentration of PM_{2.5} and PM₁₀⁵⁶. We cannot forget that pollution variables are somehow related with transport (e.g. airplanes, trains and cars)

Population and Area of the countries studied had different predictive power along the studied period. On the 21st of March (C. March), both, Population and Area had high correlation with number of infected people (p=0.00007 and p=0.0043 respectively) but on the consecutive months (C. April, C. May and C. June), while Population had no statistical significance, the Area variable retained predictive power for the number of infected people. The p values for Area were p=0.009 on C. April, p=0.0002 on C. May and p=0.0004 on C. June. Similarly, during N-C May and N-C June Population variable lost its predictive power and had no significance but Area did (N-C May: p=0.00002, R²=0,31; N-C June: p=0.00004, R²=0,30).

When we considered the relation of these two variables with the number of deaths, we found similar outputs, on the 21st of March (C. March) both had statistical significance (Population: p=0.014; Area: p=0.0053), but on the following cumulative months the Population variable was no longer significant while Area still was (p=0.04 on C. April; p=0.007 on C. May; p=0.0015 on C. June). Again, during N-C May and N-C June, Population loses its significance while Area retains it (N-C May: p=0,004; N-C June: p=0.001)). Aldibasi (2020) found a statistically significant positive association between Area(km²) and mortality rate and a negative association between Population size and incidence rate⁴⁴ (Results are shown on Table 2 and Table 3).

Table 2: $\underline{Cumulative\ Data}$. Variables that are clearly predictive of the covid-19 pandemic (statistically significant linear regression p < 0.05)

	I. CUMULATI		BLES THAT ARE CLE atistically significant lin 21 st of MAl	near regression p < 0.0		19 PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients
			Dead vs. infec	cted covid-19			
dead by covid-19	infected by covid-19	194.20	1 48	<0.00001 ***	0.90	0.80	$\alpha = -44.44$ $\beta = 0.0531$
			International Mo	bility Indicators			
infected by covid-19	Inbound tourism arrivals (in millions)	24.73	1 48	0.00001 ***	0.58	0.34	α=-3605.16 β=404.06
dead by covid-19	Inbound tourism arrivals (in millions)	15.23	1 48	0.0003 ***	0.49	0.24	α=-206 β=20.15
			Air Pollution	1 Indicators		•	,
infected by covid-19	electricity consumption	10.76	1 48	<0.00001 ***	0.69	0.47	α=1407.39 β<0.00001
dead by covid-19	electricity consumption	10.35	1 43	0.0023 ***	0.42	0.17	α=101.56 β<0.00001
infected by covid-19	Carbon dioxide emissions	55.34	1 48	<0.00001 ***	0.73	0.54	α=1604.23 β=0.00001
dead by covid-19	Carbon dioxide emissions	12.83	1 48	0.0008***	0.46	0.21	α=104.38 β<0.00001
infected by covid-19	Nitrous oxide emissions	29.32	1 48	<0.00001 ***	0.65	0.43	α=26250 β=0.4545
dead by covid-19	Nitrous oxide emissions	8.59	1 39	0.0056**	0.43	0.18	α=125.52 β= 0.0939
infected by covid-19	Methane emissions	28,62	1 40	<0.00001 ***	0.65	0.42	α=35022 β=0.1012
dead by covid-19	Methane emissions	8.62	1 39	0.0055**	0.42	0.18	α=135 β=0.00133
			Social and Geogr	aphic Indicators			
infected by covid-19	Population	10.90	1 48	0.00007***	0.53	0.28	α=2520 β=0.00003
dead by covid-19	Population	6.48	1 48	0.014*	0.34	0.12	α=134 β<00001
infected by covid-19	Area	7,70	1 48	0.0049 ***	0.39	0.1153	α=2626 β=0.0015
dead by covid-19	Area	8.51	1 48	0.0053 **	0.39	0.15	α=78.14 β=0.00009

	I. CUMULATIV		BLES THAT ARE CLE atistically significant lin 21st of API	near regression p < 0.0		9 PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients
W /	<u>, , , , , , , , , , , , , , , , , , , </u>		Dead vs. infec	ted covid-19			
dead by covid-19	infected by covid-19	136.20	1 43	<0.00001 ***	0.87	0.76	$\alpha = -35423$ $\beta = 3580$
			International Mo	bility Indicators			
infected by covid-19	Inbound tourism arrivals (in millions)	31.08	1 43	0.00001 ***	0.65	0.42	α=786 β=0.057
dead by covid-19	Inbound tourism arrivals (in millions)	73.49	1 43	0.00001 ***	0.79	0.63	α=-3136 β=285
	, ,		Air Pollution	Indicators		•	,
infected by covid-19	electricity consumption	20.76	1 43	0.00001 ***	0.57	0.33	α=18774 β<0.00001
dead by covid-19	electricity consumption	8.32	1 43	0.006 ***	0.40	0.16	α=2162 β<0.00001
infected by covid-19	Carbon dioxide emissions	8.71	1 43	0.005 ***	0.41	0.17	α=28961 β=0.00003
dead by covid-19	Carbon dioxide emissions	3.37	1 43	0.0735	0.27	0.07	α=2690 β<0.00001
infected by covid-19	Nitrous oxide emissions	7.99	1 43	0.007 **	0.40	0.16	α=26250 β=0.4545
dead by covid-19	Nitrous oxide emissions	3.24	1 43	0.08 n.s.	0.26	0.07	α =2558 β = 0.01974
infected by covid-19	Methane emissions	2.85	1 43	0.10 n.s.	0.25	0.06	α=35022 β=0.1012
dead by covid-19	Methane emissions	0.90	1 43	0.34 n.s.	0.14	0.02	α=3017 β=0.00377
	1		Social and Geogra	aphic Indicators		-	•
infected by covid-19	Population	1.09	1 43	0.30 n.s.	0.157	0.024	α=40190 β=0.00007
dead by covid-19	Population	0.32	1 43	0.57 n.s.	0.086	0.007	α=3221 β<00001
infected by covid-19	Area	7,45	1 43	0.009 **	0.384	0.147	α=22664 β=0.01185
dead by covid-19	Area	4.31	1 43	0.04 *	0.30	0.09	α=2211 β=0.00006

	I. CUMULATIV		BLES THAT ARE CLE atistically significant lin 21st of MA	near regression p < 0.0		9 PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients
W 7	. , ,		Dead vs. infec	ted covid-19			
dead by covid-19	infected by covid-19	4912108	1 48	<0.00001 ***	0.99	0.99	α=-0.22617 β=0.02910
			International Mo	bility Indicators			
infected by covid-19	Inbound tourism arrivals (in millions)	19.52	1 48	0.00006 ***	0.54	0.29	α =-43061 β =05702
dead by covid-19	Inbound tourism arrivals (in millions)	40.22	1 48	<0.00001 ***	0.67	0.46	α=-4900 β=485
	, , ,		Air Pollution	Indicators		•	•
infected by covid-19	electricity consumption	22.47	1 48	0.00002 ***	0.56	0.32	α=34430 β<0.00001
dead by covid-19	electricity consumption	14.00	1 48	0.00049 ***	0.47	0.23	α=3116 β<0.00001
infected by covid-19	Carbon dioxide emissions	9.20	1 48	0.004 **	0.40	0.16	α=53388 β=0.00005
dead by covid-19	Carbon dioxide emissions	5.25	1 48	0.026*	0.31	0.10	α=4383 β<0.00001
infected by covid-19	Nitrous oxide emissions	3.31	1 39	0.08 n.s.	0.28	0.08	α=48735 β=0.2236
dead by covid-19	Nitrous oxide emissions	0.51	1 39	0.48 n.s.	0.11	0.013	α=3930 β= 0.009
infected by covid-19	Methane emissions	4.91	1 39	0.03*	0.33	0.11	α=47540 β=0.09
dead by covid-19	Methane emissions	0.21	1 39	0.65 n.s.	0.07	0.005	α=4083 β=0.0019
	-		Social and Geogra	aphic Indicators		-	,
infected by covid-19	Population	1.62	1 48	0.20 n.s.	0.18	0.032	α=70907 β=0.00015
dead by covid-19	Population	0.68	1 48	0.41 n.s.	0.12	0.014	α=5450 β<0.00001
infected by covid-19	Area	15,38	1 48	0.0002 ***	0.49	0.243	α=29559 β=0.02954
dead by covid-19	Area	7.86	1 48	0.007 **	0.37	0.14	α=3202 β=0.0015

	I. CUMULATIV		LES THAT ARE CLE tistically significant lin 21st of JU	near regression p < 0.0		9 PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients
			Dead vs. infec	ted covid-19			•
dead by covid-19	infected by covid-19	269.83	1 48	<0.00001 ***	0.92	0.85	α=977.02 β=0.05189
			International Mo	bility Indicators			
infected by covid-19	Inbound tourism arrivals (in millions)	965	1 48	0.0003 ***	0.41	0.17	α=-10030 β=6857
dead by covid-19	Inbound tourism arrivals (in millions)	25.49	1 48	=0.00001 ***	0.59	0.35	α=-4108 β=556
			Air Pollution	Indicators		•	
infected by covid-19	electricity consumption	18.57	1 48	0.00008 ***	0.53	0.28	α=68670 β<0.00001
dead by covid-19	electricity consumption	14.43	1 48	0.00041 ***	0.48	0.23	α=4588 β<0.00001
infected by covid-19	Carbon dioxide emissions	7.77	1 48	0.0075 **	0.37	0.14	α=96950 β=0.00007
dead by covid-19	Carbon dioxide emissions	5.41	1 48	0.024*	0.32	0.10	α=6196 β<0.00001
infected by covid-19	Nitrous oxide emissions	5.49	1 39	0.024 *.	0.35	0.12	α=81526 β=0.0683
dead by covid-19	Nitrous oxide emissions	2.42	1 39	0.13 n.s.	0.24	0.06	α=5218 β=0.0262
infected by covid-19	Methane emissions	5.74	1 39	0.021*	0.36	0.13	α=82736 β=0.2293
dead by covid-19	Methane emissions	1.37	1 39	0.25 n.s.	0.18	0.003	α=5544 β=0.0067
			Social and Geogra	aphic Indicators			
infected by covid-19	Population	2.92	1 48	0.09 n.s.	0.24	0.06	α=112655 β=0.00031
dead by covid-19	Population	1.45	1 48	0.23 n.s.	0.17	0.03	α=7215 β=0.00001
infected by covid-19	Area	20.40	1 48	0.00004 ***	0.55	0.30	α=45752 β=0.5175
dead by covid-19	Area	11.24	1 48	0.0015 **	0.43	0.19	α=4051 β=0.00232

Table 3: Non-Cumulative Data. Variables that are clearly predictive of the covid-19 pandemic (statistically significant linear regression p < 0.05)

	II. NON-CUMULA			CLEARLY PREDICT		D-19 PANDEMIC	
		(sta	itistically significant li Non-Cumul	near regression p < 0.0 lative MAY	05)		
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R2	α and β coefficients
* '			Dead vs. infec	cted Covid-19			•
dead by covid-19	infected by covid-19	277	1 48	<0.000001***	0.92	0.85	α=161 β=0.067
			International Mo	bility Indicators			
infected by covid-19	Inbound tourism arrivals (in millions)	8.55	1 48	0.0052 ***	0.39	0.15	α=7108 β=2155
dead by covid-19	Inbound tourism arrivals (in millions)	17.61	1 48	0.0001 ***	0.52	0.27	α=-3136 β=285
	, , , , , , , , , , , , , , , , , , ,		Air Pollution	n Indicators		•	
infected by covid-19	electricity consumption	17.77	1 48	0.0001 ***	0.52	0.27	α=16750 β=0.00001
dead by covid-19	electricity consumption	15.16	1 48	0.0003 ***	0.49	0.24	α=1274 β=0.00001
infected by covid-19	Carbon dioxide emissions	7.26	1 48	0.009 ***	0.36	0.13	α=26171 β=0.00002
dead by covid-19	Carbon dioxide emissions	5.39	1 48	0.02*	0.32	0.10	α=1967 β=0.00001
infected by covid-19	Nitrous oxide emissions	2.42	1 39	0.13 n.s.	0.24	0.06	α=21943 β=0.1315
dead by covid-19	Nitrous oxide emissions	1.51	1 39	0.22 n. s.	0.19	0.037	α =1512 β = 0.00594
infected by covid-19	Methane emissions	4,59	1 39	0.04*	0.32	0.10	α=20458 β=0.059
dead by covid-19	Methane emissions	0.69	1 39	0.41 n. s.	0.13	0.018	α=1604 β=0.00138
			Social & Geogra	aphic Indicators		_	
infected by covid-19	Population	1.72	1 48	0.19 n. s.	0.19	0.034	α=-33384 β=0.00008
dead by covid-19	Population	0.80	1 48	0.37 n. s.	0.128	0.016	α=2542 β<0.00001
infected by covid-19	Area	21,94	1 48	0.00002 ***	0.56	0.31	α=7912 β=0.0175
dead by covid-19	Area	8.81	1 48	0.004 **	0.39	0.16	α=1236 β=0.0009

	II. NON-CUMULA			CLEARLY PREDICT near regression p < 0.0 ative JUNE		D-19 PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients
(4/ /			Dead vs. infec	ted Covid-19		•	•
dead by covid-19	infected by covid-19	269	1 48	<0.000001***	0.92	0.85	α=6197 β=16.35
			International Mo	bility Indicators			
infected by covid-19	Inbound tourism arrivals (in millions)	9.65	1 48	0.0003 ***	0.41	0.17	α=-10030 β=6858
dead by covid-19	Inbound tourism arrivals (in millions)	25.49	1 48	0.00001 ***	0.59	0.35	α=-4108 β=556
			Air Pollution	n Indicators		•	,
infected by covid-19	electricity consumption	18.57	1 48	0.0008 ***	0.53	0.27	α=66670 β<0.00001
dead by covid-19	electricity consumption	14.43	1 48	0.0004 ***	0.48	0.23	α=4588 β<0.00001
infected by covid-19	Carbon dioxide emissions	7.76	1 48	0.007 ***	0.37	0.14	α=96950 β=0.00007
dead by covid-19	Carbon dioxide emissions	5.40	1 48	0.02 *	0.32	0.10	α=6198 β<0.00001
infected by covid-19	Nitrous oxide emissions	5.49	1 39	0.02 *	0.35	0.12	α=81526 β=0.6683
dead by covid-19	Nitrous oxide emissions	2.42	1 39	0.12 n.s.	0.24	0.058	α =5218 β = 0.026225
infected by covid-19	Methane emissions	5,73	1 39	0.02 *	0.36	0.13	α=82776 β=0.22929
dead by covid-19	Methane emissions	1.37	1 39	0.25 n.s.	0.18	0.033	α=5544 β=0.00672
			Social & Geogra	phic Indicators			
infected by covid-19	Population	2.92	1 48	0.09 n.s.	0.24	0.006	α=-112655 β=0.00031
dead by covid-19	Population	0.25	1 48	0.61 n.s.	0.08	0.006	α=4541 β=130
infected by covid-19	Area	20,40	1 48	0.00004 ***	0.55	0.30	α=45752 β=0.05175
dead by covid-19	Area	11.23	1 48	0.001 ***	0.43	0.19	α=4051 β=0.0023

2. Variables that are not predictive of the covid-19 pandemic (non-statistically significant linear regression p > 0.05)

None of the variables included in this group had statistical significance during the time period studied, from 21st of March to the 21st of June. The regression analysis with Hospital Bed Density (beds per 1000 people), Number of Physicians (per 1000 people) had no correlation with number of infected nor number of deaths at any moment in our study, either using cumulative data or noncumulative data. When confronted with this new pandemic, health care systems showed little efficiency in preventing infections and death counts, due to probably lack of knowledge. We thought this should change with time, their predictive power had to grow as health care systems dealt with SARS-CoV-2, but astonishingly, after several months since the outbreak, all this relevant health parameters remained irrelevant. What proved to be more effective was early preventive measures. Lowental et al (2020) point in the same direction, countries that implemented early measures to limit population mixing limited the number of deaths, in this same study they explain that a delay of 7,49 days in initialling social distancing would double the number of deaths.⁵⁷. For Papadopoulus et al (2020) early introduction of these policies are of greater importance than its stringency.46

Researches in R&D (per million people) or Infant Mortality Rate (indicator of a countries health and care system) showed similar behaviour. In the same line, Yao et al. (2020) did not find significance in the association between hospital beds per capita and COVID-19 death rate⁵⁶. Sorci et al. (2020) found that the Number of hospital beds per 1,000 inhabitants was negatively associated with Case Fatality Rate ⁵⁸. Qiu et al (2020) in their study in China, found that COVID-19 transmission is negatively moderated by the number of doctors at the city level⁵⁹

The logic of any disease is that good health and research parameters of a country should have a positive effect over the number of infected or the number of deaths, but our results indicate no significance at all. It is necessary to highlight that the best predictor we found to death count by COVID-19 is the number of infected by COVID-19 (p>0,00001), either with cumulative data or with non-cumulative data, there is a cause-effect relation. Despite the increase in knowledge, despite improvements in health care systems... the percentage of deaths for a given number of infected could not be reduced. What it really shows

is that prevention, and not treatment, is the best way to face the COVID-19 pandemic.

Unemployment rate, Inflation rate and Military expenditure, categorised as economic indicators fell also in this group of non-predictive variables. We did not find statistically significant regressions between the different economic indicators we selected and the dependent variables (number of infected and number of deaths). Richness of a country does not predict its infected or dead by COVID-19, with the exception of those that showed easiness to travel across the country (i.e. Inbound tourism). Yao et al. among other variables they also studied Gross Domestic Product (GDP) per capita as a measurement of a country's wealth and in their work, GDP per capita was found not to have significance in the association with COVID-19 death rate⁵⁶. According to Qiu et al. (2020) COVID-19 transmission is positively moderated by GDP per capita⁵⁹

In our study, Urban population (%) and Population Growth rate were also not statistically significant in relation with number of infected people and number of deaths by SARS-CoV-2.

Government expenditure on education (as % of GDP) had some correlation with number of infected and number of deaths by SARS-CoV-2 on the 21st of March but not the following two months.

The Population ages 65 and above in our study had no relation with the number of infected on the 21st of March, gained significance on the 21st of April (p=0.06) and went back on the 21st of May to be not statistically significant. When we studied Population ages 65 and above against number of deaths it showed no correlation in March or April, but it did in May (p=0.09).

Messner also studied the effect of education and countries' median age on the COVID-19 outbreak and found that the quality of a country's education system is positively associated with the outbreak and that countries with an older population are more affected⁶⁰

Shagam (2020) using multivariate linear regression found significant correlation between incidence and mortality rates with GDP per capita (p = 2.6×10^{-15} and 7.0×10^{-4} , respectively), country-specific duration of the outbreak (2.6×10^{-4} and 0.0019), fraction of citizens over 65 years old (p = 0.0049 and 3.8×10^{-4}) and level of press freedom (p = 0.021 and 0.019). Table 4 and table 5 can be found in complementary material.

Table 4.- $\underline{Cumulative\ Data}$. Variables that are not predictive of the covid-19 pandemic (non-statistically significant linear regression p > 0.05)

	III: CUMULA		IABLES THAT ARE statistically significant 21st of MAI	linear regression p >		PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	\mathbb{R}^2	α and β coefficients
			Health & Resea	rch Indicators			
infected by covid-19	Hospital Bed Density	0.07	1 36	0.78 n.s.	0.05	0.00	α=5059 β=-248
dead by covid-19	Hospital Bed Density	0.16	1 35	0.70 n.s.	0.07	0.004	α =380.10 β =-21.19
infected by covid-19	Physicians (per 1,000 people)	0.19	1 39	0.66 n.s.	0.07	0.001	α=24452 β=8703
dead by covid-19	Physicians (per 1,000 people)	0.16	1 39	0.69 n.s.	0.06	0.004	α=179 β=43.44
infected by covid-19	Researches in R & D	0.52	1 43	0.48 n.s.	0.22	0.05	α=174 β=4946
dead by covid-19	Researches in R & D	0.51	1 43	0.46 n.s.	0.20	0.04	$\alpha = 176$ $\beta = 4427$
			Economic 1	ndicators			
infected by covid-19	Unemployment rate	0.04	1 48	0.83 n.s.	0.03	0.001	α=4995 β=85
dead by covid-19	Unemployment rate	0.84	1 48	0.36 n.s.	0.13	0.02	α=96.92 β=21.71
infected by covid-19	Inflation rate	0.52	1 48	0.47 n.s.	0.10	0.01	α=6671 β=-293
dead by covid-19	Inflation rate	0.19	1 48	0.66 n.s.	0.06	0.004	α=291 β=-10.47
infected by covid-19	Expenditure on education	3.78	1 39	0.06	0.30	0.09	α=21704 β=-3270
dead by covid-19	Expenditure on education	3.17	1 43	0.08	0.27	0.07	α=1146 β=-180
infected by covid-19	Military expenditures	0.16	1 48	0.69 n.s.	0.06	0.001	α=6743 β=-587
dead by covid-19	Military expenditures	0.21	1 48	0.65 n.s.	0.07	0.004	α=329 β=-39

			Social and Geogr	aphic Indicators			
infected by covid-19	Urban Population	0.06	1 39	0.81 n.s.	0.04	0.001	α=-8756 β=-34
dead by covid-19	Urban Population	0.07	1 39	0.79 n.s.	0.04	0.001	$\alpha = -459$ $\beta = -2.24$
infected by covid-19	Median age	2.80	1 48	0.10 n.s.	0.23	0.05	α=-13175 β=503
dead by covid-19	Median age	1.88	1 48	0.17 n.s.	0.19	0.04	α=-667 β=24.66
infected by covid-19	Population ages 65 and above	0.72	1 39	0.39 n.s.	0.13	0.02	α=1959 β=312
dead by covid-19	Population ages 65 and above	1.87	1 48	0.18 n.s.	0.19	0.04	α=-667 β=24.66
infected by covid-19	Proportion of women in national parliaments	0.22	1 39	0.64 n.s.	0.07	0.005	α=3475 β=104
dead by covid-19	Proportion of women in national parliaments	0.21	1 39	0.65 n.s.	0.07	0.005	α=-134 β=6.02
infected by covid-19	Population Growth rate	1.1	1 48	0.20 n.s.	0.15	0.022	$\alpha = 7756$ $\beta = -3411$
dead by covid-19	Population Growth rate	0.65	1 48	0.42 n.s.	0.11	0.013	α=351 β=-156
infected by covid-19	Infant Mortality Rate	0.78	1 48	0.38 n.s.	0.13	0.016	α=7572 β=-201
dead by covid-19	Infant Mortality Rate	0.65	1 48	0.42 n.s.	0.12	0.013	α=351 β=-156
infected by covid-19	Life expectancy at birth	1.06	1 48	0.31 n.s.	0.15	0.021	α=-28178 β=434
dead by covid-19	Life expectancy at birth	0.70	1 48	0.40 n.s.	0.12	0.01	α=-1388 β=21.12
infected by covid-19	Obesity - Adult prevalence rate	1.70	1 39	0.20 n.s.	0.20	0.004	α=-14009 β=-372
dead by covid-19	Obesity - Adult prevalence rate	0.64	1 39	0.43 n.s.	0.13	0.016	α=583 β=-13.80

II: CUMULATIVE DATA, VARIABLES THAT ARE NOT PREDICTIVE OF THE COVID-19 PANDEMIC (non-statistically significant linear regression p > 0.05) 21st of APRIL 2020 dependent variable predictive variable d. f. regression \mathbb{R}^2 P F value R α and β coefficients (v) (x) residue Health & Research Indicators $\alpha = 55993$ infected by covid-19 Hospital Bed Density 0.03 0.85 n.s. 0.03 0.0006 43 $\beta = -1195$ dead by $\alpha = 4105$ Hospital Bed Density 0.03 0.84 n.s. 0.03 0.0010 covid-19 43 $\beta = -86.81$ Physicians (per 1,000 $\alpha = 24452$ infected by covid-19 0.44 0.51 n.s. 0.10 0.01 43 people) $\beta = 8703$ Physicians (per 1,000 dead by $\alpha = 918$ 0.03 1.30 0.26 n.s. 0.17 covid-19 people) 43 $\beta = 961$ 1 $\alpha = 1461$ infected by covid-19 Researches in R & D 0.53 0.47 n.s. 0.11 0.01 43 $\beta = 5.71$ dead by $\alpha = 1313$ 0.19 0.04 Researches in R & D 1.74 0.19 n.s. covid-19 43 $\beta = 1427$ **Economic Indicators** α=47418 infected by covid-19 0.0002 0.98 n.s. 0.002 0.023 Unemployment rate 43 $\beta = 48.05$ dead by $\alpha = 2240$ Unemployment rate 0.43 0.43 n.s. 0.119 0.014 covid-19 43 $\beta = 176$ α=53445 infected by covid-19 Inflation rate 0.24 0.63 n.s. 0.074 0.005 43 $\beta = 1768$ dead by 1 $\alpha = 3970$ Inflation rate 0.40 0.53 n.s. 0.096 0.009 covid-19 43 $\beta = -147$ Expenditure on $\alpha = 42032$ infected by covid-19 0.21 0.65 n.s. 0.015 0.123 education 43 $\beta = -3265$ dead by Expenditure on $\alpha = 1281$ 0.0008 0.97 n.s. 0.007 0.0001 covid-19 43 education $\beta=10$ 1 $\alpha = 30854$ infected by covid-19 Military expenditures 0.54 0.46 n.s. 0.111 0.012 43 $\beta = 8962$ dead by $\alpha = 3.327$ Military expenditures 0.012 0.91 n.s. 0.016 0.000 covid-19 43 $\beta = 87$

			Social & Geogra	phic indicators			
infected by covid-19	Urban Population	0.85	1 43	0.36 n.s.	0.139	0.019	α=-20436 β=941
dead by covid-19	Urban Population	1.28	1 43	0.26 n.s.	0.169	0.028	α=-1901 β=74
infected by covid-19	Median age	1.36	1 43	0.25 n.s.	0.175	0.030	α=-64791 β=3004
dead by covid-19	Median age	2.99	1 43	0.09 n.s.	0.255	0.065	α=-7165 β=284
infected by covid-19	Population ages 65 and above	1.47	1 43	0.23 n.s.	0.162	0.053	α=4325 β=5172
dead by covid-19	Population ages 65 and above	3.83	1 43	0.06 n.s.	0.286	0.081	α=-935 β=323
infected by covid-19	Proportion of women in national parliaments	0.241	1 43	0.63 n.s.	0.074	0.005	α=26128 β=804
dead by covid-19	Proportion of women in national parliaments	2.77	1 43	0.10 n.s.	0.246	0.060	α=-1128 β=172
infected by covid-19	Population Growth rate	0.21	1 43	0.65 n.s.	0.069	0.004	α=56011 β=-12647
dead by covid-19	Population Growth rate	0.46	1 43	0.50 n.s.	0.103	0.011	α=4286 β=-1220
infected by covid-19	Infant Mortality Rate	1.27	1 43	0.26 n.s.	0.169	0.028	α=4286 β=-1220
dead by covid-19	Infant Mortality Rate	2.36	1 43	0.13 n.s.	0.228	0.052	α=5284 β=-186
infected by covid-19	Life expectancy at birth	1.44	1 43	0.23 n.s.	0.179	0.032	α=-282391 β=4237
dead by covid-19	Life expectancy at birth	3.31	1 43	0.07 n.s.	0.268	0.072	α=-28426 β=409
infected by covid-19	Obesity - Adult prevalence rate	3.09	1 43	0.09 n.s.	0.259	0.007	α=-25123 β=3440
dead by covid-19	Obesity - Adult prevalence rate	2.15	1 43	0.15 n.s.	0.218	0.048	α=-506 β=188

II: CUMULATIVE DATA, VARIABLES THAT ARE NOT PREDICTIVE OF THE COVID-19 PANDEMIC (non-statistically significant linear regression p > 0.05) 21st of MAY 2020 dependent variable predictive variable d. f. regression P \mathbb{R}^2 F value R α and β coefficients residue (v) (x) Health & Research Indicators $\alpha = 55993$ infected by covid-19 Hospital Bed Density 0.09 0.76 n.s. 0.05 0.0003 36 $\beta = -1195$ dead by $\alpha = 3755$ Hospital Bed Density 0.005 0.94 n.s. 0.01 0.0001 covid-19 35 $\beta = -30.12$ Physicians (per 1,000 $\alpha = 35276$ infected by covid-19 0.87 0.35 n.s. 0.15 0.02 39 people) $\beta = 8492$ Physicians (per 1,000 dead by $\alpha = 1084$ 0.04 1.77 0.19 n.s. 0.21 covid-19 people) 39 $\beta = 1186$ $\alpha = 71910$ infected by covid-19 Researches in R & D 0.53 0.44 n.s. 0.25 0.06 43 $\beta = -779$ dead by $\alpha = 3978$ Researches in R & D 0.10 0.19 n.s. 0.10 0.009 covid-19 40 $\beta = -0.222$ **Economic Indicators** α=91570 infected by covid-19 0.01 0.92 n.s. 0.0001 Unemployment rate 0.01 48 $\beta = -634$ dead by $\alpha = 5226$ Unemployment rate 0.10 0.04 0.002 0.76 n.s. covid-19 48 $\beta = 130$ α=95252 infected by covid-19 Inflation rate 0.13 0.05 0.002 0.71 n.s. 43 $\beta = -2281$ dead by 1 $\alpha = 7130$ Inflation rate 0.40 0.53 n.s. 0.09 0.008 covid-19 48 $\beta = -266$ Expenditure on $\alpha = 72682$ infected by covid-19 0.11 0.05 0.002 0.74 n.s. education 39 $\beta = -3021$ dead by Expenditure on $\alpha = 1281$ 0.0008 0.97 n.s. 0.0001 0.007 covid-19 43 $\beta=10$ education $\alpha = 37127$ infected by covid-19 Military expenditures 1.37 0.25 n.s. 0.16 0.027 48 $\beta = 25781$ dead by $\alpha = 3.327$ 0.012 Military expenditures 0.91 n.s. 0.016 0.000 covid-19 43 $\beta = 87$

			Social & Geogra	phic Indicators			
infected by covid-19	Urban Population	0.83	1 39	0.37 n.s.	0.14	0.020	α=-11100 β=656
dead by covid-19	Urban Population	1.17	1 43	0.29 n.s.	0.17	0.029	α=-1233 β=77
infected by covid-19	Median age	0.36	1 48	0.55 n.s.	0.09	0.007	α=-18867 β=2836
dead by covid-19	Median age	1.44	1 48	0.23 ns	0.17	0.03	α=-8026 β=380
infected by covid-19	Population ages 65 and above	0.39	1 39	0.39 n.s	0.10	0.010	α=42120 β=1176
dead by covid-19	Population ages 65 and above	2.98	1 39	0.09	0.27	0.071	α=-42 β=314
infected by covid-19	Proportion of women in national parliaments	<0.01	1 39	0.99 n.s.	0.00	0.0001	α=58525 β=-2,66
dead by covid-19	Proportion of women in national parliaments	2.98	1 39	0.09	0.27	0.071	α=-42.20 β=314
infected by covid-19	Population Growth rate	0.06	1 48	0.81 n.s.	0.03	0.001	α=94458 β=-12003
dead by covid-19	Population Growth rate	0.12	1 48	0.72 n.s.	0.05	0.003	α=6925 β=-1207
infected by covid-19	Infant Mortality Rate	0.49	1 48	0.49 n.s.	0.10	0.010	α =110832 β =-2449
dead by covid-19	Infant Mortality Rate	1.49	1 48	0.23 n.s.	0.17	0.030	α=8956 β=-287
infected by covid-19	Life expectancy at birth	0.41	1 48	0.52 n.s.	0.09	0.008	α=-236851 β=4166
dead by covid-19	Life expectancy at birth	2.27	1 48	0.14	0.21	0.045	α=-44583 β=652
infected by covid-19	Obesity - Adult prevalence rate	0.16	1 39	0.68 ns	0.06	0.0004	α=45936 β=605
dead by covid-19	Obesity - Adult prevalence rate	0.13	1 39	0.72 n.s.	0.06	0.003	α=3207 β=54

III: CUMULATIVE DATA, VARIABLES THAT ARE NOT PREDICTIVE OF THE COVID-19 PANDEMIC (non-statistically significant linear regression p > 0.05) 21st of JUNE 2020 dependent variable predictive variable d. f. regression P \mathbb{R}^2 F value R α and β coefficients residue (v) (x) Health & Research Indicators $\alpha = 139284$ infected by covid-19 Hospital Bed Density 0.38 0.54 n.s. 0.001 0.10 36 $\beta = -6798$ dead by $\alpha = 7600$ Hospital Bed Density 0.39 0.54 n.s. 0.10 0.01 covid-19 35 $\beta = -373$ Physicians (per 1,000 $\alpha = 135554$ infected by covid-19 0.18 0.68 n.s. 0.07 0.004 39 people) $\beta = -9150$ Physicians (per 1,000 dead by $\alpha = 4818$ 0.21 0.65 n.s. 0.07 0.005 covid-19 people) 39 $\beta = 564$ $\alpha = 244006$ infected by covid-19 Researches in R & D 1.14 0.31 n.s. 0.32 0.10 43 $\beta = -3856$ dead by $\alpha = 10205$ Researches in R & D 0.52 0.49 n.s. 0.22 0.05 covid-19 43 $\beta = -1.27$ **Economic Indicators** $\alpha = 132051$ infected by covid-19 0.04 0.03 0.001 Unemployment rate 0.84 n.s. 48 $\beta = 1986$ dead by $\alpha = 7214$ Unemployment rate 0.12 0.73 n.s. 0.05 0.02 covid-19 48 $\beta = 188$ α=153325 infected by covid-19 Inflation rate 0.04 0.84 n.s. 0.3 0.0008 48 $\beta = -1922$ dead by 1 $\alpha = 9496$ Inflation rate 0.21 0.64 n.s. 0.07 0.004 covid-19 48 $\beta = -255$ Expenditure on 1 $\alpha = 77890$ infected by covid-19 0.10 0.05 0.003 0.75 n.s. education 39 $\beta = 6940$ dead by Expenditure on 1 $\alpha = 2744$ 0.40 0.01 0.53 n.s. 0.10 covid-19 39 education $\beta = 767$ $\alpha = 60594$ infected by covid-19 Military expenditures 1.48 0.23 n.s. 0.17 0.003 48 $\beta = 42273$ dead by $\alpha = 2744$ 0.40 Military expenditures 0.53 n.s. 0.10 0.001 covid-19 48 $\beta = 766$

			Social and Geogr	aphic Indicators			
infected by covid-19	Urban Population	0.38	1 39	0.54 n.s.	0.10	0.001	α=34166 β=1059
dead by covid-19	Urban Population	1.06	1 39	0.31 n.s.	0.16	0.003	α=-890 β=100
infected by covid-19	Median age	0.005	1 48	0.94 n.s.	0.01	0.0001	α =166722 β =545
dead by covid-19	Median age	0.37	1 48	0.54 n.s.	0.9	0.08	α=-1017 β=257
infected by covid-19	Population ages 65 and above	0.77	1 39	0.38 n.s.	0.14	0.02	α=164989 β=-3920
dead by covid-19	Population ages 65 and above	0.26	1 39	0.61 n.s.	0.08	0.007	α=4514 β=131
infected by covid-19	Proportion of women in national parliaments	1.14	1 39	0.29 n.s.	0.17	0.03	α=187929 β=-2851
dead by covid-19	Proportion of women in national parliaments	1.28	1 39	0.26 n.s.	0.18	0.003	α=1686 β=172
infected by covid-19	Population Growth rate	0.002	1 48	0.96 n.s.	0.006	0.00004	α=144141 β=3540
dead by covid-19	Population Growth rate	0.02	1 48	0.90 n.s.	0.02	0.0003	α=8924 β=-561
infected by covid-19	Infant Mortality Rate	0.0002	1 48	0.99 n.s.	0.002	0	α=147091 β=-74
dead by covid-19	Infant Mortality Rate	0.50	1 48	0.48 n.s.	0.10	0.01	α=10708 β=-219
infected by covid-19	Life expectancy at birth	0.002	1 48	0.96 n.s.	0.007	0.00005	α=184296 β=-487
dead by covid-19	Life expectancy at birth	0.93	1 48	0.34 n.s.	0.14	0.02	α=-34683 β=556
infected by covid-19	Obesity - Adult prevalence rate	0.10	1 39	0.75 n.s.	0.05	0.003	α=87007 β=1140
dead by covid-19	Obesity - Adult prevalence rate	0.19	1 39	0.66 n.s.	0.07	0.005	α=5524 β=89

Table 5.- Non-Cumulative Data. Variables that are not predictive of the covid-19 pandemic (non-statistically significant linear regression p > 0.05)

	II. NON-CUM		. VARIABLES THAT I -statistically significant Non-Cumula	linear regression p >		PANDEMIC	
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients
			Health and Resea	arch Indicators			•
infected by covid-19	Hospital Bed Density	0.011	1 35	0.92 n. s.	0.018	0.0003	α=30277 β=-337
dead by covid-19	Hospital Bed Density	0.128	1 35	0.72 n. s.	0.06	0.003	α=1954 β=-62
infected by covid-19	Physicians (per 1,000 people)	0.005	1 39	0.94 n. s.	0.011	0.0001	α=29968 β=-443
dead by covid-19	Physicians (per 1,000 people)	0.45	1 39	0.50 n. s.	0.107	0.011	α=1123 β=237
infected by covid-19	Researches in R & D	0.75	1 10	0.41 n. s.	0.264	0.069	α=49560 β=-7186
dead by covid-19	Researches in R & D	0.24	1 10	0.63 n. s.	0.15	0.024	α =2830 β =-0.28
			Economic I	Indicators			
infected by covid-19	Unemployment rate	0.053	1 48	0.81 n. s.	0.033	0.001	α=47410 β=-745
dead by covid-19	Unemployment rate	0.008	1 48	0.93 n. s.	0.013	0.0001	α=3127 β=-21
infected by covid-19	Inflation rate	0.049	1 48	0.82 n. s.	0.032	0.001	α=46646 β=-720
dead by covid-19	Inflation rate	0.20	1 48	0.65 n. s.	0.065	0.004	α=3363 β=-107
infected by covid-19	Expenditure on education	0.004	1 39	0.95 n. s.	0.010	0.0001	α=25895 β=374
dead by covid-19	Expenditure on education	1.020	1 39	0.32 n. s.	0.159	0.025	α=136 β=346
infected by covid-19	Military expenditures	0.38	1 48	0.54 n. s.	0.088	0.009	α=1975 β=517
dead by covid-19	Military expenditures	0.37	1 48	0.54 n. s.	0.088	0.007	α=1913 β=524

			Social & Geogra	phic Indicators			
infected by covid-19	Urban Population	0.37	1 39	0.54 n. s.	0.097	0.009	α=5913 β=301
dead by covid-19	Urban Population	1.73	1 39	0.19 n. s.	0.206	0.042	α=-829 β=27
infected by covid-19	Median age	0.0006	1 48	0.98 n. s.	0.003	0.0001	α=39814 β=59.67
dead by covid-19	Median age	0.35	1 48	0.55 n. s.	0.085	0.007	α=-972 β=105
infected by covid-19	Population ages 65 and above	0.61	1 39	0.44 n. s.	0.124	0.015	α=41596 β=-1004
dead by covid-19	Population ages 65 and above	0.52	1 39	0.47 n. s.	0.114	0.013	α=1042 β=52
infected by covid-19	Proportion of women in national parliaments	1.97	1 39	0.17 n. s.	0.22	0.05	α=55438 β=-1061
dead by covid-19	Proportion of women in national parliaments	0.98	1 39	0.33 n. s.	0.156	0.024	α=610 β=43
infected by covid-19	Population Growth rate	0.002	1 48	0.95 n. s.	0.007	0.0001	α=42879 β=-1332
dead by covid-19	Population Growth rate	0.011	1 48	0.91 n. s.	0.015	0.0001	α=3102 β=-200
infected by covid-19	Infant Mortality Rate	0.06	1 48	0.80 n. s.	0.035	0.001	α=46375 β=-445
dead by covid-19	Infant Mortality Rate	0.61	1 48	0.44 n. s.	0.112	0.012	α=3987 β=-103
infected by covid-19	Life expectancy at birth	0.0003	1 48	0.99 n. s.	0.002	0.000	α=46719 β=-60
dead by covid-19	Life expectancy at birth	0.95	1 48	0.33 n. s.	0.139	0.019	α=-15620 β=239
infected by covid-19	Obesity - Adult prevalence rate	0.22	1 39	0.64 n. s.	0.07	0.006	α=17756 β=479
dead by covid-19	Obesity - Adult prevalence rate	0.36	1 39	0.55 n. s.	0.096	0.009	α=-1058 β=34

II. NON-CUMULATIVE DATA. VARIABLES THAT NOT PREDICTIVE OF THE COVID-19 PANDEMIC									
(non-statistically significant linear regression p > 0.05) Non-Cumulative JUNE									
dependent variable (y)	predictive variable (x)	F value	d. f. regression residue	P	R	R ²	α and β coefficients		
Health and Research Indicators									
infected by covid-19	Hospital Bed Density	0.38	1 35	0.54 n.s.	0.10	0.001	α=139284 β=-6798		
dead by covid-19	Hospital Bed Density	0.39	1 35	0.54 n.s.	0.10	0.011	α=7600 β=-373		
infected by covid-19	Physicians (per 1,000 people)	0.17	1 39	0.67 n.s.	0.07	0.004	α =135554 β =-9150		
dead by covid-19	Physicians (per 1,000 people)	0.20	1 39	0.65 n. s	0.07	0.005	α=4818 β=564		
infected by covid-19	Researches in R & D	1.31	1 10	0.31 n.s.	0.32	0.10	α=211006 β=-38		
dead by covid-19	Researches in R & D	0.005	1 10	0.94 n.s.	0.023	0.000	α=7025 β =-311		
			Economic 1	Indicators					
infected by covid-19	Unemployment rate	0.041	1 48	0.84 n.s.	0.003	0.001	α=132050 β=-1983		
dead by covid-19	Unemployment rate	0.12	1 48	0.73 n.s.	0.050	0.002	α=7214 β=188		
infected by covid-19	Inflation rate	0.038	1 48	0.85 n.s.	0.028	0.001	α=153325 β=-1922		
dead by covid-19	Inflation rate	0.21	1 48	0.64 n.s.	0.067	0.004	α=9496 β=-255		
infected by covid-19	Expenditure on education	0.169	1 39	0.69 n.s.	0.129	0.017	α=187429 β=-35773		
dead by covid-19	Expenditure on education	0.039	1 39	0.53 n.s.	0.100	0.01'	α=2744 β=768		
infected by covid-19	Military expenditures	1.48	1 48	0.23 n.s.	0.17	0.029	α=64594 β=42273		
dead by covid-19	Military expenditures	0.12	1 48	0.72 n.s.	0.05	0.002	α=7210 β=703		

			Social & Geogra	phic Indicators			
infected by covid-19	Urban Population	0.38	1 39	0.38 n.s	0.098	0.009	α=34166 β=1059
dead by covid-19	Urban Population	1.07	1 39	0.30 n.s	0.16	0.003	α=-890 β=100
infected by covid-19	Median Age	0.005	1 48	0.94 n.s.	0.01	0.0001	α=166722 β=-545
dead by covid-19	Median Age	0.37	1 48	0.54 n.s.	0.088	0.008	α=-1017 β=256
infected by covid-19	Population ages 65 and above	0.76	1 39	0.39 n.s	0.139	0.019	α=164989 β=-3920
dead by covid-19	Population ages 65 and above	1.45	1 48	0.23 n.s	0.171	0.029	α=7215 β=0.00001
infected by covid-19	Proportion of women in national parliaments	1.14	1 39	0.29 n.s.	0.17	0.03	α=187929 β=-2851
dead by covid-19	Proportion of women in national parliaments	1.28	1 39	0.26 n.s.	0.18	0.032	α=1686 β=172
infected by covid-19	Population Growth rate	0.002	1 48	0.96 n.s.	0.006	0.000	α=144141 β=3540
dead by covid-19	Population Growth rate	0.015	1 48	0.90 n.s.	0.002	0.000	α=8924 β=-561
infected by covid-19	Infant Mortality Rate	0.0002	1 48	0.98 n.s.	0.035	0.001	α=147091 β=-74
dead by covid-19	Infant Mortality Rate	0.50	1 48	0.48 n.s.	0.10	0.01	α=10708 β=-219
infected by covid-19	Life expectancy at birth	0.002	1 48	0.96 n.s.	0.006	0.000	α=4186296 β=-488
dead by covid-19	Life expectancy at birth	0.93	1 48	0.34 n.s.	0.138	0.019	α=-30683 β=556
infected by covid-19	Obesity - Adult prevalence rate	0.10	1 39	0.75 n.s.	0.05	0.002	α=87007 β=1140
dead by covid-19	Obesity - Adult prevalence rate	0.19	1 39	0.66 n.s.	0.07	0.005	α=4525 β=88

Statistically, with these 50 countries (nearly 70% world's population) number of dead depends only on number of infected according to:

Number of dead= $\beta*N^\circ$ infected+ α

With α values ranging from -35423 to 6197 and β from 0.02910 to 16.35. The best predictor to death count by SARS-CoV2 was the number of infected (p>0,00001), either with cumulative data or with non-cumulative data, there is a cause effect relation.

The best predictive variable was inbound tourism, and it did not change throughout the time period studied, it retained its predictive strength using cumulative data or non-cumulative data.

Air pollution predictive variables are of great significance, their ability to predict number of infected and dead should make us think about the importance of One Health. Environmental health and Animal health are essential to Human health.

Despite every effort done, the increase in knowledge, the improvements implemented in health care systems, etc. yet is not enough. None of the health & research indicators had significance at all during the four-month cumulative data regressions, nor when non-cumulative data was used. As stated previously, what it really shows is that prevention, and not treatment, is the best way to confront SARS-CoV-2 outbreak, and by extension any future epidemics that may arise unknown to medicine. Some lessons must be learnt from this pandemic. Those countries that conducted more tests to isolate infectious promptly and impose to their citizens policies to prevent the spread of the virus at early stages were, by far, least affected.

According to Loewenthal et al. (2020), to reduce the number of deaths is essential to implement confinement policies at a very early stage. If social distancing is not adopted, death incidents are doubled every 7.49 days.⁵⁷ Piguillem et al. using Italy's early figures to calibrate their model they estimate that without lockdowns there could be as much as 800, 000 fatalities. With a very early intervention that number reduces to about 120 – 230 fatalities, while if the intervention is started 60 days later, the number of fatalities is between 7, 200 and 9, 000⁶¹. Italy had, as to 21st of June, 238,275 infected people and 34,610 dead according to the World Health Organisation COVID-19 dashboard (https://covid19.who.int).

Early detection of cases through surveillance and aggressive contact tracing around known cases has helped to contain spread of the outbreak in Singapore. Together with other healthcare, border and community measures, they allow the COVID-19 outbreak to be managed without major disruption to daily living. Countries could consider these measures for a proportionate response to the risk of COVID-19⁶².

Our study shows, with very high significance, what variables predict SARS-CoV-2 infected and dead counts In order to minimise the impact of future unknown pathogens arising, is paramount that we learn the lesson from this pandemic. Prevention is the key.

ACKNOWLEDGEMENTS

We thank to Dr. C. García-Balboa, H. Díaz-Alejo, P. Martinez-Alesón for their help and advice on this paper.

BIBLIOGRAFY

- 1. Li, K. et al. The Clinical and Chest CT Features Associated with Severe and Critical COVID-19 Pneumonia. *Invest. Radiol.* (2020).
- 2. Zhou, G., Chen, S. & Chen, Z. Back to the spring of Wuhan: facts and hope of COVID-19 outbreak. Frontiers of Medicine (2020). doi:10.1007/s11684-020-0758-9
- 3. Wang, J., Tang, K., Feng, K. & Lv, W. High Temperature and High Humidity Reduce the Transmission of COVID-19. SSRN Electron. J. (2020). doi:10.2139/ssrn.3551767
- 4. Pirouz, B., Haghshenas, S. S., Pirouz, B., Haghshenas, S. S. & Piro, P. Development of an assessment method for investigating the impact of climate and urban parameters in confirmed cases of COVID-19: A new challenge in sustainable development. *Int. J. Environ. Res. Public Health* 17, (2020).
- 5. Liu, J. et al. Impact of meteorological factors on the COVID-19 transmission: A multi-city study in China. *Sci. Total Environ.* **726**, 138513 (2020).
- Guha, A., Bonsu, J., Dey, A. & Addison, D. Community and Socioeconomic Factors Associated with COVID-19 in the United States: Zip code level cross sectional analysis. medRxiv 2020.04.19.20071944

- (2020). doi:10.1101/2020.04.19.20071944
- 7. Whittle, R. S. & Diaz-Artiles, A. An ecological study of socioeconomic predictors in detection of COVID-19 cases across neighborhoods in New York City. *medRxiv* 2020.04.17.20069823 (2020). doi:10.1101/2020.04.17.20069823
- 8. Cheng, Y. et al. Kidney impairment is associated with in-hospital death of COVID-19 patients. *MedRxiv* (2020).
- 9. Zhang, F. et al. Myocardial injury is associated with in-hospital mortality of confirmed or suspected COVID-19 in Wuhan, China: A single center retrospective cohort study. *MedRxiv* (2020).
- 10. Chen, J. et al. Clinical progression of patients with COVID-19 in Shanghai, China. J. Infect. (2020).
- 11. Du, R.-H. et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. Eur. Respir. J. (2020).
- 12. Li, Z. et al. Caution on kidney dysfunctions of COVID-19 patients. (2020).
- 13. Cai, Q. et al. Experimental treatment with favipiravir for COVID-19: an open-label control study. *Engineering* (2020).
- 14. Shi, S. et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol.* (2020).
- 15. Chen, C., Yan, J. T., Zhou, N., Zhao, J. P. & Wang, D. W. Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19. Zhonghua Xin Xue Guan Bing Za Zhi 48, E008–E008 (2020).
- 16. Cheng, Y. et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int.* (2020).
- 17. Li, J.-W. et al. The impact of 2019 novel coronavirus on heart injury: A systemic review and Meta-analysis. *Prog. Cardiovasc. Dis.* (2020). doi:10.1016/j.pcad.2020.04.008
- 18. Huang, I., Lim, M. A. & Pranata, R. Diabetes mellitus is associated with increased mortality and severity of disease in COVID-19 pneumonia A systematic review, meta-analysis, and meta-regression: Diabetes and

- COVID-19. Diabetes Metab. Syndr. Clin. Res. Rev. 14, 395-403 (2020).
- 19. Guan, W. et al. Comorbidity and its impact on 1590 patients with Covid-19 in China: A Nationwide Analysis. Eur. Respir. J. (2020).
- 20. Meng, Y. et al. Sex-specific clinical characteristics and prognosis of coronavirus disease-19 infection in Wuhan, China: A retrospective study of 168 severe patients. *PLOS Pathog.* **16**, e1008520 (2020).
- 21. Zhang, J. et al. Risk factors for disease severity, unimprovement, and mortality of COVID-19 patients in Wuhan, China. Clin. Microbiol. Infect. (2020). doi:10.1016/j.cmi.2020.04.012
- 22. Chen, R. et al. Risk factors of fatal outcome in hospitalized subjects with coronavirus disease 2019 from a nationwide analysis in China. Chest (2020). doi:10.1016/j.chest.2020.04.010
- 23. Zhou, F. et al. Articles Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. (2020). doi:10.1016/S0140-6736(20)30566-3
- 24. Aifen Lin et al. Early risk factors for the duration of SARS-CoV-2 viral positivity in COVID-19 patients. Clinical infectious diseases: an official publication of the Infectious Diseases Society of America (2020).
- 25. Aifen Lin et al. Early risk factors for the duration of SARS-CoV-2 viral positivity in COVID-19 patients. Clinical infectious diseases: an official publication of the Infectious Diseases Society of America (2020).
- 26. Du, R. H. et al. Predictors of Mortality for Patients with COVID-19 Pneumonia Caused by SARS-CoV-2: A Prospective Cohort Study. Eur. Respir. J. (2020). doi:10.1183/13993003.00524-2020
- 27. Yan, C. H., Faraji, F., Prajapati, D. P., Ostrander, B. T. & DeConde, A. S. Self-reported olfactory loss associates with outpatient clinical course in Covid-19. *Int. Forum Allergy Rhinol.* (2020). doi:10.1002/alr.22592
- 28. Vardavas, C. I. & Nikitara, K. COVID-19 and smoking: A systematic review of the evidence. *Tob. Induc. Dis.* **18**, (2020).
- 29. Shi, P. et al. Impact of temperature on the dynamics of the COVID-19 outbreak in China. Sci. Total Environ. **728**, (2020).
- 30. Bukhari, Q. & Jameel, Y. Will Coronavirus Pandemic Diminish by Summer?

- SSRN Electron. J. (2020). doi:10.2139/ssrn.3556998
- 31. Ficetola, G. F. & Rubolini, D. Climate affects global patterns of COVID-19 early outbreak dynamics. *medRxiv* 2020.03.23.20040501 (2020). doi:10.1101/2020.03.23.20040501
- 32. Mollalo, A., Vahedi, B. & Rivera, K. M. GIS-based spatial modeling of COVID-19 incidence rate in the continental United States. *Sci. Total Environ.* **728**, 138884 (2020).
- 33. Triplett, M. Evidence that higher temperatures are associated with lower incidence of COVID-19 in pandemic state, cumulative cases reported up to March 27, 2020. *medRxiv* 2020.04.02.20051524 (2020). doi:10.1101/2020.04.02.20051524
- 34. Wang, M. et al. Temperature significant change COVID-19 Transmission in 429 cities. *medRxiv* 2020.02.22.20025791 (2020). doi:10.1101/2020.02.22.20025791
- 35. Tobías, A. & Molina, T. Is temperature reducing the transmission of COVID-19? Environmental Research **186**, (2020).
- 36. Shi, P. et al. Impact of temperature on the dynamics of the COVID-19 outbreak in China. Sci. Total Environ. 138890 (2020). doi:10.1016/J.SCITOTENV.2020.138890
- 37. Mollalo, A., Vahedi, B. & Rivera, K. M. GIS-based spatial modeling of COVID-19 incidence rate in the continental United States. *Sci. Total Environ.* 138884 (2020). doi:10.1016/j.scitotenv.2020.138884
- 38. Luo, W. et al. The role of absolute humidity on transmission rates of the COVID-19 outbreak. (2020).
- 39. Hisato Takagi, MD, PhD, Toshiki Kuno, MD, PhD, Yujiro Yokoyama, MD, Hiroki Ueyama, MD, Takuya Matsushiro, MD, Yosuke Hari, MD, Tomo Ando, M. Risk and protective factors of SARS-CoV-2 infection Metaregression of data from worldwide nations. 1–11 (2020).
- 40. Travaglio, M., Popovic, R., Yu, Y., Leal, N. & Martins, L. M. Links between air pollution and COVID-19 in England. *medRxiv* 2020.04.16.20067405 (2020). doi:10.1101/2020.04.16.20067405
- 41. World Health Organisation Coronavirus Disease (COVID-19) Dashboard.

- Available at: https://covid19.who.int/region/euro/country/al. (Accessed: 21st April 2020)
- 42. World Tourism Organization. Country profile inbound tourism | OMT.
- 43. Walach, H., Hockertz, S. & Sciences, M. What association do political interventions, environmental and health variables have with the number of Covid-19 cases and deaths? (2020).
- 44. Aldibasi, O. S. The Association of Country-Level Factors with Outcomes of COVID-19: Analysis of the pandemic after one million cases CURRENT STATUS: POSTED. 1–19 doi:10.21203/rs.3.rs-25834/v1
- 45. Daon, Y., Thompson, R. N. & Obolski, U. Estimating COVID-19 outbreak risk through air travel. *medRxiv* 2020.04.16.20067496 (2020). doi:10.1101/2020.04.16.20067496
- 46. Papadopoulos, D. I., Donkov, I., Charitopoulos, K. & Bishara, S. The impact of lockdown measures on COVID-19: a worldwide comparison. medRxiv 2020.05.22.20106476 (2020). doi:10.1101/2020.05.22.20106476
- 47. Chinazzi, M. et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science* (80-.). **368**, 395–400 (2020).
- 48. Peirlinck, M., Linka, K., Sahli Costabal, F. & Kuhl, E. Outbreak dynamics of COVID-19 in China and the United States. *Biomech. Model. Mechanobiol.* 1–15 (2020). doi:10.1007/s10237-020-01332-5
- 49. Kubota, Y., Shiono, T., Kusumoto, B. & Fujinuma, J. Multiple drivers of the COVID-19 spread: role of climate, international mobility, and region-specific conditions. *medRxiv* 2020.04.20.20072157 (2020). doi:10.1101/2020.04.20.20072157
- 50. Coelho, M. T. P. et al. Exponential phase of covid19 expansion is not driven by climate at global scale. medRxiv 2020.04.02.20050773 (2020). doi:10.1101/2020.04.02.20050773
- 51. Gross, B. et al. Spatio-temporal propagation of COVID-19 pandemics. 1–7 (2020).
- 52. Tuli, S., Tuli, S., Verma, R. & Tuli, R. Modelling for prediction of the spread and severity of COVID-19 and its association with socioeconomic

- factors and virus types. 1-13 (2020).
- 53. Koh, W. C., Naing, L. & Wong, J. Estimating the impact of physical distancing measures in containing COVID-19: an empirical analysis. medRxiv 2020.06.11.20128074 (2020). doi:10.1101/2020.06.11.20128074
- 54. Anzai, A., Kobayashi, T., Linton, N. M. & Kinoshita, R. Assessing the Impact of Reduced Travel on Exportation Dynamics of Novel Coronavirus Infection (COVID-19).
- 55. Merler, S., Pastore, A., Mu, K., Rossi, L. & Sun, K. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. **400**, 395–400 (2020).
- 56. Yao, Y. & Ph, D. Spatial Correlation of Particulate Matter Pollution and Death Rate of COVID-19 /China/ PM2.5, PM10/multiple linear regression. (2020).
- 57. Loewenthal, G. et al. COVID-19 pandemic-related lockdown: response time is more important than its strictness. medRxiv 2020.06.11.20128520 (2020). doi:10.1101/2020.06.11.20128520
- 58. Sorci, G., Faivre, B. & Morand, S. Why does COVID-19 case fatality rate vary among countries? *medRxiv* 2020.04.17.20069393 (2020). doi:10.1101/2020.04.17.20069393
- 59. Qiu, Y., Chen, X. & Shi, W. Impacts of social and economic factors on the transmission of coronavirus disease 2019 (COVID-19) in China. *J. Popul. Econ.* **2019**, 1–27 (2020).
- 60. Messner, W. The Institutional and Cultural Context of Cross-National Variation in COVID-19 Outbreaks. *medRxiv* **2019**, 2020.03.30.20047589 (2020).
- 61. Piguillem, F. The Optimal COVID-19 Quarantine and Testing. (2020).
- 62. Lee, V. J., Chiew, C. J. & Khong, W. X. Interrupting transmission of COVID-19: lessons from containment efforts in Singapore. 1–5 (2020). doi:10.1093/jtm/taaa039