

## **Coblation versus Bipolar Diathermy Hemostasis in Pediatric Tonsillectomy**

### **Patients: Systematic Review and Meta-Analysis**

Mohammad Karam<sup>1\*</sup>, Ahmad Abul<sup>1\*</sup>, Abdulwahab Althuwaini<sup>1</sup>, Talal Alenezi<sup>1</sup>, Ali Aljadi<sup>1</sup>,

Abdulredha Almuhanha<sup>1</sup>, Abdulrahman AlNaseem<sup>2</sup>, Abdulmalik Alsaif<sup>1</sup>, Athari Alwael<sup>3</sup>

<sup>1</sup> School of Medicine, University of Leeds, UK

<sup>2</sup> School of Medicine, University of Manchester, UK

<sup>3</sup> Department of ENT, Consultant Otorhinolaryngology Head and Neck Surgery, Al Jahra

Hospital, Jahra, Kuwait

\*contributed equally as first co-authors

#### **Corresponding Author:**

Mohammad Karam

Address: 129 Candle House, 1 Wharf Approach, Leeds, United Kingdom.

Phone: +44 7480644489

Email: [mhk130@outlook.com](mailto:mhk130@outlook.com)

## Abstract

### Objective

To compare the outcomes of coblation versus bipolar in pediatric patients undergoing tonsillectomy.

### Methods

A systematic review and meta-analysis were performed as per the Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) Guidelines and an electronic search of information was conducted to identify all Randomized Controlled Trials (RCTs) comparing the outcomes of coblation versus bipolar in pediatric patients undergoing tonsillectomy. Intraoperative bleeding, reactionary hemorrhage, delayed hemorrhage and post-operative pain were primary outcome measures. Secondary outcome measures included return to normal diet, effects on tonsillar bed, operation time and administration of analgesia. Fixed and random effects models were used for the analysis.

### Results

Seven studies enrolling 1328 patients were identified. There was a significant difference between coblation and bipolar groups in terms of delayed hemorrhage (Odds Ratio [OR] = 0.25, P = 0.0007) and post-operative pain (standardized mean difference [MD] = -2.13, P = 0.0007). Intraoperative bleeding (MD = -43.26, P = 0.11) and reactionary hemorrhage did not show any significant difference. For secondary outcomes, coblation group had improved outcomes in terms of administration of analgesia, diet and tonsillar tissue recovery and thermal damage. No significant difference was reported in terms of operation time.

### Conclusions

Coblation is a superior option when compared to bipolar technique for pediatric patients undergoing tonsillectomy as it improves post-operative pain and delayed hemorrhage and does not worsen intraoperative bleeding and reactionary hemorrhage.

## Key words

Tonsillectomy; Coblation; Bipolar Diathermy; Post-operative Pain; Hemorrhage

## Highlights

- Coblation tonsillectomy was less painful than bipolar diathermy.
- Coblation tonsillectomy was associated with less delayed hemorrhage than bipolar diathermy.
- Tonsillar tissue recovery, thermal damage and return to normal diet were better in the coblation group.

## Introduction

Tonsillectomy is one of the oldest surgical operations in medicine and one of the most common operations performed by otolaryngologists<sup>1,2</sup>. Some potential indications for pediatric tonsillectomy include recurrent tonsillitis, sleep apnea as well as PFAPA Syndrome (Periodic Fever, Aphthous Stomatitis, Pharyngitis, Adenitis) and it is therefore one of the most common operations during childhood<sup>3</sup>. Several techniques are used to perform tonsillectomies, including blunt dissection, guillotine, bipolar diathermy dissection, laser dissection, and the more recent coblation method<sup>4</sup>. Despite the range of available techniques, postoperative pain, primary or reactionary hemorrhage, and postoperative infection associated with the hemorrhage continue to present as the main post-tonsillectomy complications<sup>5</sup>. Therefore, studies continue to debate the optimal tonsillectomy technique.

Coblation (cold ablation) is a relatively new tonsillectomy technique that has earned increased popularity due to the decreased postoperative pain and reduced intraoperative bleeding that comes with its use<sup>6</sup>. Coblation is operated at surface temperatures (40-70°C) much lower than those of more traditional techniques<sup>7</sup>. Instead of relying on heat, coblation applies radiofrequency energy to a conductive natural salt solution, forming a plasma membrane comprised of highly ionised particles that hold enough energy to break the molecular bonds holding the tissue, thereby safely removing the target tissue<sup>7-9</sup>. The use of coblation eliminates the risk of causing thermal damage that comes with heat and minimizes necrosis of surrounding healthy tissue, therefore resulting in minimal pain and faster recovery<sup>8,9</sup>.

Bipolar diathermy is an electrosurgery technique that functions by passing an alternating current at a high-frequency through a pair of forceps to cut the tissue and coagulate the blood vessels<sup>10</sup>. Compared to monopolar diathermy, bipolar diathermy provides more control over the targeted area and uses less energy, thus causing less damage<sup>10</sup>. The use of bipolar diathermy to perform tonsillectomy is first described in a paper<sup>11</sup> written in 1994. Bipolar diathermy tonsillectomy

is proven to be safe for both adults and children<sup>11,12</sup>. Although the pain and morbidity rates are similar to other tonsillectomy techniques like cold dissection tonsillectomy, the bipolar diathermy technique provides significant advantages such as shorter operative time and lower blood loss levels<sup>11,12</sup>.

Several published studies assessed the effectiveness of coblation compared with bipolar diathermy techniques in pediatric patients undergoing tonsillectomy<sup>13-19</sup> including a previous meta-analysis comparing coblation to several techniques used in tonsillectomy<sup>20</sup>. However, this is the first study to perform a meta-analysis on these two specific techniques focusing on the pediatric population.

## Methods

A systematic review and meta-analysis were conducted as per the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines<sup>21</sup>.

### Eligibility criteria

All randomized control trials and observational studies comparing coblation versus bipolar diathermy hemostasis techniques for tonsillectomy were included. Coblation was the intervention group of interest and bipolar diathermy was the comparator. The study also included patients having bipolar scissors, bipolar forceps as well as cold steel and bipolar diathermy hemostasis as the comparator. All patients were included irrespective of gender or co-morbidity status as long as they belonged to either a study or control group and were pediatric patients. All case reports, cohort studies without a comparison group, and studies not written in English were excluded. The study also excluded the following techniques: bipolar molecular resonance coagulation, tonsillotomy, unipolar or monopolar diathermy, adenotonsillectomy and conventional or traditional tonsillectomy without explicitly stating bipolar diathermy as the primary method for hemostasis.

### Primary Outcomes

The primary outcomes are intraoperative bleeding, reactionary hemorrhage (within 24 hours after the operation), delayed hemorrhage (bleeding after 24 hours of the operation) and post-operative pain on day 7.

### Secondary Outcomes

The secondary outcomes included return to normal diet, effects on tonsillar bed (degree of healing in tonsillar fossae and thermal damage to tonsillar tissue), operation time and administration of analgesia.

### Literature search strategy

Two authors AA and MK independently searched the following electronic databases: MEDLINE, EMBASE, EMCARE, CINAHL, and the Cochrane Central Register of Controlled Trials (CENTRAL). The

last search was run on the 12<sup>th</sup> of April 2020. Thesaurus headings, search operators and limits in each of the above databases, were adapted accordingly. In addition, World Health Organization International Clinical Trials Registry (<http://apps.who.int/trialsearch/>), ClinicalTrials.gov (<http://clinical-trials.gov/>), and ISRCTN Register (<http://www.isrctn.com/>) were searched for details of ongoing and unpublished studies. No language restrictions were applied in our search strategies. The search terminologies included “coblation”, “bipolar”, and “tonsillectomy”. The bibliographic lists of relevant articles were also reviewed.

### Selection of Studies

The title and abstract of articles identified from the literature searches were assessed independently by two authors (AA and MK). The full texts of relevant reports were retrieved and articles that met the eligibility criteria of our review were selected. Any discrepancies in study selection were resolved by discussion between the authors.

### Data Extraction and Management

An electronic data extraction spreadsheet was created in line with Cochrane’s data collection form for intervention reviews. The spreadsheet was pilot tested in randomly selected articles and adjusted accordingly. Our data extraction spreadsheet included study-related data (first author, year of publication, country of origin of the corresponding author, journal in which the study was published, study design, study size, clinical condition of the study participants, type of intervention, and comparison), baseline demographics of the included populations (age and gender), and primary and secondary outcome data. The authors cooperatively collected and recorded the results and any disagreements were solved via discussion.

### Data synthesis

Data synthesis was conducted using the Review Manager 5.3 software. The extracted data was entered into Review Manager by three independent authors. The analysis involved used was

based on fixed and random effects modelling. The results were reported in forest plots with 95% Confidence Intervals (CIs).

For dichotomous outcomes, the Odds Ratio (OR) was calculated between the 2 groups. The OR is the odds of an event in the coblation group compared with the bipolar group. An OR of greater than 1 for the delayed hemorrhage would favor the coblation group, an OR of less than 1 would favor the bipolar group, and a OR of 1 would favor neither groups.

For continuous outcomes, the Mean Difference (MD) was calculated between the 2 groups. A positive MD for the post-operative pain score by day 7 and intraoperative bleeding would favor the coblation group, a negative MD would favor the bipolar group, and an MD of 0 would favor neither groups.

#### Assessment of Heterogeneity

Heterogeneity among the studies was assessed using the Cochran Q test ( $\chi^2$ ). Inconsistency was quantified by calculating  $I^2$  and interpreted using the following guide: 0% to 25% may represent low heterogeneity, 25% to 75% may represent moderate heterogeneity, and 75% to 100% may represent high heterogeneity.

#### Methodological quality and risk of bias assessment


Multiple authors independently assessed the methodological quality as well as the risk of bias for articles matching the inclusion criteria. For randomized trials, the Cochrane's tool for evaluating the risk of bias was used. Domains assessed included selection bias, performance bias, detection bias, attrition bias, reporting bias, and other possible sources of bias. Randomized controlled trial (RCT) studies are classified into low, unclear, or high risk of bias. For non-randomized studies, the Newcastle-Ottawa Scale was used<sup>22</sup>. It uses a star grading system to assess studies in terms of three domains: selection, comparability, and exposure. The total maximum score for each study is nine stars. The overall rating of either good, fair or poor quality was based on the Agency for Healthcare Research and Quality (AHRQ) standards<sup>22</sup>.



## Results

### Literature search results

The search strategy retrieved 546 studies and after a thorough screening of the retrieved articles using the inclusion and exclusion criteria, the authors identified seven studies in total which met the eligibility criteria (Figure 1).



**Figure 1: Prisma Flow Diagram.** The PRISMA diagram details the search and selection processes applied during the overview. PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

## Description of Studies:

Table 1 summarises the baseline characteristics of included studies published from 2001 to 2018. All included population were pediatric patients under the age of 18 who underwent tonsillectomy.

### *Temple et al. 2001*

Temple et al. conducted a single center prospective RCT that included 38 pediatric patients listed for a routine tonsillectomy with a history of chronic tonsillitis or obstructive tonsils. Patients were randomized via closed opaque envelope technique to either have bilateral coblation tonsillectomy (using an ArthroCare CoVac™ 70 ArthroWand®) or bilateral standard bipolar dissection tonsillectomy. All tonsillectomies were extracapsular.

### *Mitic et al. 2007*

Mitic et al. performed a single center prospective RCT that included 40 patients with standard indication for tonsillectomy. Randomizing was achieved by closed opaque envelope technique to either have coblation tonsillectomy or dissection tonsillectomy with bipolar diathermy hemostasis. Bipolar dissection was done using standard technique (set at 4/10 and 50 Watts power). All tonsillectomies were extracapsular.

### *Parker et al. 2009*

Parker et al. performed a single center prospective RCT that included 60 pediatric patients undergoing tonsillectomy by either cold steel dissection or coblator dissection. The trial was double blinded and a computer-generated random sequence, in sealed opaque envelopes, was used for allocation of procedure technique. All tonsillectomies were extracapsular.

### *Roje et al. 2009*

Roje et al. performed a single center, prospective RCT that included 72 pediatric patients listed for tonsillectomy. Randomization was fulfilled by using computer-generated random numbers that separated the patients into one of two groups that would undergo either coblation

tonsillectomy or conventional cold-steel tonsillectomy with bipolar diathermy. All tonsillectomies were extracapsular.

*Belloso et al. 2010*

Belloso et al. performed a single center prospective observational cohort study which included 1008 participants from July 2001 to January 2003; 526 patients received a coblation tonsillectomy and 482 patients received a blunt dissection tonsillectomy. Participants data was extracted from a tonsillectomy audit. All tonsillectomies were extracapsular. Bipolar dissection was done using standard technique (set at 4/10- and 50-Watts power).

*Konsolov et al. 2017*

Konsolov et al. carried out a single center prospective cohort study which included 60 children aged 3-18 years. The children were divided into two equal groups. One group received traditional blunt dissection with bipolar diathermy hemostasis and the other underwent coblation tonsillectomy using the ArthroCare CoVac™ 70 ArthroWand®. All tonsillectomies were extracapsular.

*Bhardwaj et al. 2018*

Bhardwaj et al. conducted a single center prospective RCT that included 100 pediatric patients undergoing tonsillectomy. Patients were given random numbers and they got randomly allocated to one of two groups: one group undergoing bipolar diathermy and the other undergoing coblation. Bipolar dissection was done using 12 watts of power. All tonsillectomies were extracapsular.

Study (Year)	Country, Journal	Study Design	Sex (Male:Female)	Mean Age ± SD (Range)	Total Study Sample (Control:Intervention)	Interventions Compared
Temple et al. (2001)	International Journal of Paediatric otorhinolaryngology, UK	RCT‡	19:19	5.6 years	38 (18:20)	Standard bipolar versus coblation tonsillectomy
Mitic et al. (2007)	Clinical Otolaryngology, Norway	RCT	NR†	NR (4-12 years)	40 (20:20)	Coblator or steel dissecting instruments versus bipolar diathermy
Parker et al. (2009)	Clinical Otolaryngology, UK	RCT	35:44	8.2 years (4-15 years)	70 (35:35)	Cold steel dissection with bipolar hemoostasis versus coblator dissection
Roje et al. (2009)	Collegium antropologicum, Croatia	RCT	41:31	6 years (3-16 years)	72 (36:36)	Conventional cold steel tonsillectomy with bipolar diathermy coagulation versus coblator
Konsulov et al.	International Journal of Otorhinolaryngology, Bulgaria	Prospective Comparative Study	NR	NR (3-18 years) Coblation: 8.16 ± 4.74 Bipolar: 6.87 ± 3.01	60 (30:30)	Coblator II system ArthroCare (Smith and Nephew) Evac 70 wand for extracapsular dissection vs. extracapsular blunt dissection with bipolar diathermy
Belloso et al. (2010)	The Laryngoscope, UK	Cohort	NR	Coblation: 7.76 ± 3.56 years Bipolar: 7.60 ± 3.44 years	1008 (526:482)	Coblation or blunt dissection versus bipolar diathermy hemostasis
Bhardwaj et al. (2018)	Indian Journal of Otolaryngology and Head & Neck Surgery, India	RCT	NR	NR (4-14 years)	100 (50:50)	Coblation Assissted versus Bipolar Diathermy

Table 1. Baseline Characteristics of the Included Studies. NR: not reported.

† NR: Not Reported

‡ RCT: Randomized Control Trial

## Primary Outcomes:

### *Bleeding*

Three outcomes of bleeding were reported in the included studies, namely intraoperative hemorrhage, reactionary hemorrhage and delayed hemorrhage.


Intraoperative bleeding was reported in two studies included 132 patients as shown in Figure 2. The mean difference analyses showed no statistically significant difference however a trend is demonstrated favoring the coblation group (MD = -43.26, CI = -96.33 to 9.80, P = 0.11). A high level of heterogeneity was found amongst the studies ( $I^2 = 99\%$ ,  $P < 0.00001$ ). Mitic et al. also reported less intraoperative bleeding in the coblation group (28.25 mL) than the bipolar group (62.25 mL). However, Hasan et al. reported no significant difference between both groups.

Reactionary or primary hemorrhage was reported by Roje et al. who did not record any case of primary hemorrhage in both groups. However, Kunsulov et al. reported two cases of reactionary hemorrhage in the bipolar group, compared to none in those that underwent coblation tonsillectomy.


Delayed or secondary hemorrhage was reported in four studies enrolling 1210 patients as demonstrated in Figure 3. There was a statistically significant difference seen in the odd ratio analyses showing a lower rate of delayed hemorrhage for the coblation group (OR = 0.25, CI = 0.11 to 0.55, P = 0.0007). A low level of heterogeneity was found amongst the studies ( $I^2 = 7\%$ , P = 0.34).

Post-operative Pain by Day 7


In Figure 4, post-operative pain by day 7 was reported using different pain scales in three studies enrolling 200 patients. There was a statistically significant difference seen in the standardized mean difference analyses showing less pain for the coblation group (standardized MD = -2.13, CI = -3.37 to -0.90, P = 0.0007). A high level of heterogeneity was found amongst the studies ( $I^2 = 91\%$ ,  $P < 0.0001$ ). Temple et al. also reported mean post-operative pain scores however did not include standard deviation; hence it wasn't possible to quantitatively assess them in the forest plots. Temple reported a significant difference favoring coblation ( $p < 0.0001$ ).



**Figure 2: Forest Plot of Coblation versus Bipolar Tonsillectomy – Intraoperative Bleeding.** Quantitative analysis showing the mean difference in the intraoperative bleeding reported as median by Roje et al. (2009) and Consulov et al. (2017).



**Figure 3: Forest Plot of Coblation versus Bipolar – Delayed Haemorrhage.** Quantitative analysis showing the odd ratio in delayed haemorrhage reported by Parker et al. (2009), Roje et al. (2009), Belloso et al. (2010) and Consulov et al. (2017).



**Figure 4: Forest Plot of Coblation versus Bipolar – Post-operative Pain by Day 7.** Quantitative analysis showing the odd ratio in delayed haemorrhage reported by Mitic et al. (2007), Consulov et al. (2017) and Bhardwaj et al. (2018).

Secondary outcomes:

#### *Return to Normal Diet*

Three studies with two different types of assessment of diet were included. Temple et al. reported a statistically significant difference in the days it took to return to normal diet between the two groups, with an average of 2.4 days for patients who had coblation tonsillectomy versus an average of 7.6 days for patients who had routine bipolar dissection. On the other hand, Parker et al. reported no significant difference in the number of days taken by the two groups to return to normal diet, with a steady increase from day 6 onwards in both groups. Mitic et al. reported diet using a nutrition score during a 10-day postoperative period and found that there was a statistically significant difference between the score of patients who had coblation tonsillectomy and those who had dissection tonsillectomy across the 10 days, favoring the former.

#### *Effect on Tonsillar Bed*

A significant difference in the degree of healing in tonsillar fossae between the two interventions was reported by Temple et al., with nearly all coblation fossae healed 9 days post-operatively, while bipolar dissection patients had considerable slough. Roje et al. reported a statistically significant mean difference of the depth of thermal damage ( $t = -40,1$ ;  $p < 0.001$ ) to tonsillar tissue, where the coblation technique caused damage two times shallower than that caused by bipolar diathermy hemostasis ( $428.58 \pm 47.4 \mu\text{m}$  and  $841.17 \pm 39.7 \mu\text{m}$ , respectively).

#### *Operation time (min)*

Operation time was defined as point of knife to skin contact according to Mitic et al.. This study reported a statistically insignificant difference between coblation and bipolar groups (26.6 min and 25.6 min, respectively).

#### *Administration of Analgesia*

Four studies have assessed analgesics administration in different ways. Parker et al identified less analgesic requirements by patients that underwent coblation tonsillectomy in the first 12 hours post-operatively. In addition, Roje et al and Konsulov agreed that patients undergoing

coblation required lower number of days on analgesics for the coblation group compared to those that received the bipolar technique. Roje also identified that the coblation group required a lower number of analgesic applications (4 vs. 8). Mitic et al has reported a lower medication intake score for the coblation group however did not specify how the scale works. Other than Roje, the studies have mentioned using the same analgesics (ibuprofen and paracetamol).

#### Methodological Quality and Risk of Bias Assessment

The Cochrane Collaboration's Tool was used to assess the quality of the RCTs included in the study (Table 2). The Newcastle-Ottawa scale was used to assess the quality of the non-randomized studies (Table 3) which offers a star system for analysis<sup>22</sup>. The quality of the included non-randomized study was deemed to be high in selection and exposure but low in comparability. Overall, both studies were of good quality based on the AHRQ standards<sup>22</sup>.



First Author	Bias	Authors' Judgement	Support for Judgement
<b>Temple et al. (2001)</b>	Random sequence generation (selection bias)	Unclear Risk	No information given
	Allocation concealment (selection bias)	Low Risk	Closed opaque envelope technique
	Blinding of participants and personnel (performance bias)	High Risk	Single surgeon – not blinded.
	Blinding of outcome assessment (detection bias)	Low Risk	Outcome measurement is not likely to be influenced by lack of blinding
	Incomplete outcome data (attrition bias)	High Risk	Standard deviation for pain is not reported
	Selective reporting (reporting bias)	Unclear Risk	No information given
	Other bias	Low Risk	Similar baseline characteristics in both groups.
<b>Mitic et al. (2007)</b>	Random sequence generation (selection bias)	Low Risk	A randomised sequence was generated by a statistician of the two interventions.
	Allocation concealment (selection bias)	Low Risk	Allocation of treatment was envelope sealed by a statistician and was then opened in the surgery room in sequential order by a nurse.
	Blinding of participants and personnel (performance bias)	High Risk	Surgeons were not blinded
	Blinding of outcome assessment (detection bias)	Low Risk	Outcome assessment was done by parents and nurses whom all were blinded for the operation method. The data from parents and nurses were compared and no difference was identified.
	Incomplete outcome data (attrition bias)	Low Risk	All questionnaires were complete.
	Selective reporting (reporting bias)	High Risk	Return to normal diet mentioned in abstract was not reported.
	Other bias	Low Risk	Similar baseline characteristics in both groups.
<b>Parker et al. (2009)</b>	Random sequence generation (selection bias)	Low Risk	Random sequence generated by envelope
	Allocation concealment (selection bias)	Low Risk	Sealed opaque envelope technique
	Blinding of participants and personnel (performance bias)	High risk	Single surgeon.
	Blinding of outcome assessment (detection bias)	Low Risk	Double blinded study. The children, the parents and the nursing staff doing the pain assessments and prescribing the analgesia were not informed which technique had been used. The surgeon took no part in the pain assessments.
	Incomplete outcome data (attrition bias)	Low Risk	Missing outcome data balanced in numbers across intervention groups, with similar reasons for missing data across groups

	Selective reporting (reporting bias)	Low Risk	All outcome data reported
	Other bias	Low Risk	Similar baseline characteristics in both groups.
<b>Roje et al. (2009)</b>	Random sequence generation (selection bias)	Low Risk	Computer-generated randomisation was used for selection of children and separating them into groups from a large ENT database.
	Allocation concealment (selection bias)	Unclear Risk	No information given
	Blinding of participants and personnel (performance bias)	High Risk	Single blinded (parents). Single surgeon not blinded
	Blinding of outcome assessment (detection bias)	Unclear Risk	Parents assessing secondary outcomes did not know which procedure was used.
	Incomplete outcome data (attrition bias)	High Risk	Participants lost to follow up (15 total- 7 coblation and 8 bipolar group)
	Selective reporting (reporting bias)	Unclear Risk	Insufficient information.
	Other bias	Low Risk	Similar baseline characteristics in both groups.
	<b>Bhardwaj et al. (2018)</b>	Random sequence generation (selection bias)	Unclear Risk
Allocation concealment (selection bias)		Unclear Risk	Random numbers but not explicitly concealed
Blinding of participants and personnel (performance bias)		Unclear Risk	Insufficient information
Blinding of outcome assessment (detection bias)		Unclear Risk	Insufficient information
Incomplete outcome data (attrition bias)		Low Risk	No missing information
Selective reporting (reporting bias)		Low Risk	No published protocol but all the expected and pre-specified outcomes are reported
Other bias		Low Risk	No significant demographic difference between the 2 groups.

Table 2. Bias analysis of the Randomized Trials using the Cochrane Collaboration's Tool

<b>Study</b>	<b>Selection</b>	<b>Comparability</b>	<b>Exposure</b>
<b>Belloso et al. (2010)</b>	****	*	***
<b>Konsulov et al. (2017)</b>	****	*	**

Table 3. Newcastle-Ottawa Scale to Assess the Quality of Non-randomised Studies.

## Discussion:

Coblation showed a superior effect when compared with bipolar diathermy in pediatric patients undergoing tonsillectomy in terms of delayed hemorrhage and postoperative pain, as shown by the results of the analyses. Although there were no significant differences between the coblation and bipolar groups in terms of intraoperative bleeding ( $P = 0.11$ ) and reactionary hemorrhage, delayed hemorrhage was significantly ( $P = 0.0007$ ) lower in the former group. Similarly, a significant ( $P = 0.0007$ ) improvement in post-operative pain was noted for the coblation group. Regarding the between-study heterogeneity, it was low for delayed hemorrhage ( $I^2 = 7\%$ ) and high for post-operative pain ( $I^2 = 91\%$ ) and intraoperative bleeding ( $I^2 = 99\%$ ), based on the assessment mentioned in the methodology.

Along with the outcomes mentioned above, the findings of this study reported several secondary outcomes that proved coblation to be a more effective technique than bipolar. Both tonsillar tissue recovery and thermal damage were significantly better in the coblation group. This correlates with analgesic administration whereby coblation required less doses compared to bipolar dissection. Generally, return to normal diet was quicker for the coblation group. There was no significant difference between the two groups in terms of operation time.

Currently, there is a debate in the literature with regards to the most efficient technique used for tonsillectomy in children<sup>23</sup>. The results of the current study are comparable to some studies where coblation tonsillectomy was found to be less painful than bipolar tonsillectomy in the immediate and overall postoperative period, which resulted in a swifter recovery and reduced analgesic requirements<sup>24, 25</sup>. A randomized control trial which included 80 adolescents found that pain and otalgia post-operatively was very slightly lower in the coblation group; this difference was, however, considered clinically insignificant<sup>25</sup>.

A more recent study in Iraq<sup>26</sup>, which compared intraoperative efficiency and postoperative recovery between bipolar electrocautery and coblation tonsillectomy in children, recorded

statistically and clinically significant higher amounts of intra-operative blood loss in bipolar technique than using coblation (Bipolar 1.43 ml vs. Coblation 15.37 ml,  $P < 0.001$ ). Although this outcome conflicts with the results of this study, the coblation technique was associated with lower mean pain scores, which is the direction of this study<sup>26</sup>.

A study in the United States included 7,562 patients under 12 years of age compared the costs of treatment and management of children undergoing tonsillectomy by either coblation or electrocautery technique<sup>27</sup>. The study reported statistically significant lower costs for coblation compared to electrocautery surgery (\$1,009 versus \$1,162;  $P < 0.0001$ ) and pharmacy (\$102.40 versus \$117.20;  $P < 0.0001$ ) costs. However, when central supply is put into consideration the total cost of coblation is slightly higher<sup>27</sup> (\$2,646 versus \$2,591;  $P = 0.0011$ ) with a difference of \$55.94. However, because coblation is less likely to result in severe hemorrhage, it was found that using coblation tonsillectomy could save the National Health Service in the United Kingdom an incremental cost of £2000 for every avoided hemorrhage as opposed to using cold dissection with bipolar diathermy for haemostasis<sup>28</sup>. In a retrospective audit of 1,336 patients who had undergone coblation, it was reported that this technique had an increased requirement for operative intervention to manage secondary haemorrhage<sup>29</sup>.

This study supports the use of coblation technique over bipolar diathermy for pediatric patients undergoing tonsillectomy. To further support the current conclusion, the authors suggest using coblation tonsillectomy under rigorous and well-designed RCTs to appropriately record data for primary and secondary outcomes.

A systematic review and meta-analysis were performed on available studies in order to provide an evidence-based conclusion. The studies and their results were described and summarized, and the risk of bias in each study was appraised. The study designs and populations were standardized, with inclusion and exclusion criteria pre-defined. The interventions were homogenous across the included studies, with patients undergoing either coblation tonsillectomy or

bipolar diathermy. Although some variation was observed in the control group such as using bipolar scissors as the primary cutting tool, bipolar diathermy was used as the primary technique for hemostasis. Despite these strengths, a number of inherent limitations should be considered. The current meta-analysis enrolled only seven studies with a total sample size of 1328 patients. There was a relatively large range in their sample sizes (1008 and 38 participants for Belloso et al. and Temple et al., respectively). Further clinical trials should therefore be completed to support the current study findings of the superior effect of coblation technique for pediatric patients undergoing tonsillectomy.

## Conclusions

Although the evidence is limited, with only seven studies comparing coblation and bipolar techniques, the results of this meta-analysis suggest that coblation improves the outcomes in paediatric patients undergoing tonsillectomy as it decreases delayed hemorrhage and post-operative pain and does not worsen intraoperative bleeding and reactionary hemorrhage. Further clinical studies are required to support the efficiency of coblation tonsillectomy technique.

## Declarations

Ethics Approval and Consent to Participate

Not Applicable.

Consent for Publication

Not Applicable.

Availability of Data and Materials

The datasets generated and analysed during the current study are available from the corresponding author on reasonable request.

Competing Interests

The author(s) declared that they have no competing interests.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

Author Contributions

Study concept and design: Mohammad Karam and Ahmad Abul

Acquisition of Data: Abdulredha Almuhanha, Abdulrahman AlNaseem, Abdulmalik Alsaif

Analysis of Data: Mohammad Karam, Ahmad Abul, and Ali Aljadi

Interpretation of Data: Mohammad Karam, Ahmad Abul, Abdulwahab Althuwaini, Ali Aljadi, and

Talal Alanezi

All authors contributed in the drafting of the manuscript.

Study Supervision: Athari Alwael

All authors have read and approved the final draft manuscript.

Acknowledgements

Not Applicable.

## References

1. Lowe D, van der Meulen J, Cromwell Det al. Key messages from the National Prospective Tonsillectomy Audit. *The Laryngoscope* 2007; 117:717-724.
2. Curtin JM. The history of tonsil and adenoid surgery. *Otolaryngologic clinics of North America* 1987; 20:415-419.
3. Gysin C. Indications of pediatric tonsillectomy. *ORL; journal for oto-rhino-laryngology and its related specialties* 2013; 75:193-202.
4. Back L, Paloheimo M, Ylikoski J. Traditional tonsillectomy compared with bipolar radiofrequency thermal ablation tonsillectomy in adults: a pilot study. *Archives of otolaryngology--head & neck surgery* 2001; 127:1106-1112.
5. Johnson LB, Elluru RG, Myer CM, 3rd. Complications of adenotonsillectomy. *The Laryngoscope* 2002; 112:35-36.
6. V N, Dutta A, Sabarigirish K. A comparative study of coblation assisted adenotonsillectomy and cold dissection adenotonsillectomy in children. *International Journal of Otorhinolaryngology and Head and Neck Surgery* 2016; 3:122.
7. Shapiro NL, Bhattacharyya N. Cold dissection versus coblation-assisted adenotonsillectomy in children. *The Laryngoscope* 2007; 117:406-410.
8. Paramasivan VK, Arumugam SV, Kameswaran M. Randomised comparative study of adenotonsillectomy by conventional and coblation method for children with obstructive sleep apnoea. *International journal of pediatric otorhinolaryngology* 2012; 76:816-821.
9. Sergeev VN, Belov SV. Coblation Technology: a New Method for High-Frequency Electrosurgery. *Biomedical Engineering* 2003; 37:22-25.
10. McCauley G. Understanding electrosurgery. *Bovie Med Corp.* 2010; 4:4-15.
11. Pang YT, El-Hakim H, Rothera MP. Bipolar diathermy tonsillectomy. *Clinical Otolaryngology & Allied Sciences* 1994; 19:355-357.



12. Hesham A. Bipolar diathermy versus cold dissection in paediatric tonsillectomy. International journal of pediatric otorhinolaryngology 2009; 73:793-795.
13. Temple RH, Timms MS. Paediatric coblation tonsillectomy. International journal of pediatric otorhinolaryngology 2001; 61:195-198.
14. Mitic S, Tvinnereim M, Lie E, Šaltytė BJ. A pilot randomised controlled trial of coblation tonsillectomy versus dissection tonsillectomy with bipolar diathermy haemostasis. Clinical Otolaryngology 2007; 32:261-267.
15. Parker D, Howe L, Unsworth V, Hilliam R. A randomised controlled trial to compare postoperative pain in children undergoing tonsillectomy using cold steel dissection with bipolar haemostasis versus coblation technique. Clinical Otolaryngology 2009 ; 34:225-231.
16. Roje Ž, Račić G, Đogaš Z, Pešutić Pisac V, Timms M. Postoperative morbidity and histopathologic characteristics of tonsillar tissue following coblation tonsillectomy in children: a prospective randomised single-blind study. Collegium antropologicum 2009; 33:293-298.
17. Belloso A, Chidambaram A, Morar P, Timms MS. Coblation tonsillectomy versus dissection tonsillectomy: postoperative hemorrhage. The Laryngoscope 2003; 113:2010-2013.
18. Bhardwaj B, Singh J. Comparative Post-operative Pain Analysis Between Coblator Assisted and Bipolar Diathermy Tonsillectomy in Paediatric Patients. Indian Journal of Otolaryngology and Head & Neck Surgery 2019; 71:90-94.
19. Konsulov S, Konsulov S, Dzhabazov K, Kopanov P. Comparison Between Coblation Assisted Tonsillectomy Versus Conventional Tonsillectomy Regarding the Postoperative Pain and Bleeding. International Journal of Otorhinolaryngology 2017; 3:1.
20. Pynnonen M, Brinkmeier JV, Thorne MC, Chong LY, Burton MJ. Coblation versus other surgical techniques for tonsillectomy. Cochrane Database of Systematic Reviews 2017; 8.
21. Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. Annals of internal medicine 2009; 151:264-269.

22. Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses  
[www.ohri.ca/programs/clinical\\_epidemiology/oxford.htm](http://www.ohri.ca/programs/clinical_epidemiology/oxford.htm)
23. Wiltshire D, Cronin M, Lintern N, Fraser-Kirk K, Anderson S, Barr R, Bennett D, Bond C. The debate continues: a prospective, randomised, single-blind study comparing Coblation and bipolar tonsillectomy techniques. *The Journal of Laryngology & Otology* 2018; 132:240-245.
24. Metcalfe C, Muzaffar J, Daultrey C, Coulson C. Coblation tonsillectomy: a systematic review and descriptive analysis. *European Archives of Oto-Rhino-Laryngology* 2017; 274:2637-2647.
25. Hong SM, Cho JG, Chae SW, Lee HM, Woo JS. Coblation vs. electrocautery tonsillectomy: a prospective randomised study comparing clinical outcomes in adolescents and adults. *Clinical and experimental otorhinolaryngology* 2013; 6:90.
26. Saeed HI, Emin A, Yaseen MA, Saleh AM. A comparative study between coblation and bipolar electrocautery tonsillectomy in children. *Zanco Journal of Medical Sciences* 2019; 23:81-89.
27. Adeyemi A, Delhougne G. PMD51-COMPARISON OF COSTS AND OUTCOMES BETWEEN COBLATION TECHNOLOGY AND ELECTROCAUTERY FOLLOWING TONSILLECTOMY AND ADENOIDECTOMY PROCEDURES. *Value in Health* 2018; 21:S251.
28. Guest JF, Rana K, Hopkins C. Cost-effectiveness of Coblation compared with cold steel tonsillectomies in the UK. *The Journal of Laryngology & Otology* 2018; 132:1119-27.
29. Praveen CV, Parthiban S, Terry RM. High incidence of post-tonsillectomy secondary haemorrhage following coblation tonsillectomy. *Indian Journal of Otolaryngology and Head & Neck Surgery* 2013; 65:24-28.