

Supplementary Information: Analysis of the potential for a malaria vaccine to reduce gaps in malaria intervention coverage

Table S1: Countries eligible for inclusion in the analysis and surveys from which data were extracted. All surveys are available for download from the Demographic and Health Surveys website [1].

Country	Surveys used
Angola (AO)	2015 DHS
Benin (BJ)	2012 DHS
Burkina Faso (BF)	2010 DHS, 2014 MIS
Burundi (BU)	2016 DHS
Cameroon (CM)	2011 DHS
Cote d'Ivoire (CI)	2012 DHS
Democratic Republic of Congo (CD)	2013 DHS
Ghana (GH)	2014 DHS, 2016 MIS
Guinea (GN)	2012 DHS
Kenya (KE)	2014 DHS, 2015 MIS
Liberia (LB)	2013 DHS, 2016 MIS
Malawi (MW)	2015 DHS, 2017 MIS
Mali (ML)	2012 DHS, 2015 MIS
Mozambique (MZ)	2015 AIS
Nigeria (NG)	2013 DHS, 2015 MIS
Sierra Leone (SL)	2013 DHS, 2016 MIS
Tanzania (TZ)	2015 DHS, 2017 MIS
Togo (TG)	2013 DHS, 2017 MIS
Uganda (UG)	2016 DHS
Zambia (ZM)	2013 DHS, 2015 MIS




Figure S1: Coverage of the DTP3 vaccine at the country level, compared to measles vaccine coverage.

Spearman's correlation coefficient: 0.950. The red line represents correlation of 1.0. Countries shown: AO – Angola, BF – Burkina Faso, BJ – Benin, BU – Burundi, CD - Democratic Republic of Congo, CI – Cote d'Ivoire, CM – Cameroon, GH – Ghana, GN – Guinea, KE – Kenya, LB – Liberia, ML – Mali, MW – Malawi, MZ – Mozambique, NG – Nigeria, SL – Sierra Leone, TG – Togo, TZ – Tanzania, UG - Uganda and ZM – Zambia.




Figure S2: Yearly clinical incidence of malaria as a function of the *P. falciparum* parasite prevalence in children aged 12–59 months (*PfPR*_{12–59}). Clinical incidence is the yearly number of events per child aged 12–59 months. This relationship was computed using a deterministic mathematical model for malaria transmission [2].

Table S2: Numbers of children and proportions of multinomial regression predictors in each country.

Country code	Number of children (thousands)	Sex		Age		Urban/ Rural		Mother's education		Wealth Index	
		Male	Female	One	Two	Urban	Rural	None & Primary	Secondary & Tertiary	Bottom 60%	Top 40%
Angola	5451	50%	50%	52%	48%	56%	44%	71%	29%	78%	23%
Burkina Faso	4817	51%	49%	52%	48%	37%	63%	89%	12%	64%	36%
Benin	5461	51%	49%	51%	49%	24%	76%	94%	6%	61%	39%
Burundi	4979	50%	50%	52%	48%	16%	84%	87%	13%	60%	39%
Democratic Republic of Congo	2293	49%	51%	53%	47%	42%	58%	65%	35%	64%	36%
Cote d'Ivoire	2814	50%	50%	50%	50%	34%	66%	89%	10%	70%	30%
Cameroon	6831	50%	50%	50%	50%	30%	70%	65%	35%	69%	31%
Ghana	2496	52%	48%	52%	48%	31%	69%	89%	11%	63%	37%
Guinea	2255	52%	48%	50%	50%	41%	59%	54%	46%	71%	28%
Kenya	8055	51%	49%	50%	50%	32%	68%	73%	27%	71%	29%
Liberia	2699	52%	48%	53%	47%	32%	68%	79%	20%	83%	17%
Mali	6492	49%	51%	50%	50%	16%	84%	79%	22%	65%	36%
Malawi	3608	50%	50%	51%	49%	26%	74%	90%	10%	58%	41%
Mozambique	1977	48%	52%	51%	49%	36%	64%	79%	21%	56%	44%
Nigeria	11154	51%	49%	52%	48%	34%	66%	65%	35%	64%	36%
Sierra Leone	4012	49%	51%	51%	49%	30%	70%	81%	19%	64%	36%
Togo	4032	50%	50%	53%	47%	24%	76%	81%	19%	62%	37%
Tanzania	2676	50%	50%	52%	48%	30%	70%	78%	22%	68%	32%
Uganda	5814	51%	49%	50%	50%	19%	81%	75%	25%	66%	33%
Zambia	5092	50%	50%	51%	49%	37%	63%	66%	34%	71%	29%

References

1. ICF. Data. The DHS Program website. Funded by USAID. <https://dhsprogram.com/Data/>. Accessed 23 Mar 2020.
2. Griffin JT, Ferguson NM, Ghani AC. Estimates of the changing age-burden of *Plasmodium falciparum* malaria disease in sub-Saharan Africa. *Nat Commun.* 2014;5:3136. doi:10.1038/ncomms4136.