

Supplementary Material:

**Onset, duration, and persistence of taste and smell changes and other COVID-19
symptoms: longitudinal study in Israeli patients**

Hadar Klein^{1#}, Kim Asseo^{1#}, Noam Karni^{2#}, Yuval Benjamini³, Ran Nir-Paz⁴, Mordechai
Muszkat², Sarah Israel^{2§}, Masha Y. Niv^{1§*}

#equal contributions as first authors

§equal contributions as senior authors

*corresponding author

¹The Institute of Biochemistry, Food and Nutrition, The Hebrew University, Rehovot, Israel

²Department of Medicine, Hadassah University Hospital, Mt. Scopus Campus, Jerusalem,
Israel

³Department of Statistics and Data Science, The Hebrew University, Mt. Scopus Campus,
Jerusalem, Israel

⁴Department of Clinical Microbiology and Infectious Diseases, Hadassah-Hebrew University
medical center

Supplementary material

27 symptoms:		
≥50%	<50%	
<u>Taste change</u>	<u>Productive cough</u>	Bad taste in mouth
<u>Smell change</u>	<u>Runny nose</u>	Voice change
<u>Fever</u>	<u>Sore throat</u>	Eyes burn
<u>Dry cough</u>	Chemesthesis change	Dizziness
<u>Muscle aches</u>	<u>Diarrhea</u>	Vision changes
<u>Headache</u>	Chest pain	Coated tongue
<u>Lack of appetite</u>	Abdominal pain	Ears pressure
	<u>Breath difficulty</u>	Lacrimation
	<u>Vomit/nausea</u>	Hearing change
	<u>Fatigue</u>	Eyes discharge

Figure S1. 27 symptoms of the questionnaire. Symptoms with prevalence higher than 50% are presented with orange background, and those with prevalence lower than 50% are presents with light blue background. Official CDC symptoms are bolded. Symptoms that were included in the final analysis are underlined.




Figure S2. Distributions of symptoms' order of appearance. Y axis shows number of patients (n) and X-axis is order of appearance of the specified symptom (first symptom to appear or followed other symptoms).

Table S1. Unresolved symptoms at 6-months follow-up. Number of patients experiencing each symptom is listed.

Symptom	Unresolved
Fatigue	23
Smell change	15
Breath difficulty	10
Taste change	8
Memory disorders	6
Muscle aches	5
Headache	4
Hair loss	3
Nose blockage	2
Legs pain	2
Eye disorders	2
Low physical performance	2
Vomit	1
Throat ache	1
Chest pain	1
Dizziness	1
Knee pain	1
Mouth sores	1
Hands pain	1
Ear pain	1
Paresthesia	1
Palpitations	1
Anemia	1
Anxiety	1
Dry cough	1
Hearing disorder	1
Rib pain	1
Depression	1
Concentration Disorders	1
Runny nose	1
Abdominal pain	1
Diarrhea	1




Figure S3. Correlation between taste change severity and duration (n=76). Y axis shows taste change duration (days) and X axis shows taste change severity.




Figure S4. Smell and taste ratings over 6-weeks follow-ups. A-C. Smell scores of patients recovered at the initial (A), second (B), and third (C) interviews. D-F. Taste scores of patients recovered at the initial (D), second (E), and third (F) interviews. We plot the mean and the confidence interval for the mean. Y axis shows rating of smell/taste ability and X axis depicts 5 time points ratings: before, during and after their illness, rated at initial questionnaire (blue), and at two following questionnaires (3-weeks (light blue) and 6-weeks (turquoise) follow-ups).