

## The Influence of Light Pollution and Light-at-Night on the Circadian Clock: Supplement

### Authors

Dennis Khodasevich<sup>1</sup>, Susan Tsui<sup>1</sup>, Darwin Keung<sup>1</sup>, Debra J. Skene<sup>2</sup>, and Micaela E. Martinez<sup>1\*</sup>

### Affiliations

1. Environmental Health Sciences, Mailman School of Public Health, Columbia University, USA

\*correspondence: mem2353@cumc.columbia.edu

2. Chronobiology, Faculty of Health and Medical Sciences, University of Surrey, UK. \*corresponding author



**Figure S1:** Example of the light sensor (worn around the neck) and wrist temperature sensor (worn around the wrist).



**Figure S2: Outdoor light sensor.** All outdoor light sensors were hung 1 meter off the ground facing North.

	Difference	Lower Est.	Upper Est.	p-value
<b>Outdoor Comparisons</b>				
Spring-Fall	716.55	249.44	1183.67	***
Summer-Fall	1030.92	563.81	1498.04	***
Winter-Fall	-274.93	-742.04	192.18	ns
Summer-Spring	314.37	-152.74	781.48	ns
Winter-Spring	-991.48	-1458.60	-524.37	***
Winter-Summer	-1305.85	-1772.96	-838.74	***
<b>Individual Comparisons</b>				
Spring-Fall	5.62	-212.85	224.09	ns
Summer-Fall	100.48	-95.66	296.62	ns
Winter-Fall	-145.94	-351.55	59.67	ns
Summer-Spring	94.86	-117.96	307.68	ns
Winter-Spring	-151.56	-373.14	70.01	ns
Winter-Summer	-246.42	-446.01	-46.84	**
<b>Individual vs Outdoor</b>				
Fall	1045.30	685.37	1405.24	***
Winter	916.32	554.49	1278.14	***
Spring	1756.24	1386.95	2125.52	***
Summer	1975.74	1619.22	2332.27	***

**Table 1:** Tukey's Honest Significant Difference test results for the mean comparisons of total light exposure AUC summaries. Seasonal comparisons between Outdoor Sensors are shown in the Outdoor section. Seasonal comparisons between Individual Sensors are shown in the Individual section. Comparisons between Outdoor and Individual Sensors within the same season are shown in the Outdoor vs Individual section.

**Random Effects**

	Variance	Std. Dev.
Individual Number	1.621	1.273
Residual	5.571	2.360

**Fixed Effects**

	Estimate	Std. Error	df	t-value	Pr(> t )
(Intercept)	1.65E+01	5.97E-01	1.16E+02	27.65	2.00E-16
Morning Light	-3.95E-03	9.125E-04	2.65E+02	-4.33	2.11E-05
Nighttime Light	2.76E-03	1.14E-03	2.51E+02	2.41	0.0165
Afternoon Light	8.16E-04	8.60E-04	2.68E+02	0.95	0.3435

**Scaled Residuals**

	Min	1Q	Median	3Q	Max
	-2.798	-0.576	0.027	0.470	3.270

**Table S2: Linear Mixed Model Summary Table.** Linear mixed model fit by REML, t-tests use Satterthwaite's method. Accompanying p-values for the fixed effects are under the "Pr(>|t|)" column. Random effects allow for y-intercept to vary between individuals. All input variables had a variance inflation factor below 1.47, suggesting no multicollinearity among the variables.

Season	Individual	Outdoor
Summer	June 21 - August 7	July 2 - July 11
Fall	September 6 - October 10	October 3 - October 13
Winter	January 3 - January 18	January 5 - January 14
Spring	March 25 - April 18	April 5 - April 14

Table S3: Observation periods for individual participants and outdoor sensors.




Figure S3: Fitted periodic temperature data. Predicted wrist temperature from the fitted periodic function. Arrangement of rows is matched to Figure 1c.