Functional Genetic Severity	Score		1	2	3	4	5	6	Statistics
N (% total)	Number of patients		16	0	4	10	8	14	
					- /				
Tumor load	N (%)	Bilateral VS	11 (68,7%)	0 (0%)	3 (75%)	10 (100%)	8 (100%)	14 (100%)	$X^{2}(1) = 8.9, p = 0.002$
		Unilateral VS	5 (31,2%)	0 (0%)	1 (25%)	0 (0%)	0 (0%)	0 (0%)	X ² (1) = 8.9, p = 0.002
		Intracranial meningioma	9 (56,2%)	0 (0%)	3 (75%)	3 (30%)	5 (62,5%)	10 (71,4%)	X ² (1) = 0.35, p = 0.55
		Intracranial Schwannoma	3 (18,7%)	0 (0%)	3 (75%)	3 (30%)	3 (37,5%)	7 (50%)	X ² (1) = 1.8, p = 0.17
		Spinal meningioma	5 (31,2%)	0 (0%)	1 (25%)	1 (10%)	3 (37,5%)	5 (45,5%)	X ² (1) = 0.10, p = 0.74
		Spinal Schwannoma	8 (50%)	0 (0%)	4 (100%)	6 (60%)	6 (75%)	11 (100%)	X ² (1) = 1.7, p = 0.17
		Spinal ependymoma	4 (25%)	0 (0%)	0 (0%)	2 (20%)	3 (37,5%)	5 (45,5%)	X ² (1) = 0.8, p = 0.35
Ocular Features	N (%)	Epiretinal membranes	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	NA
		Cataract	3 (37,5%)	0 (0%)	1 (50%)	3 (30%)	1 (12,5%)	3 (42,9%)	X ² (1) = 0, p = 1
		Combined hamartoma	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	2 (28,6%)	X ² (1) = 3.1, p = 0.07
		Optic nerve meningioma	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (12,5%)	0 (0%)	X ² (1) = 0.3, p = 0.53
	Mean (SD)	Total eye features		0 (0%)					r _s (49) = 0.06, p = 0.67
Hearing Outcomes	N (%)	Hearing grade	1 12 (75%)	0 (0%)	4 (100%)	3 (30%)	3 (37,5%)	6 (42,9%)	X ² (1) = 5.4, p = 0.019
			2 1 (6,2%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	X ² (1) = 1.5, p = 0.20
		3 o 4	0 (0%)	0 (0%)	0 (0%)	0 (0%)	2 (25%)	1 (7,1%)	X ² (1) = 2.1, p = 0.14
			5 2 (12,5%)	0 (0%)	0 (0%)	3 (30%)	1 (12,5%)	4 (28,6%)	X ² (1) = 1.6, p = 0.20
			6 2 (12,5%)	0 (0%)	0 (0%)	4 (40%)	2 (25%)	3 (21,4%)	X ² (1) = 0.6, p = 0.42
		Ago of loss of usoful boaring		0 (00)	15 0 (10 10)	22.22 (0.42)	20.20 (44.0		(40) 0.50 0.004
	Mean (SD)	Age of loss of useful hearing	3 48,31 (17,16)	0 (0%)	46,8 (18,43)	32,38 (8,43)	29,38 (11,8	9) 26,23 (12,58)	r _s (48) = -0.58, p < 0.001
Cutaneus manifestations	N (%)		4 (25%)	0 (0%)	0 (0%)	0 (0%)	4 (50%)	9 (75%)	X ² (1) = 6.2, p = 0.012

* For group 1 and 6, only 4 out 16 and 11 out of 14 patients, respectively, have been tested by a spinal magnetic resonance