

Supplementary Table 2. Major interventions in relation to Genetic Severity Score

Genetic Severity			1 Tissue Mosaic	2A Mild	2B Moderate	3 Severe	Statistics	
N (%)	Proportion of patients whithin	VS surgery	7 (36,84%)	9 (100%)	5 (50%)	10 (71,4%)	$\chi^2(1) = 4.08, p = 0.043$	
		Non-VS intracranial surgery	2 (10,5%)	0 (0%)	3 (30%)	5 (35,7%)	$\chi^2(1) = 3.99, p = 0.045$	
		Spinal surgery	6 (31,58%)	1 (11,11%)	4 (40%)	2 (14,3%)	$\chi^2(1) = 0.72, p = 0.39$	
		Shunt surgery	0 (0%)	0 (0%)	0 (0%)	1 (7,1%)	$\chi^2(1) = 1.73, p = 0.18$	
		Radiotherapy	1 (5,26%)	3 (33,33%)	3 (30%)	8 (57,1%)	$\chi^2(1) = 8.9, p = 0.002$	
		Bevacizumab	0 (0%)	1 (11,11%)	4 (40%)	6 (42,9%)	$\chi^2(1) = 10.2, p = 0.001$	
Total number of major interventions per person, grouped			7 (36,84%)	1 (11,11%)	0 (0%)	2 (14,3%)	$\chi^2(1) = 4.08, p = 0.043$	
Mean (SD)	grouped	0		5 (26,31%)	2 (22,22%)	3 (30%)	$\chi^2(1) = 0.05, p = 0.82$	
		1		2 (10,52%)	4 (44,44%)	2 (20%)	$\chi^2(1) = 0, p = 1$	
		2		1 (5,26%)	3 (33,33%)	1 (10%)	$\chi^2(1) = 0.15, p = 0.69$	
		3		3 (15,79%)	0 (0%)	4 (40%)	$\chi^2(1) = 4.10, p = 0.042$	
		4 or more						
Total number of major interventions per person,			1,53 (1,91)	1,90 (0,99)	3,09 (2,34)	3,79 (3,36)	$r_s(50) = 0.35, p = 0.01$	
Number of total surgeries			1,53 (2,07)	1,50 (0,71)	1,91 (1,64)	2,64 (2,98)	$r_s(50) = 0.18, p = 0.19$	
Age at first radiotherapy session			23	37 (9,64)	30,33 (6,66)	24,29 (8,16)	$r_s(12) = -0.42, p = 0.13$	
Age started bevacizumab			-	32	26,67 (14,57)	23,67 (8,48)	$r_s(8) = -0.26, p = 0.46$	
Age at first surgery			35,44 (13,36)	29,78 (12,05)	26,67 (7,57)	20,80 (7,73)	$r_s(35) = -0.47, p = 0.003$	
Age at first major intervention			34,20 (13,20)	29,78 (12,05)	25,55 (8,55)	20,55 (7,89)	$r_s(39) = -0.47, p = 0.001$	
Ratio of total number of major interventions to current age			0,03 (0,04)	0,04 (0,02)	0,08 (0,06)	0,13 (0,11)	$r_s(50) = 0.49, p < 0.001$	