

eTable 1. Associations of baseline clinical characteristics with the endpoint of death in patients with and without respiratory support (oxygen or intubation).

	Full population	With the end-point event	Without the end-point event	Crude analysis	Multivariable analysis	Collinearity diagnosis (VIF)
	N (%)	N (%)	N (%)	HR (SE) / p-value	HR (SE) / p-value	
<i>With respiratory support</i>	1,192 (100%)	308 (25.8%)	884 (74.2%)			
Sex						1.02
<i>Women</i>	325 (27.3%)	78 (25.3%)	247 (27.9%)	Ref.	Ref.	
<i>Men</i>	867 (72.7%)	230 (74.7%)	637 (72.1%)	1.08 (0.13) / 0.563	1.23 (0.18) / 0.254	
Age						1.01
<i>18 to 50 years</i>	249 (20.9%)	32 (10.4%)	217 (24.5%)	Ref.	Ref.	
<i>51 to 70 years</i>	646 (54.2%)	149 (48.4%)	497 (56.2%)	1.73 (0.19) / 0.005*	1.03 (0.36) / 0.934	
<i>More than 70 years</i>	297 (24.9%)	127 (41.2%)	170 (19.2%)	3.79 (0.20) / <0.001*	2.92 (0.38) / 0.005*	
Obesity ^a						1.02
<i>Yes</i>	345 (28.9%)	92 (29.9%)	253 (28.6%)	1.04 (0.12) / 0.735	0.83 (0.21) / 0.379	
<i>No</i>	847 (71.1%)	216 (70.1%)	631 (71.4%)	Ref.	Ref.	
Smoking ^b						1.01
<i>Yes</i>	181 (15.2%)	38 (12.3%)	143 (16.2%)	0.75 (0.17) / 0.095	0.60 (0.35) / 0.142	
<i>No</i>	1011 (84.8%)	270 (87.7%)	741 (83.8%)	Ref.	Ref.	
Any medical condition ^γ						1.03
<i>Yes</i>	793 (66.5%)	257 (83.4%)	536 (60.6%)	3.47 (0.15) / <0.001*	4.35 (0.20) / <0.001*	
<i>No</i>	399 (33.5%)	51 (16.6%)	348 (39.4%)	Ref.	Ref.	
Clinical severity of Covid-19 at admission ^μ						1.02

<i>Yes</i>	552 (46.3%)	147 (47.7%)	405 (45.8%)	1.21 (0.13) / 0.157	1.53 (0.17) / 0.011*	
<i>No</i>	387 (32.5%)	89 (28.9%)	298 (33.7%)	Ref.	Ref.	
<i>Missing</i>	253 (21.2%)	72 (23.4%)	181 (20.5%)	1.38 (0.16) / 0.041*	1.92 (0.23) / 0.004*	
Biological severity of Covid-19 at admission ^κ						1.01
<i>Yes</i>	768 (64.4%)	193 (62.7%)	575 (65.0%)	0.89 (0.13) / 0.390	1.06 (0.20) / 0.778	
<i>No</i>	287 (24.1%)	79 (25.6%)	208 (23.5%)	Ref.	Ref.	
<i>Missing</i>	137 (11.5%)	36 (11.7%)	101 (11.4%)	1.00 (0.20) / 0.998	1.20 (0.37) / 0.654	
<i>Without respiratory support</i>	11,018 (100%)	1,100 (10.0%)	9,918 (90.0%)			
Sex						1.03
<i>Women</i>	5770 (52.4%)	450 (40.9%)	5320 (53.6%)	Ref.	Ref.	
<i>Men</i>	5248 (47.6%)	650 (59.1%)	4598 (46.4%)	1.50 (0.06) / <0.001*	1.14 (0.09) / 0.134	
Age						1.08
<i>18 to 50 years</i>	4169 (37.8%)	19 (1.73%)	4150 (41.8%)	Ref.	Ref.	
<i>51 to 70 years</i>	3392 (30.8%)	159 (14.5%)	3233 (32.6%)	8.18 (0.24) / <0.001*	4.46 (0.35) / <0.001*	
<i>More than 70 years</i>	3457 (31.4%)	922 (83.8%)	2535 (25.6%)	35.71 (0.23) / <0.001*	21.42 (0.34) / <0.001*	
Obesity ^α						1.02
<i>Yes</i>	1301 (11.8%)	200 (18.2%)	1101 (11.1%)	1.36 (0.08) / <0.001*	1.09 (0.10) / 0.403	
<i>No</i>	9717 (88.2%)	900 (81.8%)	8817 (88.9%)	Ref.	Ref.	
Smoking ^β						1.03
<i>Yes</i>	921 (8.36%)	166 (15.1%)	755 (7.61%)	1.48 (0.08) / <0.001*	0.93 (0.10) / 0.485	
<i>No</i>	10097 (91.6%)	934 (84.9%)	9163 (92.4%)	Ref.	Ref.	
Any medical condition ^γ						1.14
<i>Yes</i>	2712 (24.6%)	637 (57.9%)	2075 (20.9%)	4.66 (0.06) / <0.001*	3.19 (0.09) / <0.001*	
<i>No</i>	8306 (75.4%)	463 (42.1%)	7843 (79.1%)	Ref.	Ref.	

Clinical severity of Covid-19 at admission ^h						1.13
<i>Yes</i>	2039 (18.5%)	480 (43.6%)	1559 (15.7%)	2.53 (0.08) / <0.001*	2.07 (0.11) / <0.001*	
<i>No</i>	2801 (25.4%)	260 (23.6%)	2541 (25.6%)	Ref.	Ref.	
<i>Missing</i>	6178 (56.1%)	360 (32.7%)	5818 (58.7%)	0.75 (0.08) / <0.001*	1.75 (0.13) / <0.001*	
Biological severity of Covid-19 at admission ^k						1.14
<i>Yes</i>	3323 (30.2%)	670 (60.9%)	2653 (26.7%)	2.54 (0.08) / <0.001*	1.72 (0.10) / <0.001*	
<i>No</i>	2925 (26.5%)	243 (22.1%)	2682 (27.0%)	Ref.	Ref.	
<i>Missing</i>	4770 (43.3%)	187 (17.0%)	4583 (46.2%)	0.61 (0.10) / <0.001*	1.15 (0.14) / 0.321	

^a Defined as having a body-mass index higher than 30 kg/m² or an International Statistical Classification of Diseases and Related Health Problems (ICD-10) diagnosis code for obesity (E66.0, E66.1, E66.2, E66.8, E66.9).

^b Current smoking status was self-reported.

^γ Assessed using ICD-10 diagnosis codes for diabetes mellitus (E11), diseases of the circulatory system (I00-I99), diseases of the respiratory system (J00-J99), neoplasms (C00-D49), and diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D5-D8)

^h Defined as having at least one of the following criteria: respiratory rate > 24 breaths/min or < 12 breaths/min, resting peripheral capillary oxygen saturation in ambient air < 90%, temperature > 40°C, or systolic blood pressure < 100 mm Hg.

^k Defined as having at least one of the following criteria: high neutrophil-to-lymphocyte ratio, low *lymphocyte-to-C-reactive protein* (both variables were dichotomized at the median of the values observed in the full sample), and plasma *lactate* levels *higher than 2 mmol/L*.

* p-value is significant (p<0.05).

Abbreviations: HR, hazard ratio; SE, standard error; VIF, variance inflation factor.