

eTable 3. Interaction of baseline characteristics with dexamethasone use on the endpoint of death among adult inpatients with COVID-19.

	With respiratory support	Without respiratory support
	Multivariable Cox regression analysis	Multivariable Cox regression analysis
	HR (SE) / p-value	HR (SE) / p-value
<i>Characteristics</i>		
Sex		
Women	Ref.	Ref.
Men	1.86 (0.22 – 15.99; 0.573)	0.91 (0.18 – 4.57; 0.907)
Age		
18 to 50 years	Ref.	Ref.
51 to 70 years	0.64 (0.11 – 3.66; 0.619)	NA
More than 70 years	0.24 (0.03 – 1.84; 0.172)	NA
Obesity ^a		
Yes	2.54 (0.59 – 11.02; 0.213)	3.90 (1.13 – 13.44; 0.031*)
No	Ref.	Ref.
Smoking		
Yes	1.28 (0.12 – 13.51; 0.839)	0.22 (0.02 – 2.64; 0.233)
No	Ref.	Ref.
Any medical condition ^b		
Yes	0.35 (0.08 – 1.62; 0.181)	1.11 (0.28 – 4.34; 0.880)
No	Ref.	Ref.
Clinical severity of Covid-19 at admission ^h		
Yes	1.35 (0.30 – 6.05; 0.694)	0.30 (0.04 – 2.55; 0.788)
No	Ref.	Ref.
Missing	NA	0.46 (0.12 – 1.78; 0.263)
Biological severity of Covid-19 at admission ^k		
Yes	0.69 (0.10 – 4.70; 0.701)	0.31 (0.09 – 1.01; 0.052)
No	Ref.	Ref.
Missing	3.24 (0.30 – 35.00; 0.332)	0.35 (0.04 – 3.35; 0.362)

^a Defined as having a body-mass index higher than 30 kg/m² or an International Statistical Classification of Diseases and Related Health Problems (ICD-10) diagnosis code for obesity (E66.0, E66.1, E66.2, E66.8, E66.9).

^b Assessed using ICD-10 codes for diabetes mellitus, diseases of the circulatory system, diseases of the respiratory system, neoplasms, and diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism based on ICD-10 classification.

^h Defined as having at least one of the following criteria: respiratory rate > 24 breaths/min, oxygen saturation < 92%, temperature ≥ 40°C, and systolic blood pressure < 100 mm Hg.

^k Defined as having at least one of the following criteria: neutrophil-to-lymphocyte ratio higher than the median in the full sample, lymphocyte-to-C-reactive protein ratio lower than the median in the full sample, and plasma lactate level higher than 2 mmol/L.

* p-value is significant (p<0.05).