



Supplementary Figures

Supplementary Figure 1. Correlation between models. We tested the association of SNPs across the genome with Δ NIHSS using an additive linear model adjusted by sex, age, the two Principal Components SNP genotyping batch, TOAST classification and baseline NIHSS. To test the effect of tPA on the results, we perform a joint analysis in the complete GENISIS cohort including tPA as covariate and compare it to the same model without tPA. The figure show how in both models, the results are highly correlated. The X-axis shows the negative \log_{10} -transformed p-values for the model without tPA; the Y-axis shows the negative \log_{10} -transformed p-values for the model with tPA as covariate

Delta P-value correlation




Supplementary Figure 2. Δ NIHSS Distribution. **A.** Distribution of Δ NIHSS (blue), baseline NIHSS (green) and 24h NIHSS (red) in the complete GENESIS cohort. **B.** Distribution of Δ NIHSS by ethnic background; non-Hispanic white (blue), Hispanic (green), Korea (orange) and US participants with African descent (US AfA – yellow).




Supplementary Figure 3. Association results. Local Manhattan plots for **A.** rs58763243, **C.** rs13403787, **E.** rs12641856, **G.** rs6930598 and **I.** rs10807797 along with the corresponding forest plots, **B,** **D,** **F,** **H** and **J,** showing the contribution of each population to the overall signal.


A.




B.




C.





D.





E.**F.**


Population	Beta	St. Error	P Value
Spain	-0.185	0.239	0.438
Finland	-0.136	0.491	0.782
Poland	0.499	0.596	0.403
US EuA	-0.659	0.528	0.212
Costa Rica	0.872	2.144	0.685
Mexico	MAF<0.03		
Korea	3.715	0.686	1.378E-07
US AfA	MAF<0.03		

Summary 0.110**G.****H.**


Population	Beta	St. Error	P Value
Spain	-0.311	0.203	0.126
Finland	-0.396	0.568	0.486
Poland	0.395	0.629	0.531
US EuA	-0.139	0.498	0.781
Costa Rica	-0.561	1.444	0.698
Mexico	-4.864	1.539	0.003
Korea	2.615	0.529	1.33E-06
US AfA	-0.512	0.583	0.381

Summary -0.070**I.****J.**


Population	Beta	St. Error	P Value
Spain	-0.531	0.131	4.804E-05
Finland	-0.634	0.268	0.018
Poland	-0.130	0.337	0.701
US EuA	-0.530	0.291	0.069
Costa Rica	-1.880	0.793	0.019
Mexico	0.881	0.838	0.298
Korea	0.487	0.288	0.092
US AfA	0.425	0.481	0.378





Summary -0.370





Supplementary Figure 4. Single-nuclei data on brain cortex. Expression patterns across the brain cortex cell populations derived from single nuclei RNAseq data for for **A.** *GALNT10*, candidate gene along with *GRIA1* for the genome wide locus in chr.5 lead by rs114248865; **B.** *DFNB59* also known as *PJKV*, candidate gene for the genome wide locus in chr.2 lead by rs13403787, **C.** *TWISTNB*, candidate gene for the genome wide locus in chr6 along with **D.** *ABCB5* lead by rs10807797.




Supplementary Figure 5. PRSice results. We used the software PRSice2 to perform comparisons using polygenic risk scores at different p-value thresholds between the genetic architecture of Δ NIHSS and **A and B**. stroke risk, **C and D** age at death, **E and F** total cholesterol, **G and H** triglycerides, **I and J** high density lipoproteins, **K and L** low density lipoproteins and **M and N** body mass index.



E.**F.****G.****H.**

I.**J.****K.****L.**

M.**N.**