

Supplementary 1

The analysis of COVID-19 and Age among the never smokers and those with history of smoking (current smokers and former smokers)

Supplementary 1: Table S1

Table S1: Baseline characteristics of never smokers

	COVID-19 positive 71 (36.60%)	COVID-19 Negative 123 (63.4 %)
Demographics		
Age, median (IQR) [range]	60 (47 -70) [20 - 90]	65 (48-78) [24-97]
Males no (%)	37 (52.11)	58 (47.54)
Labs		
Lymphopenia ≤ 1500	52 (74.29)	71 (60.17)
Lymphopenia ≤ 1000	38 (54.29)	43 (36.44) *
Medical Comorbidities No (%)		
Diabetes Mellitus	14 (19.72)	38 (31.15)
Dementia	1 (1.41)	7 (5.74)
Stroke	5 (7.04)	12 (9.84)
Parkinson's	1 (1.41)	3 (2.46)
Multiple sclerosis	0 (0)	1 (0.82)
Demyelinating Disease	1 (1.41)	0 (0)
End Stage Renal Disease	0 (0)	5 (4.13)
CKD ≥ Stage 3	3 (4.23)	9 (7.38)
CHF history	3 (4.23)	17 (13.93) *
COPD	1 (1.41)	7 (5.74)
Asthma	7 (9.86)	17 (13.93)
SLE	0 (0)	2 (1.65)
Organ Transplant	0 (0)	3 (2.46)
Hypertension	27 (38.03)	58 (47.54)
CAD	6 (8.57)	17 (14.05)
BMI ≥ 30	28 (48.28)	45 (45.45)
HIV	0 (0.00)	0 (0.00)
Non-COVID-19 respiratory viral illness	0 (0)	17 (15.6)

Abbreviations: COPD chronic obstructive pulmonary disease; CAD coronary artery disease; BMI body mass index; SLE systemic lupus erythromatosus; CHF congestive heart failure; CKD chronic kidney disease; IQR Interquartile range; COVID-19 corona virus disease 2019 ; HIV human immunodeficiency virus.

The asterisk indicates that the P value is less than 0.05 in the comparison of COVID-19 positive and negative column; Values in the COVID-19 negative columns without an asterisk have p values ≥0.05; P values were calculated using the independent t-test.

Supplementary 1: Table S2

Table S2: Baseline characteristics of those with history of smoking (current and former smokers)

	COVID-19 positive 63 (27.75%)	COVID-19 Negative 164 (72.25 %)
Demographics		
Age, median (IQR) [range]	61 (44 -76) [20 - 88]	66 (55-76) [20-97]
Males no (%)	43 (68.25)	83 (50.61) *
Labs		
Lymphopenia \leq 1500	46 (73.02)	101 (62.73)
Lymphopenia \leq 1000	28 (44.44)	77 (47.83)
Medical Comorbidities No (%)		
Diabetes Mellitus	16 (25.40)	51 (31.29)
Dementia	4 (6.35)	13 (7.98)
Stroke	5 (7.94)	24 (14.72)
Parkinson's	3 (4.76)	2 (1.23)
Multiple sclerosis	0 (0)	6 (3.70)
Demyelination Disease	1 (1.59)	3 (1.84)
End Stage Renal Disease	0 (0)	7 (4.38)
CKD \geq Stage 3	12 (19.05)	19 (11.66)
CHF history	7 (11.11)	32 (19.63)
COPD	8 (12.70)	61 (37.42) *
Asthma	8 (12.70)	19 (11.66)
SLE	0 (0)	4 (2.45)
Organ Transplant	0 (0)	1 (0.62)
Hypertension	29 (46.03)	93 (57.06)
CAD	11 (17.46)	35 (21.47)
BMI \geq 30	27 (51.92)	66 (46.48)
HIV	0 (0)	2 (1.23)
Non-COVID-19 respiratory viral Illness	0 (0)	32 (21.33)

Abbreviations: COPD chronic obstructive pulmonary disease; CAD coronary artery disease; BMI body mass index; SLE systemic lupus erythromatosus; CHF congestive heart failure; CKD chronic kidney disease; IQR Interquartile range; COVID-19 corona virus disease 2019 ; HIV human immunodeficiency virus.

The asterisk indicates that the P value is less than 0.05 in the comparison of COVID-19 positive and negative column; Values in the COVID-19 negative columns without an asterisk have p values \geq 0.05; P values were calculated using the independent t-test.

Supplementary 1: Table S3

Table S3 below shows homogeneity test. The difference between never smokers and those with history of smoking (current plus former smokers) among the COPD, CHF, Parkinson disease and the 40 to 49-year olds is significantly different.

Table S3: The homogeneity test				
	COVID-19 Status	P-Value		RR (Confidence Interval)
Smoking Status	Never Smokers	0.524		1.023 (0.954-1.097)
Demographics	Age Categories	20	Reference	1
		40	0.027	1.292 (1.030-1.620)
		50	0.048	1.089 (1.001-1.184)
		70	0.343	1.040 (0.959-1.128)
	AgeCat6#NeverSmokers	20 2	Reference	1
		40 2	0.034	0.779 (0.618-0.982)
		50 2	0.740	0.980 (0.872-1.102)
		70 2	0.734	1.021 (0.904-1.154)
	Gender	0.018		1.074 (1.012-1.140)
	Medical comorbidities	COPD#c.NeverSmokers	No	Reference
Yes			0.018	0.950 (0.910-0.991)
CHF#c.NeverSmokers		No	Reference	1
		Yes	0.009	0.958 (0.928-0.989)
Diabetes		0.712		0.987 (0.921-1.058)
Hypertension		0.068		1.067 (0.995-1.143)
Non-COVID-19 respiratory viral illness		0.000		0.910 (0.866 - 0.955)
BMI4		0.136		1.004 (0.999-1.008)
Lymphocyt1e		0.525		1.020 (0.961-1.082)
Asthma		0.354		1.054 (0.943-1.179)
CKD#c.NeverSmokers		No	Reference	1
		Yes	0.902	0.995 (0.923-1.073)
SLE		0.119		0.891 (0.771-1.030)
Organ Transplant		0.107		0.898 (0.789-1.023)
CAD		0.690		1.017 (0.936-1.105)
HIV		0.001		0.860 (0.785-0.943)
Parkinsons#c.NeverSmoker		No	Reference	1
		Yes	0.023	0.935 (0.883-0.991)
Multiple sclerosis		0.078		0.912 (0.823-1.011)
ESRD		0.024		0.916 (0.849-0.989)
Stroke#c.NeverSmoker		No	Reference	1
		Yes	0.229	0.972 (0.927-1.018)
Demyelinatingdisease		0.291		0.929 (0.809-1.066)

Supplementary 1: Table S4

Table S4 below shows the results of the association between COVID-19 status and age without adjusting for never smoking. The 50 to 69-year olds had a higher risk of testing positive for COVID-19. Besides that, males had a higher risk of testing positive for COVID-19, while patients with CHF history, ESRD, HIV positive and non-COVID-19 respiratory viral illness patients had a lower risk of testing positive for COVID-19.

	COVID-19 Status	P-value		RR (Confidence Interval)	
Age Categories	20 to 39 years	Reference		1	
	40 to 49 years	0.056		1.152 (0.997 - 1.331)	
	50 to 69 years	0.017		1.082 (1.014 - 1.154)	
	70 to 99 years	0.289		1.046 (0.963 - 1.135)	
Smoking Status	Never Smokers	0.808		0.992 (0.930 - 1.059)	
Demographics	Gender	Female	Reference		1
		Male	0.008		1.084(1.021 - 1.151)
Labs	Lymphopenia (≤ 1500)	0.757		1.009 (.951 - .0715)	
Medical Comorbidities	COPD	0.083		0.950 (0.896 - 1.007)	
	Non-COVID-19 respiratory viral illness	Absent	Reference	1	
		Present	0.000		0.905 (0.860 - 0.953)
	Hypertension	0.109		1.061 (0.987 - 1.141)	
	Diabetes	0.573		0.980 (0.915 - 1.051)	
	Asthma	0.422		1.047 (0.936 - 1.171)	
	SLE	0.301		0.945 (0.848 - 1.052)	
	Organ Transplant	0.090		0.898 (0.792 - 1.017)	
	CKD \geq Stage 3	0.570		1.035 (0.919 - 1.166)	
	CHF history	Absent	Reference	1	
		Present	0.018		0.946 (0.904 - 0.990)
	CAD	0.726		1.015(0.935-1.101)	
	Parkinson's Disease	0.082		0.911(0.819 - 1.012)	
	Multiple sclerosis	0.183		0.946(0.872- 1.027)	
	ESRD	Absent	Reference	1	
		Present	0.011		0.908(0.843- 0.978)
	Stroke	0.355		0.966(0.897 - 1.040)	
	Demyelinating disease	0.296		0.930(0.811 - 1.066)	
	HIV	Absent	Reference	1	
		Present	0.001		0.870 (0.804 - 0.941)
BMI \geq 30	0.193		1.003 (0.998 - 1.008)		

Supplementary 1: Table S5

Table S5 below shows the results of the association between COVID-19 status and age without adjusting for never smoking. The 50 to 69-year olds and males were more likely to test positive for COVID-19, while CHF history, ESRD, HIV positive and patients with non-COVID-19 respiratory viral illness were less likely to test positive for COVID-19.

Table S5: The Association between COVID-19 Status and Age with adjusting for never smokers				
	COVID-19 Status	P-value		RR (Confidence Interval)
Age Categories	20 to 39 years	Reference		1
	40 to 49 years	0.048		1.002(1.002 - 1.325)
	50 to 69 years	0.015		1.081(1.016 - 1.150)
	70 to 99 years	0.312		1.041 (0.963 - 1.126)
Demographics	Gender	0.007		1.085(1.022 - 1.151)
Labs	Lymphopenia (≤ 1500)	0.771		1.009 (0.951 - 0.070)
Medical Comorbidities	COPD	0.111		0.955 (0.902 - 1.011)
	Non-COVID-19 respiratory viral illness	Absent	Reference	1
		Present	0.000	0.907 (0.863 - 0.953)
	Hypertension	0.095		1.062 (0.989 - 1.140)
	Diabetes	0.568		0.980 (0.916 - 1.049)
	Asthma	0.411		1.048 (0.937 - 1.172)
	SLE	0.167		0.945 (0.873 - 1.024)
	Organ Transplant	0.063		0.894 (0.793 - 1.006)
	CKD \geq Stage 3	0.539		1.039 (0.920 - 1.173)
	CHF history	Absent	Reference	1
		Present	0.018	0.947 (0.905 - 0.990)
	CAD	0.711		1.015(0.937-1.008)
	Parkinson's Disease	0.080		0.912 (0.822 - 1.011)
	Multiple sclerosis	0.148		0.949 (0.884- 1.019)
	ESRD	Absent	Reference	1
		Present	0.012	0.909(0.843- 0.980)
	Stroke	0.383		0.968 (0.899 - 1.042)
	Demyelinating disease	0.297		0.932(0.817 - 1.064)
	HIV	Absent	Reference	1
		Present	0.001	0.876 (0.813 - 0.944)
	BMI \geq 30	0.195		1.003 (0.998 - 1.008)

Supplementary 1: Table S6

Table S6 below shows the analysis results the COVID-19 status and age among those with history of smoking (current and former smokers). The 40 to 49-year olds were more likely to test positive for COVID-19. While the HIV positive and non-COVID-19 respiratory viral illness patients were less likely to test positive for COVID-19.

Table S6: The Association between COVID-19 Status and Age among current and former smokers				
	COVID-19 Status	P-value		RR (Confidence Interval)
Age Categories	20 to 39 years	Reference		1
	40 to 49 years	0.018		1.309 (1.047 - 1.635)
	50 to 69 years	0.143		1.063 (0.980 - 1.153)
	70 to 99 years	0.817		0.990 (0.907 - 1.080)
Demographics	Gender	0.053		1.064 (0.999 - 1.132)
Labs	Lymphopenia (≤ 1500)	0.270		1.035 (0.974 - 1.100)
Medical Comorbidities	COPD	0.707		.989 (.933 - 1.048)
	Non-COVID-19 respiratory viral illness	Absent	Reference	1
		Present	0.005	0.907 (0.846 - 0.972)
	Hypertension	0.104		1.080 (0.984 - 1.186)
	Diabetes	0.150		0.944 (0.874 - 1.021)
	Asthma	0.758		1.021 (.894 - 1.167)
	SLE	0.188		0.918 (0.808 - 1.043)
	Organ Transplant	Omitted		1
	CKD \geq Stage 3	0.122		1.126 (0.969 - 1.308)
	CHF history	Absent	Reference	1
		Present	0.061	0.943 (0.887 - 1.003)
	CAD	0.592		1.023 (0.941 - 1.113)
	Parkinson's disease	0.555		1.032 (0.930 - 1.144)
	Multiple sclerosis	0.098		0.898 (0.791 - 1.020)
	ESRD	0.092		0.936 (0.867 - 1.011)
	Stroke	0.680		0.980 (0.892 - 1.077)
	Demyelinating disease	0.147		0.929 (0.841 - 1.026)
	HIV	Absent	Reference	1
		Present	0.006	0.863 (0.775 - 0.959)
	BMI ≥ 30	0.359		1.003 (.997 - 1.008)

Supplementary 1: Table S7

Table S7 below shows the results of the association between COVID-19 and age among the never smokers. Age was not associated with testing positive for COVID-19. Stroke, COPD, CHF history, CKD, Parkinson's disease and non-COVID-19 respiratory viral illness patients were less likely to test positive for COVID-19.

Table S7: The Association between COVID-19 Status and Age among never smokers				
	COVID-19 Status	P-value		RR (Confidence Interval)
Age Categories	20 to 39 years	Reference		1
	40 to 49 years	0.614		0.976 (0.890- 1.071)
	50 to 69 years	0.207		1.049 (0.974 - 1.130)
	70 to 99 years	0.490		1.051 (0.913 - 1.209)
Demographics	Gender	0.107		1.091 (0.981 - 1.213)
Lab	Lymphopenia (≤ 1500)	0.795		1.015 (0.906 - 1.137)
Medical Comorbidities	COPD	Absent	Reference	1
		Present	0.006	0.842 (0.745 - 0.952)
	Non-COVID-19 respiratory viral illness	Absent	Reference	1
		Present	0.008	0.902 (0.837 - 0.973)
	Hypertension	0.212		1.063 (0.966 - 1.171)
	Diabetes	0.453		1.040 (0.939 - 1.151)
	Asthma	0.370		1.087 (0.905 - 1.306)
	SLE	0.252		0.914 (0.785 - 1.065)
	Organ Transplant	0.219		0.906 (0.773 - 1.061)
	CKD \geq Stage 3	Absent	Reference	1
		Present	0.040	0.879 (0.777 - 0.994)
	CHF history	Absent	Reference	1
		Present	0.032	0.907 (0.830 - 0.991)
	CAD	0.921		1.007 (0.875 - 1.159)
	Parkinson's Disease	Absent	Reference	1
		Present	0.011	0.823 (0.708 - 0.957)
	Multiple sclerosis			Omitted 1
	ESRD	0.174		0.896 (0.764 - 1.050)
	Stroke	Absent	Reference	1
		Present	0.017	0.903 (0.831 - 0.982)
	Demyelinating disease			Omitted 1
	HIV			Omitted 1
	BMI \geq 30	0.541		1.003 (0.995 - 1.010)

Abbreviations

ACE-2 angiotensin converting enzyme 2

CAD coronary artery disease

CDC center for disease control

COPD chronic obstructive pulmonary disease

BMI body mass index

SLE systemic lupus erythromatosus

CHF congestive heart failure

ESRD end stage renal disease

CKD chronic kidney disease

COVID-19 corona virus disease 2019

HIV human immunodeficiency virus

Interquartile range IQR

Polymerase chain reaction PCR

RSV respiratory syncytial virus

USA united states of America

UK United Kingdom