

Supplementary 2

The analysis of COVID-19 and Age among the never smokers and current smokers (excluding the former smokers)

Table T1 below shows the association between COVID-19 and age adjusting for smoking status. The 50 to 69 year old patients were more likely to test positive for COVID-19, while COPD, Parkinson’s disease and demyelinating disease patients were less likely to test positive for COVID-19.

Table T1: The Association between COVID-19 Status and Age adjusting for smoking status				
	COVID-19	P-Value	RR (Confidence Interval)	
Smoking status	Never-Current Smokers	0.485	0.973 (0.901-1.051)	
Demographics	Age Categories	20-39 years	Reference 1	
		40-49 years	0.289 1.090 (0.930-1.278)	
		50-69 years	0.019 1.084 (1.013-1.159)	
		70-100 years	0.076 1.130 (0.987-1.293)	
	Gender	0.060	1.081 (0.997-1.173)	
Medical comorbidities	COPD	Absent	Reference 1	
		Present	0.010 0.874 (0.789-0.968)	
	CHF		0.096	0.943 (0.879-1.011)
	Diabetes		0.829	1.010 (0.925-1.102)
	Hypertension		0.396	1.037 (0.954-1.128)
	Non-COVID-19 respiratory viral illness	Absent	Reference 1	
		Present	0.005 0.904 (0.842- 0.971)	
	BMI4		0.090	1.006 (0.999-1.014)
	Lymphocyte 1500		0.874	0.994 (0.918-1.075)
	Asthma		0.565	1.035 (0.920-1.164)
	CKD		0.072	0.915 (0.831-1.008)
	SLE		0.759	0.968 (0.789-1.189)
	Organ Transplant		0.393	0.946 (0.834-1.074)
	CAD		0.501	0.967 (0.878-1.065)
	HIV			1
	Parkinson’s disease	Absent	Reference 1	
		Present	0.008 0.816 (0.702-0.905)	
	Multiple sclerosis		0.812	0.983 (0.853-1.133)
	ESRD		0.212	0.938 (0.849-1.037)
	Stroke		0.211	0.956 (0.891-1.026)
Demyelinating disease	Absent	Reference 1		
	Present	0.036 0.860 (0.747-0.990)		
	_cons	0.026	1.636 (1.06-2.523)	

Table T2 below shows the association between COVID-19 and age without adjusting for smoking status. The males and 50 to 69 year old patients were more likely to test positive for COVID-19, while CHF, HIV positive, ESRD and demyelinating disease patients were less likely to test positive for COVID-19.

Table T2: The Association between COVID-19 Status and Age without adjusting for Smoking status				
	COVID-19		P-Value	RR (Confidence Interval)
Demographics	Age Categories	20-39 years	Reference	1
		40-49 years	0.048	1.152 (1.002-1.325)
		50-69 years	0.015	1.081 (1.016-1.150)
		70-100 years	0.312	1.041 (0.963-1.126)
	Gender	Female	Reference	1
		Male	0.007	1.085 (1.022-1.151)
Medical comorbidities	COPD		0.111	0.955 (0.902-1.010)
	CHF	Absent	Reference	1
		Present	0.018	0.947 (0.905-0.991)
	Diabetes		0.568	0.980 (0.916-1.049)
	Hypertension		0.095	1.062 (0.990-1.140)
	Non-COVID-19 respiratory viral illness	Absent	Reference	1
		Present	0.000	0.908 (0.863-0.953)
	BMI		0.195	1.003 (0.998-1.008)
	Lymphocyte		0.771	1.009 (0.951-1.070)
	Asthma		0.411	1.048 (0.937-1.172)
	CKD		0.539	1.039 (0.920-1.173)
	SLE		0.167	0.945 (0.872-1.024)
	Organ Transplant		0.063	0.894 (0.793-1.006)
	CAD		0.711	1.015 (0.937-1.101)
	HIV	Absent	Reference	1
		Present	0.001	0.876 (0.813-0.944)
	Parkinson's disease		0.080	0.911 (0.822-1.011)
	Multiple sclerosis		0.148	0.949 (0.884-1.019)
	ESRD	Absent	Reference	1
		Present	0.012	0.909 (0.843-0.979)
Stroke		0.383	0.968 (0.899-1.042)	
Demyelinating disease		0.297	0.932 (0.817-1.064)	
_cons		0.145	1.343 (0.904-1.997)	

Table T3 below shows the association between COVID-19 and age among never smokers. Male never smokers were more likely to test positive for COVID-19, while CHF, organ transplant, HIV positive, ESRD, non-COVID-19 respiratory viral illness patients were less likely to test positive for COVID-19.

Table T3: The Association between COVID-19 Status and Age among the never smokers				
	COVID-19		P-Value	RR (95% Conf. Interval)
Demographics	Age Categories	20-39 years	Reference	1
		40-49 years	0.212	1.101 (0.946-1.281)
		50-69 years	0.045	1.078 (1.002-1.160)
		70-100 years	0.829	1.009 (0.927-1.098)
	Gender	Female	Reference	1
		Male	0.008	1.010 (1.025-1.180)
Medical comorbidities	COPD		0.149	0.955 (0.896-1.017)
	CHF	Absent	Reference	1
		Present	0.007	0.931 (0.884-0.981)
	Diabetes		0.843	0.993 (0.922-1.068)
	Hypertension		0.084	1.074 (0.991-1.163)
	Non-COVID-19 respiratory viral illness	Absent	Reference	1
		Present	0.000	0.896 (0.847-0.949)
	BMI		0.704	1.001 (0.996-1.006)
	Lymphocyte		0.646	1.017 (0.945-1.096)
	Asthma		0.261	1.083 (0.943-1.243)
	CKD		0.529	1.042 (0.917-1.184)
	SLE		0.092	0.938 (0.870-1.011)
	Organ Transplant	Absent	Reference	1
		Present	0.024	0.860 (0.755-0.980)
	CAD		0.533	1.030 (0.937-1.134)
	HIV	Absent	Reference	1
		Present	0.001	0.875 (0.809-0.947)
	Parkinson's disease		0.171	0.917 (0.811-1.038)
	Multiple sclerosis		0.735	0.981 (0.879-1.095)
	ESRD	Absent	Reference	1
		Present	0.007	0.888 (0.814-0.969)
	Stroke		0.232	0.946 (0.865-1.036)
	Demyelinating disease		0.757	1.018 (0.908-1.142)
_cons		0.308	1.243 (0.819-1.886)	

Table T4 below shows the association between COVID-19 and age among the current smokers. The 40 to 49 year old current smokers were more likely to test positive for COVID-19, while HIV positive, Non-COVID-19 respiratory viral illness current smokers while less likely to test positive for COVID-19.

Table T4: The Association between COVID-19 Status and Age among the current smokers				
	COVID-19		P-Value	RR [95% Conf. Interval]
Demographics	Age	20-39 years	Reference	1
		40-49 years	0.018	1.309 (1.047-1.635)
		50-69 years	0.143	1.063 (0.980-1.153)
		70-100 years	0.817	0.990 (0.907-1.080)
	Gender		0.053	1.064 (0.999-1.132)
Medical comorbidities	COPD		0.707	0.989 (0.933-1.048)
	CHF		0.061	0.943 (0.887-1.003)
	Diabetes		0.150	0.945 (0.874-1.021)
	Hypertension		0.104	1.080 (0.984-1.186)
	Non-COVID-19 respiratory viral illness	Absent	Reference	1
		Present	0.005	0.907 (0.846 -0.972)
	BMI4		0.359	1.003 (0.997-1.008)
	Lymphocyte		0.270	1.035 (0.974-1.100)
	Asthma		0.758	1.021 (0.894-1.167)
	CKD		0.122	1.125 (0.969-1.308)
	SLE		0.188	0.918 (0.808-1.043)
	Organ Transplant			1
	CAD		0.592	1.023 (0.941-1.113)
	HIV	Absent		
		Present	0.006	0.863 (0.775-0.959)
	Parkinson's disease		0.555	1.032 (0.930-1.144)
	Multiple sclerosis		0.098	0.898 (0.791-1.020)
	ESRD		0.092	0.936 (0.867-1.011)
	Stroke		0.680	0.980 (0.892-1.077)
	Demyelinating disease		0.147	0.929 (0.841-1.026)
_cons		0.906	1.025 (0.678-1.549)	

Table T5 below shows the homogeneity test. The difference between never smokers and current smokers among the COPD, CHF, non-COVID-19 respiratory viral illness and Parkinson's disease is significantly different as shown by the homogeneity test in Supplementary table S3.

Table T5: The homogeneity test				
	COVID-19		P-Value	RR [95% Conf. Interval]
Smoking Status	Never-Current Smokers		0.434	0.973 (0.908- 1.042)
Demographics	Age Categories	20-39	Reference	1
		40-49	0.146	1.267 (0.921- 1.742)
		50-59	0.046	1.063 (1.001- 1.128)
		70-100	0.047	1.120 (1.001- 1.252)
	2.NevCurSmokers		Reference	1
	AgeCat6#NevCurSmokers	20 2	Reference	1
		40 2	0.142	0.782 (0.563- 1.086)
		50 2	0.837	1.011 (0.911- 1.122)
		70 2	0.728	0.975 (0.844- 1.126)
	Gender#c.NevCurSmokers		Reference	1
Male		0.096	1.044 (0.992- 1.098)	
Medical comorbidities	COPD#c.NevCurSmokers		1	Reference
		Yes	0.006	0.921 (0.868- 0.977)
	CHF#c.NevCurSmokers		1	Reference
		Yes	0.029	0.959 (0.923- 0.996)
	Diabetes		0.602	1.023 (0.940- 1.112)
	Hypertension		0.541	1.026 (0.945- 1.113)
	Viral1#c.NevCurSmokers		Reference	1
		2 Yes	0.003	0.950 (0.918 - 0.983)
	BMI4		0.086	1.006 (0.999- 1.013)
	Lymphocyt1e		0.988	1.001 (0.925- 1.082)
	Asthma		0.553	1.038 (0.918- 1.172)
	CKD		0.058	0.907 (0.820- 1.003)
	SLE		0.247	0.817 (0.580- 1.150)
	Organ Transplant		0.262	0.930 (0.820- 1.056)
	CAD		0.891	0.993 (0.902- 1.094)
	HIV		Reference	1
	Parkinson's disease		0.006	0.819 (0.709- 0.945)
	Multiple sclerosis		0.605	1.034 (0.911- 1.174)
	ESRD#c.NevCurSmokers		Reference	1
		Yes	0.311	0.966 (0.904- 1.033)
	Stroke		0.110	0.945 (0.882- 1.013)
	Demyelinating disease		0.153	0.923 (0.826- 1.030)

Abbreviations

ACE-2 angiotensin converting enzyme 2

CAD coronary artery disease

CDC center for disease control

COPD chronic obstructive pulmonary disease

BMI body mass index

SLE systemic lupus erythromatosus

CHF congestive heart failure

ESRD end stage renal disease

CKD chronic kidney disease

COVID-19 corona virus disease 2019

HIV human immunodeficiency virus

Interquartile range IQR

Polymerase chain reaction PCR

RSV respiratory syncytial virus

USA united states of America

UK United Kingdom