

## Surgery & COVID-19: A rapid scoping review of the impact of COVID-19 on surgical services during public health emergencies

Connor M O’Rielly MSc,<sup>1</sup> Joshua S Ng-Kamstra MD,<sup>2</sup> Ania Kania-Richmond PhD,<sup>3</sup> Joseph C Dort MD,<sup>1,4</sup> Jonathan White MD, PhD<sup>5</sup> Jill Robert BScN,<sup>3,6</sup> Mary Brindle MD,<sup>7</sup> Khara M Sauro PhD<sup>1,4</sup>

<sup>1</sup> Department of Community Health Sciences & O’Brien Institute for Public Health, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>2</sup> Department of Critical Care Medicine, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>3</sup> Bone and Joint Health Strategic Clinical Network, Alberta Health Services, Alberta, Canada

<sup>4</sup> Departments of Surgery & Oncology, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

<sup>5</sup> Department of Surgery, Faculty of Medicine & Dentistry, University of Alberta, Edmonton, Alberta, Canada

<sup>6</sup> Surgery Strategic Clinical Network, Alberta Health Services, Alberta, Canada

<sup>7</sup> Department of Pediatrics, Cumming School of Medicine, University of Calgary, Calgary, Alberta, Canada

Corresponding Author:

Khara Sauro

3280 Hospital Dr. NW, Room 3D41 | Calgary, AB | T2N 4Z6

T: 403-210-5463

E: [kmsauro@ucalgary.ca](mailto:kmsauro@ucalgary.ca)

Word Count: 3556 words

Keywords: public health emergency, surgical services, healthcare resource utilization, health services

**ABSTRACT** (275/300 words)

**Background:** Healthcare systems globally have been challenged by the COVID-19 pandemic, necessitating the reorganization of surgical services to free capacity within healthcare systems.

**Objectives:** To understand how surgical services have been reorganized during and following public health emergencies, and the consequences of these changes for patients, healthcare providers and healthcare systems.

**Methods:** This rapid scoping review searched academic databases and grey literature sources to identify studies examining surgical service delivery during public health emergencies including COVID-19, and the impact on patients, providers and healthcare systems. Recommendations and guidelines were excluded. Screening was completed in partial (title, abstract) or complete (full text) duplicate following pilot reviews of 50 articles to ensure reliable application of eligibility criteria.

**Results:** One hundred and thirty-two studies were included in this review; 111 described reorganization of surgical services, 55 described the consequences of reorganizing surgical services and six reported actions taken to rebuild surgical capacity in public health emergencies. Reorganizations of surgical services were grouped under six domains: case selection/triage, PPE regulations and practice, workforce composition and deployment, outpatient and inpatient patient care, resident and fellow education, and the hospital or clinical environment. Service reorganizations led to large reductions in non-urgent surgical volumes, increases in surgical wait times, and impacted medical training (i.e., reduced case involvement) and patient outcomes (e.g.,

increases in pain). Strategies for rebuilding surgical capacity were scarce, but focused on the availability of staff, PPE, and patient readiness for surgery as key factors to consider before resuming services.

**Conclusions:** Reorganization of surgical services in response to public health emergencies appears to be context-dependent and has far-reaching consequences that must be better understood in order to optimize future health system responses to public health emergencies.

## ARTICLE SUMMARY

### **Strengths and limitations of the study:**

- This rapid scoping review provides an exhaustive and rigorous summary of the academic and grey literature regarding modifications to surgical services in response to public health emergencies, especially COVID-19.
- This study did not limit studies based on location or language of publication to ensure a worldwide pandemic had contributions from worldwide voices.
- Both quantitative and qualitative outcomes were included, with a mix of inductive and deductive data abstraction approaches to provide a comprehensive understanding of surgical services during public health emergencies.
- Studies with potential relevance to this question are emerging at an unprecedented rate in response to the COVID-19 pandemic and as such, some may not be included in the current review.

**Original protocol for the study:** As requested, the original unpublished protocol for this study is included as a supplementary file.

**Funding statement:** This study did not receive grant from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interest statement:** All authors declare that they have no competing interests in accordance with the International Committee of Medical Journal Editors uniform declaration of competing interests.

## INTRODUCTION

The novel SARS-CoV-2 (COVID-19) virus has spread across the globe with unrelenting speed. At the time of writing, over 16 million cases have been confirmed with 650,000 fatalities.<sup>1</sup> To protect the most vulnerable in our societies, efforts to curb further escalation (e.g., travel restrictions, physical distancing) have had a focal objective: to prevent surges that could overwhelm healthcare including shortages in personal protective equipment (PPE), ventilators, and hospital capacity.

Medical institutions have taken steps to maximize staff, PPE, ventilators, and intensive care unit (ICU) capacity in case public health efforts to ‘flatten the curve’ are insufficient. Most notably, surgical programs have suspended non-urgent (or ‘elective’) surgical procedures. Non-urgent surgeries are often defined as procedures for which a delay of three months or longer would not result in significant adverse effects to the patient.<sup>2,3</sup> These changes have thrust patients, providers, and healthcare organizations into previously unexplored territory.

While governing bodies such as colleges and academies of surgery have made recommendations to alter surgical service delivery in response to COVID-19, they have not always provided explicit instructions on how programs should operationalize the recommendations. As such, approaches to surgical triage and service delivery remain unclear: who has done what where, and why? Further, the impacts of adopting these recommendations on surgical programs, and more importantly, the physical and psychological well-being of patients and healthcare providers have only been hypothesized.<sup>4</sup> Lastly, as COVID-19 begins to release its grip on the world and the post-pandemic recovery begins, programs will be tasked with rebuilding the surgical capacity necessary to reschedule and resume the backlog of postponed procedures. Evidence distilled from the experiences of others in the context of COVID-19 and

other public health emergencies (i.e., H1N1, Ebola, SARS) is needed to guide approaches to surgical service delivery.

To enable evidence-informed reorganization and resumption of non-urgent surgeries post-COVID-19 and for future public health emergencies, we conducted a rapid scoping review to synthesize relevant and available literature. Our aim was to understand how surgical services were reorganized in response to COVID-19 and other public health emergencies; how reorganization impacted patients, healthcare providers, and health systems; and what approaches have been taken to resume surgical service delivery.

## **METHODS**

### **Study Design**

This scoping review followed the Joanna Briggs Institute methodology and Preferred Reporting Items for Systematic reviews and Meta-Analysis extension for Scoping Reviews (PRISMA-ScR) checklist.<sup>5 6</sup> The rapidly evolving situation of the current COVID-19 pandemic demanded a similarly rapid evidence synthesis. Therefore, methodological concessions recommended by the World Health Organization and Cochrane guidance for rapid reviews were made.<sup>7 8</sup> This review addressed three questions: 1) How have surgical services been reorganized in response to public health emergencies? 2) What are the patient-, healthcare provider-, and system-level consequences of reorganizing surgical services? and 3) What approaches were used for resuming surgical services?

### **Search Strategy**

The search strategy was developed by two investigators (CO, KS) and refined by others with context expertise in surgery and literature review methodology (JNK, AKR). The search strategy included subject headings, keywords, and synonyms identifying public health emergencies in general and specific public health emergencies (Ebola, SARS-CoV1, H1N1, MERS), and surgery; and were tailored for each database (Appendix A). Given the exploratory nature of the review we did not filter by study design or publication type, and since the impacts of a pandemic spans many countries there were no language restrictions.

We used the search strategy to search MEDLINE (including Epub Ahead of Print, In-Process & Other Non-Indexed Citations) and Embase from inception until May 8, 2020. Anticipating pertinent information may not be published (i.e., joint statements, recommendations, and guidelines from surgical colleges) we supplemented the database search with a structured grey literature search including targeted website searching, advanced and general Google searching, and contact with knowledge experts (Appendix A).<sup>9</sup> The reference lists of included studies were screened for relevant studies not otherwise captured.

## **Study Selection**

Titles and abstracts were reviewed by one of two independent reviewers with a third, independent reviewer screening 25% of randomly selected references in duplicate. Full texts of studies considered potentially eligible at title/abstract screening phase by at least one reviewer were reviewed in duplicate by two reviewers for eligibility. Any disagreement between reviewers at the full text screening phase was resolved through discussion and did not necessitate a third reviewer. If studies were excluded at the full text screening phase, the reason for exclusion was noted. Full text articles meeting eligibility criteria were included and data were

abstracted using a standardized data abstraction form (Appendix B). At both stages of screening, a pilot sample of 50 articles were jointly reviewed by both reviewers to ensure reliable application of eligibility criteria between reviewers.

## **Study Eligibility**

Studies were eligible for inclusion if they discussed alterations to surgical services during public health emergencies and reported: 1) reorganization of surgical services, 2) impact of reorganizing surgical services on patients, healthcare providers, or healthcare system or 3) approaches to resuming surgical capacity. Studies of any design or publication date were eligible for inclusion. Studies in any language were eligible, but consistent with rapid review methods, studies not easily translated by authors were excluded from the data synthesis, although citations are still provided. Studies were excluded if they described: only urgent interventions arising during a hospital admission (e.g., emergency tracheostomy, caesarean section), settings beyond in-patient acute care (e.g., outpatient clinics including dental clinics), surgical services beyond a public health emergency without comparison to public health emergencies, healthcare services not specifically related to surgical service.

Notably, our intention was to include guidelines that made recommendations regarding provision of surgical services; however, a high-quality review of guidelines was published<sup>10</sup> during the preparation of this review and as such, we chose to exclude guidelines.

## **Data Extraction**

Data were abstracted by one reviewer, and verified by a second reviewer, using a standardized data abstraction form (Appendix B). Data included: date of publication, country,



study design, definition of non-urgent surgery, characteristics of study sample (if applicable), outcomes of interest for the three research questions, detailed below.

### **Outcomes of Interest**

Our primary outcomes were reorganization of surgical services, impact of reorganization and resuming surgical services. We intentionally included a broad array of outcomes and used an inductive approach to data abstraction to gain a comprehensive understanding of surgical services and the impact during public health emergencies.

We collected qualitative data from studies reporting on changes to surgical programming, conceptualized into five categories: changes to triage criteria or case selection, changes to PPE practices, workforce changes, changes to patient care, changes to resident and fellow education, and environmental changes. Qualitative and quantitative data on the impact of reorganization of surgical services was organized by impact on: patients, providers and healthcare system. To illustrate temporal changes, data preceding, during and after the precipitating event were collected whenever possible. Quantitative variables of interest included: adverse events (including morbidity and mortality), primary care and emergency department visits, number of hospital and ICU admissions, length of hospital and ICU stay, number of surgical procedures performed and number of procedures cancelled, care costs, and wait times for non-urgent surgery. Qualitative variables included narrative description of patient or physician experience, written descriptions of changes to physician remuneration, or comments surrounding surgical waitlist composition. Qualitative data was also collected on details of efforts to rebuild capacity to surgical services.

## **Study Quality (Risk of Bias) Assessment**

Given the aim of a rapid scoping review is not to appraise evidence but to map the available literature,<sup>11</sup> quality appraisal of included studies was not performed.

## **Data Synthesis, Analysis and Reporting**

Study and sample data were summarized narratively in a data table and using descriptive statistics where appropriate. We decided *a priori* to use a random effects model for meta-analysis if there was sufficient data on the impact of changes to surgical services to pool, however this was not feasible. Instead, descriptive statistics were used to synthesize the quantitative outcomes. Data were synthesized and presented separately for each of the three research questions.

## **Patient and Public Partnership**

Patients and the public were not involved in the conception or analysis of this review.

## **RESULTS**

### **Search Results**

A total of 3 013 unique scholarly articles and 106 sources of grey literature were identified, of which 702 were considered eligible for full text review. After full text review, 120 studies and five documents from the grey literature were included. Screening of the reference lists of included articles led to seven additional studies being included for a total of 132 included studies. Thirty-seven studies contributed data to more than one of the research questions resulting in the qualitative synthesis of 111 studies assessing alterations to service delivery, 55 studies evaluating the consequences of these changes, and six studies enumerating their

procedures for rebuilding capacity (Table 1). The flow of evidence sources within the study is detailed in Figure 1. One Spanish language study was translated for inclusion,<sup>12</sup> but two studies could not be readily translated therefore they are not included in the synthesis.<sup>13 14</sup>

## **Description of Studies**

The majority of included studies were published in 2020 about COVID-19 (87.9%, n=116); fewer studies were related to other public health emergencies: SARS (7.58%, n=10), Ebola (2.27%, n=3), H1N1(1.52%, n=2), and MERS (0.76%, n=1). Over two thirds of the included studies (74.2%) emerged from the countries hit earliest by COVID-19; China (14.4%, n=19), Singapore (8.33%, n=11), Italy (19.7%, n=26), and the USA (31.8%, n=41). While many studies described the experiences of their surgical departments as a whole, oncology (15.9%, n=21), orthopedics (13.6%, n=18), and neurosurgery (11.4%, n=15) were the specialties most prominently represented. Summaries of descriptive study information are shown in Figure 2.

## **Reorganization of Surgical Service**

A number of themes emerged from the 108 studies describing reorganization of surgical services. Nearly all studies reported partial, with most reporting full cessation of non-urgent surgeries at their centre, albeit with varying definitions of “non-urgent” (e.g., can be safely postponed for 3 months) and “urgent” (e.g., patient would have adverse outcome if not completed within 7 days). Changes to service delivery were focused on six domains: case selection/triage, PPE regulations and practice, workforce composition and deployment, outpatient and inpatient patient care, resident and fellow education, and the hospital or clinical

environment (Table 2). The three domains that were most frequently reported (case selection/triage, patient care, and workforce) are described in greater detail below.

*1. Changes to Case Selection and Triage Procedures.* The “where” of surgery is described above; however, the issue of what patients can safely undergo which surgical procedures was also discussed in the included studies. We identified cancelling or postponing “non-urgent” surgeries was almost universal. Most often hospitals cancelled surgeries via telephone or text message, but some studies identified that patients initiated their own surgical cancellations due to concerns with safety and contamination. While urgent surgeries were triaged according to routine practice, new triage decisions were made for non-urgent (including oncology) procedures. Methods for triaging non-urgent procedures varied across studies, from the use of guideline supported checklists of eligible procedures to virtual multidisciplinary meetings where the treating surgeon presented details of the case (e.g., patient characteristics, acuity, imaging) to a larger group representing many surgical specialties to reach consensus on each case.

*2. Changes to Patient Care.* Sixty-two studies reported complete cessation or marked reduction of in-person, non-urgent outpatient clinic visits. In these studies, only urgent patients and those requiring post-operative suture or staple removal were granted in-person visits under strict conditions including mask wearing, negative symptom check, history or temperature pre-screening. Studies specific to COVID-19 almost universally filled the resulting care gap for patients deemed “non-urgent” using telephone or video-based telemedicine. Interfaces used include, but were not limited to Zoom, WeChat, Facetime, telephone, and SMS text messaging. A reported advantage of telemedicine was the ability to not only follow-up with returning

patients but to also continue consultations and establish contact with new patients who would require care when non-urgent surgeries resumed. While some admitted a historical reluctance to transition to video-based telemedicine and reported early concerns with their ability to establish secure connections with patients, frequently their worries faded with use and many reported telemedicine would remain integrated in their practices beyond the pandemic.

*3. Changes to the Workforce.* Fourteen of the included studies describe changing the workforce into a minimum of two teams; a “contaminated” team providing care to infected patients and a “clean” team managing those not infected. When these teams were kept separate from one another both inside and outside of the hospital setting, surgical departments were able to continue managing the inevitable emergencies (as well as non-urgent procedures in some settings) without cross contamination during the public health emergencies. New work rotations and shift schedules were created to ensure this structure was sustainable, often with extra healthcare providers designated to replace those with exposures and to provide adequate time off to prevent burnout. This practice was only possible with wards, operating rooms, and pathways (i.e. corridors, elevators) that are separated under the same “clean” and “contaminated” designation. In the most extreme case, entire hospitals were designated for each patient group, as was done by Singapore during SARS<sup>15</sup> and Italy during COVID-19.<sup>16</sup>

### **Impact of Reorganizing Surgical Services**

Of the 55 studies with data relevant to this question, 42 were focused on changes in surgical volumes with six reporting changes to surgical waitlist time or composition, four underlining changes to resident and fellow involvement in surgery, and two showing changes in

patient pain, anxiety, and depression. These recurring outcome measures are summarized below with data for all studies relevant to this question shown in Appendix C.

*Changes in Surgical Volumes.* Thirty-seven studies provided data for this outcome, with 37.8% (n=14) reporting a greater than 75% reduction and 70.3% (n=26) reporting a greater than 50% reductions in their overall or site specific non-urgent surgical volumes (Figure 3a). Not all studies reported reductions; as one study from an oncology “hub” hospital in Italy reported a 20% increase in their surgical volumes, likely due to more cases being diverted to their hospital during the COVID-19 pandemic.<sup>17</sup>

*Changes in Resident/Fellow Involvement in Surgical Activities.* Four studies<sup>18-21</sup> reported on this outcome; two survey-based case series, one resident-level case study and one study containing both survey and case log data. The reductions in surgical involvement for residents are shown by quartile in Figure 3b.

*Changes to Waitlist Length and Composition.* Five studies<sup>22-26</sup> reported data for this outcome. One centre reported a 64% increase in length of their minor colorectal surgery waitlist<sup>25</sup> and another centre (head and neck oncological surgery program) reported a 500% increase in latency from diagnosis to surgery.<sup>26</sup> One study reported no waitlist deaths during the COVID-19 pandemic<sup>24</sup> while another saw a small decrease in the number of weekly waitlist deaths.<sup>23</sup> A single study identified more patients leaving their renal transplantation waitlist due to mortality or clinical deterioration.<sup>22</sup>

*Changes in Patient Pain, Anxiety, and Depression.* Two studies<sup>27 28</sup> reported pain, anxiety, and depression among more than half of waitlist patients; 42.1% experienced anxiety, and 26.3% experienced depression (Figure 3c). The leading reported cause of patient anxiety was a lack of knowledge about when their surgeries would be rescheduled. Other than a single study describing the negative financial effects of the COVID-19 pandemic,<sup>29</sup> impacts on healthcare providers and their practices were rarely discussed.

### **Rebuild Surgical Capacity**

A total of seven studies reported the experience of rebuilding surgical capacity in their departments, hospitals, or systems; all studies referred to the COVID-19 pandemic. One study from China reported reopening non-urgent surgeries with close consideration of risk for imported transmission but did not provide further detail of triage or prioritization.<sup>30</sup> Among studies that changed their surgical triage practices, patients were prioritized for surgery based on procedure acuity or urgency (i.e., risk to patients if surgery were further delayed), resource intensity, and procedural complexity. Four studies<sup>31-34</sup> noted that prior to resuming non-urgent surgeries, availability of the staff, (Operating Room) ORs, PPE, and testing was necessary to prepare for a large and complicated surgical backlog.

## **DISCUSSION**

This review identified over 3,000 evidence sources, 132 of which were included. Approaches to reorganizing surgical services varied between studies and centers, but the cancellation or postponement of non-urgent surgeries such as arthroplasty surgeries for chronic joint pain, coronary artery bypass graft surgery for asymptomatic individuals, and primary

gastric bypass surgery was nearly universal.<sup>2</sup> The most frequently reported change to surgical services was modified triage criteria for surgical cases, workforce, and approach to patient care. Many studies reported a decrease in surgical volumes due to public health emergencies, while a few reported the non-surgical impacts such as patient wellbeing or changes in healthcare utilization beyond the surgical wards. Very few studies described their experience resuming surgical services after a public health emergency.

The varied approaches to providing surgical services during a public health emergency identified in this review illustrate that a “one fits all” approach does not exist. Changes to surgical services likely depends on the characteristics of specific centers and their patients. While several guidelines have been published with recommendations on how to provide surgical care during COVID-19, we chose to exclude guidelines and recommendations from this review for two reasons: 1) a high quality review of surgical recommendations for the response to COVID-19 was published by one of the authors just prior to this study<sup>10</sup> and 2) because there is abundant evidence suggesting guidelines and recommendations for practice are frequently not implemented into clinical practice.<sup>35-41</sup> Some of the guideline recommendations in the review by Søreide et al.<sup>10</sup> were implemented within the included studies in the present review; such as creating areas within-hospital for ‘clean’ and ‘contaminated’ cases and workforce redeployment to critical care. However, other recommendations were infrequently noted, such as the dedicated use of isolated, negative pressure ORs for patients with COVID-19. These resource intensive practices may not have been attainable under the pressures of managing public health emergencies and may not be feasible in low-resource settings.



Changes to surgical services, such as cancelling or postponing non-urgent surgeries may be necessary to manage public health emergencies to reduce the risk of contamination and increase capacity within hospitals. However, the impact of these changes remains poorly understood. Many studies reported decreases in surgical volumes, but few other variables were explored with regards to the impact on patients, providers, and healthcare systems. Five studies examined the impact of changes to surgical services among physicians and trainees, and found that training was compromised in some specialties.<sup>18-21</sup> The finding that medical training was compromised is particularly important for understanding the downstream and long-term repercussions of the response to public health emergencies; decreases in surgical volumes and clinical hours for trainees could have negative and unintended effects on the future quality and safety of patient care.<sup>42</sup> Studies examining the effects of surgical service alterations on patients noted negative effects on mental health outcomes,<sup>27 28</sup> pain,<sup>27</sup> and an increased incidence of death among surgical patients.<sup>22 23 43</sup>

Very few studies described specific actions undertaken to rebuild and resume pre-public health emergencies surgical capacity. This may be due to the fact that most included studies examined the ongoing COVID-19 pandemic, or because few places have implemented specific plans to date. Included studies did describe consideration of system-level factors like availability of PPE and ORs. However, more patient-centric considerations such as organizing child care and requesting time away from their job during a pandemic, are needed. Interestingly, one study reported 14% of surgical patients initiated the cancellation of their surgery,<sup>27</sup> which suggests patient readiness for surgery during- and post-COVID-19 should be considered. For evidence to inform policy, additional research is needed to understand the impacts of different approaches for resuming surgical services.

This study is, to our knowledge, the first comprehensive scoping review of evidence around reallocation of surgical services during public health emergencies. While this study has several strengths, including a comprehensive search of academic and grey literature sources, and a mix of inductive and deductive data abstraction approaches, there are some limitations that should be considered when interpreting our findings. We modified the Joanna Briggs methodology for scoping reviews,<sup>5</sup> according to the World Health Organization and Cochrane's guidance on conducting rapid reviews,<sup>7,8</sup> with the intent of balancing rigor with a timely and policy-responsive review of the literature. Also, given that the evidence around the COVID-19 pandemic is growing at an unprecedented rate, it is possible that additional studies have been published since we ran our search strategy, especially around resuming surgical services. In order to mitigate this limitation, an ongoing effort to pivot this study into a living review is underway to ensure the data presented is up-to-date. Notably, this review did not identify evidence from any low or middle-income countries who may face unique challenges during a pandemic compared to high income countries described in our review. It is also likely that during the global pandemic, many healthcare institutions have been focused on coping with COVID-19 instead of publishing their experiences; we hope more organizations will add their experience to the literature.

In conclusion, we report early evidence of the operational changes that have occurred internationally in response to public health emergencies which could inform the ongoing response to COVID-19 and future public health emergencies. This study identified a gap in our understanding of the impact of these changes on patients, providers, and the healthcare system

which should be the focus of research moving forward to provide an evidence-based approach to managing surgical patients in future public health emergencies.

### **Author Contributions**

CO contributed to the design and conceptualization of the review, analysis and interpretations of the data, and drafting and revising the manuscript; JSN contributed to the design of the review, interpretation of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript; AKR contributed to the design of the review, interpretation of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript; JCD contributed to the interpretation of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript; JW contributed to the interpretation of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript; JR contributed to the interpretation of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript; MB contributed to the interpretation of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript; KMS contributed to the design and conceptualization of the review, analysis and interpretations of the data, providing feedback on the manuscript, and gave approval of the final version of the manuscript.









## References

1. Johns Hopkins Center for Systems Science and Engineering (CSSE). COVID-19 Dashboard. [Available from: <https://coronavirus.jhu.edu/map.html>] accessed July 27, 2020.
2. American College of Surgeons. COVID-19: Recommendations for management of elective surgical procedures. [Available from: [https://www.facs.org/-/media/files/covid19/recommendations\\_for\\_management\\_of\\_elective\\_surgical\\_procedure\\_s.ashx](https://www.facs.org/-/media/files/covid19/recommendations_for_management_of_elective_surgical_procedure_s.ashx)] accessed July 2, 2020.
3. Shared Health Manitoba. COVID-19: The New Normal. [Available from: <https://sharedhealthmb.ca/files/covid-19-memo-the-new-normal.pdf>] accessed May 8, 2020.
4. COVIDSurg Collaborative. Elective surgery cancellations due to the COVID-19 pandemic: global predictive modelling to inform surgical recovery plans. *British Journal of Surgery* 2020 doi: doi:10.1002/bjs.11746 [published Online First: 2020 May 12]
5. Peters MDJ, Godfrey CM, McInerney P, et al. Chapter 11: Scoping Reviews. In: E. A, Z. M, eds. Joanna Briggs Institute Reviewer's Manual: The Joanna Briggs Institute, 2017.
6. Tricco AC, Lillie E, Zarin W, et al. PRISMA Extension for Scoping Reviews (PRISMA-ScR): Checklist and Explanation. *Ann Intern Med* 2018;169(7):467-73. doi: 10.7326/m18-0850 [published Online First: 2018/09/05]
7. Tricco AC, Langlois EV, Straus SE, Editors. Rapid reviews to strengthen health policy and systems: a practical guide. Geneva: World Health Organization, 2017. Licence: CC BY-NC-SA 3.0 IGO
8. Garritty C, Gartlehner G, Kamel C, et al. Cochrane Rapid Reviews. Interim Guidance from the Cochrane Rapid Review Methods Group., 2020.
9. Godin K, Stapleton J, Kirkpatrick SI, et al. Applying systematic review search methods to the grey literature: a case study examining guidelines for school-based breakfast programs in Canada. *Syst Rev* 2015;4:138. doi: 10.1186/s13643-015-0125-0 [published Online First: 2015/10/27]
10. Soreide K, Hallet J, Matthews JB, et al. Immediate and long-term impact of the COVID-19 pandemic on delivery of surgical services. *Br J Surg* 2020 doi: 10.1002/bjs.11670 [published Online First: 2020/05/01]
11. Munn Z, Peters MDJ, Stern C, et al. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Medical Research Methodology* 2018;18(1):143. doi: 10.1186/s12874-018-0611-x
12. Alvarez Gallego M, Gortazar de Las Casas S, Pascual Miguelanez I, et al. SARS-CoV-2 pandemic on the activity and professionals of a General Surgery and Digestive Surgery Service in a tertiary hospital. *Cir Esp* 2020 doi: 10.1016/j.ciresp.2020.04.001 [published Online First: 2020/04/28]
13. Wang X, Wang Z, Yao C, et al. Management of ophthalmic perioperative period during 2019 novel coronavirus disease outbreak. *Zhonghua Shiyan Yanke Zazhi/Chinese Journal of Experimental Ophthalmology* 2020;38(3):200-03. doi: <http://dx.doi.org/10.3760/cma.j.issn115989-20200224-00100>
14. Zhen L, Lin T, Zhao ML, et al. [Management strategy for the resumption of regular diagnosis and treatment in gastrointestinal surgery department during the outbreak of coronavirus disease 2019 (COVID-19)]. *Zhonghua Wei Chang Wai Ke Za Zhi* 2020;23(4):321-26.





- doi: 10.3760/cma.j.issn.1671-0274.2020-0316-00146 [published Online First: 2020/04/21]
15. Chee VW, Khoo ML, Lee SF, et al. Infection control measures for operative procedures in severe acute respiratory syndrome-related patients. *Anesthesiology* 2004;100(6):1394-8. doi: 10.1097/00000542-200406000-00010 [published Online First: 2004/05/29]
  16. Fontanella MM, Saraceno G, Lei T, et al. Neurosurgical activity during COVID-19 pandemic: an expert opinion from China, South Korea, Italy, United States of America, Colombia and United Kingdom. *J Neurosurg Sci* 2020 doi: 10.23736/S0390-5616.20.04994-2 [published Online First: 2020/04/30]
  17. Vicini E, Galimberti V, Naninato P, et al. COVID-19: The European institute of oncology as a "hub" centre for breast cancer surgery during the pandemic in Milan (Lombardy region, northern Italy) - A screenshot of the first month. *Eur J Surg Oncol* 2020 doi: 10.1016/j.ejso.2020.04.026 [published Online First: 2020/05/04]
  18. Amparore D, Claps F, Cacciamani GE, et al. Impact of the COVID-19 pandemic on urology residency training in Italy. *Minerva Urol Nefrol* 2020 doi: 10.23736/S0393-2249.20.03868-0 [published Online First: 2020/04/08]
  19. Cai Y, Jiam NT, Wai KC, et al. Otolaryngology Resident Practices and Perceptions in the Initial Phase of the U.S. COVID-19 Pandemic. *Laryngoscope* 2020 doi: 10.1002/lary.28733 [published Online First: 2020/05/06]
  20. Givi B, Moore MG, Bewley AF, et al. Advanced head and neck surgery training during the COVID-19 pandemic. *Head Neck* 2020 doi: 10.1002/hed.26252 [published Online First: 2020/05/10]
  21. Hemingway JF, Singh N, Starnes BW. Emerging practice patterns in vascular surgery during the COVID-19 pandemic. *J Vasc Surg* 2020 doi: 10.1016/j.jvs.2020.04.492 [published Online First: 2020/05/04]
  22. de Vries APJ, Alwayn IPJ, Hoek RAS, et al. Immediate impact of COVID-19 on transplant activity in the Netherlands. *Transpl Immunol* 2020:101304. doi: 10.1016/j.trim.2020.101304 [published Online First: 2020/05/07]
  23. Dominguez-Gil B, Coll E, Fernandez-Ruiz M, et al. COVID-19 in Spain: Transplantation in the midst of the pandemic. *Am J Transplant* 2020 doi: 10.1111/ajt.15983 [published Online First: 2020/05/03]
  24. Lauterio A, De Carlis R, Belli L, et al. How to guarantee liver transplantation in the north of Italy during the COVID-19 pandemic: A sound transplant protection strategy. *Transpl Int* 2020 doi: 10.1111/tri.13633 [published Online First: 2020/04/30]
  25. Bradford IM. Tales from the frontline: the colorectal battle against SARS. *Colorectal Dis* 2004;6(2):121-3. doi: 10.1111/j.1462-8910.2004.00600.x [published Online First: 2004/03/11]
  26. Lee AKF, Cho RHW, Lau EHL, et al. Mitigation of head and neck cancer service disruption during COVID-19 in Hong Kong through telehealth and multi-institutional collaboration. *Head Neck* 2020 doi: 10.1002/hed.26226 [published Online First: 2020/05/02]
  27. Brown TS, Bedard NA, Rojas EO, et al. The Effect of the COVID-19 Pandemic on Electively Scheduled Hip and Knee Arthroplasty Patients in the United States. *J Arthroplasty* 2020 doi: 10.1016/j.arth.2020.04.052 [published Online First: 2020/05/08]
  28. Wan IYP, Wat KHY, Ng CSH, et al. Evaluation of the emotional status of patients on a waiting list for thoracic surgery during the outbreak of Severe Acute Respiratory Syndrome(SARS). *Stress and Health* 2004;20(4):209-12. doi: 10.1002/smi.1013

29. Vaccaro AR, Getz CL, Cohen BE, et al. Practice Management During the COVID-19 Pandemic. *J Am Acad Orthop Surg* 2020;28(11):464-70. doi: 10.5435/JAAOS-D-20-00379 [published Online First: 2020/04/15]
30. Shen Y, Cui Y, Li N, et al. Emergency Responses to Covid-19 Outbreak: Experiences and Lessons from a General Hospital in Nanjing, China. *Cardiovasc Intervent Radiol* 2020;43(6):810-19. doi: 10.1007/s00270-020-02474-w [published Online First: 2020/04/29]
31. Busin M, Yu AC, Ponzin D. Coping with COVID-19: An Italian Perspective on Corneal Surgery and Eye Banking in the Time of a Pandemic and Beyond. *Ophthalmology* 2020 doi: 10.1016/j.ophtha.2020.04.031 [published Online First: 2020/05/08]
32. Pelt CE, Campbell KL, Gililand JM, et al. The Rapid Response to the COVID-19 Pandemic by the Arthroplasty Divisions at Two Academic Referral Centers. *J Arthroplasty* 2020 doi: 10.1016/j.arth.2020.04.030 [published Online First: 2020/05/02]
33. Walker JP. Resuming Elective Surgery at UTMB Predicated on Patient and Staff Well-Being, 2020.
34. Health BMO. A Commitment to Surgical Renewal in BC. Canada, 2020.
35. Ben-Menachem E, French JA. Guidelines--are they useful? *Epilepsia* 2006;47 Suppl 1:62-4. doi: 10.1111/j.1528-1167.2006.00663.x [published Online First: 2006/10/19]
36. Cabana MD, Rand CS, Powe NR, et al. Why don't physicians follow clinical practice guidelines? A framework for improvement. *Jama* 1999;282(15):1458-65. [published Online First: 1999/10/27]
37. Farquhar C, Kofa E, Slutsky J. Clinicians' attitudes to clinical practice guidelines: a systematic review. *Med J Aust* 2002;177:502-06.
38. Grimshaw JM, Russell IT. Effect of clinical guidelines on medical practice: a systematic review of rigorous evaluations. *Lancet* 1993;342(8883):1317-22. [published Online First: 1993/11/27]
39. Grimshaw JM, Thomas RE, MacLennan G, et al. Effectiveness and efficiency of guideline dissemination and implementation strategies. *Health technology assessment (Winchester, England)* 2004;8(6):iii-iv, 1-72. [published Online First: 2004/02/13]
40. Lomas J, Anderson G, Domnick-Pierre K, et al. Do practice guidelines guide practice? The effect of a consensus statement on the practice of physicians. *N Engl J Med* 1989;321(19):1306-11.
41. Lugtenberg M, Burgers JS, Westert GP. Effects of evidence-based clinical practice guidelines on quality of care: a systematic review. *Quality & safety in health care* 2009;18(5):385-92. doi: 10.1136/qshc.2008.028043 [published Online First: 2009/10/09]
42. Ahmed N, Devitt KS, Keshet I, et al. A systematic review of the effects of resident duty hour restrictions in surgery: impact on resident wellness, training, and patient outcomes. *Ann Surg* 2014;259(6):1041-53. doi: 10.1097/sla.0000000000000595 [published Online First: 2014/03/26]
43. D'Apolito R, Faraldi M, Ottaiano I, et al. Disruption of Arthroplasty Practice in an Orthopedic Center in Northern Italy During the Coronavirus Disease 2019 Pandemic. *J Arthroplasty* 2020 doi: 10.1016/j.arth.2020.04.057 [published Online First: 2020/05/07]

















**Table 1.** Description of included studies
















Author	Year	Country	Public Health Emergency	Surgical Specialty	Changes to Surgical Services	Impact of Changes Examined	Resumption of Services
Alvarez-Gallego	2020	Spain	COVID-19	General			
Ammar	2020	USA	COVID-19	Neurosurgery			
Amparore	2020	Italy	COVID-19	Urology		Changes in clinical and surgical resident involvement	
Ansarin	2020	Italy	COVID-19	Oncology (Head & Neck)			
Bashir	2020	UK	COVID-19	Vascular			
Ben Abdallah	2020	France	COVID-19	Vascular			
Bernucci	2020	Italy	COVID-19	Neurosurgery		Changes in surgical volume	
Bettinelli	2020	Italy	COVID-19	Orthopedics			
Bolkan	2014	Norway	Ebola	Obstetrics		Changes in non-Ebola admissions and surgical volumes	
Bolkan	2018	Norway	Ebola	Obstetrics			Changes in non-Ebola admissions and surgical volumes
Bourlon	2009	Mexico	H1N1	—		Number of surgical cancellations	

































Bradford	2003	China	SARS	GI		Changes in admissions and surgical volumes	
Brethauer	2020	USA	COVID-19	GI			Operative cases placed in “depot” to be rescheduled alongside new teleconsults
Brown	2020	USA	COVID-19	Orthopedics		Patient pain, anxiety and physical function	
Buckstein	2020	USA	COVID-19	Radiation Oncology			
Bundu	2016	Sierra Leone	Ebola	—		Changes in ED/ward admissions and surgical activity	
Burke	2020	USA	COVID-19	Neurosurgery			
Busin	2020	Italy	COVID-19	Ophthalmology		Changes in demand/donations for cornea bank	Set reasonable timelines for patients requiring low acuity surgery. Surgical work schedule extended into evenings and weekends
Cai	2020	USA	COVID-19	Otolaryngology		Changes in resident	













						educational programming
Cakmak	2020	Turkey	COVID-19	Oncology (Breast)		Changes in surgical volume
Carenzo	2020	Italy	COVID-19	—		Changes in surgical volume
Cenzato	2020	Italy	COVID-19	Neurosurgery		
Chan	2006	China	SARS	Ophthalmology		
Chee	2004	Singapore	SARS	—		
Chew	2020	Singapore	COVID-19	General		
Chisci	2020	Italy	COVID-19	Vascular		Changes in surgical volume
Civantos	2020	USA	COVID-19	Oncology (Head & Neck)		Number of surgical cancellations
D'Apolito	2020	Italy	COVID-19	Orthopedics		Changes in surgical volume
de Vries	2020	Netherlands	COVID-19	Transplant		Changes in transplantation volumes
Ding	2020	Singapore	COVID-19	Orthopedics		
Dominguez-Gil	2020	Spain	COVID-19	Transplant		Changes in transplantation volumes
Doussot	2020	France	COVID-19	Oncology		Changes in surgical volume
Dowdell	2020	USA	COVID-19	Orthopedics		
Ducournau	2020	France	COVID-19	Plastics		
Eichberg	2020	USA	COVID-19	Neurosurgery		
Ficarra	2020	Italy	COVID-19	Urology		Proportion of surgical cancellations

Fontanella	2020	Italy	COVID-19	Neurosurgery		initiated by patients Changes in surgical volume
Fontanella	2020	Italy	COVID-19	Neurosurgery		
Giorgi	2020	Italy	COVID-19	Spinal		
Givi	2020	USA	COVID-19	Oncology (Head & Neck)		Changes in fellow educational programming
Gomez-Barrena	2020	Spain	COVID-19	Orthopedics		Changes in surgical volume
Gouveia	2020	Portugal	COVID-19	Vascular		
Guerci	2020	Italy	COVID-19	General		
Gupta	2020	India	COVID-19	Oncology (Head & Neck)		
Haines	2003	China	SARS	Obstetrics		
Hemingway	2020	USA	COVID-19	Vascular		Changes in clinical and surgical volumes
Hormati	2020	Iran	COVID-19	GI		
Hu	2020	China	COVID-19	Oncology		
Jean	2020	USA	COVID-19	Neurosurgery		Changes in surgical volume
Kempa	2020	Poland	COVID-19	Electrophysiology		Changes in surgical volume
Kessler	2020	USA	COVID-19	Neurosurgery		
Konda	2020	USA	COVID-19	Orthopedics		
Kuo	2010	Argentina	H1N1	Ophthalmology		Changes in clinical and surgical volumes
Lai	2020	China	COVID-19	Ophthalmology		
Lancaster	2020	USA	COVID-19	—		Changes in surgical volume






Langer	2020	USA	COVID-19	Plastics		
Lauterio	2020	Italy	COVID-19	Transplant		Changes in transplantation volumes
Lee	2020	China	COVID-19	Oncology (Head & Neck)		Changes in surgical wait times
Leong Tan	2020	Singapore	COVID-19	Vascular		
Li	2020	China	COVID-19	Transplant		
Liebensteiner	2020	Austria	COVID-19	Orthopedics		
Liu	2003	Singapore	SARS	Anesthesia		
Mak	2020	China	COVID-19	Plastics		
Marti	2020	Spain	COVID-19	Oncology (Breast)		
Maurizi	2020	Italy	COVID-19	Thoracic		Changes in surgical volume
McBride	2020	Australia	COVID-19	—		
McMillan	2020	Canada	COVID-19	—		Prioritization first of patients who would be at increased risk with further delay, followed by those waiting longest
Meneghini	2020	USA	COVID-19	Orthopedics		
Meyer	2020	France	COVID-19	Neurosurgery		Changes in surgical volume
Morgan	2020	UK	COVID-19	Orthopedics		
Nair	2020	India	COVID-19	Ophthalmology		
Nassar	2020	USA	COVID-19	General		
Park	2020	USA	COVID-19	Otolaryngology		









Park	2020	South Korea	MERS	—		Changes in surgical volume	
Patel	2020	USA	COVID-19	Oncology (Head & Neck)			
Patel	2020	USA	COVID-19	Otolaryngology			
Pelt	2020	USA	COVID-19	Orthopedics		Number of surgical postponements	Surgeries prioritized based on complexity and predicted LOS, scheduling only completed if appropriate PPE and screening available.
Pittet	2020	Switzerland	COVID-19	—		Changes in surgical volume	
Prachand	2020	USA	COVID-19	—		Number of surgical cancellations	
Price	2020	USA	COVID-19	Dermatology			
Qadan	2020	USA	COVID-19	Oncology (GI/Hepatobiliary)			
Ralli	2020	Italy	COVID-19	Otolaryngology			
Rampinelli	2020	Italy	COVID-19	Oncology (Head & Neck)		Changes in surgical volume	
Randelli	2020	Italy	COVID-19	Orthopedics			
Ricciardi	2020	Italy	COVID-19	Neurosurgery			
Ross	2020	USA	COVID-19	—		Changes in clinical volumes	

Rubin	2020	USA	COVID-19	Electrophysiology		Changes in lab capacity and consultation volumes
Rubin	2020	USA	COVID-19	Electrophysiology		Changes in surgical volume
Salengar	2020	USA	COVID-19	Cardiac		
Sarpong	2020	USA	COVID-19	Orthopedics		Changes in surgical volume
Schull	2007	Canada	SARS	—		
Schwarzkopf	2020	USA	COVID-19	Orthopedics		Changes in surgical volume
Scullen	2020	USA	COVID-19	Neurosurgery		
Seese	2020	USA	COVID-19	Cardiac		
Sethi	2020	USA	COVID-19	GI		Changes in surgical and consultation volumes
Shen	2020	China	COVID-19	—		Scheduling resumed following consideration of reduced risk of imported transmission and growing waitlist
Shih	2020	China	COVID-19	Ophthalmology		Changes in surgical volume
Shokri	2020	USA	COVID-19	Plastics		Changes in surgical volume
Sobel	2020	USA	COVID-19	Urology		







Sun	2020	China	COVID-19	Neurosurgery		Emergency surgeries performed	
Tan	2004	Singapore	SARS	Anesthesia			
Tan	2020	Singapore	COVID-19	Urology		Changes in surgical and consultation volumes	
Tan	2020	China	COVID-19	Neurosurgery			
Tay	2020	Singapore	COVID-19	Orthopedics		Changes in surgical volumes	
Tay	2020	Singapore	COVID-19	Orthopedics			
Thaler	2020	Austria	COVID-19	Orthopedics			
Tolone	2020	Italy	COVID-19	—			
Too	2020	Singapore	COVID-19	Interventional Radiology			
Topf	2020	USA	COVID-19	Oncology (Head & Neck)			
Tsui	2005	China	SARS	Cardiac			
Tzeng	2020	USA	COVID-19	Oncology			
Unal	2020	Turkey	COVID-19	Vascular			
Vaccaro	2020	USA	COVID-19	Orthopedics		Changes in physician remuneration and staffing	
Valenza	2020	Italy	COVID-19	Oncology		Changes in surgical volumes	
van de Haar	2020	Netherlands	COVID-19	Oncology			
Various	2020	Canada	COVID-19	—		Number of surgical postponements	Calling patients to assess their ability to reschedule,

contracting private facilities with focus on urgent surgeries, patients waiting twice their clinical benchmarks or surgeries with minimal LOS

Various	2020	Ireland	COVID-19	—			
Vicini	2020	Italy	COVID-19	Oncology (Breast)		Changes in surgical volumes	
Vlantis	2004	China	SARS	Otolaryngology		Changes in outpatient and surgical volumes	
Walker	2020	USA	COVID-19	—		Number of surgical cancellations	Assess readiness of staff to safely resume high volumes of surgery and ensured availability of rapid in-house testing
Wan	2004	China	SARS	Thoracic		Patient anxiety and depression	
Wasser	2020	Israel	COVID-19	Ophthalmology			

Williams	2020	USA	COVID-19	Ophthalmology		Number of surgeries rescheduled
Wong	2020	Singapore	COVID-19	Anesthesia		
Wu	2020	China	COVID-19	Oncology (Head & Neck)		
Xiaolei	2020	China	COVID-19	Ophthalmology		
Zangrillo	2020	Italy	COVID-19	—		
Zarzaur	2020	USA	COVID-19	—		
Zeng	2020	China	COVID-19	Oncology (Head & Neck)		
Zizzo	2020	Italy	COVID-19	—		
Zoia	2020	Italy	COVID-19	Neurosurgery		

Domains of change to surgical services, represented numerically:

-  Changes to case selection and surgical triage
-  Changes to PPE protocols and practices
-  Changes to the surgical workforce.
-  Changes to inpatient and outpatient care
-  Changes to resident and fellow education
-  Changes to the environment

*Abbreviations:* — =data not provided, COVID-19=Coronavirus Disease 2019, ED=Emergency department, GI=Gastrointestinal; LOS=Length of stay, PPE=Personal protective equipment, SARS=Severe Acute Respiratory Syndrome



**Table 2.** Reorganization of surgical services, by domain.

<b>Change Domain</b>	<b>Number of Studies (%)</b>	<b>Examples of Change</b>
Triage or Case Selection	80 (74.7)	<ol style="list-style-type: none"> <li>1. Prioritization of patients based on pre-defined levels of acuity;</li> <li>2. Virtual multidisciplinary meetings or tumor boards;</li> <li>3. Creation of specialty-specific lists outlining surgery-eligible and ineligible ailments, often with inclusion of case-by-case category.</li> <li>4. Postponement based on high-risk patient characteristics (i.e., older age, multimorbidity) and expected need for ICU.</li> </ol>
PPE	63 (58.3)	<ol style="list-style-type: none"> <li>1. Hospital wide surgical mask mandate for staff and attendees;</li> <li>2. Standard level of PPE outlined for all patient encounters with enhanced PPE (e.g., addition of N95 or PAPR, head and shoe covering) protocol for specific procedures or care of infected patients;</li> <li>3. Refresher instruction courses provided to all hospital staff;</li> <li>4. Trained observer supervising all perioperative donning and doffing of PPE to ensure safety and compliance.</li> </ol>
Workforce	70 (64.8)	<ol style="list-style-type: none"> <li>1. Separation of clinical staff into rotating “clean” and “dirty” teams caring for exclusively for non-infected and infected patients, respectively;</li> <li>2. Temperature and symptom screening of staff with mandated quarantine periods in cases of unprotected exposure;</li> <li>3. Case discussions, handover and clinical staff meetings transitioned to virtual format;</li> <li>4. Redeployment of staff to hospital areas requiring support (e.g., ICU), often paired with virtual training to ensure comfortable transition.</li> </ol>

Patient Care	95 (88.0)	<ol style="list-style-type: none"> <li>1. Complete cancellation or transition to telemedicine for all non-urgent and routine perioperative clinical visits;</li> <li>2. Patient temperature, symptom and travel history screening before entry to clinic (relevant for urgent surgical patients);</li> <li>3. Preference for endovascular or minimally invasive surgical approaches when possible, use of conservative care when possible (oncology);</li> <li>4. Restrictions on number of accompanying persons or visitors (often zero with some allowing maximum of 1).</li> </ol>
Resident/Fellow Education	35 (32.4)	<ol style="list-style-type: none"> <li>1. Changes to resident/fellow team structure and rotation schedules to ensure continued coverage of department and maximize resident/fellow safety;</li> <li>2. Redeployment of residents to non-specialty areas requiring clinical support;</li> <li>3. Curriculum and conferences shifted to online format to allow continued e-learning for off-duty trainees;</li> <li>4. Trainees involvement in surgical care of infected persons ceased or altered (e.g., only admitted to OR during low-risk/non-aerosolizing procedures).</li> </ol>
Environment	70 (64.8)	<ol style="list-style-type: none"> <li>1. Dedication of wards (hallways, elevators), ORs, or entire hospitals to treat for only those infected or not infected;</li> <li>2. Use of negative-pressure OR when possible;</li> <li>3. Transformation of surgical wards, ORs and outpatient clinics into patient care areas to increase surge capacity;</li> <li>4. Double occupancy patient rooms reduced to single occupancy, or</li> </ol>

physical measures (e.g., cubicles,  
distanced waiting room chairs)  
implemented.


*Abbreviations:* ICU= Intensive care unit, PPE= Personal protective equipment, PAPR= Powered air purifying respirator, OR= Operating room.

Figure 1. PRISMA Flow diagram.

Figure 2. Summary of study characteristics.

Figure 3. Summary of leading impacts of changes to surgical programming

**Figure 1.** Flow of studies in the scoping review



\*Includes guidelines, recommendation-based reviews, projections or estimations without mention of true changes to surgical programming.

## Figure 2. Study characteristics




Figure 2. **A.** Country of publication, **B.** Public health emergency discussed, and **C.** Surgical specialty addressed ('Other' includes Cardiac (n=3), Anesthesia (n=3), Electrophysiology (n=3), Obstetrics and Gynecology (n=3), Thoracic (n=2), Interventional Radiology (n=1), and Dermatology (n=1)).



**Figure 3.** Summary of the impacts of alterations to surgical services




Figure 3. A summary of the impacts of alterations to surgical services during public health emergencies on **A.** Overall surgical activity (n=37 studies), **B.** Resident and fellow involvement in surgery (n=5 studies) where circle size represents the number of studies contributing to that quartile, and **C.** Patient experience (n=2 studies).