

Online Data Supplement:

An epithelial microRNA upregulates airway IL-25 and TSLP expression in type 2-high asthma via targeting CD39-extracellular ATP axis


Kan Zhang^{1,2,3*}, Yuchen Feng^{1,2*}, Yuxia Liang^{1,2}, Wenliang Wu^{1,2}, Chenli Chang^{1,2}, Dian Chen^{1,2}, Shengchong Chen^{1,2}, Jiali Gao^{1,2}, Gongqi Chen^{1,2}, Lingling Yi^{1,2}, Dan Cheng⁴, Guohua Zhen^{1,2†}

¹Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China; ²Key Laboratory of Respiratory Diseases, National Health Commission of People's Republic of China, and National Clinical Research Center for Respiratory Diseases, Wuhan, China; ³Department of Respiratory and Critical Care Medicine, The Affiliated Suzhou Hospital of Nanjing Medical University, Suzhou, China; ⁴Department of Respiratory and Critical Care Medicine, Renmin Hospital of Wuhan University, Wuhan, China.

Supplementary Table. Primers for quantitative PCR

Gene	Species	Type	Sequence
<i>β-actin</i>	Human	Forward	GCAAGCAGGACTATGACGAG
		Reverse	CAAATAAAGCCATGCCAATC
<i>CD39</i>	Human	Forward	ACTATCGAGTCCCCAGATAATGC
		Reverse	CCTGATCCTTCCCATAGCACAA
<i>CLCA1</i>	Human	Forward	ATGGCTATGAAGGCATTGTCTG
		Reverse	TGGCACATTGGGGTCTGATTG
<i>GAPDH</i>	Human	Forward	AAGGTGAAGGTCGGAGTCAAC
		Reverse	GGGGTCATTGATGGCAACAATA
<i>POSTN</i>	Human	Forward	GACCGTGTGCTTACACAAATTG
		Reverse	AAGTGACCGTCTCTTCCAAGG
<i>SERPINB2</i>	Human	Forward	TCCTGGGTCAAGACTCAAACC
		Reverse	CATCCTGGTATCCCCATCTACA
<i>β-actin</i>	Mouse	Forward	GGCTGTATTCCCCTCCATCG
		Reverse	CCAGTTGGTAACAATGCCATGT
<i>Gapdh</i>	Mouse	Forward	TGGCCTTCCGTGTTCTTAC
		Reverse	GAGTTGCTGTTGAAGTCGCA
<i>Cd39</i>	Mouse	Forward	AGATGAAATCGGTGCGTACCT
		Reverse	GAGTCTGGTGATGCTTGGATG

Supplementary Figure and Figure Legend:



Supplementary Figure. Overexpression of airway miR-206 expression aggravates HDM-induced AHR, airway inflammation, mucus overproduction and type 2 response in mice. (A) Respiratory resistance in response to different concentration of intravenous acetylcholine at 24 h after the last HDM or saline challenge in mice intranasally administered with control or miR-206 agomir. (B) Inflammatory scores of lung sections from mice intranasally administered with control or miR-206 agomir and

challenged with HDM or saline were calculated as described in Methods. (C) Counts for macrophages, eosinophils, lymphocytes and neutrophils in BALF. (D) H&E staining of representative lung sections. (E) PAS staining for mucus in representative lung sections. (F) The numbers of PAS-staining-positive cells were counted in four random fields for each lung section at $\times 200$ magnification. (G-I) The protein levels of IL-4 (G), IL-5 (H), IL-13 (I) in BALF were determined by ELISA. (J) Plasma IgE levels in peripheral blood were determined by ELISA. $n = 6-10$ mice per group combined from 2 experiments. Data are mean \pm SD. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$ (one-way ANOVA with Bonferroni's post hoc test). Scale bar = 50 μm .