

1 **Clinical and Virological Characteristics of Hospitalized COVID-19**
2 **Patients in a German Tertiary Care Center during the First Wave of**
3 **the SARS-CoV-2 Pandemic**

4 **Running title: Berlin COVID-19 cohort**

5 **Author list:**

6 Charlotte Thibeault^{1,2}, Barbara Mühlemann^{1,3,4}, Elisa T. Helbig^{1,2}, Mirja Mittermaier^{1,2,5},
7 Tilman Lingscheid^{1,2}, Pinkus Tober-Lau^{1,2}, Lil A. Meyer-Arndt^{1,6}, Leonie Meiners^{1,3,4},
8 Paula Stubbemann^{1,2}, Sascha S. Haenel^{1,2}, Laure Bosquillon de Jarcy^{1,2,3}, Lena
9 Lippert^{1,2}, Moritz Pfeiffer^{1,2}, Miriam S. Stegemann^{1,2}, Robert Roehle^{1,5,7,9}, Janine
10 Wiebach^{1,5,7}, Stefan Hippenstiel^{1,2}, Thomas Zoller^{1,2}, Holger Müller-Redetzky^{1,2},
11 Alexander Uhrig^{1,2}, Felix Balzer^{1,8}, Christof von Kalle^{1,9}, Norbert Suttorp^{1,2,10}, Terry C.
12 Jones^{1,3,4,11}, Christian Drosten^{1,3,4}, Martin Witzernath^{1,2,10}, Leif E. Sander^{1,2}, **Pa-COVID**
13 **Study Group**, Victor M. Corman^{1,3,4*}, Florian Kurth^{1,2,12*}

14 **Affiliations:**

15 ¹Charité–Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-
16 Universität zu Berlin, and Berlin Institute of Health, 10117 Berlin, Germany

17 ²Department of Infectious Diseases and Respiratory Medicine, Charité – Universitätsmedizin
18 Berlin, 13353 Berlin, Germany

19 ³Institute of Virology, Charité - Universitätsmedizin Berlin, 10117 Berlin, Germany

20
21 ⁴German Center for Infection Research (DZIF), Associated Partner Site, 10117 Berlin,

22 Germany

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

1 ⁵Berlin Institute of Health (BIH), 10178 Berlin, Germany

2 ⁶Department of Neurology, Charité – Universitätsmedizin Berlin, 10117 Berlin, Germany

3 ⁷Institute of Biometry and Clinical Epidemiology, Charité – Universitätsmedizin Berlin, 10117
4 Berlin, Germany

5 ⁸Department of Anesthesiology and Intensive Care Medicine, Charité – Universitätsmedizin
6 Berlin, 13353 Berlin, Germany

7 ⁹Clinical Study Center (CSC), Berlin Institute of Health, and Charité - Universitätsmedizin
8 Berlin, 10117 Berlin, Germany

9 ¹⁰German Center for Lung Research (DZL), 35392 Gießen, Germany

10 ¹¹Centre for Pathogen Evolution, Department of Zoology, University of Cambridge, Downing
11 St., Cambridge, CB2 3EJ, U.K.

12 ¹²Department of Tropical Medicine, Bernhard Nocht Institute for Tropical Medicine, and
13 Department of Medicine I, University Medical Center Hamburg-Eppendorf, 20359 Hamburg,
14 Germany

15 * these authors contributed equally

16

17

18

19

20

21

22

23

24

1 **Pa-COVID-19 study group:**

2 Linda Jürgens, Malte Kleinschmidt, Sophy Denker, Christoph Ruwwe-Glösenkamp, Bettina
3 Temmesfeld-Wollbrück, Katrin M. Heim, Dirk Schürmann, Andreas Hocke, Bastian Opitz,
4 Belén Millet Pascual-Leone, Rosa C. Schuhmacher, Nadine Olk, David Hillus, Felix Machleidt,
5 Sebastian Albus, Felix Bremer, Jan-Moritz Doehn, Carmen Garcia, Philipp Knape, Philipp M.
6 Krause, Liron Lechtenberg, Yaosi Li, Panagiotis Pergantis, Teresa Ritter, Berna Yedikat,
7 Christian Zobel, Friederike L. Hefele, Ute Kellermann, Mariana Schürmann, Lisa-Marie
8 Wackernagel, Anne Wetzel, Daniel Grund, Jens K. Haumesser, Johannes Hodes, Johannes
9 Rein, Peter Radünzel, Astrid Breitbart, Sergej Münzenberg, Dominik Soll, Tamar
10 Zhamurashvili, Ralf-Harto Hübner, Florian Alius, Tim Andermann, Thomas Cronen, Simon
11 Fraumann, Nikolaj Frost, Dominik Geus, Gisele J. Godzick-Njomgang, Anne Herholz, Vera
12 Hermanns, Moritz Hilbrandt, Till Jacobi, Ye-Ji Kim, Elena Madlung, Luise Martin, Nikolai
13 Menner, Agata Mikolajewska, Luisa Mrziglod, Nadine Muller, Michaela Niebank, Eva Pappé,
14 Frieder Pfäfflin, Lennart Pfannkuch, Matthias Raspe, Nicola Reck, Anne Ritter, Jacopo
15 Saccomanno, Laura K. Schmalbrock, Fridolin Steinbeis, Christoph Tabeling, Markus
16 Vogtmann, Susanne Weber, Markus Brack, Matthias Felten, Sein Schmidt, Maria Rönnefarth,
17 Georg Schwanitz, Alexander Krannich, Saskia Zvorc, Uwe D. Behrens, Lucie Kretzler, Linna
18 Li, Isabelle Wirsching, Chantip Dang-Heine, Michael Hummel, Dana Briesemeister, Denise
19 Treue, Martin Möckel, Samuel Knauß, Matthias Endres, Claudia Spies, Steffen Weber-
20 Carstens, Jan M. Kruse, Daniel Zickler, Andreas Edel, Britta Stier, Philipp Enghard, Roland
21 Körner, Kai-Uwe Eckardt, Lucas Elbert, Christopher Neumann, Marius A. Eckart, Thuy N.
22 Pham, Solveig Schönberger, Alexander Wree, Frank Tacke, Josef Mang, Nadia A. de Vries,
23 Marcel Wittenberg, Jana Riecke

24

25

26

27

1 **Notes**

2 **Conflict of interest**

3 CT, BM, ETH, MM, TL, PTL, LAMA, LM, PS, SSH, LBJ, LL, MP, MSS, RR, JW, SH, TZ, HMR,
4 AU, FB, CK, NS, TCJ, CD, MW, LES, PSG, VMC and FK have no conflicts of interest to
5 declare.

6 **Funding**

7 CT is a participant in the academic program “clinical studies in infectiology” funded by the
8 German Research Foundation (DFG).

9 MM is a participant in the BIH-Charité Digital Clinician Scientist Program funded by the Charité
10 – Universitätsmedizin Berlin, the Berlin Institute of Health (BIH), and the German Research
11 Foundation (DFG).

12 This study was supported by the German Federal Ministry of Education and Research
13 (NaFoUniMedCovid19 – COVIM, FKZ: 01KX2021 and PROVID - FKZ 01KI20160A) to LES,
14 MW, CD, and VMC and PROVID to SH.

15 The work of CD and VMC is supported by the Berlin Institute of Health, Charite -
16 Universitätsmedizin Berlin and by the German Ministry of Health (Konsiliarlabor für
17 Coronaviren).

18

19

20

21

1 **Abstract:**

2 **Background**

3 Adequate patient allocation is pivotal for optimal resource management in strained
4 healthcare systems, and requires detailed knowledge of clinical and virological disease
5 trajectories.

6 **Methods**

7 A cohort of 168 hospitalized adult COVID-19 patients enrolled in a prospective
8 observational study at a large European tertiary care center was analyzed.

9 **Results**

10 Forty-four percent (71/161) of patients required invasive mechanical ventilation (IMV).
11 Shorter duration of symptoms before admission (aOR 1.22 per day less, 95%CI 1.10-
12 1.37, $p < 0.01$), age 60-69 as compared to 18-59 years (aOR 4.33, 95%CI 1.07-20.10,
13 $p = 0.04$), and history of hypertension (aOR 5.55, 95%CI 2.00-16.82, $p < 0.01$) were
14 associated with need for IMV. Patients on IMV had higher maximal concentrations,
15 slower decline rates, and longer shedding of SARS-CoV-2 than non-IMV patients (33
16 days, IQR 26-46.75, vs 18 days, IQR 16-46.75, respectively, $p < 0.01$). Median duration
17 of hospitalization was 9 days (IQR 6-15.5) for non-IMV and 49.5 days (IQR 36.8-82.5)
18 for IMV-patients.

19 **Conclusion**

20 Our results indicate a short duration of symptoms before admission as a risk factor for
21 severe disease and different viral load kinetics in severely affected patients.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18

Key words: Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2); Coronavirus disease 2019 (COVID-19); Viral concentration; COVID-19 Nucleic Acid Testing; Respiratory Distress Syndrome; Mechanical ventilation; Artificial respiration; prospective study; symptom assessment

1 **Background**

2 The ongoing Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)
3 pandemic places an unprecedented burden on healthcare systems worldwide. Host
4 factors predictive of severe clinical course and adverse outcome in patients with
5 Coronavirus disease 2019 (COVID-19) include older age, male gender, and pre-
6 existing chronic comorbidities [1-6]. Several risk scores containing clinical
7 characteristics, laboratory assessments, and biomarkers have been proposed for
8 improved patient management and resource allocation [5, 7] .

9 Reported proportions of hospitalized patients requiring invasive mechanical ventilation
10 (IMV) vary considerably between 2.3% [8] and 23.6% [2]. In-hospital case fatality rates
11 of 20-30% have been described in China, Germany, Italy, the UK, and the US [1, 6, 9-
12 11]. In Germany, shortages of inpatient beds and intensive care unit (ICU) capacity
13 were largely avoided during the first pandemic wave [12], contributing to a
14 comparatively low overall case fatality rate [1, 13].

15 Here, we report clinical characteristics, laboratory and virological parameters, clinical
16 course, and outcome of 168 COVID-19 patients included in a prospective
17 observational cohort study conducted at Charité – Universitätsmedizin Berlin,
18 Germany. The study was designed for deep clinical, molecular, and immunological
19 phenotyping of COVID-19 [14-18].

20 The data reflect the situation in a tertiary care referral center for the treatment of
21 patients with acute respiratory distress syndrome (ARDS), including veno-venous
22 extracorporeal membrane oxygenation (vvECMO) therapy at an associated certified
23 weaning center during the first months of the COVID-19 pandemic, before treatment

1 with dexamethasone became standard of care [19]. Specific aims of this work were to
2 identify risk factors associated with need for IMV, to analyze viral kinetics in patients
3 with and without IMV, and to provide a comprehensive description of clinical course
4 and outcome.

5 **Methods**

6 **Study cohort and data collection**

7 Data collection was performed within the Pa-COVID-19 study, a prospective
8 observational cohort study conducted at Charité – Universitätsmedizin Berlin, as
9 described [14]. Adult patients admitted between March 1st and June 30th, 2020, with
10 PCR-confirmed SARS-CoV-2 infection were included if patients or their legal
11 representatives gave informed consent. We recorded epidemiological and
12 demographic data, medical history, history of present illness, symptoms, clinical
13 course, treatment, and outcomes upon enrolment and longitudinally during
14 hospitalization. The study was approved by the ethics committee of Charité –
15 Universitätsmedizin Berlin (EA2/066/20), conducted according to the Declaration of
16 Helsinki and Good Clinical Practice principles (ICH 1996) and is registered in the
17 German and WHO international clinical trials registry (DRKS00021688) [14].

18 Comorbidities were classified using the Charlson comorbidity Index (CCI) [20]. ARDS
19 was defined according to the Berlin definition of ARDS [21]. Sequential Organ Failure
20 Assessment (SOFA) score [22] was calculated from data recorded in the ICU data
21 management system. The following predefined events were assessed in all patients:
22 1) Sepsis (defined according to sepsis-3 criteria [23], 2) venous thromboembolic
23 events (VTE; pulmonary embolism or deep vein thrombosis), 3) neurological events

1 (hemorrhagic/ischemic stroke, delirium, critical illness polyneuropathy (CIP), critical
2 illness myopathy (CIM), epileptic seizure, meningitis and encephalitis). Treatment was
3 unaffected by participation in the study. Patient allocation was performed according to
4 structured regional processes [24] and management of critically ill patients following
5 national and international recommendations [25, 26].

6 All laboratory assessments were carried out in accredited laboratories at Charité –
7 Universitätsmedizin Berlin. SARS-CoV-2 viral concentration was measured in
8 respiratory samples (nasal- or oropharyngeal swabs) by real-time RT-PCR (32). Viral
9 concentration is given as \log_{10} genome copies per swab or initial 1 mL sampling buffer.

10 **Statistical analyses**

11 Distribution of continuous variables was summarized by median and interquartile range
12 (IQR) values or mean and standard deviation (sd), as appropriate. The differences of
13 continuous variables between groups were examined by Welch's t-test or, in absence
14 of normal distribution, by Mann-Whitney *U* test. Categorical variables were compared
15 using chi-square tests. For all analyses, complete cases were used for the respective
16 evaluation.

17 We conducted a multiple logistic regression with “need for invasive mechanical
18 ventilation” as a binary dependent variable and age, BMI, hypertension, diabetes, and
19 time between symptom onset and admission to hospital as independent variables. The
20 covariates were chosen taking into account current evidence, results from univariate
21 testing, and sample size. We performed univariate tests regarding all available patient
22 factors and association with organ support treatment, complications, and outcome.
23 Patients with therapy limitations (Do Not Intubate (DNI) or Do Not Resuscitate (DNR)
24 orders) at the respective time point were excluded for comparison between non-IMV

1 and IMV patients, and for analyses of course-of-organ support and mortality. Non-
2 survivors were excluded from comparison between short- (< 15 days) or long-term (\geq
3 15 days) IMV. For analysis of SOFA score, the highest score per day was included for
4 calculation of means. The neurological component of the SOFA score was not taken
5 into consideration due to patient sedation. Patients who died or were transferred to
6 other centers were excluded from calculation of length of hospital stay. For analyses
7 of routine laboratory parameters within 72 hours of first admission, first-available
8 parameters were included.

9 We compared viral concentration between non-IMV and IMV patients, and regressed
10 viral concentration on the duration from symptom onset to admission using both the
11 first positive RT-PCR result and the RT-PCR with the highest viral concentration. For
12 the calculation of viral concentration decline, we estimated the slope parameter from a
13 linear regression of at least four viral concentration measurements over time for each
14 patient. If available, the first of at least two final negative RT-PCR results was included,
15 in which case the viral concentration of the negative RT-PCR was set to 2.0 in
16 accordance with the RT-PCR limit of detection and sample dilution factor of ~ 20 [27,
17 28]. We calculated shedding duration as the time from symptom onset to the date of
18 the first of at least two final negative RT-PCR results.

19 Analyses were conducted with R (version 3.6.1) [29], JMP (version 14.2.0) [30], and
20 statsmodels (version 0.12.0) in Python 3.7.9 [31]. A p-value < 0.05 indicates statistical
21 significance, although all results have to be considered as non-confirmatory. For this
22 reason, no adjustment for multiple testing was done.

23

24

1 **Results**

2 **Baseline characteristics**

3 Between March 1st and June 30th 2020, a total of 347 adult patients with COVID-19 were
4 hospitalized at Charité – Universitätsmedizin Berlin. This analysis includes 168 adult
5 patients who consented to participation in the prospective observational study (**Figure**
6 **1**). Sixty-five percent (110/168) were directly admitted to our center, whereas 29.8%
7 (50/168) were referred due to ARDS or other conditions requiring tertiary care. Four
8 percent (7/168) were hospitalized for other reasons and coincidentally diagnosed with
9 SARS-CoV-2 infection during routine screening, and one patient (0.6%) was admitted
10 due to a late complication of COVID-19.

11 Baseline characteristics are shown in **Table 1**. Median patient age was 61 years (IQR
12 49.3-72), and 66.1% (111/168) were male. Median CCI was 3 (IQR 1-4). Most
13 prevalent comorbidities were hypertension (53.6%, 90/168), diabetes (19.6%, 33/168),
14 and chronic pulmonary disease (16.7%, 28/168). Median time from symptom onset
15 until diagnosis was 4 days (IQR 1-7), and from symptom onset until admission to
16 hospital was 6 days (IQR 3-10).

17 **ARDS and organ support treatment**

18 Fifty-two percent (88/168) of patients developed ARDS. One of them was already
19 dependent on long-term intermittent invasive ventilation and six had DNI orders in
20 place. Of the remaining 81 patients with ARDS, 87.6% (71/81) required IMV whereas
21 12.3% (10/81) could be managed with HFNC oxygen only (**Figure 1**). Median time
22 from hospital admission to intubation was 2 days (IQR 0-4). Among all patients without
23 therapeutic limitations, 44.1% (71/161) required IMV, 9.9% (16/161) could be treated

1 with HFNC oxygen only, 24.2% (39/161) required oxygen via nasal prongs, and 22%
2 (35/161) were not in need of supplemental oxygen.

3 Age between 60 and 69 years as compared to 18 to 59 years (adjusted OR 4.33, 95%
4 CI 1.07-20.10, $p=0.05$) and pre-existing hypertension (adjusted OR 5.55, 95% CI 2.00-
5 16.82, $p<0.01$) were independent risk factors for IMV requirement in multivariable
6 analysis. Requirement for IMV increased with shorter duration of symptoms before
7 hospital admission (adjusted OR 1.22 per day less, 95% CI 1.10-1.37, $p<0.01$, **Table**
8 **1**).

9 Seventy-nine percent (56/71) of all intubated patients required long IMV. Need for long
10 IMV was associated with a short (≤ 1 day) duration from admission until intubation
11 (20.0% (3/15) of patients with short IMV vs 53.6% (30/56) patients with long IMV,
12 unadjusted OR (uOR) 4.6, 95% CI 1.17-18.16, $p=0.02$). Other factors associated with
13 long IMV were transferral from other centers (28.6% (4/14) of patients with short IMV
14 were transferred vs 69.6% (39/56) of patients with long IMV, uOR 5.73, 95% CI: 1.58-
15 20.87, $p<0.01$), a higher total mean SOFA score during the second week after initial
16 admission (8.4, 95% CI 5.83-10.9 in patients with short IMV vs 11.2, 95% CI 10.1-12.3,
17 $p<0.05$ in patients with long IMV) as well as differences in SOFA score components
18 (coagulation, hepatic impairment; for details see **Supplementary table 1**.)

19 IMV patients had a significantly higher risk of death from COVID-19 compared to non-
20 IMV patients (1.1% (1/90) of non-IMV patients died vs 29.6% (21/71) of IMV patients,
21 uOR 37.38, 95% CI 4.88-286.26, $p<0.01$). Sixty-three percent (44/70) of IMV patients
22 underwent tracheotomy at a median time of 15 days (IQR 12-20) from intubation.
23 Weaning from IMV was successfully concluded in 76.0% (38/50) of surviving IMV
24 patients after a median of 42 days (IQR 16-66) from intubation. Eighteen percent (9/50)

1 of patients remained dependent on intermittent IMV and 6.0% (3/50) on long-term
2 oxygen therapy (LTOT) upon discharge or transferral.

3 Thirty-one percent (22/71) of all IMV patients required vvECMO (hereafter termed
4 “ECMO”) treatment. ECMO was initiated a median of nine days (IQR 5.3-18.5) from
5 intubation and continued for a median of 18 days (IQR 7.8-35.5). All ECMO patients
6 required hemodialysis, 86.4% (19/22) underwent tracheotomy, 50% (11/22) had a
7 VTE, and 50% (11/22) died. Seventy-five percent (53/71) of all patients with IMV and
8 all (22/22) patients on ECMO received proning therapy. Details of the subgroup of
9 patients receiving proning therapy have been reported elsewhere [32].

10 Hemodialysis was initiated in 30.4% (51/168) of patients, and in 66.2% (47/71) of
11 patients with IMV. Hemodialysis was initiated after a median of eight days (IQR 4.5-14
12 days) following hospital admission and five days (IQR 2-8.8 days) after intubation.
13 Thirty-eight percent (18/47) of IMV patients with hemodialysis died. For details on
14 tracheotomy, ECMO and hemodialysis see **Supplementary table 2**.

15 As no evidence of efficacy of antiviral and anti-inflammatory treatments was available
16 at the time, these were only used in a small subgroup of patients of this cohort
17 **(Supplementary table 3)**.

18 **Virological and routine laboratory data**

19 Viral concentration data was available for 166/168 patients. On average, each patient
20 had seven RT-PCR tests (sd: 5.3, min=1, max=29, including positive and negative
21 result) from the day of symptom onset (or 10 days from first admission, if the date of
22 symptom onset was not available) to the end of hospitalization, with tests performed
23 every 8.4 days on average (sd: 8.8). Eighty-six patients had two final negative RT-PCR
24 tests at the end of the disease course. Median first-measured viral concentration

1 differed by 0.68 log₁₀ viral copies between IMV and non-IMV patients (5.9, IQR 4.68-
2 7.28 vs 5.22 log₁₀ viral copies, IQR 4.49-7.28, respectively; p=0.12, **Figure 2A**), and
3 median highest viral concentration by 1.19 log₁₀ viral copies (6.7, IQR 5.35-7.62 vs 5.51
4 log₁₀ viral copies, IQR 4.7-7.62, respectively; p=0.02, **Figure 2B**). Decline of viral
5 concentration (**Supplementary figure 1**) was significantly slower in IMV versus non-
6 IMV patients (−0.13, IQR −0.19- −0.08 vs −0.22 log₁₀ viral concentration decrease / day,
7 IQR −0.3- −0.08, respectively; p<0.01) (**Figure 2C, Supplementary figure 2A-D**). The
8 duration of shedding was significantly longer in IMV patients than in non-IMV patients
9 (median 33, IQR 26-46.75 vs 18 days, IQR 16-46.75, p=<0.01). We found no
10 association between viral concentration and the duration from symptom onset to
11 admission and no difference in first or in highest viral concentrations in patients
12 requiring long versus short IMV (**Supplementary figure 3A-D, Supplementary table**
13 **1**).

14 A statistically significant difference between non-IMV and IMV patients was observed
15 in the levels of C-reactive protein (CRP), procalcitonin (PCT), Interleukin-6 (IL-6),
16 lactate dehydrogenase, ferritin, white blood cell count, lymphocyte count, neutrophil-
17 to-lymphocyte ratio, creatinine, urea, aspartate aminotransferase, creatine kinase, N-
18 terminal prohormone of brain natriuretic peptide, and troponin (**Supplementary table**
19 **4**). The course of 12 routine laboratory parameters over time in non-IMV and IMV
20 patients is shown in **Figure 3**.

21 **Complications during treatment**

22 Eighteen percent (31/168) of patients developed sepsis. Occurrence of sepsis was
23 associated with organ replacement therapies, but not with patient-related risk factors
24 (**Supplementary table 5**). The first sepsis episode occurred after a median of 16 days

1 (IQR 8-21) from hospital admission and 11.5 days (IQR 3-18.3 days) from intubation.
2 In 15/31 (48.4%) of patients with sepsis, ECMO was initiated during or soon after
3 occurrence of sepsis (median time from occurrence of sepsis and initiation of ECMO,
4 0 days, IQR 0-6.25 days).

5 Nineteen percent (32/168) of patients were diagnosed with at least one VTE. Due to
6 evidence for increased risk of thromboembolic events in COVID-19 (36), therapeutic
7 anticoagulation was introduced in all critically ill patients from April 2020. A quarter of
8 VTEs (8/32) occurred without anticoagulatory treatment, 18.8% (6/32) under
9 prophylactic anticoagulation, and 46% (15/32) under therapeutic anticoagulation. In
10 9.3% (3/32) of patients, VTE was diagnosed at autopsy and anticoagulation status at
11 onset was unclear.

12 Twenty-four percent of patients (41/168) had at least one neurologic event during
13 hospitalization, including hemorrhagic stroke (9/41, 22%), ischemic stroke (3/41,
14 7.3%), delirium (17/41, 41.5%), CIP/CIM (17/41, 41.5%), and epileptic seizure (3/41,
15 7.3%). Details of sepsis episodes, VTE and neurologic events are shown in
16 **Supplementary table 5.**

17 **Outcome**

18 Median time of hospital stay was 14 days (IQR 7-35) for all, 9 days (IQR 6-15.5) for
19 non-IMV, and 49.5 days (IQR 36.8-82.5) for IMV patients.

20 Seventeen percent (29/168) of all patients died. Of all patients without therapy
21 limitations (DNR/DNI), 13.6% (22/161) died, 8.7% (14/161) were transferred to other
22 centers, and 77.6% (125/161) were discharged. Median time from first hospitalization
23 until death was 33 days (IQR 16-98).

1 In univariate analyses, CCI ≥ 3 (uOR 4.35, 95% CI 1.52-12.45, $p < 0.01$), short duration
2 of symptoms (uOR 6.08, 95% CI 1.88-19.68, $p < 0.01$), occurrence of sepsis (uOR
3 16.47, 95% CI 5.85-46.5, $p < 0.01$), occurrence of VTE (uOR 7.58, 95% CI 2.88-19.97,
4 $p < 0.01$) and higher SOFA score during 1st and 2nd week after intubation were associated
5 with death. There was a difference in the median first and highest viral concentration
6 between survivors and non-survivors (6.84, IQR 4.99-7.91 vs 5.38, IQR 4.54-7.91,
7 $p = 0.05$; and 7.14, IQR 5.39-7.91 vs 5.86, IQR 4.77-7.91, $p = 0.04$, respectively)
8 **(Supplementary table 6).**

9 Of all discharged or transferred patients, 6.5% (9/139) still required IMV, 2.9% (4/139)
10 LTOT, and 1.4% (2/139) new hemodialysis.

11 **Discussion**

12 We analyzed a detailed clinical and virological dataset from a prospective
13 observational cohort of COVID-19 patients hospitalized in a tertiary, ECMO/ARDS
14 referral and weaning center in Germany during the first wave of the SARS-CoV-2
15 pandemic, before dexamethasone became standard of care. One of our major findings
16 is that rapid clinical deterioration – as reflected by short duration of symptoms before
17 hospital admission – is a highly relevant risk factor for both need of IMV in a
18 multivariable risk model and death in univariate analyses in our cohort.

19 A shorter duration of symptoms was not associated with higher initial viral
20 concentration. There was also no difference in initial viral concentration in more
21 severely ill patients, i.e., those requiring IMV, compared to mildly ill patients. Yet, we
22 found a significant increase of inflammatory markers such as CRP, PCT, and IL-6 at
23 presentation and over time in IMV compared to non-IMV patients. Moreover, IMV

1 patients had higher maximal concentrations, a slower decline as well as longer duration
2 of shedding of SARS-CoV-2 compared to non-IMV patients. Higher maximal viral
3 concentrations were also found in patients who died, compared to survivors.

4 Our findings underline that the early inflammatory host response to SARS-CoV-2
5 determines the course of disease more than pathogen factors such as initial SARS-
6 CoV-2 RNA concentration [2, 39-41]. A short duration of symptoms before admission
7 possibly reflects a rapid increase of the level of inflammation and might serve as an
8 easy to assess risk factor for clinical deterioration [42]. During the further course of
9 COVID-19, severe disease is characterized by the inability to rapidly reduce viral
10 particles. A possible explanation for this phenotype – rapid deterioration after symptom
11 onset, high inflammatory markers, and lack of efficient clearing of viral particles – might
12 be the various kinds of immune dysregulation, as reported by our group and others
13 [17, 43, 44].

14 Conflicting results have been published regarding the association between clinical
15 severity and viral concentration for SARS-CoV-2 so far [33-38]. This could possibly be
16 explained by a rather unbalanced proportion of mildly and severely affected patients in
17 the respective studies as compared to our cohort [38]. However, it is undetermined
18 whether higher maximal viral concentration and longer duration of shedding are a
19 possible cause or an indicator of more severe organ damage and disease.

20 Regarding invasive treatment methods and complications, although it is difficult to
21 separate cause from effect, onset of sepsis before or around the time of initiation of
22 ECMO may indicate that sepsis drives clinical deterioration rather than ECMO driving
23 sepsis, a topic requiring further investigation. In line with this, sepsis was a strong risk
24 factor for death in our study.

1 Although a remarkable proportion (29.8%) of patients were transferred to our center
2 due to severe ARDS, and overall 44.1% needed IMV, we report a comparatively low
3 in-hospital mortality of 13.6% in patients without limitations of therapy. In comparison,
4 Karagiannadis et al. reported in-hospital mortality of 22.2% [1] and Rieg et al. of 23.9%
5 [45] in cohorts with 17.2% [1] and 32.9% [45] ventilated patients, respectively. Of note,
6 our data show a high median length of hospital stay of 49.5 days for patients requiring
7 IMV. By comparison, the median length of hospital stay for non-COVID ARDS patients
8 was 17 days in a recent global multi-center prospective study [46]. There is growing
9 evidence that the length of IMV-, ICU-, and inpatient treatment of patients with COVID-
10 19 ARDS exceeds that of patients with ARDS unrelated to COVID-19 [42, 45].

11 Despite the long median duration of hospital stay, a considerable percentage of
12 patients could not successfully be weaned off the ventilator (18%), and 6% required
13 ongoing oxygen therapy following discharge. The mere number of deceased patients
14 therefore depicts the burden of disease of COVID-19 only very incompletely,
15 particularly with respect to long-term morbidity. The prospective approach of our study
16 will allow us to evaluate long-term complications in the aftermath of COVID-19.

17 Prospective observational studies are often hampered by selection of patients with
18 relatively mild disease courses due to need for informed consent. In our study, a
19 deferred-consent model was applied, allowing for inclusion of critically ill patients. The
20 high proportion of severely affected patients indicates that this selection bias does not
21 apply to our data. On the other hand, one fifth of patients were only mildly affected and
22 did not require oxygen therapy, representing a sub-cohort admitted for clinical
23 observation or due to lacking the possibility of self-isolation.

1 In conclusion, we report a short duration of symptoms before clinical deterioration as
2 a risk-factor for severe COVID-19, a finding that merits further exploration as an easily
3 assessable prognostic factor. Second, we show that severely ill patients have higher
4 maximal viral concentrations and a slower decline of viral concentration compared to
5 mildly-affected patients. Third, our results demonstrate a comparatively long duration
6 of inpatient treatment for patients with IMV, largely exceeding that described for non-
7 COVID-19 ARDS patients. Our data give a comprehensive description of the clinical
8 course, virological characteristics, organ support treatment, complications, and
9 outcome of a representative cohort of patients in an unrestricted tertiary care
10 healthcare setting with comparatively low mortality. The reported findings will be of
11 value for both scientific progress in the fight against SARS-CoV-2 as well as for clinical
12 management of critical care and weaning resources in the ongoing pandemic.

13

14

15

16

17

18

19

1 **Figure legends:**

2 **Figure 1: Study cohort flowchart.** A total of 347 adult patients were hospitalized with
3 COVID-19 during the study period from March 1st until June 30th at Charité-
4 Universitätsmedizin Berlin. Of these, 168 patients could be enrolled in the prospective
5 observational study, whereas 179 denied. Among the included patients, 88 had acute
6 respiratory distress syndrome (ARDS). One patient with ARDS was already invasively
7 ventilated and six of them had DNI/DNR (do not intubate/do not resuscitate) orders in
8 place, resulting in 81 patients requiring respiratory support. Of those, 71 patients were
9 intubated and ten required only high-flow nasal cannula oxygen therapy.

10

11 **Figure 2: Comparison of viral concentration between patients with invasive**
12 **mechanical ventilation (IMV) and those without IMV (non-IMV).** A) First measured
13 viral concentration: Median \log_{10} viral concentration and (IQR) are 5.9 (4.68-7.28) for
14 IMV patients and 5.22 (4.49-7.28) for non-IMV patients. B) Highest viral concentration:
15 Median \log_{10} viral concentrations and (IQR) are 6.7 (5.35-7.62) for IMV patients and
16 5.51 (4.7-7.62) for non-IMV patients. C) Differences in the slopes of \log_{10} viral
17 concentration decline rates were estimated using a linear regression of viral
18 concentration from the full disease course of a patient and days since symptom onset
19 (n=63) or admission (n=10) for patients with and without IMV. Only patients with at
20 least four viral concentration measurements were included. D) Duration from symptom
21 onset to the first of at least two final negative RT-PCR results for ventilated and non-
22 ventilated patients. Median 33 days (IQR: 26-46.75) for IMV vs 18 days (IQR: 16-46.75)
23 for non-IMV patients, p<0.01) Pairwise comparisons were performed using a Mann-
24 Whitney *U* test. Grey horizontal lines indicate the median.

1 **Figure 3 A-L:** Comparison of laboratory parameters during the course of disease in
2 IMV (red) versus non-IMV patients (blue). X-axis: days post admission. The boxes and
3 lines are median 25th and 75th percentiles, Whiskers indicate the 1st and 99th
4 percentile. A Welch's t-test was used: * $p < 0.05$, ** $p < 0.01$.

5

6 **Supplementary figure legends:**

7 **Supplementary Figure 1:** Courses of viral concentration over time for A) IMV and B)
8 non-IMV patients. Only patients with viral concentration measurements on at least four
9 different days were included. If a patient had multiple viral concentration
10 measurements on the same day, only the highest viral concentration was included for
11 that day. If available, the first of at least two final negative PCRs is included with a viral
12 concentration of 2.0 assigned. The x-axis displays the number of days since symptom
13 onset, if available ($n=63$), or number of days since admission ($n=10$).

14 **Supplementary Figure 2:** Viral concentration decline rates, taking into account full
15 and partial viral concentration courses for each patient. Decline rates were calculated
16 using a linear regression for patients with at least four RT-PCR results. A, B) include
17 positive RT-PCR results and the first of at least two final negative RT-PCR tests for a
18 patient. C, D) only include positive RT-PCR test results. A, C) are based on all RT-
19 PCR test results for a patient, while B, D) only take into account results from RT-PCR
20 tests performed within 30 days of symptom onset ($n=63$), or date of admission ($n=10$)
21 if date of symptom onset was unknown. Pairwise comparisons were performed using
22 a Mann-Whitney U test. Grey horizontal lines indicate the median.

23

24

25

1 **Supplementary Figure 3:** Log₁₀ viral concentration plotted against time in days from
2 symptom onset to admission. A) and B) include data from invasive mechanically
3 ventilated (IMV) patients, and C) and D) from non-IMV patients. A and C) show the
4 first-measured viral concentration per patient, and B and D) illustrate the highest viral
5 concentration per patient. Shaded areas indicate the 95% confidence interval. p – p-
6 value, R – correlation coefficient.

7

8

9

10

11

12

13

14

15

16

17

18

19

20

1 **References :**

- 2 1. Karagiannidis C, Mostert C, Hentschker C, et al. Case characteristics, resource use, and
3 outcomes of 10 021 patients with COVID-19 admitted to 920 German hospitals: an
4 observational study. *The Lancet Respiratory Medicine* **2020**; 8:853-62.
- 5 2. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospital admission and critical
6 illness among 5279 people with coronavirus disease 2019 in New York City: prospective
7 cohort study. *BMJ* **2020**; 369:m1966.
- 8 3. Deng G, Yin M, Chen X, Zeng F. Clinical determinants for fatality of 44,672 patients with
9 COVID-19. *Critical Care* **2020**; 24:179.
- 10 4. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related
11 death using OpenSAFELY. *Nature* **2020**; 584:430-6.
- 12 5. Knight SR, Ho A, Pius R, et al. Risk stratification of patients admitted to hospital with
13 covid-19 using the ISARIC WHO Clinical Characterisation Protocol: development and
14 validation of the 4C Mortality Score. *BMJ* **2020**; 370:m3339.
- 15 6. Docherty AB, Harrison EM, Green CA, et al. Features of 20 133 UK patients in hospital
16 with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: prospective
17 observational cohort study. *BMJ* **2020**; 369:m1985.
- 18 7. Liu S, Yao N, Qiu Y, He C. Predictive performance of SOFA and qSOFA for in-hospital
19 mortality in severe novel coronavirus disease. *Am J Emerg Med* **2020**; 38:2074-80.
- 20 8. Guan W-j, Ni Z-y, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in
21 China. *New England Journal of Medicine* **2020**; 382:1708-20.

- 1 9. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting Characteristics, Comorbidities,
2 and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City
3 Area. *JAMA* **2020**; 323:2052-9.
- 4 10. Grasselli G, Zangrillo A, Zanella A, et al. Baseline Characteristics and Outcomes of 1591
5 Patients Infected With SARS-CoV-2 Admitted to ICUs of the Lombardy Region, Italy. *JAMA*
6 **2020**; 323:1574-81.
- 7 11. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients
8 with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet* **2020**; 395:1054-
9 62.
- 10 12. Stang A, Stang M, Jöckel K-H. Estimated Use of Intensive Care Beds Due to COVID-19
11 in Germany Over Time. *Dtsch Arztebl International* **2020**; 117:329-35.
- 12 13. Bauer J, Brüggmann D, Klingelhöfer D, et al. Access to intensive care in 14 European
13 countries: a spatial analysis of intensive care need and capacity in the light of COVID-19.
14 *Intensive Care Medicine* **2020**; 46:2026-34.
- 15 14. Kurth F, Roennefarth M, Thibeault C, et al. Studying the pathophysiology of coronavirus
16 disease 2019: a protocol for the Berlin prospective COVID-19 patient cohort (Pa-COVID-19).
17 *Infection* **2020**; 48:619-26.
- 18 15. Chua RL, Lukassen S, Trump S, et al. COVID-19 severity correlates with airway
19 epithelium-immune cell interactions identified by single-cell analysis. *Nat Biotechnol* **2020**;
20 38:970-9.
- 21 16. Messner CB, Demichev V, Wendisch D, et al. Ultra-High-Throughput Clinical Proteomics
22 Reveals Classifiers of COVID-19 Infection. *Cell Systems* **2020**; 11:11-24.e4.
- 23 17. Schulte-Schrepping J, Reusch N, Paclik D, et al. Severe COVID-19 Is Marked by a
24 Dysregulated Myeloid Cell Compartment. *Cell* **2020**; 182:1419-40.e23.

- 1 18. Braun J, Loyal L, Frentsch M, et al. SARS-CoV-2-reactive T cells in healthy donors and
2 patients with COVID-19. *Nature* **2020**; 587:270-4.
- 3 19. Group TRC. Dexamethasone in Hospitalized Patients with Covid-19 — Preliminary
4 Report. *New England Journal of Medicine* **2020**.
- 5 20. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying
6 prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*
7 **1987**; 40:373-83.
- 8 21. Ranieri VM, Rubenfeld GD, Thompson BT, et al. Acute respiratory distress syndrome:
9 the Berlin Definition. *Jama* **2012**; 307:2526-33.
- 10 22. Vincent JL, Moreno R, Takala J, et al. The SOFA (Sepsis-related Organ Failure
11 Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on
12 Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive*
13 *Care Med* **1996**; 22:707-10.
- 14 23. Singer M, Deutschman CS, Seymour CW, et al. The Third International Consensus
15 Definitions for Sepsis and Septic Shock (Sepsis-3). *JAMA* **2016**; 315:801-10.
- 16 24. Wiesner B, Bachmann M, Blum TG, et al. Responsibilities of Weaning Centers during the
17 COVID-19 Pandemic Outbreak - Recommendations for the Assignment of ICU Capacities in
18 COVID-19 Patients as shown by the Berlin-Brandenburg POST-SAVE-Model. *Pneumologie*
19 **2020**; 74:358-65.
- 20 25. Kluge S, Janssens U, Welte T, et al. Recommendations for treatment of critically ill
21 patients with COVID-19. *Der Anaesthesist* **2020**.
- 22 26. Brower RG, Matthay MA, Morris A, Schoenfeld D, Thompson BT, Wheeler A. Ventilation
23 with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and
24 the acute respiratory distress syndrome. *N Engl J Med* **2000**; 342:1301-8.

- 1 27. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV)
2 by real-time RT-PCR. *Euro Surveill* **2020**; 25.
- 3 28. Wölfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized
4 patients with COVID-2019. *Nature* **2020**; 581:465-9.
- 5 29. Team RC. A language and environment for statistical computing. R Foundation for
6 Statistical computing. **2018**:<https://www.R-project.org/>.
- 7 30. Sall JL, Ann; Stephens, Mia; Loring, Sheila. JMP® Start Statistics: A Guide to Statistics
8 and Data Analysis Using JMP®, Sixth
9 Edition. **2017**:[https://support.sas.com/content/dam/SAS/support/en/books/jmp-start-](https://support.sas.com/content/dam/SAS/support/en/books/jmp-start-statistics/70199_excerpt.pdf)
10 [statistics/70199_excerpt.pdf](https://support.sas.com/content/dam/SAS/support/en/books/jmp-start-statistics/70199_excerpt.pdf).
- 11 31. Seabold SP, Josef. statsmodels: Econometric and statistical modeling with python.
12 Proceedings of the 9th Python in Science Conference **2010**.
- 13 32. Mittermaier M, Pickerodt P, Kurth F, et al. Evaluation of PEEP and prone positioning in
14 early COVID-19 ARDS. *EClinicalMedicine*.
- 15 33. To KK-W, Tsang OT-Y, Leung W-S, et al. Temporal profiles of viral load in posterior
16 oropharyngeal saliva samples and serum antibody responses during infection by SARS-CoV-
17 2: an observational cohort study. *The Lancet Infectious Diseases* **2020**; 20:565-74.
- 18 34. Magleby R, Westblade LF, Trzebucki A, et al. Impact of Severe Acute Respiratory
19 Syndrome Coronavirus 2 Viral Load on Risk of Intubation and Mortality Among Hospitalized
20 Patients With Coronavirus Disease 2019. *Clinical Infectious Diseases* **2020**.
- 21 35. Liu Y, Yan L-M, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *The*
22 *Lancet Infectious Diseases* **2020**; 20:656-7.

- 1 36. Argyropoulos KV, Serrano A, Hu J, et al. Association of Initial Viral Load in Severe Acute
2 Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Patients with Outcome and Symptoms.
3 *Am J Pathol* **2020**; 190:1881-7.
- 4 37. Cevik M, Tate M, Lloyd O, Maraolo AE, Schafers J, Ho A. SARS-CoV-2, SARS-CoV, and
5 MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic
6 review and meta-analysis. *The Lancet Microbe*.
- 7 38. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of
8 COVID-19. *Nature Medicine* **2020**; 26:672-5.
- 9 39. Wu X, Liu L, Jiao J, Yang L, Zhu B, Li X. Characterisation of clinical, laboratory and
10 imaging factors related to mild vs. severe covid-19 infection: a systematic review and meta-
11 analysis. *Ann Med* **2020**; 52:334-44.
- 12 40. Khinda J, Janjua NZ, Cheng S, van den Heuvel ER, Bhatti P, Darvishian M. Association
13 between markers of immune response at hospital admission and COVID-19 disease severity
14 and mortality: A meta-analysis and meta-regression. *J Med Virol* **2020**.
- 15 41. Shao M, Li X, Liu F, Tian T, Luo J, Yang Y. Acute kidney injury is associated with severe
16 infection and fatality in patients with COVID-19: A systematic review and meta-analysis of 40
17 studies and 24,527 patients. *Pharmacol Res* **2020**; 161:105107.
- 18 42. Schmidt M, Hajage D, Demoule A, et al. Clinical characteristics and day-90 outcomes of
19 4244 critically ill adults with COVID-19: a prospective cohort study. *Intensive Care Medicine*
20 **2020**.
- 21 43. Lucas C, Wong P, Klein J, et al. Longitudinal analyses reveal immunological misfiring in
22 severe COVID-19. *Nature* **2020**; 584:463-9.
- 23 44. Fajgenbaum DC, June CH. Cytokine Storm. *New England Journal of Medicine* **2020**;
24 383:2255-73.

- 1 45. Rieg S, von Cube M, Kalbhenn J, et al. COVID-19 in-hospital mortality and mode of
2 death in a dynamic and non-restricted tertiary care model in Germany. PLoS One **2020**;
3 15:e0242127.

- 4 46. Bellani G, Laffey JG, Pham T, et al. Epidemiology, Patterns of Care, and Mortality for
5 Patients With Acute Respiratory Distress Syndrome in Intensive Care Units in 50 Countries.
6 JAMA **2016**; 315:788-800.

- 7

- 8





